


HOSTED BY



Contents lists available at ScienceDirect

International Journal of Pediatrics and Adolescent Medicine

journal homepage: <http://www.elsevier.com/locate/ijpam>

Original article

Gut microbiome pattern in adolescents with functional gastrointestinal disease

Andrew R. Nafarin^a, Badriul Hegar^a, Hikari A. Sjakti^a, Yvan Vandenplas^{b,*}^a Department of Child Health, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo Hospital, Jakarta, Indonesia^b KidZ Health Castle, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel, Brussels, Belgium

ARTICLE INFO

Article history:

Received 19 July 2018

Received in revised form

13 January 2019

Accepted 27 January 2019

Available online 2 February 2019

Keywords:

Functional gastrointestinal disease

Adolescents

Microbiome

ABSTRACT

Background: Functional gastrointestinal disease (FGID) has a worldwide prevalence of 10–45%, and is one of the most common causes of recurrent abdominal pain in children. FGID is characterized with abdominal discomfort and changes in bowel movement. Alteration in gut microbiota is associated with FGID, but data are limited, and there are no data from Indonesia.

Methods: A case–control study was conducted in 22 FGID children and 28 healthy subjects aged 13–18 years at the junior high school and senior high school in Central Jakarta. FGID was diagnosed using Rome IV criteria. Age, sex, and level of education were recorded. Stool samples were collected and investigated for *Bifidobacterium* spp. and Enterobacteriaceae.

Results: Most of the FGID subjects were females (17/22), with a median age of 16 years. The median values of *Bifidobacterium* spp. were 138.95 (range: 0.2–22,735.8) CFU/gram for the FGID subjects and 232.5 (range: 1.9–38,985.6) CFU/gram in healthy subjects, which showed no statistically significant difference ($P = .49$). The median values of Enterobacteriaceae were 58.9 (range: 2.5–9577.8) CFU/gram in FGID subjects and 85 (range: 12.1–3139.4) CFU/gram in healthy subjects, which showed no statistically significant difference ($P = .94$). Our findings indicate that the gut microbiome of adolescents with FGIDs is characterized by a huge variability in levels of *Bifidobacterium* spp. and Enterobacteriaceae.

Conclusion: Because of the wide range detected in the number of *Bifidobacterium* spp. and Enterobacteriaceae in FGID and healthy subjects, no statistically significant difference was observed. More studies in larger groups of selected patients may be needed.

© 2019 Publishing services provided by Elsevier B.V. on behalf of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Functional gastrointestinal disease (FGID) is defined using the Rome IV criteria and must, by definition, include all of the following: i) abdominal discomfort (an uncomfortable sensation not described as pain) or pain associated with 2 or more of the following: at least 25% of the time improved with defecation and/or onset associated with a change in the frequency of stool and/or onset associated with a change in the form (appearance) of stool and ii) no evidence of an inflammatory, anatomic, metabolic, or neoplastic process that explains the subject's symptoms [1].

Data of the prevalence of FGID in Asian children and adolescents

are limited. According to Rome II criteria, the prevalence of FGID in the outpatient clinic of the Pediatric Gastrohepatology Clinic, Cipto Mangunkusumo Hospital, Indonesia, in 2001 was estimated at 4% [2]. In a study based on Rome III criteria in 2014, 14 out of 232 high school students in Jakarta were classified as suffering from FGID, a prevalence of 6.0% [3].

Different mechanisms have been suggested for the pathophysiology of FGID, including psychological disorder, visceral hypersensitivity, and gut microbiome composition [4]. Recent studies suggest that FGID is associated with changes in the gut microbiome [5–8]. The goal of this study was to analyze the *Bifidobacterium* and Enterobacteriaceae microbiome in children with and without FGID.

2. Materials and methods

2.1. Subjects

From May 2016 to September 2016, 22 Rome IV criteria-positive

* Corresponding author. Kidz Health Castle, UZ Brussel, Vrije Universiteit Brussel, Laarbeeklaan, 101, 1090, Brussels, Belgium.

E-mail address: yvan.vandenplas@uzbrussel.be (Y. Vandenplas).

Peer review under responsibility of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia.

adolescents for FGID aged 13–18 years were recruited from the 216th Junior High School and 68th Senior High School in Central Jakarta. Clinical symptoms of recurrent abdominal pain occurring at least 1 day per week in the last 3 months are associated with two or more of the following symptoms (1) related to defecation, (2) associated with a change in the frequency of stool, and (3) associated with a change in the form (appearance) of stool. Participants were excluded if they had taken antibiotics during the 3 months before inclusion or had suffered any organic disease during 6 months before inclusion in the study. During the same period, 28 healthy age-matched volunteers without any sign or symptoms of organic disease and without intake of antibiotics were included as control. Briefly, the patient and control groups were comparable with similar characteristics, except for the presence of symptoms of an FGID. Approval of the Ethical Committee of the Medical College of Indonesia University was obtained (number 875/UN2.F1/ETIK/2015).

2.2. Study protocol

Stools were collected and transported in an icebox to the Prodia Research and Esoteric Laboratory in Central Jakarta, Indonesia within 1 h after defecation. Once arrived in the laboratory, samples were frozen at -70°C before DNA extraction. The laboratory was blinded about the origin of the stools, i.e., patients versus controls. Real-time polymerase chain reaction (PCR) was performed using CFX96 from Bio-Rad Laboratories, Inc. The bacterial reference strains were *Bifidobacterium breve* (ATCC 15700) and *Escherichia coli* (ATCC 11775) by Remel™. The sequences for *Bifidobacterium* are CGCGTCYGGTGTGAAAG and CCCACATCCAGCATCCA. The sequences of *Enterobacteriaceae* are TGCCGTAATTCGGGAGAAGGCA and TCAAGGACCAGTGTTCAGTGTC. DNA extraction was performed using QIAamp DNA Stool Mini Kit by Qiagen.

2.3. Statistical analysis

All data were collected and analyzed with SPSS version 20.0. The Mann–Whitney *U* test was used to analyze the data. *P* value less than 0.05 was considered statistically significant. Data are expressed as median (minimum to maximum value).

3. Results

A total of 50 subjects (22 subjects with FGIDs and 28 healthy controls; 37 females and 13 males) were included. The median age of the FGID group was 16 years, and the median age of the control group was 14 years (Table 1).

The median value of *Bifidobacterium* spp. was 138.95 (0.2–22,735.8) CFU/gram in the FGID group and 232.5 (1.9–38,985.6) CFU/gram in the control group (*p* 0.49). The median value of *Enterobacteriaceae* is 58.9 (2.5–9577.8) CFU/gram in the FGID group and 85 (12.1–3139.4) CFU/gram in the control group (*p*

0.94). No statistically significant difference was found for *Bifidobacterium* spp. and *Enterobacteriaceae* between the FGID and control groups (Table 2).

The data of *Bifidobacterium* spp. and *Enterobacteriaceae* were transformed, which produced a normal curve (Shapiro–Wilk test > 0.05). The mean value of log *Bifidobacterium* spp. was 1.94 (95% CI 1.29–2.59) in the FGID group and 2.31 (95% CI 1.89–2.73) in the control group (*P* 0.31). The mean value of log *Enterobacteriaceae* was 2.09 (95% CI 1.63–2.55) in the FGID group and 2.0 (95% CI 1.72–2.28) in the control group (*P* 0.73). No statistically significant difference was found for *Bifidobacterium* spp. and *Enterobacteriaceae* between the FGID and control groups (Table 3).

These data indicate that the gut microbiome of adolescents with FGID is characterized by a huge variability in levels of *Bifidobacterium* and *Enterobacteriaceae*, which result in a statistically nonsignificant trend for lower levels of *Bifidobacterium* spp. and higher levels of *Enterobacteriaceae* in adolescents with FGIDs.

4. Discussion

Risk factors for FGID include differences according to age, race, sex, diet, region, psychological insults and familial history [9]. The etiology of FGID is not fully understood. Recent studies showed that the gut microbiota pattern of patients with FGID differs significantly from that of healthy individuals, with an increase in potentially pathogenic bacteria and a decrease in beneficial bacteria [9–12]. Probiotics given during management induce significant changes in the gut microbiota pattern and reduce FGID symptoms [13,14].

This study uses PCR to analyze *Bifidobacterium* spp., which represents beneficial bacteria, and *Enterobacteriaceae*, which represents potentially pathogenic bacteria. The two types of bacteria were selected, as there are studies that show significant differences between both types [12,13,15].

This is the first study conducted on Indonesian children and adolescents with FGID and their gut microbiota pattern. This study did not compare food and drinking habits or socioeconomic status. A previous study performed in the same school showed no significant differences between gender, age, academic records, parental educational level, number of siblings, socioeconomic status, use of antibiotics during the previous two months, and consumption of bread and drinking milk/coffee/energy drinks [3]. The prevalence of FGID in this study is 6%, which is the same as that observed in the previous study. Devanarayana et al. found in a meta-analysis that FGID prevalence in Asia is approximately 2.8–25.7%, with a median of 12.4% [16].

This study also did not exclude subjects who sometimes consumed commercial probiotic dairy products such as Yakult™ containing the *Lactobacillus casei* Shirota strain. A previous study conducted in 39 subjects with FGID showed that *L. casei* Shirota, by consuming Yakult™ twice daily for 8 weeks (6.5×10^9 CFU), failed to show significant improvement [17]. *L. casei* Shirota strain or placebo administered to 10 healthy subjects thrice daily as 100 ml (minimum 10^9 CFU/ml) for 4 consecutive weeks showed a significant increase in *Lactobacillus* but no increase in *Bifidobacterium*. [18] To date, no trial has been carried out in patients with FGID along with a comparable amount of *L. casei* Shirota strain.

Generally, the gut microbiota is composed of beneficial and potentially pathogenic bacteria. In healthy individuals, there is a balance between beneficial bacteria such as *Bifidobacterium* and *Lactobacillus* and potentially pathogenic bacteria such as *Enterobacteriaceae*, *Streptococcus*, *Enterococcus*, *Bacteroides*, *Clostridium*, *Staphylococcus*, *Escherichia*, and *Proteus* [19]. The healthy balance in the gut microbiota is the fundamental for a normal gut function, while any dysbiosis may produce gut symptoms.

Table 1
Subject characteristics.

	FGID group (n = 22)	Control group (n = 28)
Sex		
Male	5	8
Female	17	20
School		
Junior High	9	18
Senior High	13	10
Age (years old) ^a	16 (13–17)	14 (13–17)

^a Data expressed as median (minimum to maximum value).

Table 2
Bacterial count results.

Bacterial count (CFU/gram)	FGID (N = 22)	Control (N = 28)	<i>P</i> ^a
<i>Bifidobacterium</i> spp.	138.95 (0.2–22,735.8)	232.5 (1.9–38,985.6)	.493
<i>Enterobacteriaceae</i>	58.9 (2.5–9577.8)	85 (12.1–3139.4)	.938

Legend: FGID: Functional Gastrointestinal Disorder; Data are expressed as median (minimum to maximum value); ^a Mann–Whitney *U* Test.

Table 3
Log bacterial count results.

Log Bacterial count	FGID (N = 22)	Control (N = 28)	<i>P</i> ^a
<i>Bifidobacterium</i> spp.	1.94 (1.46)	2.31 (1.08)	.312
<i>Enterobacteriaceae</i>	2.09 (1.03)	2.00 (0.71)	.726

Legend: FGID: Functional Gastrointestinal Disorder; Data are expressed as mean (±SD); ^a Unpaired *t*-Test.

Bifidobacterium spp. is a dominant microorganism in the human intestine that produces essential nutrients for the mucosa, such as short-chain fatty acids (SCFAs) and lactic acid. They eliminate toxins and unnecessary substances by decreasing the intraluminal pH and inhibiting the growth of potential pathogenic organisms such as *E. coli*, *Salmonella*, and *Staphylococcus aureus* [20,21]. *Bifidobacterium* also stimulates gut epithelial cell turnover, hence correlating positively with secretory IgA, and prevents bacterial translocation, hence correlating negatively with *E. coli* counts [22,23]. This study found a trend for higher counts of *Bifidobacterium* in healthy subjects, although the difference was not statistically significant. Previous studies also showed higher counts of *Bifidobacterium* spp. such as *Bifidobacterium catenulatum* in healthy subjects. [9] The tendency of lower counts of *Bifidobacterium* in subjects with FGID might be a marker for the gut microbiota pattern in Indonesia. The composition and diversity of the microbiome are established on the basis of the luminal environment (physical, chemical, and biological exposome) and host surveillance. The chemical exposome derived from nutrients and other xenobiotics can influence the dynamics of the microbiome community (the stability, diversity, or resilience) [24].

Enterobacteriaceae, the main cause of endotoxin production, comprises many species such as *E. coli*, *Klebsiella pneumoniae*, *Salmonella typhimurium*, *Shigella*, *Proteus*, *Enterobacter*, *Serratia*, and *Citrobacter*. These microorganisms in Enterobacteriaceae may produce ammonia and sulfured hydrogen and can impair gut absorptive function in the colon and jejunum, with decreased absorption of water, glucose, and electrolytes [25]. Pimentel et al. found that 78% of the subjects with FGID had small intestinal bacterial overgrowth. Bacterial eradication revealed an improvement in symptoms in these patients; 48% of eradicated subjects did no longer meet the Rome criteria for FGID [26]. Our study found a trend for higher counts of Enterobacteriaceae in subjects with FGID, although not statistically significant. Previous studies found a statistically significant increase in subjects with FGID [12,22,27–29].

5. Conclusion

Because of the wide range observed in the number of *Bifidobacterium* spp. and Enterobacteriaceae in healthy adolescents and those with FGID, no statistically significant difference was observed. The wide range may be due to environment, such as dietary factors. Consequently, more data are needed in large patient groups under strict control of environmental factors.

Conflicts of interest

There is no conflict of interest. This research was funded by the

researcher.

Ethical statement

Approval of the Ethical Committee of the Medical College of Indonesia University was obtained (number 875/UN2.F1/ETIK/2015).

CRediT authorship contribution statement

Andrew R. Nafarin: Data curation, Project administration, Writing - original draft, Conceptualization. **Badriul Hegar:** Data curation, Project administration, Writing - original draft, Conceptualization. **Hikari A. Sjakti:** Data curation, Project administration, Writing - original draft. **Yvan Vandenplas:** Writing - original draft, Data curation, Conceptualization.

References

- [1] Benninga MA, Faure C, Hyman PE, St James Roberts I, Schechter NL, Nurko S. Childhood functional gastrointestinal disorders: neonate/toddler. *Gastroenterology* 2016;150:1443–551.
- [2] Hegar B. Rome II Criteria in developing country. *Acta Gastroenterol Belg* 2003;8:11.
- [3] Lara Filleskes AP, Fatima SA, Hegar B. Irritable bowel syndrome and its associated factors in adolescents. *Paediatr Indones* 2014;54:344–50.
- [4] Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006;130:1480–91.
- [5] Quigley EM. Bacterial flora in irritable bowel syndrome: role in pathophysiology, implications for management. *J Dig Dis* 2007;8:2–7.
- [6] Lin HC. Small intestinal bacterial overgrowth: a framework for understanding irritable bowel syndrome. *J Am Med Assoc* 2004;292:852–8.
- [7] Pimentel M, Park S, Mirocha J, Kane SV, Kong Y. The effect of a nonabsorbed oral antibiotic (rifaximin) on the symptoms of the irritable bowel syndrome: a randomized trial. *Ann Intern Med* 2006;145:557–63.
- [8] O'Mahony L, McCarthy J, Kelly P, Hurley G, Luo F, Chen K, dkk. Lactobacillus and bifidobacterium in irritable bowel syndrome: symptom responses and relationship to cytokine profiles. *Gastroenterology* 2005;128:541–51.
- [9] Zhu X, Chen W, Zhu X, Shen Y. A cross-sectional study of risk factors for irritable bowel syndrome in children 8–13 years of age in Suzhou, China. *Gastroenterol Res Pract* 2014;2014:198461.
- [10] Malinen E, Rinttilä T, Kajander K, Matto J, Kassinen A, Krogius L, dkk. Analysis of the fecal microbiota of irritable bowel syndrome patients and healthy controls with real-time PCR. *Am J Gastroenterol* 2005;100:373–82.
- [11] Kassinen A, Krogius-Kurikka L, Makivuokko H, Rinttilä T, Paulin L, Corander J, dkk. The fecal microbiota of irritable bowel syndrome patients differs significantly from that of healthy subjects. *Gastroenterology* 2007;133:24–33.
- [12] Kerckhoffs AP, Samsom M, van der Rest ME, de Vogel J, Knol J, Ben-Amor K, dkk. Lower Bifidobacteria counts in both duodenal mucosa-associated and fecal microbiota in irritable bowel syndrome patients. *World J Gastroenterol* 2009;15:2887–92.
- [13] Fan YJ, Chen SJ, Yu YC, Si JM, Liu B. A probiotic treatment containing Lactobacillus, Bifidobacterium and Enterococcus improves FGID symptoms in an open label trial. *J Zhejiang Univ - Sci B* 2006;7:987–91.
- [14] Francavilla R, Miniello V, Magista AM, De Canio A, Bucci N, Gagliardi F, dkk. A randomized controlled trial of Lactobacillus GG in children with functional abdominal pain. *Pediatrics* 2010;126:e1445–52.
- [15] Gerritsen J, Smidt H, Rijkers GT, de Vos WM. Intestinal microbiota in human health and disease: the impact of probiotics. *Genes Nutr* 2011;6:209–40.
- [16] Devanarayana NM, Rajindrajith S, Pathmeswaran A, Abegunasekara C, Gunawardena NK, Benninga MA. Epidemiology of irritable bowel syndrome in children and adolescents in Asia. *J Pediatr Gastroenterol Nutr* 2015;60:792–8.
- [17] Thijssen AY, Clemens CH, Vankerckhoven V, Goossens H, Jonkers DM, Masclee AA. Efficacy of Lactobacillus casei Shirota for patients with irritable bowel syndrome. *Eur J Gastroenterol Hepatol* 2016;28:8–14.
- [18] Spanhaak S, Havenaar R, Schaafsma G. The effect of consumption of milk fermented by Lactobacillus casei strain Shirota on the intestinal microflora

- and immune parameters in humans. *Eur J Clin Nutr* 1998;52:899–907.
- [19] Round JL, Mazmanian SK. The gut microbiota shapes intestinal immune responses during health and disease. *Nat Rev Immunol* 2009;9:313–23.
- [20] Kerckhoffs AP, Samsom M, van der Rest ME, de Vogel J, Knol J, Ben-Amor K, et al. Lower Bifidobacteria counts in both duodenal mucosa-associated and fecal microbiota in irritable bowel syndrome patients. *World J Gastroenterol* 2009;15:2887–92.
- [21] King TS, Elia M, Hunter JO. Abnormal colonic fermentation in irritable bowel syndrome. *Lancet* 1998;352:1187–9.
- [22] Jian-Min Si Y-CY, Fan Yu-Jing, Chen Shu-Jie. Intestinal microecology and quality of life in irritable bowel syndrome patients. *World J Gastroenterol* 2004;10:1802–5.
- [23] O'Sullivan DJ, Kullen MJ. Tracking of probiotic bifidobacteria in the intestine. *Int Dairy J* 1998;513–25.
- [24] Moon Y. Microbiome-linked crosstalk in the gastrointestinal exposome towards host health and disease. *Pediatr Gastroenterol Hepatol Nutr* 2016;19:221–8.
- [25] Johnson CL, Versalovic J. The human microbiome and its potential importance to pediatrics. *Pediatrics* 2012;129:950–60.
- [26] Pimentel M, Chow EJ, Lin HC. Eradication of small intestinal bacterial overgrowth reduces symptoms of irritable bowel syndrome. *Am J Gastroenterol* 2000;95:3503–6.
- [27] Carroll IM, Ringel-Kulka T, Siddle JP, Ringel Y. Alteration in composition and diversity of the intestinal microbiota in patients with diarrhea-predominant irritable bowel syndrome. *Neurogastroenterol Motil* 2012;24:521–30.
- [28] Carroll IM, Chang YH, Park J, Sartor RB, Ringel Y. Luminal and mucosal-associated intestinal microbiota in patients with diarrhea-predominant irritable bowel syndrome. *Gut Pathog* 2010;2:19.
- [29] Sobieszczanska BM, Osek J, Wasko-Czopnik D, Dworniczek E, Jermakow K. Association of enteroaggregative *Escherichia coli* with irritable bowel syndrome. *Clin Microbiol Infect* 2007;13:404–7.