

activity and CBX2 knockdown led to a decrease in ALDH activity. Examination of primary samples from the GTFB and TMA revealed CBX2 expression is increased in HGSOC and is upregulated in approximately 58% of metastases when compared to the primary tumor.

**Conclusions:** CBX2 directly impacts proliferation and is overexpressed in HGSOC, indicating CBX2 may be associated with advanced disease. Elucidation of the mechanism is ongoing, however, a stem-like phenotype seems to play a role. This work expands our understanding of HGSOC progression and identifies a novel therapeutic target.

doi:[10.1016/j.ygyno.2019.03.220](https://doi.org/10.1016/j.ygyno.2019.03.220)

#### Poster #31

##### Salvage treatment in recurrent endometrial cancer of the pelvis and peritoneal cavity

L. McAlarnen<sup>a</sup>, K. Ryan<sup>b</sup>, W. Adams<sup>c</sup>, A. Gliniewicz<sup>a</sup>, A. Winder<sup>a</sup>, M. Liotta<sup>a</sup>, R. Potkul<sup>a</sup>, W. Small Jr<sup>a</sup>, M. Harkenrider<sup>a</sup>. <sup>a</sup>Loyola University Medical Center. <sup>b</sup>Loyola University Chicago Stritch School of Medicine. <sup>c</sup>Loyola University Chicago Health Sciences Division

**Objectives:** Regional recurrence of endometrial cancer (EC) is uncommon but represents a challenging yet potentially curable group of patients. Here we seek to determine optimal methods of salvage therapy for regionally recurrent EC.

**Methods:** A single institution database was analyzed from 2007 to present with 22 cases identified of nodal, pelvic, or peritoneal cavity recurrences of EC treated with curative intent. Patient, tumor, and treatment characteristics were identified and analyzed for both initial and recurrent treatment. Univariable Cox proportional hazards models were used to estimate the risk of a second recurrence. Due to sparse event rates, conclusions were confirmed with Fisher's exact tests.

**Results:** At diagnosis, 73% were endometrioid histology, 73% stage 1, and 27% with LVSI. Of 22 cases of recurrent EC, 13 recurrences (59%) were regional including the pelvic and paraaortic nodes, while 9 recurrences (41%) were to the abdomen. Twelve patients experienced remission from last treatment to most recent follow up ranging from 20 days to over 6 years. Nine (75%) of the patients currently in remission underwent surgery, EBRT, and chemotherapy. Nine of 22 patients experienced a second pelvic or peritoneal recurrence (41%). Three of the 4 patients with distant metastases had regional or abdominal recurrences. The overall probability of survival two years after a regional or abdominal recurrence treated with salvage therapy was 69% (95% CI: 38% - 86%). The overall probability of progression-free survival at 2 years was 51% (95% CI: 26% - 72%).

**Conclusions:** In this sample, we found no meaningful association of a definitive salvage regimen and survival for recurrent EC of the pelvis and peritoneal cavity. Aggressive use of multimodality therapy with surgery followed by tumor-directed radiotherapy and chemotherapy has favorable progression-free and overall survival in this very high-risk population of recurrent EC patients.

doi:[10.1016/j.ygyno.2019.03.221](https://doi.org/10.1016/j.ygyno.2019.03.221)

#### Poster #32

##### Determining the methylation patterns of clinically normal endometrium and multiple tumor regions from uteri containing endometrial cancer

M. Bagaria, S. Kerr, E. Heinzen, M. Maurer, E. Shields, M. Lemens, V. Shridhar, J. Bakkum-Gamez

**Objectives:** Aberrant DNA methylation occurs early in carcinogenesis and is being explored as a biomarker for early detection of endometrial cancer (EC). Methylation can be present as a field effect

in histologically normal tissues and can be associated with future cancer development. Additionally, molecular diversity or intratumoral heterogeneity (ITH) within a cancer can be related with a higher risk of recurrence. Here we explore the field effect and ITH of EC through methylation analyses of normal and tumor regions.

**Methods:** Cases of hysterectomy for EC from 1/2011-12/2013 were retrospectively identified. Women with germline genetic mutations, presence of synchronous cancers, or history of chemotherapy or radiation receipt were excluded. Normal endometrium (NE), precancerous lesions (PC), and up to 3 separate tumor regions from within each hysterectomy specimen were selected by a single gynecologic pathologist. Extracted DNA from each area underwent pyrosequencing of 4 genes previously identified as methylated in type I (RASSF1A, CDH13) or type II (HTR1B, ADCYAP1) EC. Methylation percentage was evaluated individually across CpG sites and averaged across each gene. The CpG sites of each gene were noted to have consistent methylation using hierarchical clustering. Differences in methylation between NE and EC for each gene were assessed using paired t test. Patterns of methylation across the tumor regions within the patient and between patients were assessed using principal component analysis.

**Results:** Among 24 EC cases, 4 were clear cell (CC), 6 grade 1 or 2 endometrioid adenocarcinoma (EA1/2), 4 grade 3 endometrioid adenocarcinoma (EA3), and 10 serous. The mean age of this cohort was 64 years. In the hysterectomy specimens, NE areas were available in 14/24 (58%), PC lesions in 11/24 (45%) and 3 separate tumor regions in 22/24 women (91%). NE had significantly lower methylation than tumor regions for all the 4 genes (all  $p < 0.005$ ). Tumor methylation did not appear to be associated with age for any of the genes tested. Intratumoral variation in methylation was observed, though the level of magnitude was smaller than the difference in tumor vs. NE or tumor vs. PC lesions. Fig 1

**Conclusions:** Normal endometrium did not exhibit epigenetic changes identified in EC tumor regions and methylation ITH was observed in EC tumors. Both of these findings suggest molecular changes associated with EC development are focal but heterogeneous.

doi:[10.1016/j.ygyno.2019.03.222](https://doi.org/10.1016/j.ygyno.2019.03.222)

#### Poster #33

##### Clinicopathologic factors associated with increased risk of recurrence in stage IA grade 1 endometrial cancer

C. Nwachukwu, M. Baskovic, D. Fujimoto, R. Von-Eyben, D.P. English, E. Kidd. Stanford University Medical Center, Stanford, CA

**Objectives:** Endometrial cancer (EC) recurrence risk generally depends on pathologic prognostic factors such as stage, grade, depth of myometrial invasion (MI), and lymph node (LN) involvement. FIGO Stage IA grade 1 (IA G1) endometrial cancers lack most adverse features and are categorized in the low recurrence risk group, however, the factors associated with recurrences in this group are not clearly defined though recurrences have been reported in about 5% of cases. The purpose of this study was to identify clinical and pathologic factors that predict for tumor recurrence in IAG1 EC.

**Methods:** We retrospectively reviewed clinical records for EC patients diagnosed between January 1996 and July 2017 at our institution. 127 patients with FIGO 2009 Stage IA grade 1 EC who underwent surgical resection were included. Baseline characteristics were analyzed with chi-square tests. Univariate logistic regression analysis was performed to test for factors that associate with recurrence.

**Results:** Median follow up was 22 months (m). Tumor recurrence was recorded in 12 (8.6%) of patients with median time to