



Analysis of urine-specific antibiograms from veterans to guide empiric therapy for suspected urinary tract infection

Ketzela J. Marsh^{a,*}, Lesley Mundy^b, John J. Holter^b, James R. Johnson^{a,b}

^a University of Minnesota, Minneapolis, MN, USA

^b Minneapolis Veterans Affairs Medical Center, Minneapolis, MN, USA

ARTICLE INFO

Article history:

Received 24 July 2018

Received in revised form 17 July 2019

Accepted 23 July 2019

Available online 30 July 2019

Keywords:

Urinary tract infections

Antimicrobial susceptibility

Escherichia coli

Fosfomycin

ABSTRACT

Urinary tract infection (UTI) is common among patients at Veterans Affairs Medical Centers (VAMCs), many of whom are elderly men with underlying urological problems. Most UTI guidelines address uncomplicated UTI in women, and clinicians may select empiric therapy based on local hospital-wide *Escherichia coli* cumulative susceptibility (antibiogram) data. To inform selection of empiric therapy for UTI at the Minneapolis VAMC (MVAMC), we compiled antimicrobial susceptibility testing (AST) results for 1 year's urine isolates. We analyzed these AST results (bioMérieux VITEK®) for 2494 microbiologically significant urine isolates at MVAMC from June 2013 through May 2014. For antimicrobial–organism combinations that were not tested, we imputed results based on local or published data and/or expert opinion. For ambiguous antimicrobial–organism combinations, we analyzed susceptibility as both 0% and 100%. We calculated cumulative percent susceptible for 26 relevant antimicrobial agents, overall and stratified by Gram stain characteristic and clinical site. The study population included 1548 Gram-negative and 946 Gram-positive urine isolates. Species distribution varied significantly by clinical site. *E. coli* represented only 27% of isolates overall (9–37%, depending on site); also prevalent were *Enterococcus* (14%) and other Gram-positive organisms (23%). Urine-specific antibiograms varied significantly by Gram stain characteristic, between *E. coli* and other Gram-negative organisms, and by clinical site. Of the oral agents, only fosfomycin provided ≥80% susceptibility. Ultimately, *E. coli* represented urine isolates poorly with respect to species distribution and AST results. We conclude that urine-specific antibiograms, stratified by Gram stain characteristic and clinical site, may improve empirical UTI therapy for veterans.

© 2019 Elsevier Inc. All rights reserved.

1. Introduction

Empirical antimicrobial therapy for urinary tract infection (UTI) ideally should provide reliable activity against the patient's urine organism. To accomplish this, therapy must be selected with attention to the likely pathogens and their anticipated antimicrobial susceptibility patterns (Gupta et al. 2011a).

A prominent 2010 international practice guidelines document regarding UTI management addresses empirical therapy selection but is limited to acute uncomplicated cystitis and pyelonephritis in premenopausal women (Gupta et al. 2011a). However, UTI is a significant problem also among veterans, many of whom are elderly men, often with anatomical or functional urinary tract abnormalities. No available guidelines address empiric antimicrobial therapy for UTI in this distinctive population.

Existing guidelines and expert opinion often suggest reliance on the susceptibility patterns of local uropathogens when choosing empiric therapy for UTI (Gupta et al. 2011a; Colgan and Williams 2011; Hooton 2012; Gupta et al. 2011b). However, such data usually are unavailable because laboratories typically report cumulative susceptibility (antibiogram) data stratified by species, not specimen type, and for all samples combined, not stratified by clinical site. Clinicians therefore commonly rely on their local laboratory's overall antibiogram for *E. coli*, the cause of 75–95% of uncomplicated UTI episodes in women (Hooton 2012). Yet, previous studies have shown that susceptibility patterns of *E. coli* at a given institution can vary according to clinical setting such as intensive care unit vs. inpatient ward vs. outpatient sites (Williams et al. 2003; Colgan et al. 2008).

Previous studies of the antibiograms of urine isolates (hereafter, urine-specific antibiograms) from children (Tamma et al. 2014; Dahle et al. 2012; Edlin et al. 2013), inpatients versus outpatients (Saperston et al. 2014), and Emergency Department patients (Jorgensen et al. 2017) found important differences between these setting-specific urine antibiograms and the overall hospital antibiogram, which suggest that use of such data conceivably could improve clinical care. However,

* Corresponding author at: University of Minnesota, 420 Delaware Street SE, MMC #250, Minneapolis, MN 55455. Tel.: +1-612-626-9943.

E-mail address: marsh846@umn.edu (K.J. Marsh).

it remains unclear whether these differences were due to the isolates being urine-specific, setting-specific, or both. To our knowledge, no study has assessed urine-specific antibiograms stratified by clinical site among veterans.

Accordingly, we compiled and analyzed urine-specific antibiograms for the Minneapolis Veterans Affairs Medical Center (MVAMC) using 1 year's antimicrobial susceptibility testing (AST) data. Our goal was to identify appropriate empiric treatment options for UTI in our population, both overall and stratified by clinical site of origin, and to determine whether the *E. coli* antibiogram is a suitable surrogate for a urine-specific antibiogram, either overall or for individual clinical sites.

2. Materials and methods

2.1. Study population

We compiled organism identity and AST results for 2494 (96%) of the 2587 total urine isolates recovered and reported by the MVAMC clinical laboratory from June 2013 through May 2014, excluding only the 93 diphtheroid isolates as presumed contaminants. We were limited by the laboratory's criteria for microbiological significance, according to which organisms are identified only if present at $\geq 100,000$ colony-forming units (CFU) per milliliter for voided urine, and ≥ 5000 CFU/mL for catheterized urine.

Only 1 isolate of a given organism from the same patient within 30 days was analyzed, regardless of source. Of the 2494 isolates, 2109 (85%) had undergone AST by the clinical laboratory for selected agents on a bioMérieux VITEK® instrument; we used these results. For the remaining isolates, which included *Aerococcus urinae* ($n = 11$), *Aerococcus viridans* ($n = 4$), coagulase-negative *Staphylococcus* ($n = 293$; only 1 *S. saprophyticus*), viridans group *Streptococcus* ($n = 86$), *Streptococcus agalactiae* ($n = 43$), and *Streptococcus bovis* ($n = 2$), we imputed antimicrobial susceptibility results, as described below.

2.2. Susceptibility imputations

Because of this study's exploratory nature, we assessed a broadly inclusive list of antimicrobial agents ($n = 26$) that included all agents the MVAMC clinical microbiology laboratory routinely tests against urine organisms, plus fosfomycin. The laboratory performs routine AST only for certain urine organisms (depending on organism identity) and then only for certain organism-specific antimicrobials, which are selected based largely on the organism's Gram stain characteristic. Additionally, based on clinician request, the laboratory also performs AST selectively for specific organisms, agents, or combinations thereof that are not tested routinely. Consequently, for many of the antimicrobial-organism combinations relevant to the present study, few or no directly determined AST results were available. Therefore, for valid overall comparisons between agents regardless of organism identity or Gram stain characteristic, we imputed results for the antimicrobial-organism combinations that the laboratory did not test directly. Susceptibility imputations relied on published data (Masters 2015; *The Sanford guide to antimicrobial therapy* 2018; Traub and Leonhard 1997; Matesanz et al. 2015; Cattoir et al. 2010; Mohan et al. 2017; Rasmussen 2016; Deletoile et al. 2009; Falagas et al. 2009; Falagas et al. 2010a; Falagas et al. 2010b; Karageorgopoulos et al. 2012; Maraki et al. 2009; Popovic et al. 2010; Zhanel et al. 2016), opinions from expert colleagues, local aggregate data from before the study period, and/or data generated during the study period for the subset of study isolates of a given organism type that underwent AST by special request.

We chose not to report AST data for tetracyclines. For Gram-negative organisms, the MVAMC microbiology laboratory does not perform tetracycline AST unless requested by a clinician (which seldom occurs), and the literature contains few relevant data to support imputations. Therefore, despite the availability of tetracycline AST data for some Gram-positive organisms (from both the MVAMC laboratory and the

literature), tetracycline data could not be included in a cumulative report for all urine isolates combined.

Additionally, for certain antimicrobial-organism combinations, because of i) the absence of local data and ii) conflicting expert opinions and published evidence, we performed imputations by assuming both extremes of the plausible range of susceptibility prevalence values. Specifically, for *Enterococcus* species with trimethoprim-sulfamethoxazole (TMP/SMX) (Wisell et al. 2008), and *Staphylococcus saprophyticus* and *Proteus* species with fosfomycin (Zhanel et al. 2016), we imputed both 0% and 100% susceptibility. Likewise, for *Pseudomonas aeruginosa* and fosfomycin, we imputed both 0% and 50% susceptibility (Falagas et al. 2009). We chose the 50% as the upper bound because few of our *Pseudomonas* isolates were extensively multidrug resistant (e.g., 87% were susceptible to ceftipime and 95% to piperacillin/tazobactam).

Overall, our reported susceptibility data (70,632 total isolate-specific antimicrobial-organism combinations) reflect 46% direct determinations, 42% imputations based on historical local data, external published data, and expert opinion, and 12% imputations based on current local data from other isolates within the same organism category.

2.3. Statistical analysis

We calculated the cumulative percent of urine isolates susceptible to each of 26 antimicrobial agents, both overall and stratified by Gram stain characteristic and clinical site, i.e., community residential centers (CRCs), extended care center (ECC), intensive care unit (ICU), inpatient ward, or outpatient. Similar calculations were made for *E. coli* isolates only. Comparisons for percent susceptible were tested using a χ^2 test, with $P < 0.05$ considered statistically significant. Because this was an exploratory analysis, we did not adjust for multiple comparisons, a decision supported by a biostatistician colleague (personal communication, Paul Thuras).

3. Results

3.1. Species distribution

The 2494 urine isolates included 1548 Gram-negative and 946 Gram-positive organisms. *E. coli* accounted for only 27% of isolates overall (by site for 9% ICU to 37% CRC) and 44% of the Gram-negative isolates overall (by site for 13% ICU to 70% CRC). *Enterococcus* and other Gram-positive organisms were also prevalent, both overall (15% and 22%, respectively) and at each site (Table 1).

Statistical analysis showed that the 5 clinical sites differed significantly for the prevalence of each of the main microbial categories, both overall and for at least 1 site compared with the others (Table 1). By contrast, they did not differ significantly for the prevalence of Gram-positive or Gram-negative organisms collectively, or of the minor microbial subsets (i.e., <5% of the total population: *Streptococcus* species and miscellaneous Gram-positive organisms).

3.2. Urine specific-antibiograms overall

Urine-specific antibiograms (for 26 total agents) demonstrated overall percent susceptibility <80% for all oral agents except fosfomycin (fluoroquinolones, 61–68%; TMP/SMX, 57% or 78% depending on *Enterococcus*; nitrofurantoin, 64%) and <90% for all intravenous agents (ceftriaxone, 65%; ertapenem, 69%; imipenem, 89%; piperacillin/tazobactam, 86%) (Table 2). By contrast, fosfomycin exhibited 81–95% overall imputed susceptibility. The urine-specific antibiogram based solely on directly determined data (which were available for only a subset of all antimicrobial-organism combinations) differed significantly from the comprehensive antibiogram that included imputations.

Table 1

Species distribution of urine isolates by clinical site.

Organism	Prevalence of organism, column % ^a						P value ^b					
	All (n = 2494)	CRC (n = 90)	ECC (n = 98)	ICU (n = 33)	WARD (n = 541)	OTPT (n = 1732)	5 sites	All vs. CRC	All vs. ECC	All vs. ICU	All vs. WARD	All vs. OTPT
Gram-positives	37	48	30	27	37	38	0.08	0.06	0.09	0.21	0.80	0.82
<i>Enterococcus</i>	15	8	17	12	20	14	0.003	0.05	0.58	0.61	0.01	0.29
<i>Staphylococcus</i>	16	34	10	15	11	18	<0.001	<0.001	0.09	0.81	0.001	0.29
<i>Streptococcus</i>	5	6	2	0	5	6	0.38	0.90	0.16	0.18	0.97	0.68
Other Gram-positive spp.	1	0	0	0	1	1	0.68	0.46	0.44	0.66	0.40	0.93
Gram-negatives	61	52	70	73	63	62	0.08	0.06	0.09	0.21	0.80	0.82
<i>Escherichia coli</i>	27	37	22	9	17	30	<0.001	0.05	0.30	0.02	<0.001	0.02
<i>Klebsiella spp.</i>	12	7	28	15	13	12	<0.001	0.10	<0.001	0.64	0.77	0.46
<i>Pseudomonas spp.</i>	7	0	8	15	15	4	<0.001	0.01	0.61	0.06	<0.001	0.001
Other Gram-negative spp	15	9	12	33	17	15	0.01	0.08	0.37	0.005	0.35	0.78

^a Site definitions: CRC, community residential centers; ECC, extended care center; ICU, intensive care unit; WARD, inpatient ward; OTPT, outpatient (including Emergency Department and Urgent Care).

^b P values < 0.05 (by χ^2 test) are shown in boldface.

3.3. Urine-specific antibiograms by organism

Urine-specific antibiograms differed significantly between all Gram-negative organisms combined, all Gram-positive organisms combined, and all *E. coli* (Table 2). For *E. coli*, where comparisons were possible (given the limited list of agents for which the hospital laboratory reports results), the urine-specific antibiogram did not differ significantly from the hospital's overall (i.e., all sites and all specimen types) antibiogram from the same period (data not shown). By contrast, among the present urine isolates, the susceptibility results for most of the studied antimicrobials differed significantly between *E. coli* vs. all Gram-negative organisms combined (Table 2).

3.4. Urine-specific antibiograms by clinical site

Urine-specific antibiograms differed significantly across the 5 clinical sites for 22 of the 28 antimicrobial susceptibility end points assessed. The antibiogram for all sites combined differed significantly for at least 1 antimicrobial in comparison with each clinical site's antibiogram and for most antimicrobials in comparison with the antibiograms for CRC and inpatient ward (Table 3). Moreover, for each clinical site except the ICU (where small numbers limited statistical power), the antibiogram for all organisms differed significantly for 1 or more antimicrobials from the same site's *E. coli* antibiogram (data not shown).

Table 2Urine-specific antibiogram by organism type, including *Escherichia coli*.

Antimicrobial	Proportion susceptible, column %				P value ^a			
	All (n = 2494)	Gram-positives (n = 946)	Gram-negatives (n = 1548)	<i>E. coli</i> (n = 679)	All vs. Gram-positives	All vs. Gram-negatives	All vs. <i>E. coli</i>	Gram-negatives vs. <i>E. coli</i>
Amikacin	62	0	100	100	<0.001	<0.001	<0.001	0.10
Ampicillin	39	54	31	56	<0.001	<0.001	<0.001	<0.001
Ampicillin/sulbactam	59	73	51	65	<0.001	<0.001	0.006	<0.001
Penicillin	39	53	31	56	<0.001	<0.001	<0.001	<0.001
Cefazolin	55	37	65	90	<0.001	<0.001	<0.001	<0.001
Cefepime	72	37	94	95	<0.001	<0.001	<0.001	0.48
Cefoxitin	56	37	67	90	<0.001	<0.001	<0.001	<0.001
Ceftazidime	62	10	93	94	<0.001	<0.001	<0.001	0.67
Ceftriaxone	65	37	83	94	<0.001	<0.001	<0.001	<0.001
Ciprofloxacin	61	37	76	72	<0.001	<0.001	<0.001	0.04
Clindamycin	14	38	0	0	<0.001	<0.001	<0.001	N/A ^f
Ertapenem	69	37	88	100	<0.001	<0.001	<0.001	<0.001
Erythromycin	12	33	0	0	<0.001	<0.001	<0.001	N/A
Fosfomycin 100/50 ^b	95	100	92	100	<0.001	<0.001	<0.001	<0.001
Fosfomycin 0 ^c	81	86	79	100	0.003	0.03	<0.001	<0.001
Gentamicin	74	43	93	92	<0.001	<0.001	<0.001	0.22
Imipenem	89	75	97	100	<0.001	<0.001	<0.001	<0.001
Levofloxacin	65	47	75	72	<0.001	<0.001	<0.001	0.11
Linezolid	37	98	0	0	<0.001	<0.001	<0.001	N/A
Moxifloxacin	68	56	75	72	<0.001	<0.001	0.047	0.11
Nitrofurantoin	64	82	53	91	<0.001	<0.001	<0.001	<0.001
Oxacillin	13	35	0	0	<0.001	<0.001	<0.001	N/A
Piperacillin/tazobactam	86	73	94	98	<0.001	<0.001	<0.001	<0.001
Quinupristin/dalfopristin	24	62	0	0	<0.001	<0.001	<0.001	N/A
Tobramycin	58	0	94	91	<0.001	<0.001	<0.001	0.009
Trimethoprim/sulfamethoxazole 100 ^d	78	88	72	77	<0.001	<0.001	0.58	0.02
Trimethoprim/sulfamethoxazole 0 ^e	57	33	72	77	<0.001	<0.001	<0.001	0.02
Vancomycin	37	98	0	0	<0.001	<0.001	<0.001	N/A

^a P values < 0.05 (by χ^2 test) are shown in boldface.

^b Fosfomycin 100/50: *Proteus* species imputed as 100% susceptible, *Staphylococcus saprophyticus* imputed as 100% susceptible, and *Pseudomonas aeruginosa* imputed as 50% susceptible.

^c Fosfomycin 0: *Proteus* species imputed as 0% susceptible, *Staphylococcus saprophyticus* imputed as 0% susceptible, and *Pseudomonas aeruginosa* imputed as 0% susceptible.

^d Trimethoprim/sulfamethoxazole 100: *Enterococcus* species imputed as 100% susceptible.

^e Trimethoprim/sulfamethoxazole 0: *Enterococcus* species imputed as 0% susceptible.

^f N/A: not applicable (0% prevalence in both groups).

Table 3
Urine-specific antibiograms by clinical site.

Antimicrobials	Proportion susceptible, column % ^a						P value ^b					
	All (n = 2494)	CRC (n = 90)	ECC (n = 98)	ICU (n = 33)	WARD (n = 541)	OTPT (n = 1732)	5 sites	All vs. CRC	All vs. ECC	All vs. ICU	All vs. WARD	All vs. OTPT
Amikacin	62	52	68	73	62	62	0.13	0.07	0.19	0.20	0.91	0.85
Ampicillin	39	34	32	28	37	41	.007	0.34	0.12	0.15	0.25	0.33
Ampicillin/sulbactam	59	45	50	48	54	62	<0.001	0.01	0.07	0.21	0.03	0.11
Penicillin	39	34	31	28	37	41	0.06	0.35	0.08	0.16	0.24	0.35
Cefazolin	55	53	53	36	44	58	<0.001	0.83	0.78	0.04	<0.001	0.04
Cefepime	72	62	72	75	67	74	0.008	0.03	0.94	0.67	0.01	0.37
Cefoxitin	56	54	55	42	45	59	<0.001	0.82	0.92	0.13	<0.001	0.03
Ceftazidime	62	55	67	64	61	62	0.57	.23	0.27	0.83	0.72	0.97
Ceftriaxone	65	62	64	57	54	68	<0.001	0.56	0.85	0.36	<0.001	0.04
Ciprofloxacin	61	46	71	71	55	63	<0.001	0.002	0.046	0.18	0.01	0.25
Clindamycin	14	36	6	10	9	16	<0.001	<0.001	0.02	0.39	<0.001	0.22
Ertapenem	69	63	69	63	57	72	<0.001	0.29	0.87	0.54	<0.001	0.02
Erythromycin	12	35	6	5	10	13	<0.001	<0.001	0.03	0.27	0.09	0.42
Fosfomycin 100/50 ^c	95	100	96	92	90	97	<0.001	0.03	0.75	0.25	<0.001	0.04
Fosfomycin 0 ^d	81	90	84	69	72	84	<0.001	0.04	0.55	0.09	<0.001	0.04
Gentamicin	74	82	76	88	68	75	0.002	0.09	0.61	0.07	0.004	0.54
Imipenem	89	68	92	85	90	89	<0.001	<0.001	0.21	0.48	0.40	0.97
Levofloxacin	65	47	71	71	56	67	<0.001	<0.001	0.16	0.33	<0.001	0.08
Linezolid	37	48	29	27	37	38	0.06	0.045	0.08	0.23	0.94	0.89
Moxifloxacin	68	48	74	76	58	71	<0.001	<0.001	0.26	0.34	<0.001	0.053
Nitrofurantoin	64	79	51	42	52	69	<0.001	.004	0.01	0.01	<0.001	0.004
Oxacillin	13	12	6	9	10	14	0.03	0.79	0.04	0.49	0.06	0.37
Piperacillin/tazobactam	86	66	85	81	87	86	<0.001	<0.001	0.71	0.49	0.38	0.69
Quinupristin/dalfopristin	24	43	15	15	18	25	<0.001	<0.001	0.06	0.26	0.01	0.35
Tobramycin	58	48	68	73	78	58	<0.001	0.045	0.049	0.10	<0.001	0.88
Trimethoprim/sulfamethoxazole 100 ^e	78	88	79	74	74	79	0.02	0.03	0.90	0.47	0.07	0.29
Trimethoprim/sulfamethoxazole 0 ^f	57	75	73	62	49	60	<0.001	0.001	0.002	0.72	<0.001	0.16
Vancomycin	37	44	27	27	36	38	0.07	0.15	0.04	0.26	0.77	0.59

^a Site definitions: CRC, community residential centers; ECC, extended care center; ICU, intensive care unit; WARD, inpatient ward; OTPT, outpatient (including Emergency Department and Urgent Care).

^b P values < 0.05 (by χ^2 test) are shown in boldface.

^c Fosfomycin 100/50: *Proteus* species and *Staphylococcus saprophyticus* imputed as 100% susceptible; *Pseudomonas aeruginosa* as 50% susceptible.

^d Fosfomycin 0: *Proteus* species, *Staphylococcus saprophyticus*, and *Pseudomonas aeruginosa* imputed as 0% susceptible.

^e Trimethoprim/sulfamethoxazole 100: *Enterococcus* species imputed as 100% susceptible.

^f Trimethoprim/sulfamethoxazole 0: *Enterococcus* species imputed as 0% susceptible.

4. Discussion

Our analysis of urine-specific antibiograms from the MVAMC clinical microbiology laboratory showed that the laboratory's aggregate susceptibility data for *E. coli*, on which clinicians may rely for empiric antibiotic selection for UTI, represents poorly all urine isolates (either overall or by clinical site), all Gram-negative urine isolates, or even the *E. coli* urine isolates at individual clinical sites. Moreover, *E. coli*, although the leading species, accounted for only a minority of urine isolates, both overall and by site; instead, multiple other Gram-negative species, *Enterococcus*, and other Gram-positive organisms accounted collectively for most isolates. Finally, the urine-specific antibiogram differed significantly, overall and for most clinical sites, from both the urine *E. coli* antibiogram and the contemporaneous hospital *E. coli* antibiogram. Therefore, for construction of antibiograms at our center — and, likely, at other similar centers — *E. coli* is a poor surrogate for the total urine isolate population, and variation between different clinical sites is substantial.

Our analysis failed to identify optimal agents for empiric therapy of UTI at our center. For nearly all agents, the overall percent of urine isolates that were susceptible fell below the guideline-recommended threshold values of 80% for treating lower urinary tract infections and 90% for treating upper urinary tract infections (Gupta et al. 2011a). Fosfomycin, the only exception, exhibited 81–95% overall imputed susceptibility. In the United States, fosfomycin is available only as an oral formulation and is approved only for treatment of lower urinary tract infections. It may be an important empiric treatment option to consider in our veteran population, especially if the patient previously had a multidrug-resistant organism or a mix of Gram-positive and Gram-

negative organisms. Among Gram-negative organisms, the difference in susceptibility prevalence between cefepime (94%) and ceftriaxone (83%) was attributable almost entirely to *Pseudomonas aeruginosa*. This suggests that providers should still consider traditional risk factors for resistant organisms, such as hospital exposure, when choosing a parenteral agent for suspected UTI.

In that regard, the emerging concept of “patient-specific antibiograms” that incorporate patient characteristics such as age, length of hospital stay, comorbid diagnoses, previous antibiotic exposure, and prior AST results may assist with decisions regarding empirical antimicrobial therapy (Overly et al. 2018). Additionally, studies of the clinical efficacy of TMP/SMX against *Enterococcus* and of fosfomycin against urinary organisms generally could clarify the role of these agents in empiric therapy for UTI in veterans and similar patients, reducing the current uncertainty that obliged us to perform sensitivity analyses using extreme values for percent susceptible.

Our study had several limitations. First, the susceptibility data were deidentified, precluding medical record review to determine whether isolates represented symptomatic UTI versus asymptomatic bacteriuria and to assess clinical outcomes in relation to *in vitro* susceptibility. Notably, our objective was to recommend empiric antibiotic options for providers intent on treating for UTI, irrespective of clinical presentation. Many patients are treated for UTI despite not fulfilling standard criteria for symptomatic UTI. Additionally, some Gram-positive organisms (such as coagulase-negative staphylococci) that might be considered contaminants in healthy women can be important pathogens for veterans with indwelling catheters. Although sensitivity analysis showed that removal of data for coagulase-negative staphylococci significantly

altered the antibiogram (not shown), we chose to retain these data given the target patient population. By contrast, removal of data for viridans group *Streptococcus* did not meaningfully alter the antibiogram (not shown).

Second, because 22% of veterans diagnosed with UTI in the outpatient setting at our MVAMC do not have a urine culture done (Drekonja et al. 2010), the isolates studied likely do not represent all diagnosed UTIs in our population. Third, the clinical site recorded for each isolate was the origin of the urine specimen and may not reflect the acuity of the case, if for example the urine sample was collected in a clinic or the emergency department, but the patient ultimately was admitted to an inpatient ward or the ICU. Fourth, the findings might not be generalizable outside of the VA system. Fifth, the absence of susceptibility testing results for many antimicrobial–organism combinations obliged extensive use of imputation, which was limited by the available evidence and expert opinion, although these same limitations apply also in clinical practice.

Our study also had notable strengths. First, its focus on urine isolate-specific AST data may better reflect the susceptibility profiles of uropathogens as opposed to diverse-source pathogens. Second, the large number of urine isolate-specific antimicrobial–organism combinations (70,632), which represented a full year of data across all clinical sites served by MVAMC, allowed an analysis that included the full spectrum of urine culture findings across our institution. Third, MVAMC serves a population mainly of elderly men, many of whom have anatomical or functional urinary tract abnormalities. Because existing UTI guidelines address empiric antibiotic selection only for uncomplicated UTI in women, our findings provide novel important information for providers caring for elderly men.

In conclusion, our findings identify serious limitations in reliance on the existing hospital laboratory antibiogram for guidance in selecting empiric UTI therapy at our institution; contradict the assumption that *E. coli* is a reliable surrogate for urine organisms generally; and show that, at our center, system-wide data apply unreliably to individual clinical sites. Use of urine-specific antibiograms that are stratified by clinical site conceivably could improve empirical selection of UTI therapy for veterans, as could performance of urine Gram stains combined with stratification of urine antibiogram data by Gram stain characteristic.

Acknowledgments

Funding

This work was supported by the National Institute of Child Health and Human Development at the National Institutes of Health [T32 HD068229] and by the National Institute of Allergy and Infectious Diseases at the National Institutes of Health [T32 AI055433], both for K.J.M.

Prior presentations

Findings of the study were presented as posters at *IDWeek 2015* and *IDWeek 2017*. Both meetings were in San Diego, CA, in October 2015 and October 2017, respectively.

Appreciation

We thank Barbara Murray for her expert opinion regarding TMP/SMX and *Enterococcus*. We also thank Dimitri Drekonja for reviewing the manuscript prior to submission.

Potential conflicts of interest

Authors K.J.M., L.M., and J.J.H. have no conflicts. Author J.R.J. has received grants or contracts from Achaogen, Allergan, Melinta, Merck, Shionogi, Syntiron, and Tetrphase; has consultancies with Crucell/

Janssen and Syntiron; and has patent applications for tests to detect specific *E. coli* strains.

References

- Cattoir V, Kobal A, Légrand P. *Aerococcus urinae* and *Aerococcus sanguinicola*, two frequently misidentified uropathogens. *Scand J Infect Dis* 2010;42(10):775–80.
- Colgan R, Williams M. Diagnosis and treatment of acute uncomplicated cystitis. *Am Fam Physician* 2011;84(7):771–6.
- Colgan R, Johnson JR, Kuskowski M, Gupta K. Risk factors for trimethoprim-sulfamethoxazole resistance in patients with acute uncomplicated cystitis. *Antimicrob Agents Chemother* 2008;52(3):846–51.
- Dahle KW, Korgenski EK, Hersh AL, Srivastava R, Gesteland PH. Clinical value of an ambulatory-based antibiogram for uropathogens in children. *J Pediatric Infect Dis Soc* 2012;1(4):333–6.
- Deletoile A, Decre D, Courant S, Passet V, Audo J, Grimont P, et al. Phylogeny and identification of *Pantoea* species and typing of *Pantoea agglomerans* strains by multilocus gene sequencing. *J Clin Microbiol* 2009;47(2):300–10.
- Drekonja DM, Okoye NC, Kuskowski MA, Johnson JR. Appropriateness of urinary tract infection diagnosis and treatment duration. *Arch Intern Med* 2010;170(5):489–90.
- Edlin RS, Shapiro DJ, Hersh AL, Copp HL. Antibiotic resistance patterns of outpatient pediatric urinary tract infections. *J Urol* 2013;190(1):222–7.
- Falagas ME, Kastoris AC, Karageorgopoulos DE, Rafailidis PI. Fosfomycin for the treatment of infections caused by multidrug-resistant non-fermenting Gram-negative bacilli: a systematic review of microbiological, animal and clinical studies. *Int J Antimicrob Agents* 2009;34(2):111–20.
- Falagas ME, Maraki S, Karageorgopoulos DE, Kastoris AC, Kapaskelis A, Samonis G. Antimicrobial susceptibility of Gram-positive non-urinary isolates to fosfomycin. *Int J Antimicrob Agents* 2010a;35(5):497–9.
- Falagas ME, Maraki S, Karageorgopoulos DE, Kastoris AC, Mavromanolakis E, Samonis G. Antimicrobial susceptibility of multidrug-resistant (MDR) and extensively drug-resistant (XDR) Enterobacteriaceae isolates to fosfomycin. *Int J Antimicrob Agents* 2010b;35(3):240–3.
- Gupta K, Hooton TM, Naber KG, Wullt B, Colgan R, Miller LG, et al. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin Infect Dis* 2011a;52(5):e103–20.
- Gupta K, Hooton TM, Miller L. Uncomplicated UTIIGC. Managing uncomplicated urinary tract infection—making sense out of resistance data. *Clin Infect Dis* 2011b; 53(10): 1041–2.
- Hooton TM. Clinical practice. Uncomplicated urinary tract infection. *N Engl J Med* 2012; 366(11): 1028–37.
- Jorgensen S, Zurayk M, Yeung S, Terry J, Dunn M, Nieberg P, et al. Emergency department urinary antibiograms differ by specific patient group. *J Clin Microbiol* 2017;55(9): 2629–36.
- Karageorgopoulos DE, Wang R, Yu XH, Falagas ME. Fosfomycin: evaluation of the published evidence on the emergence of antimicrobial resistance in Gram-negative pathogens. *J Antimicrob Chemother* 2012;67(2):255–68.
- Maraki S, Samonis G, Rafailidis PI, Vouloumanou EK, Mavromanolakis E, Falagas ME. Susceptibility of urinary tract bacteria to fosfomycin. *Antimicrob Agents Chemother* 2009;53(10):4508–10.
- Masters BR. In: Bennett John E, Dolin Raphael, Blaser Martin J, editors. *Mandell, Douglas, and Bennett's principles and practice of infectious diseases*, (2015). Elsevier Saunders: Springer; 2015. ISBN: 13–978–1–4557–4801–3.
- Matesanz M, Rubal D, Iniguez I, Rabunal R, Garcia-Garrote F, Coira A, et al. Is *Streptococcus bovis* a urinary pathogen? *Eur J Clin Microbiol Infect Dis* 2015;34:719–25.
- Mohan B, Zaman K, Anand N, Taneja N. *Aerococcus viridans*: a rare pathogen causing urinary tract infection. *J Clin Diagn Res* 2017;11(1):DR01–DR3.
- Overly S, Mehta J, Hunt LN, Palermo B, Ketcherside WJ, Chen Y, et al. Predictive patient-specific antibiograms could improve precision of empiric antibiotic therapy: a performance evaluation in an academic health system. Society for Healthcare Epidemiology of America (SHEA) Spring Conference. Portland, Oregon. April 2018:18–20.
- Popovic M, Steinort D, Pillai S, Joukhadar C. Fosfomycin: an old, new friend? *Eur J Clin Microbiol Infect Dis* 2010;29(2):127–42.
- Rasmussen M. *Aerococcus*: an increasingly acknowledged human pathogen. *Clin Microbiol Infect* 2016;22(1):22–7.
- Saperston KN, Shapiro DJ, Hersh AL, Copp HL. A comparison of inpatient versus outpatient resistance patterns of pediatric urinary tract infection. *J Urol* 2014;191(5):1608–13. Suppl.
- Tamma PD, Sklansky DJ, Palazzi DL, Swami SK, Milstone AM. Antibiotic susceptibility of common pediatric uropathogens in the United States. *Clin Infect Dis* 2014;59(5): 750–2.
- The Sanford guide to antimicrobial therapy. 48th ed, 2018.
- Traub WH, Leonhard B. Comparative susceptibility of clinical group A, B, C, F, and G beta-hemolytic streptococcal isolates to 24 antimicrobial drugs. *Chemotherapy* 1997;43 (1):10–20.
- Williams DN, Sannes MR, Eckhoff AA, Peterson PK, Johnson JR, Sannes MR, et al. Antimicrobial resistance in *Escherichia coli* causing urinary tract infections in Costa Rica: a clinical dilemma. *Int J Antimicrob Agents* 2003;21(1):79–81.
- Wisell KT, Kahlmeter G, Giske CG. Trimethoprim and enterococci in urinary tract infections: new perspectives on an old issue. *J Antimicrob Chemother* 2008;62(1): 35–40.
- Zhanel GG, Walky AJ, Karlowsky JA. Fosfomycin: a first-line oral therapy for acute uncomplicated cystitis. *Can J Infect Dis Med Microbiol* 2016;2016, 2082693.