



Quality of life of the family of children with asthma is not related to asthma severity

Vaida Taminskiene¹ · Tomas Alasevicius² · Algirdas Valiulis³ · Egle Vaitkaitiene⁴ · Rimantas Stukas¹ · Adamos Hadjipanayis⁵ · Steve Turner⁶ · Arunas Valiulis^{1,2}

Received: 5 July 2018 / Revised: 6 December 2018 / Accepted: 11 December 2018 / Published online: 4 January 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

The quality of life for the family is an important outcome of childhood asthma. The aim of the study was to describe the quality of life in families who have a child with asthma. The Pediatric Quality of Life Inventory Family Impact Module was completed by the parents of 527 children with asthma. The median overall score was 75.0 (interquartile range 63.9, 87.5). The following factors were independently associated with lower quality of life: additional difficulties such as anxiety and financial hardship (3.81 [2.45, 5.93]), waking with asthma symptoms one or more nights a week (odds ratio 2.53 [1.34, 4.75]), regular use of symptoms reliever medication (2.47 [1.57, 3.87]), and female gender (1.97 [1.27, 3.05]). Lower socioeconomic status of the family and exposure to molds at home doubled the odds for lower quality of life. Physician's diagnosed asthma severity and control were associated with quality of life in univariate, but not multivariate analysis.

Conclusion: Multiple factors, several of which are not related to asthma, contribute to the family burden of having a child with asthma. Clinicians should be mindful of the impact of asthma on the child and the family, and consider exploring factors not directly related to childhood asthma.

What is Known:

- Childhood asthma as a chronic disease impacts the quality of life of the patient, but there is also an impact on the immediate family.
- There are relatively few studies exploring the quality of life of parents of a child with asthma; the results are heterogeneous and none has been carried out in an Eastern European country.

What is New:

- This is the first study to describe caregiver's quality of life in an Eastern European population in the context of childhood asthma.
- The quality of life of the family of asthmatic child depends not only on factors related to asthma, but also non-asthma related factors such as poverty which play even more important role.

Communicated by Mario Bianchetti

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00431-018-3306-8>) contains supplementary material, which is available to authorized users.

✉ Arunas Valiulis
arunas.valiulis@mf.vu.lt

Vaida Taminskiene
vaida.taminskiene@mf.vu.lt

Tomas Alasevicius
t.alasevicius@gmail.com

Algirdas Valiulis
algirdval@gmail.com

Egle Vaitkaitiene
egle.vaitkaitiene@ismuni.lt

Rimantas Stukas
rimantas.stukas@mf.vu.lt

Adamos Hadjipanayis
adamos@paidiatros.com

Steve Turner
s.w.turner@abdn.ac.uk

Extended author information available on the last page of the article

Keywords Asthma · Children · Family · Impact · Quality of life

Abbreviations

ACT	Asthma Control Test
C	Communication Scale
CACT	Childhood Asthma Control Test
CF	Cognitive Functioning Scale
CI	Confidence interval
DA	Daily Activities Scale
DALYs	Disability-adjusted life years
EF	Emotional Functioning Scale
FR	Family Relationships Scale
IQR	Interquartile range
OR	Odds ratio
PedsQLFIM	Pediatric Quality of Life Inventory Family Impact Module
PF	Physical Functioning Scale
SF	Social Functioning Scale
QoL	Quality of life
W	Worry scale

Introduction

Asthma is a global public health problem [18] and is one of the most common chronic diseases of childhood [2]. Asthma management in childhood is essential for better overall health in youth and in later life [3]. There is a human cost to asthma evidenced as reduced quality of life. Asthma can disturb everyday life of the patient, limits physical abilities, and causes emotional and economic consequences [2]. The burden of asthma, when measured as disability-adjusted life years (DALYs), accounts for 1.1% of the global DALYs lost [21]. Asthma is among the 20 most common conditions which affect DALYs across all ages, and in children is in the top ten conditions affecting DALYs.

Childhood asthma can also affect the life of the child's family members [24]. Children with chronic diseases such as asthma require more time, care, and attention from their parents compared to children without these diagnoses [26]. Across a wide range of conditions, parents who have children with chronic conditions report increased levels of stress and having to make changes to their personal and family life to meet their child's health needs [24].

Our group and others have previously shown that the quality of life (QoL) of children with asthma is reduced in association with increasing asthma severity and poor symptom control [22, 23]. Our understanding of how pediatric asthma impacts on QoL of the family is incomplete [12, 24] and limited to Western countries. The aim of our study was to assess in an Eastern European country the QoL in families where there is a child with asthma.

Materials and methods

Study population and data collection

This study was a part of a cross-sectional study of QoL in Lithuanian children with asthma and their parents, and our methodology is previously described [23]. Parents of children with asthma aged 2–17 years were asked to participate during the scheduled outpatient visit to pediatric pulmonologist. According to national guidelines, children with mild asthma in Lithuania visit pediatric pulmonologist at least once a year and more frequently if they have moderate or severe asthma; clinicians categorized asthma severity as mild, moderate, or severe. Study data were collected in six polyclinics in the two largest cities of Lithuania during the period between December 2014 and July 2016.

Family quality of life tool

Lithuanian version of Pediatric Quality of Life Inventory Family Impact Module (PedsQLFIM) was used to determine QoL for families [26]. This questionnaire (used with the permission of Mapi Research Institute) consists of 36 questions from which the following six subscales are derived each of which describe the disease impact on the family's QoL: Physical Functioning; Emotional Functioning; Social Functioning; Cognitive Functioning; Communication; Worry; and two for the functioning of the whole family (Daily Activities and Family Relationships). Scores ranging from 0 to 100 for each scale as well as overall score were calculated, with lower scores indicating greater impact on family life. The PedsQLFIM overall score for parents who have children with no chronic condition is typically > 80 [16].

Respiratory and demographic questionnaire was completed by parents to determine their characteristics, including socioeconomic, domestic, and environmental factors, and details of their child's asthma, associated conditions, and treatment details (Supplement 1). "Other allergies" was defined as an affirmative response to the questions related to allergic rhinitis, food allergy, and skin rashes.

Childhood Asthma Control Tests (CACT) or Asthma Control Tests (ACT) were used as appropriate to measure asthma control. ACT scores (for children aged > 11 years) range from 5 to 25, while CACT scores (for children aged 4–11 years) from 0 to 27, and scores ≤ 15 were defined as "uncontrolled asthma," scores between 16 and 19 were defined as "partly controlled asthma" and scores > 19 defined as "controlled asthma" [6].

Statistical analysis

Spearman correlation, Mann–Whitney and Kruskal–Wallis tests were performed as appropriate for statistical analysis. As previously [23], PedsQLFIM scores were divided into terciles and the lowest tercile was the reference group (scores ≤ 68.06) indicating the highest asthma impact on family life. Binary logistic regression was used to identify risk factors associated with low family QoL. Variables significantly associated with PedsQLFIM scores in the univariate analysis were selected for the multivariate analysis. The final binary logistic regression model includes only statistically significant covariates; descriptive statistics of the model were also considered.

Data analysis was performed using SPSS (version 22.0; IBM Corporation, New York, NY, USA). All p values ≤ 0.05 were considered as statistically significant.

Results

Participants of the study

There were 807 parents of children with asthma invited to participate, and questionnaires were completed by 527 (65.3%).

The median age [interquartile range (IQR)] of children with asthma was 8.0 (5.0, 12.0) years, and the majority were boys (63.2%). Most of children had mild asthma and well-controlled disease (Table 1). The majority (60.9%) of children were also diagnosed with other allergies. A total of 81.9% of mothers and 91.6% of fathers had permanent jobs. More than half of respondents had an income of less than 300 euros per month per family member. Thirty percent of respondents were exposed to mold at home, and of these, half had visible mold in the bathroom, and 20% had mold in the bedrooms, living rooms, or kitchen. Further characteristics of the study participants are presented in Table 1.

Childhood asthma impact on family life

The overall median [IQR] PedsQL Family Impact Module score was 75.0 [63.9, 87.5]. The lowest scores were for the Worry scale (60.0 [45.0, 75.0]) and the highest for Communication (91.7 [66.7, 100.0]). Parents indicated increased anxiety due to asthma treatment efficiency and side effects, child's future; see Supplement 2. They reported how family activities require more time and effort because of child's asthma. Parents rarely complained about the following: nausea, disturbed memory, difficulties in solving family problems, or inability to tell about their problems and feelings to a doctor or a nurse. Childhood asthma had a greater negative impact on parents QoL compared to functioning of the whole family ($p < 0.001$): overall mean score of six scales measuring parent self-reported functioning was 73.9 (± 17.2) and overall mean

score of two scales measuring parent-reported family functioning was 76.0 (± 18.2). Overall PedsQLFIM score was lower in parents of children aged from 2 to 9 years old compared to children aged > 9 years, but this difference was not significant, except scales of Emotional Functioning and Daily Activities.

Factors associated with childhood asthma impact on family life

In univariate analyses, PedsQLFIM overall score was associated with asthma severity and control, presence of asthma symptoms during the last year, hospitalization due to asthma within the last 6 months, use of rescue inhalers, and presence of other allergies (Table 2). Parents who reported humidity and molds at their homes had lower PedsQLFIM scores as well. Parents with lower QoL experienced more difficulties because of child's disease: additional anxiety, financial costs, and difficulties to balance their professional and personal life. Girls, asthmatic children with worse general health condition, as well as children from families getting social support and having lower income had lower PedsQLFIM score. The overall PedsQLFIM score was not associated with whether the parents lived together or separately, the child's age, or exposure to pets and second-hand smoking (Table 2). Supplement 2 presents results for the six individual scales.

In the multivariate analysis (Table 3), lower PedsQLFIM was independently linked to asthma symptoms at night during the last year, use of rescue medicine, lower socioeconomic status, female gender, exposure to molds at home, and additional difficulties in a family because of child's disease (including increased tension and anxiety in family, financial hardships, impaired balance of personal, and professional life).

Discussion

Childhood asthma is a chronic disease which is known to affect the patient's life [10], and this study evaluated how characteristics of a child's asthma impact on QoL of the child's family members using the PedsQLFIM. The overall PedsQLFIM score of parents of children with asthma was lower in our population compared to other studies of children with no illness [16], and the score was higher compared to children with other complex chronic health conditions [26], oncology [1], and chronic gastrointestinal disorders [16].

This study confirms results from other studies in families with a child with conditions other than asthma that chronic childhood disease affects the whole family [1, 26], but replicates this earlier work in an Eastern European country.

In our study, the factors associated with poorer family QoL were gender, "additional difficulties" (including increased tension and anxiety in family, financial hardships, impaired balance of personal and professional life), financial problems,

Table 1 Descriptive characteristics of study participants

Asthma severity	% mild (<i>n</i>)	71.4 (376)
	% moderate (<i>n</i>)	25.6 (135)
	% severe (<i>n</i>)	3.0 (16)
Asthma control	% controlled (<i>n</i>)	62.9 (293)
	% partly controlled (<i>n</i>)	23.4 (109)
	% not controlled (<i>n</i>)	13.7 (56)
Other allergies	% allergic rhinitis (<i>n</i>)	37.3 (196)
	% food allergy (<i>n</i>)	11.0 (58)
	% allergic rhinitis and food allergies (<i>n</i>)	12.7 (67)
	% no (<i>n</i>)	32.1 (169)
	% do not know (<i>n</i>)	6.9 (36)
Source of family income	% permanent job income (<i>n</i>)	95.1 (501)
	% social benefits (<i>n</i>)	11.8 (62)
Income per family member	% up to 450 euros (<i>n</i>)	56.7 (299)
	% 451–600 euros (<i>n</i>)	10.1 (53)
	% more than 600 euros (<i>n</i>)	10.6 (56)
	% do not want to indicate (<i>n</i>)	22.6 (119)
Child's general health	% very good (<i>n</i>)	11.4 (60)
	% good (<i>n</i>)	37.2 (196)
	% moderate (<i>n</i>)	31.5 (166)
	% satisfactory (<i>n</i>)	16.1 (85)
	% poor (<i>n</i>)	3.4 (18)
	% very poor (<i>n</i>)	0.4 (2)
	% males (<i>n</i>)	63.2 (333)
	% children living with both parents (<i>n</i>)	79.1 (417)
	% mothers with higher education (<i>n</i>)	59.3 (310)
	% fathers with higher education (<i>n</i>)	46.0 (223)
	% children who felt short of breath during the last year (<i>n</i>)	83.1 (438)
	% children who woke up at night because of asthma during the last year (<i>n</i>)	70.1 (369)
	% children who used symptom relievers during the last month (<i>n</i>)	66.8 (352)
	% children who have pets at home (<i>n</i>)	42.3 (223)
	% children exposed to smoking at home (<i>n</i>)	22.0 (116)
	% children exposed to molds at home (<i>n</i>)	30.2 (159)
	% children hospitalized due to asthma within last 6 months (<i>n</i>)	16.1 (85)
	% major difficulties due to asthma (<i>n</i>)	44.6 (235)

and the frequency of nocturnal symptoms and of reliever medication use. Some of the findings in this study are consistent with previous publications in Western populations. For example, the presence and frequency of asthma symptoms and use of rescue inhalers are related to poor parental QoL [12]. Nocturnal asthma symptoms may disturb the sleep of all family members and increase the risk of parents not attending a job the next day [8].

Childhood asthma is a multifactorial disease [17] and not unexpectedly the family's quality of life is associated with many factors, some of which are not primarily asthmatic. Poverty is recognized to be associated with asthma outcomes [27], patient's [22], and caregiver's [9] QoL. Exposure to molds at home was a determinant of higher childhood asthma impact on the family, and although not linked to family QoL

elsewhere, the presence of mold is known to be associated with asthma symptoms [14] and can also be a sign of poor quality housing and an index of poverty. Presence of concomitant allergies is also a known risk factor of asthma [4], but in this study, it was associated with family's health-related QoL only in univariate, but not in multivariate analysis. We observed an association between female sex and reduced QoL of parents, and this is consistent with studies in Spain and Greece which found slightly higher QoL scores in boys with asthma compared to girls [11, 19]. A lower QoL in girls compared to boys may be represented by different psychological response to asthma [7, 19].

We have previously reported factors associated with QoL of 5–11-year-old children with asthma [23] whose parents participated in the present study, and some findings are not

Table 2 Median PedsQL Family Impact Module scores by social, environment, and lifestyle factors

Factors		N	PedsQLFIM score
Total		527	75.0
Gender	Males	333	76.4
	Females	194	71.9*
Asthma Severity	Mild	376	77.1
	Moderate	135	72.2
	Severe	16	62.8**
Asthma Control	Controlled	293	78.5
	Partly controlled	109	70.1
	Not controlled	56	67.4**
Woke up at night because of asthma during the last year	Every week	119	68.8**
	< 1 time per week	249	74.3
	None	157	84.7
Short breath during the last 12 months	Yes	327	72.2**
	No	168	85.4
General child's health	Very good	60	87.2
	Good	196	80.6
	Moderate	166	70.1
	Satisfactory	85	71.5
	(Very) poor	20	57.9**
Use of symptom relievers everyday	Yes	160	69.4**
	No	367	77.1
Family income	≤ 450 euros	299	74.3
	451–600 euros	53	75.0
	≥ 600 euros	56	82.6*
Main source of family income	Permanent job	501	75.0
	Social benefits	61	65.3**
Major difficulties due to asthma	Yes	235	68.1**
	No	292	83.3
Hospitalization within last 6 months	Yes	85	71.5*
	No	442	75.7
Other allergies (allergic rhinitis, food allergies)	Yes	321	73.6*
	No	169	79.9
Child lives with both parents	Yes	417	75.8
	No	110	72.9
Pets at home	Yes	223	75.0
	No	304	75.0
Exposure to smoking	Yes	116	75.0
	No	411	75.0
Molds at home	Yes	159	72.9*
	No	368	76.0

PF physical functioning, *EF* emotional functioning, *SF* social functioning, *CF* cognitive functioning, *C* communication, *W* worry, *DA* daily activities, *FR* family relationships

* $p < 0.05$; ** $p < 0.001$ (Mann–Whitney and Kruskal–Wallis tests)

consistent between the two studies. For example, the presence and frequency of nocturnal asthma symptoms and use of reliever medication were related to patient's and caregiver's QoL, but the child's QoL was related to shortness breath and not nocturnal symptoms or use of reliever medication [23]. In

contrast, asthma control, severity, and general child's health condition were associated with QoL of child but not parental QoL, whilst gender, additional difficulties in family, social support, and exposure to molds were associated with parental QoL but not the child's QoL [23].

Table 3 Associations of lower PedsQL Family Impact Module scores with possible risk factors in multivariate logistic regression

	OR (95%CI)	<i>p</i>
Woke up at night because of asthma during the last 12 months		
Every week	2.53 (1.34–4.75)	0.004
< 1 time per week	1.99 (1.15–3.43)	0.014
None	1	
Gender		
Female	1.97 (1.27–3.05)	0.002
Male	1	
Use of symptom relievers (almost) every day during the last month		
Yes	2.47 (1.57–3.87)	< 0.001
No	1	
Additional difficulties caused by the child's disease		
Yes	3.81 (2.45–5.93)	< 0.001
No	1	
Social benefits as the main source of income		
Yes	1.99 (1.05–3.80)	0.035
No	1	
Molds at home		
Yes	2.03 (1.29–3.19)	0.002
No	1	

$N = 382$; Wald- $\chi^2 = 35.24$; $p < 0.001$; Hosmer-Lemeshow test $\chi^2 = 7.715$; $df = 8$; $p = 0.462$; correctly classified 65.4%; Nagelkerke $R^2 = 0.340$
CI confidence interval, *OR* odds ratio

Asthma severity and control were not related to family QoL in the multivariate analysis but were associated with the child's QoL in our previous work [23], and also other studies [15, 19]. Gent et al. found that asthma symptoms impair QoL of children and caregivers' regardless of the doctor's confirmed asthma diagnosis [25]. One reason for this apparent inconsistency may be due to two specific questions related to asthma impact on family QoL (i.e., frequency of nocturnal asthma symptoms and regular use of symptom relief) which subsumed the univariate relationships between QoL and control and severity. An alternative explanation is that the parent's perception of parental QoL is different to child's perception of their own QoL.

Our study has a number of strengths and limitations. To our knowledge, this is the first study to use the PedsQLFIM in the setting of childhood asthma. A second strength is that although parental QoL has been described in another Eastern European country [24], this remains a relatively under-researched geographical area. A limitation is that QoL was not ascertained in children without asthma in our study, so we did not have a local "control" population for comparison, but QoL scores were lower than in studies of parents whose children do not have chronic conditions [16], and our focus was on determinants of QoL within an asthma population. A second limitation is that questionnaire completion rate was relatively low (65.3%), and this may have introduced some

bias into the population upon which this study is based. Another limitation of our study was that questionnaires were completed only by one parent (usually mother), and fathers can assess problems caused by child's health condition differently to mothers with the latter indicating a higher impact on family well-being [13, 20]. One more limitation is that we did not consider ethnicity which may be related to QoL, and this may be important since language barriers in ethnical minorities may result in poorer asthma management [5].

In summary, we demonstrate that a child's asthma impacts caregiver's QoL. Better appreciation by clinicians of the impact of a child's on family life (and how this impact may be lessened) may improve outcomes for both child and family. Multiple factors contribute to the burden of having a child with asthma on immediate family members, and clinicians should be mindful not only of the impact of asthma on the child and the family, but also consider exploring factors not directly related to childhood asthma.

Acknowledgments Authors express gratitude to participants of this study (children with diagnosed asthma and their parents) who agreed to participate in a study and complete questionnaires.

Authors' contributions VT developed study protocol, collected data, performed data analysis, and wrote the manuscript. TA collected data and performed data analysis. AlgV contributed to data analysis; EV developed study protocol, collected data, and reviewed the manuscript. RS supervised the design and execution of the study. AH reviewed the manuscript. ST performed data analysis and reviewed the manuscript. ArV supervised the design and execution of the study, contributed to data collection and analysis, and reviewed the manuscript.

Compliance with ethical standards

Ethical approval The study was approved by Vilnius regional ethical committee, ref. no 158200-14-749-265.

Informed consent Informed consent forms to participate in a study were signed by parents and children from 8 years old.

Conflict of interest The authors declare that they have no conflict of interests.

References

1. Al-Gamal E, Long T (2016) Psychometric properties of the Arabic version of the PedsQLTM family impact scale. *J Res Nurs* 21:609–610. <https://doi.org/10.1177/1744987116670204>
2. Asher I, Pearce N (2014) Global burden of asthma among children. *Int J Tuberc Lung Dis* 18:1269–1278. <https://doi.org/10.5588/ijtld.14.0170>
3. Bousquet J, Addis A, Agache I, Agustí A et al (2014) Integrated care pathways for airway diseases (AIRWAYS-ICPs). European innovation partnership on active and healthy ageing, action plan B3. Mechanisms of the development of allergy (MeDALL, WP7), GARD (global Alliance against chronic respiratory diseases, WHO) research demonstration project. *Eur Respir J* 44:304–323. <https://doi.org/10.1183/09031936.00014614>

4. Bousquet J, Arnavielhe S, Bedbrook A et al (2018) MASK 2017: ARIA digitally-enabled, integrated, person-centered care for rhinitis and asthma multimorbidity using real-world-evidence. *Clin Transl Allergy* 8:45. <https://doi.org/10.1186/s13601-018-0227-6>
5. Chan KS, Keeler E, Schonlau M, Rosen M, Mangione-Smith R (2005) How do ethnicity and primary language spoken at home affect management practices and outcomes in children and adolescents with asthma? *Arch Pediatr Adol Med* 159:283–289. <https://doi.org/10.1001/archpedi.159.3.283>
6. Cloutier MM, Schatz M, Castro M, Clark N, Kelly HW, Mangione-Smith R, Sheller J, Sorkness C, Stoloff S, Gergen P (2012) Asthma outcomes: composite scores of asthma control. *J Allergy Clin Immunol* 129:S24–S33. <https://doi.org/10.1016/j.jaci.2011.12.980>
7. Di Marco F, Verga M, Santus P, Giovannelli F, Busatto P, Neri M, Girbino G, Bonini S, Centanni S (2010) Close correlation between anxiety, depression, and asthma control. *Resp Med* 104:22–28. <https://doi.org/10.1016/j.rmed.2009.08.005>
8. Diette GB, Markson L, Skinner EA, Nguyen TT, Algatt-Bergstrom P, Wu AW (2000) Nocturnal asthma in children affects school attendance, school performance, and parents' work attendance. *Arch Pediatr Adol Med* 154:923–928. <https://doi.org/10.1001/archpedi.154.9.923>
9. Everhart RS, Fiese BH, Smyth JM (2008) A cumulative risk model predicting caregiver quality of life in pediatric asthma. *J Pediatr Psychol* 33:809–818. <https://doi.org/10.1093/jpepsy/jsn028>
10. Fisch SM, Neininger MP, Prenzel F, Bernhard MK, Hornemann F, Merckenschlager A, Kiess W, Bertsche T, Bertsche A (2018) Experiences, expectations, and fears of adolescents with epilepsy or bronchial asthma. *Eur J Pediatr* 177:1451–1457. <https://doi.org/10.1007/s00431-018-3200-4>
11. Garcia-Marcos L, Carvajal Uruena I, Escribano Montaner A, Fernandez Benitez M, Garcia de la Rubia S, Tauler Toro E, Pérez Fernández V, Barcina Sánchez C (2007) Seasons and other factors affecting the quality of life of asthmatic children. *J Invest Allerg Clin* 17:249–256
12. Halterman JS, Yoos HL, Conn KM, Callahan PM, Montes G, Neely TL, Szilagyi PG (2004) The impact of childhood asthma on parental quality of life. *J Asthma* 41:645–653. <https://doi.org/10.1081/JAS-200026410>
13. Jastrowski Mano KE, Khan KA, Ladwig RJ, Weisman SJ (2009) The impact of pediatric chronic pain on parents' health-related quality of life and family functioning: reliability and validity of the PedsQL 4.0 family impact module. *J Pediatr Psychol* 36:517–527
14. Jones R, Recer GM, Hwang SA, Lin S (2011) Association between indoor mold and asthma among children in Buffalo, New York. *Indoor Air* 21:156–164. <https://doi.org/10.1111/j.1600-0668.2010.00692.x>
15. Karadeniz P, Özdoğan Ş, Ayyıldız-Emecen D, Öncül Ü (2016) Asthma control test and pediatric asthma quality of life questionnaire association in children with poor asthma control. *Turkish J Pediatr* 58:464–472. <https://doi.org/10.24953/turkjpmed.2016.05.002>
16. Knez R, Stevanovic D, Vulić-Prtorić A, Vlašić-Cicvarić I, Peršić M (2015) The Croatian version of the pediatric quality of life inventory (PedsQL™) family impact module: cross-cultural adaptation and psychometric evaluation. *J Child Fam Stud* 24:363–371. <https://doi.org/10.1007/s10826-013-9844-9>
17. Mazenq J, Dubus JC, Gaudart J, Charpin D, Nougairess A, Viudes G, Noel G (2017) Air pollution and children's asthma-related emergency hospital visits in southeastern France. *Eur J Pediatr* 176:705–711. <https://doi.org/10.1007/s00431-017-2900-5>
18. Nunes C, Pereira AM, Morais-Almeida M (2017) Asthma costs and social impact. *Asthma Res Pract* 3(1):1. <https://doi.org/10.1186/s40733-016-0029-3>
19. Petsios KT, Priftis KN, Hatziagorou E, Tsanakas JN, Antonogeorgos G, Matziou VN (2013) Determinants of quality of life in children with asthma. *Pediatr Pulmonol* 48:1171–1180. <https://doi.org/10.1002/ppul.22768>
20. Sigurdardottir AO, Garwick AW, Svavarsdottir EK (2017) The importance of family support in pediatrics and its impact on healthcare satisfaction. *Scand J Caring Sci* 31:241–252. <https://doi.org/10.1111/scs.12336>
21. Soriano JB, Abajobir AA, Abate KH, Abera SF, Agrawal A, Ahmed MB, Aichour AN, Aichour I, Aichour MTE, Alam K, Alam N, Alkaabi JM, al-Maskari F, Alvis-Guzman N, Amberbir A, Amoako YA, Ansha MG, Antó JM, Asayesh H, Atey TM, Avokpaho EFGA, Barac A, Basu S, Bedi N, Bensenor IM, Berhane A, Beyene AS, Bhutta ZA, Biryukov S, Boneya DJ, Brauer M, Carpenter DO, Casey D, Christopher DJ, Dandona L, Dandona R, Dharmaratne SD, Do HP, Fischer F, Gebrehiwot TT, Geleto A, Ghoshal AG, Gillum RF, Ginawi IAM, Gupta V, Hay SI, Hedayati MT, Horita N, Hosgood HD, Jakovljevic M(M)B, James SL, Jonas JB, Kasaeian A, Khader YS, Khalil IA, Khan EA, Khang YH, Khubchandani J, Knibbs LD, Kosen S, Koul PA, Kumar GA, Leshargie CT, Liang X, el Razek HMA, Majeed A, Malta DC, Manhertz T, Marquez N, Mehari A, Mensah GA, Miller TR, Mohammad KA, Mohammed KE, Mohammed S, Mokdad AH, Naghavi M, Nguyen CT, Nguyen G, le Nguyen Q, Nguyen TH, Ningrum DNA, Nong VM, Obi JI, Odeyemi YE, Ogbo FA, Oren E, PA M, Park EK, Patton GC, Paulson K, Qorbani M, Quansah R, Rafay A, Rahman MHU, Rai RK, Rawaf S, Reinig N, Safiri S, Sarmiento-Suarez R, Sartorius B, Savic M, Sawhney M, Shigematsu M, Smith M, Tadese F, Thurston GD, Topor-Madry R, Tran BX, Ukwaja KN, van Boven JFM, Vlassov VV, Vollset SE, Wan X, Werdecker A, Hanson SW, Yano Y, Yimam HH, Yonemoto N, Yu C, Zaidi Z, el Sayed Zaki M, Lopez AD, Murray CJL, Vos T (2017) Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet Respir Med* 5:691–706. [https://doi.org/10.1016/S2213-2600\(17\)30293-X](https://doi.org/10.1016/S2213-2600(17)30293-X)
22. Taminskiene V, Mukhopadhyay S, Palmer C, Mehta A, Ayres J, Valiulis A, Turner SW (2016) Factors associated with quality of life in children with asthma living in Scotland. *Pediatr Pulmonol* 51:484–490. <https://doi.org/10.1002/ppul.23359>
23. Taminskiene V, Vaitkaitiene E, Valiulis A, Turner S, Hadjipanayis A, Stukas R, Valiulis A (2018) The self-reported quality of life of Lithuanian children with asthma was comparable to Western populations. *Acta Paediatr* 107:333–338. <https://doi.org/10.1111/apa.14140>
24. Trzcieniecka-Green A, Bargiel-Matusiewicz K, Wilczyńska A, Omar HA (2015) Quality of life of parents of children with asthma. *Int J Child Adolesc Health* 8:351–355. https://uknowledge.uky.edu/pediatrics_facpub/187
25. van Gent R, van Essen LEM, Rovers MM, Kimpen JLL, van der Ent CK, de Meer G (2007) Quality of life in children with undiagnosed and diagnosed asthma. *Eur J Pediatr* 166:843–848. <https://doi.org/10.1007/s00431-006-0358-y>
26. Varni JW, Sherman SA, Burwinkle TM, Dickinson PE, Dixon P (2004) The PedsQL™ family impact module: preliminary reliability and validity. *Health Qual Life Out* 2:55. <https://doi.org/10.1186/1477-7525-2-55>
27. Williams DR, Sternthal M, Wright RJ (2009) Social determinants: taking the social context of asthma seriously. *Pediatrics* 123:S174–S184. <https://doi.org/10.1542/peds.2008-2233H>

Affiliations

Vaida Taminskiene¹  · Tomas Alasevicius² · Algirdas Valiulis³ · Egle Vaitkaitiene⁴ · Rimantas Stukas¹ · Adamos Hadjipanayis⁵ · Steve Turner⁶ · Arunas Valiulis^{1,2}

¹ Department of Public Health, Institute of Health Sciences, Faculty of Medicine, Vilnius University, Vilnius, M.K Ciurlionio str. 21, 03101 Vilnius, Lithuania

² Clinic of Children's Diseases, Institute of Clinical Medicine, Faculty of Medicine, Vilnius University, Antakalnio str. 57, 10207 Vilnius, Lithuania

³ Department of Rehabilitation, Physical and Sports Medicine, Institute of Health Sciences, Faculty of Medicine, Vilnius University, Vilnius, M.K Ciurlionio str. 21, 03101 Vilnius, Lithuania

⁴ Medical Academy, Lithuanian University of Health Sciences, Tilzes str. 18, 47181 Kaunas, Lithuania

⁵ European University of Cyprus, 6, Diogenis Str, 2404 Engomi, Nicosia, Cyprus

⁶ Child Health, University of Aberdeen, Aberdeen AB25 2ZG, UK