

This feasibility study, describes a novel wristed needle driver in a series of LPN patients. Certainly, the small sample size (20 patients) and the fact that all procedures were performed by a single surgeon is a limitation. The authors found that use of the novel needle driver was associated with higher blood loss (200 mL vs 50 mL,  $P = .03$ ) compared to robotic surgery and the transfusion rate (30%) and positive margin rates (15%) were higher than those reported in some other series.<sup>8,9</sup> However, to the authors credit this was their initial experience with this technology and these outcomes will certainly improve with greater experience. A promising finding was that the warm ischemia time was similar between groups (LPN: 25.5 minutes vs RPN: 18.5 minutes,  $P = .36$ ).

These authors should be congratulated for attempting to reduce the costs of PN. With healthcare consuming 18% of the gross domestic product in the United States, and 22% of the current budgetary expenses for 2018, devices like this experimental needle driver have the potential to lower the cost of minimally invasive PN.<sup>10</sup> In addition, this type of device could facilitate minimally invasive PN in regions where surgical robots are not available. Certainly, larger, multicenter, prospective trials will be required to determine whether this needle driver will provide equivalent outcomes to robotic surgery but at a lower cost. In addition, comparisons between the novel needle driver and conventional LPN will be necessary to determine whether this device truly can shorten the learning curve and improve the outcomes of LPN. Technologies such as FlexDex have potential to make PN more readily available and allow for preservation of renal function to reduce risk of chronic kidney disease among this patient population.<sup>11</sup>

**Mohammad Hajiha, D. Duane Baldwin**, Department of Urology, Loma Linda University Health, Loma Linda, CA

## References

1. Cacciamani GE, Medina LG, Gill T, et al. Impact of surgical factors on robotic partial nephrectomy outcomes: comprehensive systematic review and meta-analysis. *J Urol*. 2018;200:258–274.
2. Wu Z, Li M, Liu B, et al. Robotic versus open partial nephrectomy: a systematic review and meta-analysis. *Plos ONE*. 2014;9:e94878.
3. Biehn Stewart S, Reed SD, Moul JW. Will the future of health care lead to the end of the robotic golden years? *Eur Urol*. 2014;65:325–328.
4. Hyams E, Pierorazio P, Mullins J K, Ward M, Allaf M. Laparoscopy and robotic surgery a comparative cost analysis of robot-assisted versus traditional laparoscopic partial nephrectomy. *JEndo*. 2012;26.
5. Mottrie A, Borghesi M, Ficarra V. Is traditional laparoscopy the real competitor of robot-assisted partial nephrectomy? *Eur Urol*. 2012;62:1034–1036.
6. Poon SA, Silberstein JL, Chen LY, Ehdaie B, Kim PH, Russo P. Trends in partial and radical nephrectomy: an analysis of case logs from certifying urologists. *J Urol*. 2013;190:464–469.
7. Y. Yang D, Monn M F, Bahler C D, Sundaram C P. Does robotic assistance confer an economic benefit during laparoscopic radical nephrectomy? *JUro*. 2014. Presented at annual meeting of American Urological Association, Orlando, Florida, May 16-21.
8. Benway B M, Bhayani S B, Rogers C G, et al. Robot assisted partial nephrectomy versus laparoscopic partial nephrectomy for renal tumors: a multi-institutional analysis of perioperative outcomes is accompanied. *JUrol*. 2009;182:866–873.
9. Leow JJ, Heah NH, Chang SL, Chong YL, Png KS. Outcomes of robotic versus laparoscopic partial nephrectomy: an updated meta-analysis of 4,919 patients. *J Urol*. 2016;196:1371–1377.

10. National Health Expenditures 2017 Highlights, Center for medicare & medicaid services, Page last Modified: 12/11/2018, <https://www.cms.gov/research-statistics-data-and-systems/statistics-trends-and-reports/nationalhealthexpenddata/nationalhealthaccountshistorical.html>.
11. Lau W KO, Blute M L, Weaver A L, Torres V E, Zincke H. Matched comparison of radical nephrectomy vs nephron-sparing surgery in patients with unilateral renal cell carcinoma and a normal contralateral kidney. *Mayo Clin Proc*. 2000;75:1236–1242.

<https://doi.org/10.1016/j.urology.2019.05.051>  
UROLOGY 132: 127–128, 2019. © 2019 Elsevier Inc.

## AUTHOR REPLY



We thank the editors for their thoughtful comments regarding our study.<sup>1</sup> These authors correctly note that robotic surgery has improved postoperative outcomes including shorter length of stay, decreased blood loss while sustaining curative extirpation for partial nephrectomy (PN) as compared to an open technique.<sup>2,3</sup> The rapid adoption of robotic surgery since the Food and Drug Administration approval of the Intuitive Surgical platform in 2000 has accelerated the utilization of minimally invasive techniques for PN. However, this transition should not halt innovation and utilization with laparoscopic instruments in renal surgery.

As healthcare providers, it is important to be good stewards of healthcare resource utilization, balancing the potential benefits of novel innovations while also being cognizant of the costs associated with these new technologies.<sup>4</sup> In this study we have proposed the use of a new technology in a resource limited hospital to perform a more minimally invasive technique for PN.

The authors point out that laparoscopic PN can be very technically challenging and is associated with a significant learning curve. Given that our study represents our initial experience with this technology, it is likely that these cases represent outcomes within the learning curve for this technology. In the prostate cancer literature, 1 study of laparoscopic radical prostatectomy demonstrated that the 5-year risk of recurrence decreased from 17% to 16% to 9% for patients treated by surgeons with 10, 250, and 750 prior laparoscopic procedures.<sup>5</sup> In this study we report on the first 10 patients undergoing LPN with an articulating instrument assistance completed by 1 single surgeon. Further study to examine perioperative and postoperative outcomes with additional surgical volume will be necessary to further evaluate these initial results.

Larger and multi-institutional studies will be needed to assess the future utilization of articulating laparoscopic instruments such as this technology. We agree that a direct comparison to laparoscopic instruments would be ideal to assess the proposed benefit suggested in our study. However, in hospitals without access to a robotic surgical system there will likely remain a selection bias for patients undergoing laparoscopic PN with traditional laparoscopic instruments, in that the simple cases will undergo LPN and more complex cases will undergo open PN. Therefore, careful consideration will need to be made to match complexity parameters like RENAL Nephrometry scores used in this study to compare traditional LPN to LPN with articulating instruments.<sup>6</sup>

From this initial study, we have seen that warm ischemia times, operative times, and length of hospital stay are similar to robotic PN (RPN), however we do not suggest this LPN technique replace the utilization of RPN. The literature suggests that RPN is superior to LPN, and when available RPN should be performed.<sup>7</sup> However, in the correct hospital setting without a robotic platform, LPN with articulating instruments may both improve LPN utilization and provide a cost-effective alternative to open nephrectomy.

**Hari T. Vigneswaran, Ryan W. Dobbs, Simone Crivellaro**, University of Illinois at Chicago, Department of Urology, Chicago, IL

## References

1. Vigneswaran H, Dobbs R, Huang J, et al. Use of a novel articulating laparoscopic needle driver for partial nephrectomy: an initial experience. *Urology*. 2019. Tbd.

2. Tsai SH, Tseng PT, Sherer BA, et al. Open versus robotic partial nephrectomy: systematic review and meta-analysis of contemporary studies. *Int J Med Robot*. 2019;15:e1963.
3. Wu Z, Li M, Liu B, et al. Robotic versus open partial nephrectomy: a systematic review and meta-analysis. *PLoS One*. 2014;9:e94878.
4. Dobbs RW, Magnan BP, Abhyankar N, et al. Cost effectiveness and robot-assisted urologic surgery: does it make dollars and sense. *Minerva Urol Nefrol*. 2017;69:313–323.
5. Vickers AJ, Savage CJ, Hruza M, et al. The surgical learning curve for laparoscopic radical prostatectomy: a retrospective cohort study. *Lancet Oncol*. 2009;10:475–480.
6. Kutikov A, Uzzo RG. The R.E.N.A.L. nephrometry score: a comprehensive standardized system for quantitating renal tumor size, location and depth. *J Urol*. 2009;182:844–853.
7. Mottrie A, Borghesi M, Ficarra V. Is traditional laparoscopy the real competitor of robot-assisted partial nephrectomy? *Eur Urol*. 2012;62:1034–1036. discussion 1038-1039.

<https://doi.org/10.1016/j.urology.2019.05.052>  
UROLOGY 132: 128–129, 2019. © 2019 Elsevier Inc.