



# Resting vector volume measured before ileostomy reversal may be a predictor of major fecal incontinence in patients with mid or low rectal cancer: a longitudinal cohort study using a prospective clinical database

Min Hyun Kim<sup>1</sup> · Myong Hun Ihn<sup>2</sup> · Jung Rae Cho<sup>1</sup> · Myung Jo Kim<sup>1</sup> · Sung Il Kang<sup>1</sup> · Heung-Kwon Oh<sup>1</sup> · Duck-Woo Kim<sup>1</sup> · Sung-Bum Kang<sup>1</sup>

Accepted: 3 April 2019 / Published online: 17 April 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

## Abstract

**Purpose** Despite a high incidence of fecal incontinence following sphincter-preservation surgery (SPS), there are no definitive factors measured before ileostomy reversal that predict fecal incontinence. We investigated whether vector volume anorectal manometry before ileostomy reversal predicts major fecal incontinence following SPS in patients with mid or low rectal cancer.

**Methods** This longitudinal prospective cohort study comprised 173 patients who underwent vector volume anorectal manometry before ileostomy reversal. The Fecal Incontinence Severity Index was measured 1 year after the primary SPS and classified as major incontinence (FISI score  $\geq 25$ ) or continent/minor incontinence (FISI score  $< 25$ ). Multivariable logistic regression analysis was used to identify predictors of major incontinence.

**Results** Ninety-two patients (53.1%) had major incontinence. Although tumor stage, location, and neoadjuvant chemoradiotherapy were comparable, the major incontinence group had lower resting pressure (28.4 vs. 34.3 mmHg,  $P = 0.027$ ), greater asymmetry at rest (39.1% vs. 34.1%,  $P = 0.002$ ) and squeezing (34.2% vs. 31.4%,  $P = 0.046$ ), shorter sphincter length (3.3 vs. 3.7 cm,  $P = 0.034$ ), and lower resting vector volume (143,601 vs. 278,922 mmHg<sup>2</sup> mm,  $P < 0.001$ ) compared with the continent/minor incontinence group. Resting vector volume was the only independent predictor of major incontinence (odds ratio = 0.675 per 100,000 mmHg<sup>2</sup> mm, 95% confidence interval, 0.532–0.823;  $P = 0.006$ ).

**Conclusions** This study revealed that resting vector volume before ileostomy reversal may predict major fecal incontinence. We suggest that the physiology of the anorectum should be discussed with patients before ileostomy reversal in patients at high risk of fecal incontinence.

**Keywords** Anorectal manometry · Ileostomy reversal · Fecal incontinence

---

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00384-019-03293-3>) contains supplementary material, which is available to authorized users.

---

✉ Sung-Bum Kang  
kangsb@snuh.org

<sup>1</sup> Department of Surgery, Seoul National University Bundang Hospital, 82 Gumi-ro 173 Beon-gil, Bundang-gu, Seongnam-si, Gyeonggi-do 13620, South Korea

<sup>2</sup> Department of Surgery, Soonchunhyang University Gumi Hospital, Gumi, South Korea

## Introduction

Sphincter-preservation surgery is now widely performed since the introduction of total mesorectal excision [1], the development of circular surgical staplers [2], and the application of preoperative chemoradiotherapy [3], which do not compromise the patient's oncologic outcomes. However, fecal incontinence is common after sphincter-preservation surgery for mid or low rectal cancer [4] and is a major issue owing to its significant negative impact on the patient's quality of life [5, 6]. The development of fecal incontinence may be related to radiotherapy, proximity of the tumor to the anal verge, advanced stage, and history of a diverting ostomy [7–9], because

these factors may alter the morphology and physiology of the anorectum and sphincter complex.

Anorectal manometry may be a useful tool to objectively assess the function of the anal canal. However, none of the manometric parameters are definitive predictors of major fecal incontinence before ileostomy reversal. Although some patients with fecal incontinence after rectal surgery show changes in manometry pressure, rectal capacity, or rectal compliance or asymmetry of the anal sphincter [10–15], these findings are not consistent among studies. Vector volume manometry, a measurement utilizing the three-dimensional profilometric pressure measurements across the anal sphincter, may be more advantageous than conventional manometry due to its assessment of the vector component across the anal sphincter [16, 17]. The purpose of this study was to investigate the diagnostic value of vector volume manometry for predicting major fecal incontinence before ileostomy reversal in patients with mid or low rectal cancer.

## Methods

### Patients

This was a longitudinal study of a prospective database comprising of patients who underwent ileostomy reversal at Seoul National University Bundang Hospital between March 2005 and May 2016. The cohort included patients who initially underwent sphincter-preservation surgery for mid (5–10 cm from the anal verge) or low (0–5 cm from the anal verge) rectal cancer and who underwent vector volume anorectal manometry before ileostomy reversal. Ileostomy reversal was scheduled for patients with no evidence of the disease after complete surgical recovery and/or termination of adjuvant chemotherapy. Patients were evaluated for anastomosis integrity and ileostomy reversal was not performed if there were any signs of leak or threatening signs. Patients with locally advanced rectal cancer underwent neoadjuvant chemoradiotherapy in accordance with the National Comprehensive Cancer Network guidelines [18]. Neoadjuvant therapy was normally performed for patients with a threatened mesorectal fascia shown on anorectal magnetic resonance imaging (T4 lesions, lymph node in proximity to mesorectal fascia) or clinically overt lymph node–positive tumors (N2). Sphincter-preservation surgery was done either laparoscopically or using an open method. The decision to undergo sphincter-preservation surgery was made by the clinician with the patient's consent. Patients who underwent abdominoperineal resection for lesions invading the anal sphincter were excluded from the cohort. Anastomosis was made using an endoluminal stapler or by hand-sewn methods. During the stapling procedure, the surgeon took care to avoid wedging any sphincter

muscles or pelvic floor muscles during approximation of the anvil, resulting in near-circular anastomosis.

All patients completed the translated version of the Fecal Incontinence Severity Index (FISI) questionnaire via individual interviews at 1 year after rectal cancer resection [19]. Patients were divided into two groups according to their FISI score: major incontinence (FISI score  $\geq 25$ ) or continent/minor incontinence (FISI score  $< 25$ ) [5, 20–25]. Although there is no consensus on the exact definition of major fecal incontinence based on the FISI score, we used FISI score  $\geq 25$  to define major incontinence partly based on a previous study where patients with FISI score  $\geq 25$  showed a significant decrease in their quality of life [5] and partly based on the median FISI score in our cohort. We compared the manometric parameters recorded before ileostomy reversal between the two groups. General demographic and surgical data were compared between the two groups, including the following: age, sex, body mass index (BMI), tumor location, tumor stage, radiotherapy, chemotherapy, sphincter-preservation surgery approach, operation type, and anastomotic technique. Patients with fecal incontinence were treated with various medications and dietary/exercise education methods; however, they did not undergo any means of invasive treatments such as nerve stimulation. This study was approved by the Institutional Review Board of our institution (Protocol no. B-1808-484-105).

### Anorectal manometry


Anorectal manometry was conducted in an outpatient clinic before the patient was admitted for ileostomy reversal, as in our prior study [14]. All manometry measurements were performed with a water-perfusion technique using an eight-channel Micro Tip catheter (Medtronic, Minneapolis, MN) connected to a perfusion pump. Patients were held in the left-lateral position with their hips flexed at 90°. The probe was positioned 6 cm from the anal verge and pressure was measured using a continuous pull-through method at 0.25 cm/s. The following anorectal parameters were recorded: mean resting pressure, percentage asymmetry at rest, maximal squeezing pressure, percentage asymmetry at squeezing, sustained duration, sphincter length, high-pressure zone (HPZ) length, defecation index, coughing reflex, and vector volume at rest. Anorectal sensation tests of compliance were not performed after restorative proctectomy because of the risk of bowel perforation [26]. Vector volume parameters were defined as previously described [27]. The percentage asymmetry of the sphincter was defined as the percentage of the deviation of the integrated cross-sections from a perfect circle. The HPZ length was defined as the length of the sphincter at which pressures of  $\geq 50\%$  above the basal pressure were recorded during a sustained squeeze. The resting vector volume is the vector volume of the anal canal measured at rest, which

is calculated by adding up the segmental vector volumes across the anal canal (Fig. 1). Each segmental vector volume is the product of the radial area multiplied by each segmental  $z$ -axis length. The radial area is the area of the measurements made by the profilometric profile of a given segment. Typically, for an eight-channel probe, it would be the sum of the multiplied product of each adjacent probe and then multiplied by 0.354 [27]. The resulting vector volume was calculated by the Polygram software (Medtronic GI, Minneapolis, MN) and was given in  $\text{mmHg}^2 \text{ mm}$ .

## Statistical analysis

Quantitative variables are expressed as the mean  $\pm$  standard deviation for normally distributed variables or as the median (interquartile range [IQR]) for non-normally distributed variables. The  $\chi^2$  test and Student's  $t$  test (the Mann–Whitney  $U$  test for non-normally distributed variables) were used to compare categorical and continuous variables, respectively, between the two groups. Anorectal manometry parameters were compared using Student's  $t$  test between the major and minor incontinence groups. Univariable and multivariable logistic regression analyses were also used to assess which manometric parameters were predictors of major incontinence. Parameters with  $P < 0.25$  in univariable analyses were included in the multivariable analysis, in which  $P < 0.05$  was considered statistically significant. A predictive cutoff value of resting vector volume was then calculated using receiver-operating characteristic (ROC) curve analyses by calculating the area under the ROC curve (AUC). The optimal thresholds were determined by maximizing the Youden index. Statistical analyses were performed using SPSS version 20.0 (Chicago, IL) and MedCalc 15.6.1 (Mariakerke, Belgium).

**Fig. 1** Resting vector volume in a representative patient with major incontinence (**a**; FISI score  $\geq 25$ ) and in a continent patient (**b**; FISI score = 0). The red ring represents the segment with the maximal segmental vector volume and the blue ring represents the segment with the minimal segmental vector volume of a given measurement. P proximal portion of the anal canal, D distal end of the anal canal



## Results

### Patient characteristics

A total of 173 patients with available manometry measurements before ileostomy reversal and 1-year postoperative FISI score were included in the study cohort. Their median age was 60.0 years (range 33–82 years) and 112 (64.7%) were male. Tumor location was low in 134 (77.5%) patients and preoperative chemoradiotherapy was performed in 115 (71.6%) patients. Laparoscopic surgery was performed in 117 (67.6%) patients. Stapled anastomosis and hand-sewn anastomosis were done in 111 (64.2%) and 62 (35.8%) cases, respectively. Ileostomy repair was performed at a median of 5.6 months (IQR 4.7–6.4 months) after primary resection.

On the basis of the FISI score, there were 92 patients in the major incontinence group and 81 patients in the continent/minor incontinence group; the median FISI score was 26 (IQR 12.5–32.0) in the overall cohort, 31 (IQR 29–37) in the major incontinence group, and 12 (IQR 0–20) in the continent/minor incontinence group. Baseline clinical and tumor characteristics including sex, BMI, tumor location, and stage were comparable in both groups (Table 1). However, the mean age was greater in the major incontinence group ( $62.1 \pm 10.2$  vs.  $57.2 \pm 11.4$  years). The proportions of patients who underwent preoperative radiotherapy or chemotherapy were similar in both groups. The operation type, surgical approach, and anastomosis type were also similar in both groups. The proportion of each component of the FISI score to each corresponding group is shown in Table 2.

### Anorectal manometry analysis

Anorectal manometry was performed before ileostomy reversal at a median of 4.8 months (IQR 3.9–5.8 months) after primary rectal resection. Patients who underwent sphincter-

preservation surgery had lower resting and squeezing pressures, higher asymmetry at rest and squeezing, and lower mean resting vector volume compared with previously reported values in a historical healthy control group that were assessed using the same protocol and equipment at our institution (mean resting pressure:  $56.4 \pm 14.2$  mmHg; resting asymmetry:  $29.8 \pm 8.2\%$ ; maximum squeezing pressure:  $141.7 \pm 53.8$  mmHg, squeeze asymmetry:  $28.4 \pm 9.9\%$ ; high-pressure zone length:  $2.1 \pm 0.8$  cm; mean resting vector volume:  $460,709 \pm 410,368$  mmHg<sup>2</sup> mm) [14]. In our cohort, we found that resting pressure, resting vector volume, and sphincter length were lower in the major incontinence group than in the continent/minor incontinence group. Asymmetry at rest and asymmetry at squeezing were greater in the major

incontinence group. However, maximal squeezing pressure, sustained duration, high-pressure zone, defecation index, and coughing reflex were not significantly different between the two groups (Table 3). Similar trends were observed in subgroups of patients based on cancer location (mid or low rectal cancer), radiotherapy, and type of resection (ultra-low anterior resection and intersphincteric resection) (Supplementary Tables 1–3).

Multivariable analysis showed that resting vector volume was the only statistically significant variable for major incontinence (Table 4). Anorectal manometry parameters included in the multivariable analysis were mean resting pressure ( $P = 0.027$ ), squeezing pressure ( $P = 0.111$ ), asymmetry at rest ( $P = 0.002$ ), asymmetry at squeezing ( $P = 0.046$ ), sphincter

**Table 1** Baseline characteristics

	Major incontinence ( $n = 92$ )	Continent/minor incontinence ( $n = 81$ )	<i>P</i>
Age, years, median (range)	62 (37–80)	59 (33–82)	0.005
Sex, <i>n</i> (%)			0.437
Male	62 (67.4)	50 (61.7)	
Female	30 (32.6)	31 (38.3)	
BMI <sup>a</sup>	$23.3 \pm 3.0$	$24.1 \pm 3.1$	0.096
Tumor location (cm), from the anal verge, median (IQR <sup>b</sup> )	4.0 (3.0–5.9)	5.0 (3.0–5.0)	0.556
Low rectum (0–5 cm), <i>n</i> (%)	68 (73.9)	66 (81.5)	0.276
Mid rectum (5.1–10 cm), <i>n</i> (%)	24 (26.1)	15 (18.5)	
Pathologic tumor stage, <i>n</i> (%)			0.777
0	10 (10.9)	14 (17.3)	
1	35 (38.0)	30 (37.0)	
2	13 (14.1)	12 (14.8)	
3	30 (32.6)	22 (27.2)	
4	4 (4.3)	3 (3.7)	
Distal resection margin (cm)	$1.8 \pm 1.5$	$1.7 \pm 1.4$	0.818
Radiotherapy, <i>n</i> (%)			0.324
None	30 (32.6)	21 (25.9)	
Preoperative	57 (62.0)	58 (71.6)	
Postoperative	5 (5.4)	2 (2.5)	
Chemotherapy, <i>n</i> (%)			0.347
None	16 (17.4)	16 (19.8)	
Postoperative	19 (20.7)	9 (11.1)	
Surgical approach, <i>n</i> (%)			0.105
Open	35 (38.0)	21 (25.9)	
Laparoscopic	57 (62.0)	60 (74.1)	
Operation type, <i>n</i> (%)			0.088
Low anterior resection	16 (17.4)	6 (7.4)	
Ultra-low anterior resection	49 (53.3)	54 (66.7)	
Intersphincteric resection	27 (29.3)	21 (25.9)	
Anastomosis, <i>n</i> (%)			0.346
Hand-sewn	36 (39.1)	26 (32.1)	
Stapled	56 (60.9)	55 (67.9)	

<sup>a</sup> BMI body mass index, <sup>b</sup> IQR interquartile range

**Table 2** Frequency according to types of incontinence in continent/minor incontinence group and major incontinence group

Type of incontinence		Frequency of incontinence during the past 1 month, <i>n</i> (%)					
		≥ 2/day	1/day	≥ 2/week	1/ week	1–3/ month	Less frequent/ never
Major incontinence ( <i>n</i> = 92)	Gas	63 (68.5)	6 (6.5)	10 (10.9)	1 (1.1)	2 (2.2)	10 (10.9)
	Mucous	9 (9.8)	3 (3.3)	2 (2.2)	3 (3.3)	10 (10.9)	65 (70.7)
	Liquid	15 (16.3)	7 (7.6)	17 (18.5)	7 (7.6)	9 (9.8)	37 (40.2)
	Solid	43 (46.7)	12 (13.0)	16 (17.4)	5 (5.4)	5 (5.4)	11 (12.0)
Continent/minor incontinence ( <i>n</i> = 81)	Gas	14 (17.3)	3 (3.7)	13 (16.0)	0	11 (13.6)	40 (49.4)
	Mucous	0	0	2 (2.5)	0	7 (8.6)	72 (88.9)
	Liquid	1 (1.2)	0	5 (6.2)	5 (6.2)	13 (16.0)	57 (70.4)
	Solid	2 (2.5)	2 (2.5)	9 (11.1)	2 (2.5)	13 (16.0)	53 (65.4)

length ( $P = 0.034$ ), high-pressure zone length ( $P = 0.160$ ), and resting vector volume ( $P < 0.001$ ). Because the vector volume is normally represented in the hundred-thousand-unit range, it was not possible to show an increased risk per unit increment in this parameter. Thus, a subsequent multivariable analysis was done by reducing the variable by a factor of 100,000 which yielded an odds ratio of 0.675 per 100,000 mmHg<sup>2</sup> mm (95% confidence interval, 0.532–0.823;  $P = 0.006$ ). The AUC for resting vector volume was 0.663. A cutoff value of 242,710 mmHg<sup>2</sup> mm (decrease in vector volume increases severity) was obtained with a sensitivity of 84.8% and specificity of 50.6% by maximizing the Youden index (Fig. 2).

## Discussion

There is no definitive test before ileostomy reversal for predicting fecal incontinence, even though there is a high incidence of fecal incontinence following sphincter-

preservation surgery that usually occurs after restoration of bowel continuity. This study suggests that a decreased resting vector volume shown on anorectal manometry performed before ileostomy reversal is associated with an increased risk of major fecal incontinence after sphincter-preservation surgery. Using this cohort of patients, we determined a cutoff value for resting vector volume before ileostomy reversal as a predictor of major incontinence in patients with mid or low rectal cancer. This cutoff value should be validated in a well-designed large-scale study.

Fecal incontinence is a debilitating condition after sphincter-preservation surgery, with a significant negative impact on the patient's quality of life [6]. Although fecal incontinence usually improves during the first year after surgery [28], it may persist for longer, especially in patients with a low rectal tumor or those who undergo chemoradiotherapy [8]. The ability to predict which patients are at increased risk of fecal incontinence, before

**Table 3** Anorectal manometry measurements between groups

	Major incontinence ( <i>n</i> = 92)	Continent/minor incontinence ( <i>n</i> = 81)	<i>P</i>
Mean resting pressure (mmHg)	28.4 ± 15.8	34.3 ± 18.6	0.027
Resting asymmetry (%)	39.1 ± 11.7	34.1 ± 9.6	0.002
Max squeezing pressure (mmHg)	117.3 ± 65.9	133.3 ± 65.5	0.111
Squeeze asymmetry (%)	34.2 ± 9.3	31.4 ± 8.3	0.046
Sustained duration (s)	35.2 ± 19.8	33.6 ± 23.5	0.632
Sphincter length (cm)	3.3 ± 1.1	3.7 ± 1.0	0.034
High-pressure zone (cm)	2.2 ± 1.1	2.4 ± 0.9	0.160
Defecation index	1.7 ± 1.1	1.6 ± 1.1	0.507
Coughing reflex (%)	134.6 ± 105.7	125.9 ± 74.6	0.543
Resting vector volume (mmHg <sup>2</sup> mm)	143,601 ± 118,961	278,922 ± 289,674	<0.001

Values are expressed as the mean ± standard deviation

**Table 4** Multivariable analysis of manometric parameters for major incontinence

	OR	95% CI	P
Mean resting pressure, per mmHg	1.009	0.985–1.035	0.456
Resting asymmetry, per %	1.022	0.979–1.066	0.324
Maximal squeezing pressure, per mmHg	0.999	0.993–1.004	0.589
Squeeze asymmetry, per %	1.022	0.979–1.066	0.324
Sphincter length, per cm	0.858	0.502–1.467	0.575
High-pressure zone, per cm	1.223	0.716–2.088	0.462
Resting vector volume, per mmHg <sup>2</sup> mm	1.000	1.000–1.000	0.008

OR hazard ratio, CI confidence interval


performing ileostomy reversal, would be helpful to patients because it may be possible to prevent fecal incontinence. Objective assessment of anal canal function before undergoing ileostomy reversal is a logical step toward providing the patient with better treatment options.

Many approaches have been used to investigate fecal incontinence, including diagnostic tests of sphincter morphology, nerve integrity, and function such as magnetic resonance imaging, endoscopic ultrasonography, pudendal nerve terminal latency test, and anorectal manometry. However, imaging alone does not provide adequate information about anal function, and results of the pudendal nerve terminal latency test were controversial [29]. We believe that anal manometry is the most direct method for assessing anal function. Although several studies have performed anorectal manometry after sphincter-preservation surgery in patients with fecal incontinence, few of these studies determined a cutoff value that could be used to predict fecal incontinence after ileostomy reversal, although changes in anal sphincter pressure, symmetry, and rectal compliance have been documented [10–15]. Thus, we need better understanding of the physiologic changes caused by structural defects of the anal sphincter following sphincter-preservation surgery. The present study showed that patients with major incontinence had lower resting pressure, greater asymmetry of the anal canal, and shorter sphincter length compared with the continent/minor incontinence group, as reported in previous studies [10–15]. Maximal squeezing pressure was not low in major incontinence, consistent with prior studies [10, 13, 14]. Although reduced compliance of the neorectum and decreased sensitivity of the anus to propulsion forces were reported in prior studies [10, 13–15], we did not measure rectal compliance for safety concerns, such as the risk of rectal perforation during measurement [26].

Our study showed that the resting vector volume was inversely correlated with the severity of fecal incontinence. We believe the clinical relevance of resting vector volume was underestimated in the past compared to usual parameters such as pressure, asymmetry, or sphincter length and should be revisited. The anal canal is a three-dimensional structure and

all of its components, including the muscles and nerves, act together to maintain continence. Therefore, a more comprehensive integrated parameter, such as resting vector volume, is better qualified to assess the overall quality of anal function and continence. The resting vector volume is calculated from three-dimensional profilometric pressure measurements across the anal sphincter at the resting state. The radial area of the pressure profile is calculated across each cross-section and integrated across the z-axis of the anal canal, resulting in a lower value if there is a loss of function of the internal anal sphincter in any segment or direction [27]. Previous studies showed that the resting vector volume after rectal surgery was generally lower in continent patients than in healthy individuals [14, 15, 30]. We think that that decreases in sphincter length, resting pressure, or symmetry would result in a decreased resting vector volume, which will provide a better representation of the three-dimensional structural function of the anal canal.

In our study, we estimated a cutoff vector volume value of 242,710 mmHg<sup>2</sup> mm for predicting major incontinence by maximizing the Youden index. However, the AUC of this curve was 0.663 and is a limitation to generalizing our findings. The lack of standardization of protocol and interpatient variability has prevented widespread use of manometry parameters into daily clinical decision-making [27]. We consider that the optimal cutoff value should be evaluated with consideration of interpatient variability as well as a standardized protocol to help clinical decision. Additional well-designed studies are needed to confirm the clinical value and cutoff value of resting vector volume as a predictor for major incontinence after sphincter-preservation surgery.



**Fig. 2** Receiver-operating characteristic curve of resting vector volume for predicting major incontinence. Area under the curve = 0.663

The proportion of patients suffering from fecal incontinence after sphincter-preservation surgery vary ranging from 0 to 71%, with an increased risk for low tumors and neoadjuvant radiotherapy [6, 8]. The high prevalence of fecal incontinence in our cohort is partly due to the fact that over 95% of rectal cancer patients underwent sphincter-preservation surgery in our facility, which is one of the strengths of our study. There is a trend toward more sphincter-preservation surgeries being performed than in the past [31, 32], which subsequently will increase patients suffering from fecal incontinence. Additionally, in our cohort, 72% of patients underwent neoadjuvant chemoradiotherapy and 77.5% of patients were low rectal tumors (within 5 cm from anal verge), which also contributes to a higher risk of fecal incontinence.

Our study holds some limitations and unsolved issues to discuss. First, the decision to undergo anorectal manometry testing and FISI questionnaire survey was not prospectively controlled, introducing the possibility of selection bias. The FISI questionnaire was not administered before primary surgery or more than 1 year after surgery. Thus, the effects of sphincter-preservation surgery on further deterioration of sphincter function and/or recovery are unknown. Second, patients were not assessed for pelvic nerve function due to safety issues; thus, whether incontinence was induced by nerve damage or structural defects is unknown. Third, we did not consider the possibility of age and sex differences in anal canal physiology because of the small number of patients. Future studies that account for these subgroup differences may yield differing results. Lastly, the generalizability of the resting vector volume would be limited without a larger scale study. On subgroup analysis, although a trend was observed, some subgroups failed to show a statistically significant difference in the resting vector volume.

In conclusion, this study revealed that resting vector volume measured before ileostomy reversal may be a predictor of major incontinence in patients undergoing sphincter-preserving surgery for mid or low rectal cancer. We suggest that preoperative counseling should be offered to patients at increased risk of fecal incontinence, especially in patients with a low resting vector volume. We also suggest that the physiology of the anorectum should be discussed with patients before ileostomy reversal.

### Compliance with ethical standards

**Ethical approval** All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**Conflict of interest** The authors declare that they have no conflict of interest.

### References

1. Heald RJ, Smedh RK, Kald A, Sexton R, Moran BJ (1997) Abdominoperineal excision of the rectum—an endangered operation. *Norman Nigro Lectureship. Dis Colon Rectum* 40:747–751
2. Knight CD, Griffen FD (1980) An improved technique for low anterior resection of the rectum using the EEA stapler. *Surgery* 88:710–714
3. Sauer R, Becker H, Hohenberger W, Rodel C, Wittekind C, Fietkau R, Martus P, Tschmelitsch J, Hager E, Hess CF, Karstens JH, Liersch T, Schmidberger H, Raab R, German Rectal Cancer Study Group (2004) Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 351:1731–1740
4. Bryant CL, Lunniss PJ, Knowles CH, Thaha MA, Chan CL (2012) Anterior resection syndrome. *Lancet Oncol* 13:e403–e408
5. Bordeianou L, Rockwood T, Baxter N, Lowry A, Mellgren A, Parker S (2008) Does incontinence severity correlate with quality of life? Prospective analysis of 502 consecutive patients. *Color Dis* 10:273–279
6. Emmertsen KJ, Laurberg S, Rectal Cancer Function Study Group (2013) Impact of bowel dysfunction on quality of life after sphincter-preserving resection for rectal cancer. *Br J Surg* 100:1377–1387
7. Battersby NJ, Juul T, Christensen P, Janjua AZ, Branagan G, Emmertsen KJ, Norton C, Hughes R, Laurberg S, Moran BJ, United Kingdom Low Anterior resection Syndrome Study Group (2016) Predicting the risk of bowel-related quality-of-life impairment after restorative resection for rectal cancer: a multicenter cross-sectional study. *Dis Colon Rectum* 59:270–280
8. Alavi M, Wendel CS, Krouse RS, Temple L, Hornbrook MC, Bulkley JE, McMullen CK, Grant M, Herrinton LJ (2017) Predictors of bowel function in long-term rectal cancer survivors with anastomosis. *Ann Surg Oncol* 24:3596–3603
9. Dulskas A, Miliauskas P, Tikuisis R, Escalante R, Samalavicius NE (2016) The functional results of radical rectal cancer surgery: review of the literature. *Acta Chir Belg* 116:1–10
10. Ihnat P, Vavra P, Prokop J, Pelikan A, Ihnat Rudinska L, Penka I (2018) Functional outcome of low rectal resection evaluated by anorectal manometry. *ANZ J Surg* 88:e512–e516
11. Bordeianou L, Lee KY, Rockwood T, Baxter NN, Lowry A, Mellgren A, Parker S (2008) Anal resting pressures at manometry correlate with the Fecal Incontinence Severity Index and with presence of sphincter defects on ultrasound. *Dis Colon Rectum* 51:1010–1014
12. Kroesen AJ, Runkel N, Buhr HJ (1999) Manometric analysis of anal sphincter damage after ileal pouch-anal anastomosis. *Int J Color Dis* 14:114–118
13. Matzel KE, Bittorf B, Gunther K, Stadelmaier U, Hohenberger W (2003) Rectal resection with low anastomosis: functional outcome. *Color Dis* 5:458–464
14. Kang SB, Kim N, Lee KH, Kim YH, Kim JH, Kim JS (2008) Anal sphincter asymmetry in anal incontinence after restorative proctectomy for rectal cancer. *World J Surg* 32:2083–2088
15. Rink AD, Nagelschmidt M, Radinski I, Vestweber KH (2008) Evaluation of vector manometry for characterization of functional outcome after restorative proctocolectomy. *Int J Color Dis* 23:807–815
16. Fynes MM, Behan M, O’Herlihy C, O’Connell PR (2000) Anal vector volume analysis complements endoanal ultrasonographic assessment of postpartum anal sphincter injury. *Br J Surg* 87:1209–1214
17. Zbar AP, Kmiot WA, Aslam M, Williams A, Hider A, Audisio RA, Chiappa A, deSouza N (1999) Use of vector volume manometry and endoanal magnetic resonance imaging in the adult female for

- assessment of anal sphincter dysfunction. *Dis Colon Rectum* 42: 1411–1418
18. Benson AB, Venook AP, Al-Hawary MM, Cederquist L, Chen Y-J, Ciombor KK et al (2018) Rectal cancer, version 2.2018, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Cancer Netw* 16:874–901
  19. Rockwood TH, Church JM, Fleshman JW, Kane RL, Mavrantonis C, Wexner SD, Bliss D, Lowry AC (1999) Patient and surgeon ranking of the severity of symptoms associated with fecal incontinence: the Fecal Incontinence Severity Index. *Dis Colon Rectum* 42:1525–1532
  20. Boreham MK, Richter HE, Kenton KS, Nager CW, Gregory WT, Aronson MP, Vogt VY, McIntire DD, Schaffer JI (2005) Anal incontinence in women presenting for gynecologic care: prevalence, risk factors, and impact upon quality of life. *Am J Obstet Gynecol* 192:1637–1642
  21. Norton C, Burch J, Kamm MA (2005) Patients' views of a colostomy for fecal incontinence. *Dis Colon Rectum* 48:1062–1069
  22. Gearhart S, Hull T, Floruta C, Schroeder T, Hammel J (2005) Anal manometric parameters: predictors of outcome following anal sphincter repair? *J Gastrointest Surg* 9:115–120
  23. Colquhoun P, Kaiser R Jr, Efron J, Weiss EG, Noguera JJ, Vernava AM, Wexner SD (2006) Is the quality of life better in patients with colostomy than patients with fecal incontinence? *World J Surg* 30: 1925–1928
  24. Abbas SM, Bissett IP, Neill ME, Parry BR (2005) Long-term outcome of postanal repair in the treatment of faecal incontinence. *ANZ J Surg* 75:783–786
  25. Levack MM, Savitt LR, Berger DL, Shellito PC, Hodin RA, Rattner DW, Goldberg SM, Bordeianou L (2012) Sigmoidectomy syndrome? Patients' perspectives on the functional outcomes following surgery for diverticulitis. *Dis Colon Rectum* 55:10–17
  26. Park JS, Kang SB, Kim DW, Kim NY, Lee KH, Kim YH (2007) Iatrogenic colorectal perforation induced by anorectal manometry: report of two cases after restorative proctectomy for distal rectal cancer. *World J Gastroenterol* 13:6112–6114
  27. Schizas AM, Emmanuel AV, Williams AB (2011) Anal canal vector volume manometry. *Dis Colon Rectum* 54:759–768
  28. Liu LG, Yan XB, Shan ZZ, Yan LL, Jiang CY, Zhou J, Tian Y, Jin ZM (2017) Anorectal functional outcome following laparoscopic low anterior resection for rectal cancer. *Mol Clin Oncol* 6:613–621
  29. Saraidaridis JT, Molina G, Savit LR, Milch H, Mei T, Chin S, Kuo J, Bordeianou L (2018) Pudendal nerve terminal motor latency testing does not provide useful information in guiding therapy for fecal incontinence. *Int J Color Dis* 33:305–310
  30. Schizas AM, Emmanuel AV, Williams AB (2011) Vector volume manometry—methods and normal values. *Neurogastroenterol Motil* 23:886–e393
  31. Ricciardi R, Virnig BA, Madoff RD, Rothenberger DA, Baxter NN (2007) The status of radical proctectomy and sphincter-sparing surgery in the United States. *Dis Colon Rectum* 50:1119–1127
  32. Shahjehan F, Kasi PM, Habermann E, Day CN, Colibaseanu DT, Mathis KL, Larson DW, Merchea A (2019) Trends and outcomes of sphincter-preserving surgery for rectal cancer: a national cancer database study. *Int J Color Dis* 34:239–245

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.