



Correction to: Paclitaxel-coated balloon treatment for functionally nonsignificant residual coronary lesions after balloon angioplasty

Ae-Young Her¹ · Eun-Seok Shin² · Joo Myung Lee³ · Scot Garg⁴ · Joon-Hyung Doh⁵ · Chang-Wook Nam⁶ · Bon-Kwon Koo⁷

Published online: 14 June 2019
© Springer Nature B.V. 2019

Correction to:
The International Journal of Cardiovascular Imaging
(2018) 34:1339–1347
<https://doi.org/10.1007/s10554-018-1351-z>

In the original publication of the article Fig. 1 was published incorrectly as it showed incorrect patient numbers. The corrected Fig. 1 is given in this correction.

The original article can be found online at <https://doi.org/10.1007/s10554-018-1351-z>.

✉ Eun-Seok Shin
sesim1989@gmail.com

- ¹ Division of Cardiology, Department of Internal Medicine, Kangwon National University School of Medicine, Chuncheon, South Korea
- ² Division of Cardiology, Ulsan University Hospital, University of Ulsan College of Medicine, 877, Bangeojinsunhwando-ro, Dong-gu, Ulsan 44033, South Korea
- ³ Department of Internal Medicine and Cardiovascular Center, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea
- ⁴ East Lancashire Hospitals NHS Trust, Blackburn, Lancashire, UK
- ⁵ Division of Cardiology, Inje University Ilsan Paik Hospital, Goyang, South Korea
- ⁶ Department of Internal Medicine, Dongsan Medical Center, Keimyung University College of Medicine, Taegu, South Korea
- ⁷ Department of Internal Medicine and Cardiovascular Center, Seoul National University Hospital, Seoul, South Korea




Fig. 1 Flowchart of the study