



Dietary patterns were not associated with age-related macular degeneration: a cross-sectional analysis in the Irish Nun Eye Study

Rachel V. McCarter¹ · Charlotte E. Neville¹ · Giuliana Silvestri² · Shannon Montgomery¹ · Evelyn Moore² · Vittorio Silvestri² · Christopher R. Cardwell¹ · Ruth E. Hogg¹ · Jayne V. Woodside^{1,3} · Gareth J. McKay¹

Received: 21 May 2018 / Accepted: 13 November 2018 / Published online: 22 November 2018
© The Author(s) 2018

Abstract

Background Analysing dietary patterns (DP) evaluates overall dietary intake, taking account of its complexity, quality, variance and the interaction between different foods, providing an alternative approach for the evaluation of nutritional influences on age-related macular degeneration (AMD) risk.

Aims To evaluate the relationship between DP and AMD in an older female population.

Methods Data was analysed from the cross-sectional Irish Nun Eye Study involving 1233 older women with a restricted lifestyle (mean age 76.3 years [range, 56–100 years]). The Wisconsin Age-related Maculopathy Grading System was used to classify digital colour macular fundus images and dietary intake was assessed using a food frequency questionnaire ($n = 1033$). A posteriori DP were derived using principal component analysis. Logistic regression models examined associations between DP and AMD risk with adjustment for confounders.

Results Two DP were identified: a ‘healthy’ pattern characterised by a high intake of oily fish, wholegrains, vegetables and fruit; and an ‘unhealthy’ pattern characterised by high-fat dairy products, sugar, sweets and chips. Of the participants included within the analysis, AMD status were categorised as controls ($n = 818$, 86.9%), early AMD ($n = 83$, 8.8%) and late AMD ($n = 21$, 2.2%). Regression analysis failed to identify any significant associations between healthy or unhealthy DP and AMD risk, in unadjusted and adjusted models.

Conclusion No evidence of an association between the DP identified and AMD risk was detected in this well-characterised population. Further research is required to determine the overall dietary influences on AMD risk in general population cohorts.

Keywords Age-related macular degeneration · Diet · Dietary patterns · Nutrition

Introduction

Age-related macular degeneration (AMD) is the leading cause of vision impairment within the developed world in people aged > 50 years and treatment options for slowing the

progression of visual loss due to late AMD are limited [1]. Nutritional influences and dietary interventions have been shown to modulate AMD associated risk [2–8]. Previous dietary based analyses have largely focused on the evaluation of individual nutrients, foods or groups of foods to identify independent nutritional influences affecting AMD risk, although the associations observed, have not always been consistent [9–13]. The analysis of dietary patterns (DP) evaluates overall dietary intake, taking account of its complexity, quality, variance and the interaction between different foods, providing an alternative approach for the evaluation of nutritional influences on AMD risk [14, 15].

Some studies have outlined the potential nutritional or dietary influences on AMD risk [7, 16–19]. Data emerging from the Age-Related Eye Disease Study (AREDS), advocates a healthy diet with supplementation of zinc, lutein/zeaxanthin and vitamins C and E, as an effective strategy for reducing

Rachel V. McCarter and Charlotte E. Neville contributed equally to this work.

✉ Gareth J. McKay
g.j.mckay@qub.ac.uk

¹ Centre for Public Health, Queen’s University Belfast, Belfast, UK

² Department of Ophthalmology, Belfast Health & Social Care Trust, Belfast, UK

³ UKCRC Centre of Excellence for Public Health, School of Medicine, Dentistry and Biomedical Sciences, Queen’s University Belfast, Belfast, UK

AMD progression from its early to visually-disabling late stage [16]. Another study by Chiu et al. also noted the potential benefits of omega-3 fatty acids for reducing AMD progression [20]. In relation to dietary carbohydrates, other studies have highlighted that the quality of carbohydrates consumed as measured by the glycaemic index, modify AMD risk [17, 20–23]. However, quantification of the glycaemic index of a specific food can be problematic, as it varies with respect to the other foods consumed at the same time [24]. Overall, this cumulative evidence showing associations with different foods and nutrients could suggest that the overall diet may be important given that the consumption of one food may affect the nutritional value and bioavailability of others, and the nutritional benefits may also be influenced by the individual's health status or severity of AMD.

To our knowledge, only two studies have previously reported on the associations between a posteriori-derived DP and AMD. In the US-based AREDS study, Chiu and colleagues identified two major DP, labelled as 'Oriental' and 'Western', which were strongly associated with reduced and increased risk of AMD (both early and late) respectively, accounting for <20% of the total dietary variance assessed [14, 15]. An Australian study (the Melbourne Collaborative Cohort study [MCCS]) identified a significant association between 'processed foods' and a higher prevalence of late AMD [25]. To date, we are unaware of any study that has examined associations between DP and AMD risk in European populations. As such, the aim of this study was to evaluate a posteriori-derived DP of nutritional intake and associated risk of AMD in a well-characterised cohort of older Irish women with a restricted lifestyle.

Methods

Study population

The study population was drawn from the Irish Nun Eye Study (INES), a cross-sectional study, in which DP were analysed in association with AMD prevalence. This unique population subscribes to a set of rules that governs behaviour in terms of material possessions, emotional and physical attachment, maintenance of a daily structured religious life of abstinence and prayer with dietary and lifestyle limitations and therefore represents a relatively homogenous population. A total of 1233 females were recruited between 2007 and 2009 from 123 convents across the island of Ireland; all participants were Irish, of white ethnicity, had been residing in a convent for 25 years or more and were older than 55 years. Prior to the commencement of research, ethical approval was obtained from the Office for Research Ethics Committee Northern Ireland and informed written consent was

obtained from all participants. DP analysis was performed on 1033 participants who completed a food frequency questionnaire (FFQ) as part of the study and for whom retinal images were available. The study design and sampling procedures have been described previously [26, 27].

Age-related macular degeneration characterisation

Colour macular centred fundus images were graded independently by the Network of Ophthalmic Reading Centres UK (NetwORC UK). A standardised procedure adopted from definitions of the Wisconsin Age-related Maculopathy Grading System was used to classify features of early and late AMD [28] and the definitions were based on the International Classification for Age-related Macular Degeneration [29]. The presence of features within a 6000- μm circle centred on the fovea were recorded. Drusen were categorised according to size, characteristics of homogeneity of surface features and outline. Pigmentary changes were classified into two categories: hyperpigmentation and hypopigmentation. These features were used to assign each eye to a severity grade as follows: Controls: no features of AMD or the presence of soft distinct drusen ($> 63 \mu\text{m}$ and $\leq 125 \mu\text{m}$) or pigmentary abnormalities only. Cases were defined as early AMD: soft, indistinct ($\geq 125 \mu\text{m}$) or reticular drusen only or soft distinct drusen with pigmentary abnormalities; late AMD: either geographic atrophy (i.e. well-demarcated area of retinal pigment atrophy with visible choroidal vessels) or neovascular AMD (i.e. presence of any of the following: serous or hemorrhagic retinal or retinal pigment epithelial detachment, subretinal neovascular membrane, or periretinal fibrous scar).

Assessment of dietary intake

The semi-quantitative Scottish Collaborative Group FFQ was used to assess the dietary intake of each participant (<http://www.foodfrequency.org/>). This validated tool consisted of 19 sections comprised of 170 food items. Participants had to indicate the frequency of consumption of each food item during the previous 2–3 month period. Frequency was measured in terms of the number of standard portions per day/week/month/rarely. Reported food intakes were converted into a daily weight using food portion sizes [30].

Dietary pattern identification

The individual food items within the FFQ were manually allocated to one of 38 food groups according to food type and macronutrient content (Table 1). Principal component analysis (PCA) with orthogonal rotation (varimax) was applied to this data to generate DP data and factor loadings for each individual food group. Examination of the scree plot was used to

Table 1 Food items included in food groups for dietary pattern analysis

Food groups	Food items included (from the FFQ)
Red meat	Minced beef, beef-burgers (fried), roast beef, roast pork
Organ meat	Liver
Poultry	Roast chicken
Processed meat	Grilled pork sausages, bacon gammon joint, black pudding, ham, salami, chicken pie
White fish and shellfish	Fish fingers, battered cod, poached white fish (cod), smoked white fish (cod), fish cakes, prawns, mussels
Oily fish	Fried mackerel, grilled salmon, smoked salmon, tinned tuna, sardines
Refined grains (excludes refined cereals)	White sliced bread, soft white roll, garlic bread, pitta bread, white rice, pasta (spaghetti), noodles, scones, crackers, savoury pancakes
Wholegrains (excludes wholegrain cereals)	Brown rice, oatcakes
Potatoes without waffles and roasts	Boiled potatoes, mashed potatoes, potato salad
French fries with waffles and roasts	Home-cooked and retail French fries, potato waffles, roast potatoes
Chips	Chips, reduced fat chips, tortilla chips
Pizza	Pizza
Low fat dairy excluding Horlicks	Semi-skimmed milk, skimmed milk, low fat yogurt, low calorie yogurt, low fat cheese, cottage cheese
High fat dairy	Full fat milk, full fat yogurt, fromage frais, cream, cheddar cheese, cheese spread, Philadelphia cheese, dried milk
Eggs	Boiled hen's egg, fried egg, scrambled egg, quiche
Dressings/sauces/condiments	White sauce, ketchup, mayonnaise, oil and vinegar dressing, tomato chutney, gravy, marmite
Desserts (excluding biscuits)	Milk-based pudding, sponge puddings, cheesecake, fruit pie, mousse, custard, Cornetto, other ice cream, sponge cake, sponge cake with jam/cream/icing, fruit cake, doughnuts
Chocolate excluding hot chocolate	Dairy milk bar
Biscuits	Cereal bars, digestive biscuits, custard creams, shortbread, chocolate coated biscuits
Sugar and sweets	Toffees, boiled sweets, fruit gums, jam
Nuts	Peanuts, unsalted nuts, peanut butter
Soup	Home-made soup (vegetable), tinned soup (tomato), instant soup
Tea	Tea, herbal tea
Coffee	Instant coffee, decaffeinated coffee, cappuccino
Alcohol	Low alcohol beer, dark beer (stout), light beer (lager), white wine, red wine, sherry/port, spirits/liqueurs, alcopops, cider
Fruit juice	Pure orange juice, tomato juice
Fruit	Fresh fruit salad, tinned fruit, apples, bananas, oranges, pears, peaches, kiwi, mixed dried fruit, grapes
Vegetables	Mixed vegetable stir-fry, tinned vegetables, coleslaw, sweet peppers
Lutein-zeaxanthin-rich vegetables	Carrots, sweetcorn, lettuce
Cruciferous green leafy vegetables	Cabbage, brussel sprouts, broccoli, spinach, cauliflower
Legumes	Peas, baked beans, soya beans, kidney beans, lentils
Allium family	Leeks, onions
Tomato group	Tomatoes, tomato pasta sauce
Meat replacement products	Soya milk, vegeburgers, quom, tofu
Soft drinks	Blackcurrant squash, fruit squash, diet cola, regular fizzy drinks
Hot chocolate and Horlicks	Hot chocolate, Horlicks
Refined breakfast cereals	Cornflakes, Crunchy nut cornflakes
Wholegrain breakfast cereals	Porridge, all bran, muesli, Weetabix

determine the number of factors to be retained. To aid data interpretation, food groups which had a factor loading > 0.2 were retained for describing the DP. The PCA produced factor scores for each participant which were categorised into quartiles for each DP with the fourth quartile indicating greatest conformity to the specific DP.

Demographic, lifestyle and anthropometric data

A structured questionnaire was used to assess alcohol and smoking status, disease status (presence or absence) and medication usage. Diastolic blood pressure (DBP) and systolic blood pressure (SBP) were measured in a resting seated

position using an oscillometric blood pressure aneroid sphygmomanometer (Speider and Keller). Participant's weight and height were measured and body mass index (BMI) computed as weight (kilograms)/height (metres) squared.

Statistical analysis

IBM SPSS version 21 (SPSS Inc., Chicago, IL, USA) was used for statistical analyses. For all analyses, statistical significance was considered $P < 0.05$. Descriptive statistics were obtained for all variables of interest. Categorical and continuous variables were summarised as n (%) and mean (SD), respectively. One-way analysis of variance (ANOVA) was used to assess the differences between the quartiles for continuous variables. Chi square tests were used to investigate the differences between quartiles for categorical variables. Post-hoc pairwise comparisons were performed using Mann-Whitney U tests between quartiles to determine the location of any differences (data not shown).

Logistic regression was used to explore the relationship between the DP identified (with each DP distributed into equal quartiles) and AMD status (controls versus AMD (i.e. early or late AMD)) in both unadjusted (model 1) and adjusted analyses (models 2 and 3). The reference category was the lowest quartile for each DP, indicating lowest conformity to the pattern. Within this analysis, the confounding factors were those identified from similar previously published studies and those commonly recognised to influence early or late AMD [14, 15, 25]. Model 1 was unadjusted; Model 2 was adjusted for age and BMI; Model 3 was adjusted for model 2 plus smoking, alcohol, hypertension, diabetes mellitus (DM), ischaemic heart disease (IHD) and cerebrovascular accident (CVA).

Results

The mean age of participants was 76.3 years (standard deviation [SD] 8.0; Table 2). Based on BMI classifications, 45% ($n = 412$) of participants were of normal weight, 12% ($n = 111$) were underweight, 28% ($n = 258$) were overweight and 15% ($n = 138$) were classified as clinically obese. Of the participants included within the analysis, AMD status was categorised as controls ($n = 818$, 86.9%), early AMD ($n = 83$, 8.8%) and late AMD ($n = 21$, 2.2%). Images from 19 participants (2%) were of insufficient quality for AMD phenotyping. The majority of participants were non-consumers of alcohol ($n = 864$, 92%), and had never smoked, ($n = 905$, 96%). Approximately 3% of participants ($n = 30$) had self-reported CVA, 11% had IHD ($n = 107$), 3% had DM ($n = 31$) and hypertension was reported in 41% of participants ($n = 390$).

PCA identified two major DP labelled 'healthy' and 'unhealthy'. The factor loadings for both DP are presented in

Table 2 Health and lifestyle characteristics of the Irish Nun Eye Study participants ($n_{\max} = 941$)

Characteristic	Mean (SD) ^a	Range
Age (year)	76.3 (8.0)	56–100
BMI (kg/m ²)	24.6 (5.1)	13.3–60.1
Height (m)	1.61 (0.1)	1.25–1.82
Weight (kg)	63.8 (13.0)	33–159
MABP (mmHg)	92.5 (10.3)	57–147
AMD status:		
Controls, n (%)	818 (86.9)	
Early AMD, n (%)	83 (8.8)	
Late AMD, n (%)	21 (2.2)	
Smoking status*:		
Never, n (%)	905 (96.3)	
Ever, n (%)	35 (3.7)	
Alcohol intake*:		
None, n (%)	864 (91.8)	
1–7 measures/week, n (%)	74 (7.9)	
> 7 measures/week, n (%)	3 (0.3)	
CVA, n (%)	30 (3.2)	
Diabetes mellitus, n (%)	31 (3.3)	
IHD, n (%)	107 (11.4)	
HTN, n (%)	390 (41.5)	

AMD, age-related macular degeneration; BMI, body mass index; MABP, mean arterial blood pressure; CVA, cerebrovascular accident; HTN, hypertension; IHD, ischaemic heart disease

Values are presented as mean (SD) for continuous variables and n (%) for categorical variables

Table 3. The 'healthy' DP was characterised (in decreasing order of factor loadings) by lutein/zeaxanthin-rich vegetables, green leafy vegetables, alliums, vegetables, fruit, tomatoes, legumes, nuts, oily fish, low-fat dairy products, pizza, dressings/sauces/condiments, wholegrain breakfast cereal and red meat. The 'unhealthy' DP was characterised (in decreasing order of factor loadings) by chips, French fries, alcohol, high fat dairy products, soups, desserts, sugars and sweets, wholegrains, dressings/sauces/condiments, processed meat, potatoes, eggs, refined grains, refined breakfast cereal, chocolate, vegetables, red meat, white fish and shellfish. Together, the DP characterised accounted for approximately 16% of the total dietary variance within the population.

No significant differences in participant characteristics were observed across quartiles within the 'healthy' DP, with the exception of age ($p < 0.001$) and BMI ($p = 0.05$); females in the lowest quartile tended to be older and had a higher BMI compared to those in the other quartiles. Participants in the lowest quartile of the 'unhealthy' DP also tended to be older ($p = 0.01$) while those in the highest quartile were more likely to consume alcohol ($p < 0.001$) and smoke ($p = 0.05$) (data not shown).

Table 3 Factor loading matrix for the two major dietary patterns identified in the Irish Nun Eye Study ($n = 1033$)

Food groups	Healthy (Factor 1)	Unhealthy (Factor 2)
Red meat	0.263	0.234
Processed meat	–	0.369
White fish and shellfish	–	0.224
Oily fish	0.375	–
Refined grains	–	0.302
Wholegrains	–	0.385
Potatoes	–	0.359
French fries	–	0.511
Chips	–	0.554
Pizza	0.297	–
Low fat dairy	0.305	–
High fat dairy	–	0.466
Eggs	–	0.356
Dressings	0.29	0.382
Desserts	–	0.436
Chocolate	–	0.252
Sugars and sweets	–	0.415
Nuts	0.386	–
Soups	–	0.452
Alcohol	–	0.482
Fruit	0.517	–
Vegetables	0.558	0.243
L/Z rich vegetables	0.714	–
Green leafy vegetables	0.696	–
Legumes	0.429	–
Alliums	0.59	–
Tomato	0.454	–
Refined breakfast cereal	–	0.256
Wholegrain breakfast cereal	0.274	–

L/Z, lutein-/zeaxanthin

There was no significant difference in AMD risk (i.e. controls versus AMD) across quartiles of either the ‘healthy’ or ‘unhealthy’ DP, in both the unadjusted and adjusted analyses (Table 4). Participants who adhered closely to an ‘unhealthy’ DP had a greater risk of AMD compared to those in the other quartiles, although this failed to reach significance. In the unadjusted analysis (model 1) the odds ratio for AMD in the highest fourth of the ‘healthy’ DP (i.e. most healthy) compared to the lowest fourth (i.e. least healthy) was 0.86 (95% CI 0.48–1.55; $P = 0.62$), while the odds ratio for AMD in the highest fourth of the ‘unhealthy’ DP (i.e. most unhealthy) compared to the lowest fourth (i.e. least unhealthy) was 1.50 (95% CI 0.82–2.75; $P = 0.19$). Model 2 was adjusted for age and BMI (‘healthy’ OR = 1.51 (95% CI 0.80–2.86; $P = 0.20$), ‘unhealthy’ OR = 1.34 (95% CI 0.71–2.54; $P = 0.37$). Model 3 was adjusted for model 2 covariates plus smoking and alcohol status, hypertension, DM, IHD and CVA (‘healthy’ OR =

1.50 (95% CI 0.79–2.86; $P = 0.21$), ‘unhealthy’ OR = 1.45 (95% CI 0.76–2.78; $P = 0.26$)).

Discussion

To our knowledge, this is the first study to examine associations between a posteriori-derived DP and AMD risk in a European population. Two major DP were identified using PCA and labelled as ‘healthy’ and ‘unhealthy’ with no evidence of association with AMD risk in this older population with a restricted lifestyle. Previous evidence of the association between DP and AMD risk is limited and somewhat conflicting [14, 15]. The MCCS found no association between DP and early AMD risk. However, they noted that a ‘grains and fish’ DP, characterised by frequent consumption of boiled rice, muesli, fish (not fried), chicken (not fried), vegetables, and avoidance of white bread was associated with 51% reduced odds of late AMD while a diet characterised by red and processed meats and fried foods was significantly associated with a higher prevalence of late AMD across quartiles of adherence [25]. In contrast to our findings, results from the US-based AREDS study by Chiu et al. identified strong associations between an ‘Oriental’ and ‘Western’ DP with decreased and increased risk for both early and late AMD, respectively [14]. However, comparing DP across the various studies is challenging given the complexity of the composition of the DP and the geographical variations in dietary intake. In addition, potential confounding from known or unknown variables such as genetic data (effect modification due to genotype) or variation in the classification of AMD definitions could limit detectable associations with dietary outcomes [10].

Other studies have considered various combinations of single foods and nutrients and AMD risk [11, 31]. A retrospective case-control study examining diet quality used the Alternate Healthy Eating Index (AHEI) to evaluate scores for the intake of five different food groups (grains, fruits, vegetables, meat and milk) and both total and saturated fat intake, identifying participants in the highest quartile of dietary quality with lower risk of late AMD (OR, 0.54; 95% CI, 0.30–0.99) compared to those in the lowest quartile. However, no association was found using a HEI score [31]. The Carotenoids in Age-Related Eye Disease Study (CAREDS) used FFQ data from 1313 female participants aged 55 to 74 years to generate a modified HEI (mHEI) which considered intake of whole grains, vegetables, fruits, meat, beans, eggs, fish and milk, total fat, saturated fat, sugar and alcohol [11, 32]. Their results showed a 46% lower odds for early AMD in the highest mHEI quintile compared to the lowest quintile (OR, 0.54; 95% CI, 0.33–0.88) [11]. Merle and colleagues assessed associations between progression to late AMD and adherence to a Mediterranean diet in 2525 AREDS participants [33]. An alternate Mediterranean diet (aMeDi) score

Table 4 Association between the risk of AMD (controls versus AMD) and dietary patterns in the participants of the Irish Nun Eye Study ($n = 922$)

Dietary pattern		OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>
Dietary pattern 1							
Healthy	Q1 (least healthy)	Q2		Q3		Q4 (most healthy)	
Model 1	1.00 (ref)	1.15 (0.65, 2.04)	0.62	0.99 (0.55, 1.77)	0.97	0.86 (0.48, 1.55)	0.62
Model 2	1.00 (ref)	1.30 (0.71, 2.38)	0.40	1.43 (0.77, 2.64)	0.25	1.51 (0.80, 2.86)	0.20
Model 3	1.00 (ref)	1.29 (0.70, 2.37)	0.41	1.38 (0.74, 2.57)	0.31	1.50 (0.79, 2.86)	0.21
Dietary pattern 2:							
Unhealthy	Q1 (least unhealthy)	Q2		Q3		Q4 (most unhealthy)	
Model 1	1.00 (ref)	1.30 (0.70, 2.43)	0.41	1.78 (0.99, 3.21)	0.06	1.50 (0.82, 2.75)	0.19
Model 2	1.00 (ref)	1.22 (0.63, 2.34)	0.56	1.54 (0.83, 2.86)	0.17	1.34 (0.71, 2.54)	0.37
Model 3	1.00 (ref)	1.18 (0.61, 2.28)	0.62	1.61 (0.86, 3.01)	0.14	1.45 (0.76, 2.78)	0.26

Values summarised as odds ratio for risk of AMD according to a ‘healthy’ or ‘unhealthy’ dietary pattern

Model 1, unadjusted; Model 2, adjusted for age, BMI; Model 3, adjusted for Model 2 plus smoking status, alcohol status, diabetes mellitus, HTN, IHD and CVA. *AMD*, age-related macular degeneration; *BMI*, body mass index; *CVA*, cerebrovascular accident; *HTN*, hypertension; *IHD*, ischaemic heart disease

was used to evaluate the intake of nine food groups (whole grains, fruit, vegetables, legumes, nuts, red and processed meats, fish, alcohol) as well as a mono-unsaturated to saturated fats ratio. A 26% lower risk of progression to late AMD was significantly associated with an elevated aMeDi score after adjusting for age, sex, AMD grade, total energy intake, AREDS treatment, smoking, BMI, educational level, supplement use and 10 genetic variants (hazard ratio [HR], 0.74; 95% CI, 0.61–0.91; P -trend = 0.007) [33].

This study had a number of key strengths: the standardised collection of data including potential confounding factors, the photographic grading of AMD from high-resolution digital macula-centred images and the use of phenotypic data from both eyes of each participant for analysis. AMD classification was undertaken in an independent retinal grading centre without any knowledge of the associated nutritional data. The use of DP to evaluate a broader overview of dietary intake as opposed to the assessment of individual nutrients or foods alone provides a more comprehensive understanding of the synergistic interactions between food consumption and disease risk and reduces the influences of residual confounding. The use of an a posteriori approach for DP analysis is also considered to be an approach that is free from the constraints of a priori hypotheses [34, 35].

Limitations of this study include the cross-sectional design which prevents determination of causal influences associated with dietary intake. An ‘unhealthy’ DP characterised by a range of ‘unhealthy’ foods may be associated with increased risk of AMD, however due to the low prevalence of AMD in this population ($n = 83$ for early AMD and $n = 21$ for late AMD), and the fact that this cohort were reasonably healthy given their age (96% non-smokers, 91% non-drinkers and 45% with a normal BMI) and consuming a restricted diet, statistical power to

detect an association with AMD may have been limited. Although, we evaluated dietary intake as a whole and the analysis included known non-dietary confounders, it is possible that residual confounding still represents a concern. However, the homogeneous lifestyle of this cohort is likely to have limited residual confounding. There are also limitations associated with the assessment of dietary intake. Although FFQs are commonly used to infer DP and dietary intake in epidemiological studies, they are nonetheless prone to inherent error and are limited in their ability to assess all dietary components. Although 170 foods were considered within the FFQ, additional foods consumed may have been missed. There are also drawbacks of DP analysis including consolidation of the food items into groups, choosing the number of factors to be retained, selecting the method of rotation and naming the factors identified. As this was a cross-sectional study, AMD status and dietary intake were only assessed at a single time point and therefore we were unable to account for temporal changes over a period of time. However, other studies have used comparable methods and highlighted the reproducibility of DP over time [36, 37].

In conclusion, this study showed no evidence of an association between DP and AMD risk in an ageing and well-characterised cohort with a restricted lifestyle. Nevertheless, increasing evidence suggests unhealthier food choices may contribute to increased AMD risk [3, 11, 31]. Critically, further longitudinal assessment and intervention studies are necessary to evaluate dietary intake of specific foods and DP as a risk factor for AMD, particularly in older adults with late AMD.

Acknowledgements The authors thank all study participants and also Janet Kyle for provision of the food frequency questionnaire and providing advice regarding the set up and design of the study. The project was

funded by the Northern Ireland Health and Social Care Research and Development Office (grant number RRG project 4.41), and the Department for Education and Learning, Northern Ireland. The funding organisations had no role in the design or conduct of this research.

Funding G. Silvestri reports grants from Northern Ireland Health and Personal Social Services Research and Development Office during this study, personal fees from Bayer and non-financial support from Allergan outside the submitted work. G. McKay reports personal fees from Boehringer Ingelheim.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

This article does not contain any studies with animals performed by any of the authors.

Informed consent Informed consent was obtained from all individual participants included in the study.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Miller JW (2013) Age-related macular degeneration revisited—piecing the puzzle: the LXIX Edward Jackson memorial lecture. *Am J Ophthalmol* 155:1–35
- Seddon JM, George S, Rosner B (2006) Cigarette smoking, fish consumption, omega-3 fatty acid intake, and associations with age-related macular degeneration: the US twin study of age-related macular degeneration. *Arch Ophthalmol* 124:995–1001
- Chong EW, Simpson JA, Robman LD et al (2009) Red meat and chicken consumption and its association with age-related macular degeneration. *Am J Epidemiol* 169:867–876
- Adams MK, Chong EW, Williamson E et al (2012) 20/20—alcohol and age-related macular degeneration: the Melbourne Collaborative Cohort Study. *Am J Epidemiol* 176:289–298
- Chong EW, Robman LD, Simpson JA et al (2009) Fat consumption and its association with age-related macular degeneration. *Arch Ophthalmol* 127:674–680
- Chong EW, Wong TY, Kreis AJ, Simpson JA, Guymer RH (2007) Dietary antioxidants and primary prevention of age related macular degeneration: systematic review and meta-analysis. *BMJ* 335:755
- The Age-Related Eye Disease Study 2 (AREDS2) Research Group (2013) Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: the Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trials. *JAMA* 309:2005–2015
- Seddon JM, Rosner B, Sperduto RD et al (2001) Dietary fat and risk for advanced age-related macular degeneration. *Arch Ophthalmol* 119:1191–1199
- Smith W, Mitchell P, Leeder SR (2000) Dietary fat and fish intake and age-related maculopathy. *Arch Ophthalmol* 118:401–404
- Wang JJ, Rochtchina E, Smith W et al (2009) Combined effects of complement factor H genotypes, fish consumption, and inflammatory markers on long-term risk for age-related macular degeneration in a cohort. *Am J Epidemiol* 169:633–641
- Mares JA, Voland RP, Sondel SA et al (2011) Healthy lifestyles related to subsequent prevalence of age-related macular degeneration. *Arch Ophthalmol* 129:470–480
- San Giovanni JP, Chew EY, Clemons TE et al (2007) The relationship of dietary carotenoid and vitamin A, E, and C intake with age-related macular degeneration in a case-control study: AREDS report no. 22. *Arch Ophthalmol* 125:1225–1232
- Tan JS, Wang JJ, Flood V, Mitchell P (2009) Dietary fatty acids and the 10-year incidence of age-related macular degeneration: the Blue Mountains Eye Study. *Arch Ophthalmol* 127:656–665
- Chiu CJ, Chang ML, Li T, Gensler G, Taylor A (2017) Visualization of dietary patterns and their associations with age-related macular degeneration. *Invest Ophthalmol Vis Sci* 58:1404–1410
- Chiu CJ, Chang ML, Zhang FF, Li T, Gensler G, Schleicher M, Taylor A (2014) The relationship of major American dietary patterns to age-related macular degeneration. *Am J Ophthalmol* 158:118–127
- Age-Related Eye Disease Study Research Group (2001) A randomized, placebo-controlled, clinical trial of high-dose supplementation with vitamins C and E, beta carotene, and zinc for age-related macular degeneration and vision loss: AREDS report no. 8. *Arch Ophthalmol* 119:1417–1436
- Chiu CJ, Milton RC, Klein R, Gensler G, Taylor A (2009) Dietary compound score and risk of age-related macular degeneration in the age-related eye disease study. *Ophthalmology* 116:939–946
- Stevens R, Barlett H, Cooke R (2015) Dietary analysis and nutritional behaviour in people with and without age-related macular disease. *Clin Nutr ESPEN* 10(3):e112–e117
- Braakhuis A, Raman R, Vaghefi E (2017) The association between dietary intake of antioxidants and ocular disease. *Diseases* 5:3
- Chiu CJ, Klein R, Milton RC, Gensler G, Taylor A (2009) Does eating particular diets alter the risk of age-related macular degeneration in users of the age-related eye disease study supplements? *Br J Ophthalmol* 93:1241–1246
- Chiu CJ, Hubbard LD, Armstrong J, Rogers G, Jacques PF, Chylack LT Jr, Hankinson SE, Willett WC, Taylor A (2006) Dietary glycemic index and carbohydrate in relation to early age-related macular degeneration. *Am J Clin Nutr* 83:880–886
- Chiu CJ, Milton RC, Gensler G, Taylor A (2007) Association between dietary glycemic index and age-related macular degeneration in nondiabetic participants in the age-related eye disease study. *Am J Clin Nutr* 86:180–188
- Chiu CJ, Milton RC, Klein R, Gensler G, Taylor A (2007) Dietary carbohydrate and progression of age-related macular degeneration, a prospective study from the age-related eye disease study. *Am J Clin Nutr* 86:1210–1218
- Moghaddam E, Vogt JA, Wolever TM (2006) The effects of fat and protein on glycemic responses in nondiabetic humans vary with waist circumference, fasting plasma insulin, and dietary fiber intake. *J Nutr* 136:2506–2511
- Amirul Islam FM, Chong EW, Hodge AM, Guymer RH, Aung KZ, Makeyeva GA, Baird PN, Hopper JL, English DR, Giles GG, Robman LD (2014) Dietary patterns and their associations with age-related macular degeneration: the Melbourne collaborative cohort study. *Ophthalmology* 121:1428–1434
- McGowan A, Silvestri G, Moore E et al (2014) Retinal vascular caliber, iris color and age-related macular degeneration in the Irish Nun Eye Study. *Invest Ophthalmol Vis Sci* 56:382–387

27. McGowan A, Silvestri G, Moore E, Silvestri V, Patterson CC, Maxwell AP, McKay GJ (2015) Evaluation of the retinal vasculature in hypertension and chronic kidney disease in an elderly population of Irish Nuns. *PLoS One* 10:e0136434
28. Klein R, Davis MD, Magli YL, Segal P, Klein BEK, Hubbard L (1991) The Wisconsin age-related maculopathy grading system. *Ophthalmology* 98:1128–1134
29. Bird AC, Bressler NM, Bressler SB et al (1995) An international classification and grading system for age-related maculopathy and age-related macular degeneration. The International ARM Epidemiological Study Group. *Surveys of Ophthalmology* 39:367–374
30. Food Standards Agency (2006) Food portion sizes, 3rd edn. TSO, London
31. Montgomery MP, Kamel F, Pericak-Vance MA, Haines JL, Postel EA, Agarwal A, Richards M, Scott WK, Schmidt S (2010) Overall diet quality and age-related macular degeneration. *Ophthalmic Epidemiol* 17:58–65
32. Guenther PM, Reedy J, Krebs-Smith SM, Reeve BB, Basiotis PP (2008) Development of the healthy eating index-2005. *J Am Diet Assoc* 108:1896–1901
33. Merle BM, Silver RE, Rosner B, Seddon JM (2015) Adherence to a Mediterranean diet, genetic susceptibility, and progression to advanced macular degeneration: a prospective cohort study. *Am J Clin Nutr* 102:1196–1206
34. Kastorini CM, Papadakis G, Milionis HJ, Kalantzi K, Puddu PE, Nikolaou V, Vemmos KN, Goudevenos JA, Panagiotakos DB (2013) Comparative analysis of a-priori and a-posteriori dietary patterns using state-of-the-art classification algorithms: a case/control study. *Artif Intell Med* 59:175–183
35. Neville CE, Montgomery S, Silvestri G et al (2017) Dietary patterns and retinal vessel caliber in the Irish Nun Eye Study. *J Nutr Health Aging*:1–8
36. Hu FB, Rimm E, Smith-Warner SA, Feskanich D, Stampfer MJ, Ascherio A, Sampson L, Willett WC (1999) Reproducibility and validity of dietary patterns assessed with a food-frequency questionnaire. *Am J Clin Nutr* 69:243–249
37. Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC (2000) Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* 72:912–921