



Association of *BMP15* and *GDF9* variants to premature ovarian insufficiency

Monise Santos¹ · Emerson Barchi Cordts^{1,2} · Carla Peluso² · Mayla Dornas² · Felipe Heurre Vieira Neto² · Bianca Bianco^{1,2} · Caio Parente Barbosa^{1,2} · Denise Maria Christofolini^{1,2} 

Received: 21 January 2019 / Accepted: 26 July 2019 / Published online: 7 August 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

Purpose To identify genetic variation associated to premature ovarian insufficiency (POI).

Methods A total of 74 women with POI (group POI), 45 women with increased FSH levels (group high FSH), and 88 controls (non-POI) were studied. Genotyping of *BMP15*:c.-9C>G (rs3810682), *BMP15*:c.328+905A>G (rs3897937), and *BMP15*:c.852C>T (rs17003221); and *GDF9*:c.134-694G>A (rs4705974), *GDF9*:c.-31-951G>A (rs11748063), *GDF9*:c.-152G>C (rs30177), and *GDF9*:g.1073C>T (rs803224) was performed by the TaqMan methodology. Chi-square and Fisher's exact tests were performed to evaluate the distribution of genotypes, alleles, odds ratio, and the Hardy-Weinberg equilibrium of each variation. Haplotype analysis was performed for each gene considering the case and control groups. Bonferroni's correction was applied to chi-square and Fisher's exact test data, and *p* values < 0.007 for genotypes and alleles and < 0.006 for haplotypes were considered significant.

Results It was observed a statistically significant difference in genotype distribution of *BMP15*:c.852C>T between group POI and controls (*p* < 0.001). TT and TC genotypes were more frequently observed in group POI. Genotype distribution in case group POI, however, was not in the Hardy-Weinberg equilibrium, due to the increased number of heterozygotes in the sample. Concerning *GDF9*, no association was found among the studied genetic variants and POI or high FSH groups.

Conclusion It is concluded from the present study that the genotypes CT and TT from *BMP15*:c.852C>T variation may be risk factors for the development of POI.

Keywords Premature ovarian insufficiency · *BMP15* · *GDF9* · Infertility · Gene variants

Introduction

Women's reproductive age was postponed due to the changes in professional and social life in the last decades. This fact put in evidence of some gynecological pathology that impact on ovarian reserve decrease and premature ovarian insufficiency (POI) and menopause [1].

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10815-019-01548-0>) contains supplementary material, which is available to authorized users.

✉ Denise Maria Christofolini
Denise.morf@gmail.com

¹ Instituto Ideia Fértil, Avenida Príncipe de Gales, 821, Santo André, SP 09060-650, Brazil

² Faculdade de Medicina do ABC, Av. Lauro Gomes, 2000, Room 101, CEPES, Santo André, SP 09060-870, Brazil

In regular conditions, menopause occurs among 50–52 years old, while premature menopause is considered when menses cessation and ovarian decrease occur before 40 years old [2]. Apart from menstrual irregularity, the other criterion that must be considered to proper diagnosis of POI is FSH (follicle-stimulating hormone) dosage.

Traditionally, POI has been diagnosed through two or more FSH dosages that result in levels above 40 mIU/mL at least 1 month apart. However, in 2013, the ESHRE POI Consensus (that occurred in Utrecht, the Netherlands) determined a new cutoff level for FSH as two dosages of 25 mIU/mL or above, respecting the same period of time [3].

The etiology of POI is complex and can be attributed to autoimmune diseases, infections, iatrogenic exposition, gene mutations, and chromosome aberrations [4–7]. Therefore, adrenocortical (21-OH) and anti-thyroid antibodies (TPO), drugs, and previous surgeries affecting ovaries must be

considered in clinical evaluation. Also, karyotype, *FMRI* gene premutation must be the first investigated when a genetic cause is considered.

However, mutations in other genes have emerged as possible causes of POI, when primary causes were discarded, including genes that exert hormonal effects or affect follicular function as *BMP15* and *GDF9* [8].

BMP15 is a member of the beta-growth factor superfamily (TGFb) and is located in Xp11.2, an important chromosomal region for ovarian differentiation [9, 10]. *GDF9* also is a member of beta-growth factor superfamily (TGFb) and is located in 5q31.1. [11, 12]. The factors secreted by oocytes, *BMP15* and *GDF9* play an important role in the development of the primordial follicle, ovulation process, corpus luteum formation, proliferation of granulosa cells, and oocyte maturation through paracrine/autocrine signaling pathways [9, 10, 13]. In bovines, the *BMP15* transcript is observed from oocyte maturation to morula stage, after fertilization [14].

In view of the possible role of these genes in POI development and the small number of scientific articles available, most of them performed with European and Asiatic populations, the objective of the present study was to evaluate the association of *BMP15* and *GDF9* variations with POI in a Brazilian population and to evaluate which variants are shared by patients with increased levels of FSH, in order to contribute to elucidation of the etiology of premature ovarian insufficiency.

Methods

Casuistic

The present study is an association study, using a case-control design, developed with patients recruited from the premature ovarian insufficiency outpatient clinic of the Instituto Ideia Fértil (case group) and women recruited from the gynecology and climacteric outpatient clinic of Faculdade de Medicina do ABC (control group). The research project was approved by the FMABC's Ethics Committee and all patients read and signed the informed consent form.

The case group was composed of 119 women and subdivided into two groups:

- The case group POI consisted of 74 women with POI, with FSH levels > 25 mIU/mL and cessation of menstruation before 40 years of age. POI criteria followed The ESHRE Guideline Group on POI [3]. All of them had age up to 40 years old at the time of study inclusion.
- The case group high FSH consisted of 45 women, with FSH levels between 10 and 25 mIU/mL, irregular menses, and age up to 40 years old, possible candidates to ovarian insufficiency development. Sisters or cousins of patients were not included in the sample.

Patients with a history of surgical manipulation of the gonads and gonadal lesions and history of radio- and/or chemotherapy were excluded. Also, patients with abnormal levels of thyroid hormones and reagent anti-TPO were excluded from the sample. As inclusion criteria, all patients (POI and high FSH groups) must have normal karyotype and normal genotype for the *FMRI* gene expansion (alleles with less than 55 CGG repeats.).

The control group consisted of 88 non-POI women with 50 years of age or more, whose menopause occurred after 50 years of age naturally. All of them had at least one pregnancy and parity and never experienced irregular menses. As in menopause increased levels of FSH were normally expected, no FSH dosage was performed for the control group. Women with a familial history of POI were not included in the control group.

Genotyping

The collection of 5 mL of peripheral blood in a tube containing EDTA from all the women in the study was performed. DNA extraction was performed using the salting out methodology [15].

The studied variants of the *BMP15* and *GDF9* genes, described in Table 1, were identified by real-time PCR using the TaqMan SNP genotyping assays (C_27504454_10, C_7455457_10, C_32866291_10, C_27970131_10, C_2563478_10, C_2264894_10, C_2563470_10). PCR conditions were provided by the manufacturer: heating at 60 °C for 30 s, followed by an initial denaturation of 95 °C for 20 s, followed by 40 denaturation cycles of 95 °C for 1 s, annealing at 60 °C for 20 s. Positive controls for each genotype were included in all reactions. Variant nomenclature was described according to HGVS variant nomenclature (<https://varnomen.hgvs.org>).

Statistical analysis

For quantitative variables as age and FSH levels, Shapiro-Wilk's test was performed in order to identify if the samples have normal distribution. Data with no normal distribution was described by median and percentile. Normal data was described by average and standard deviation.

The distribution data of the genotypes and alleles, odds ratio, and the Hardy-Weinberg (HW) equilibrium were analyzed by the chi-square test and Fisher exact test using BioStat 3.0 software. Bonferroni's correction was applied to genotype and allele distribution analysis and *p* values < 0.007 were considered statistically significant.

The analysis of the haplotypes for the two genes was also performed using Haploview 4.2 software. The *p* value < 0.006 was considered statistically significant after Bonferroni's correction.

Table 1 Description of *BMP15* and *GDF9* variations according to dbSNP (accessed January 15, 2019)

<i>BMP15</i> (NM_005448.2)	rs3810682:C>G	rs3897937:A>G	rs17003221:C>T	
Alleles	C>G (rev)	A>G	C>T	
Ancestral allele	G	G	T	
HGVS name	<i>BMP15</i> :c.-9 C>G	<i>BMP15</i> :c.328+905 A>G	<i>BMP15</i> :c.852 C>T	
Functional consequence	5'UTR variant	Intronic variant	p.Ser284=	
GnomAD allele frequency	<i>G</i> = 0.16200	<i>G</i> = 0.37980	<i>T</i> = 0.05660	
AbraOM allele frequency*	<i>G</i> = 0.18182	Not reported	<i>T</i> = 0.107784	
<i>GDF9</i> (NM_001288824.2)	rs4705974:G>A	rs11748063:G>A	rs30177:G>C	rs803224:C>T
Alleles	C>T (fwd)	C>T (fwd)	G>C (rev)	C>T (rev)
Ancestral allele	T	T	C	T
HGVS name	<i>GDF9</i> :c.134-694 G>A	<i>GDF9</i> :c.-31-951G>A	<i>GDF9</i> :c.-152 G>C	rs803224:C>T
Functional consequence	Intronic variant	5'UTR variant	5'UTR variant	Intergenic
GnomAD allele frequency	<i>T</i> = 0.1703	<i>T</i> = 0.3484	<i>C</i> = 0.3082	<i>T</i> = 0.1396
AbraOM allele frequency*	Not reported	Not reported	<i>C</i> = 0.724315	Not reported

*AbraOM is a database for Brazilian genomic variants [16]

Results

The average ages of patients in case groups POI and high FSH were 38 [35–41] and 37 [35–39] years old, respectively, and in the control group was 56 [53.75–60.5] years old. The average levels of FSH hormone for POI patients were 67.20 mIU/mL [53.23–108.90] in the first dosage and 65.00 [41.19–101.40] in the second dosage and 16.15 [12.26–19.19] and 16.05 [12.65–22.18] mIU/mL for high FSH patients. Additional exams and hormone dosages can be observed in supplementary data as Table 4.

GDF9 and *BMP15* genotype and allele distributions among case groups 1 and 2 and controls were statistically evaluated as well as odds ratio and HW equilibrium data. The results can be observed in Table 2.

The statistical analysis of the *BMP15*:c.852C>T showed a statistically significant difference of group POI results for genotype distribution ($p < 0.001$) when compared with the control group. The CT and TT genotypes were more frequent in the group POI. The other *BMP15* variations evaluated showed no statistically significant difference. *GDF9* variations evaluated showed no statistically significant difference among groups.

No *BMP15* or *GDF9* haplotypes were associated to POI or high FSH. Data is available in Tables 3.

Discussion

The etiology of premature ovarian insufficiency can be attributed to several genetic alterations and is still a subject of research. According to Vilodre et al. (2007), different causes of POI may lead to either reduction in the number of follicles and/or defects in the follicle stimulus mechanism [14]. In this

project, we performed the analysis of the variations rs3810682, rs3897937, and rs17003221 of the *BMP15* gene and the variations rs4705974, rs11748063, rs30177, and rs803224 of the *GDF9* gene in two groups of patients: women with established ovarian insufficiency and women with increased FSH values, possible candidates for POI. We compared the findings in these groups to the results of a control group in order to find possible risk factors for POI in our population and to identify women at high risk for developing ovarian failure. Also, we have studied haplotypes associated to the condition to both genes.

The correlation of the *BMP15* gene with POI was observed initially in animal models. Galloway et al. in 2000 [17] performed a study with sheep that had a mutation in *BMP15*. The heterozygous models had an increased rate of ovulation and birth of twins and triplets. The homozygous sheep developed ovarian failure, originated from a compromised follicular development. Besides, a substitution in *BMP15* (Y235C) in heterozygous was identified from sisters with POI leading to further investigation of the importance of the gene for the condition [18]. Since then, some authors have investigated the association of *BMP15* variations with premature ovarian insufficiency in different populations with diverse findings, as described by Qin et al. (2015) [19] and detailed in Table 5 in the Electronic Supplementary Materials (ESM) [18, 20–32]. Some variations seem to be frequent in POI cases while some were exclusive of specific populations.

In the present study, the analysis of *BMP15* genotypes and alleles showed that the CT and TT genotypes for *BMP15*:c.852 C>T were more frequent in the case group POI than in the control group. No difference was observed in the distribution of the *BMP15* variations among patients with high FSH and controls, and no haplotype was associated to POI or high FSH groups.

Table 2 Genotype and allele distributions for *BMP15* and *GDF9* polymorphisms

Gene	SNP	Group	n	Genotypes, n (%)			p	Alleles, n (%)		p	HWE
<i>BMP15</i>	rs3810682 c.-9C>G	POI	70	GG	CG	CC		G	C		
				46 (66%)	16 (23%)	8 (11%)	0.72	108 (77%)	32 (23%)	0.69	–
		High FSH	42	29 (69%)	8 (19%)	5 (12%)	0.57	66 (79%)	18 (21%)	0.94	–
	rs3897937 c.328+ 905A>G	Control	88	58 (66%)	23 (26%)	7 (8%)		139 (79%)	37 (21%)		
				GG	GA	AA		G	A		
		POI	71	3 (4%)	47 (66%)	21 (30%)	0.06	53 (37%)	89 (63%)	0.09	–
	rs17003221 c.852C>T	High FSH	43	1 (2%)	32 (74%)	10 (23%)	0.26	34 (40%)	52 (60%)	0.28	–
		Control	88	7 (8%)	68 (77%)	13 (15%)		82 (47%)	94 (53%)		
				TT	TC	CC		T	C		
		POI	70	18 (26%)	45 (64%)	7 (10%)	< 0.001	81 (58%)	59 (42%)	0.03	0.02
<i>GDF9</i>	rs30177 c.-152G>C	High FSH	43	15 (35%)	15 (35%)	13 (30%)	0.58	45 (52%)	41 (48%)	0.31	–
		Control	88	23 (26%)	34 (38%)	31 (35%)		80 (46%)	95 (54%)		
				CC	CG	GG		C	G		
	rs11748063 c.-31-951G>A	POI	74	8 (11%)	22 (30%)	44 (59%)	0.73	38 (26%)	110 (74%)	0.64	–
		High FSH	45	7 (15%)	16 (36%)	22 (49%)	0.67	30 (33%)	60 (67%)	0.37	–
		Control	87	9 (10%)	31 (36%)	47 (54%)		49 (28%)	126 (72%)		
	rs803224 Intergenic	POI	65	15 (23%)	22 (34%)	28 (43%)	0.033	52 (40%)	78 (60%)	0.02	–
		High FSH	41	5 (12%)	14 (34%)	22 (54%)	0.118	24 (29%)	58 (71%)	0.78	–
		Control	87	7 (8%)	34 (39%)	46 (53%)		48 (28%)	126 (72%)		
	rs4705974 c.134-694G>A	POI	70	56 (80%)	13 (19%)	1 (1%)	0.33	125 (89%)	15 (11%)	0.46	–
		High FSH	45	33 (73%)	8 (18%)	4 (9%)	0.72	74 (82%)	16 (18%)	0.35	–
		Control	86	68 (79%)	13 (15%)	5 (6%)		148 (87%)	23 (13%)		
				TT	TC	CC		T	C		
		POI	65	0 (0%)	14 (22%)	51 (78%)	0.31	14 (11%)	116 (89%)	0.49	–
		High FSH	41	1 (2%)	10 (24%)	30 (73%)	0.81	12 (15%)	70 (85%)	0.78	–
		Control	86	3 (3%)	17 (20%)	66 (77%)		23 (13%)	149 (87%)		

After Bonferroni's correction, the *p* value < 0.007 was considered significant for genotypes and alleles. It was not applied to HWE that considered *p* values < 0.05 to evidence disequilibrium *n*, number of samples; %, percentage of samples; *HWE*, the Hardy-Weinberg equilibrium; Statistically significant results are in italics

Table 3 Haplotype distribution concerning *BMP15* and *GDF9* variations

SNV	Haplotype associations	Frequency ¹	Case ratios ¹	Control ratios ¹	<i>p</i> value ¹	Frequency ²	Case ratios ²	Control ratios ²	<i>p</i> value ²
<i>BMP15</i> haplotype									
rs3810682	GAC	0.292	0.276	0.304	0.579	0.248	0.212	0.265	0.352
rs3897937	GGT	0.211	0.238	0.189	0.288	0.157	0.166	0.153	0.786
rs17003221	GAT	0.171	0.202	0.147	0.199	0.199	0.236	0.182	0.310
	GGC	0.108	0.056	0.149	0.008	0.184	0.172	0.190	0.732
	CAT	0.066	0.079	0.055	0.406	0.069	0.091	0.058	0.328
	CGT	0.062	0.060	0.063	0.904	0.055	0.043	0.061	0.541
	CGC	0.047	0.025	0.064	0.099	0.046	0.012	0.062	0.072
	CAC	0.044	0.065	0.028	0.107	0.041	0.068	0.028	0.132
<i>GDF9</i> haplotype									
rs4705974	CCGT	0.496	0.447	0.538	0.102	0.527	0.488	0.547	0.367
rs11748063	CTGT	0.189	0.241	0.144	0.025	0.135	0.135	0.136	0.981
rs30177	CCCC	0.105	0.094	0.115	0.557	0.129	0.158	0.115	0.317
rs803224	TTCT	0.104	0.094	0.113	0.588	0.116	0.121	0.114	0.875
	CCCT	0.057	0.060	0.054	0.804	0.043	0.043	0.045	0.806
	TTGT	0.024	0.037	0.013	0.167	0.016	0.024	0.012	0.468

After Bonferroni's correction, the *p* value < 0.006 was considered significant

¹ Case group IOP

² Case group 2 High FSH

In a previous study performed by our research group, Peluso et al. (2017) studied 186 infertile women with regular FSH levels, in order to correlate the same *BMP15* variations investigated here to ovarian response. It was observed that none of the alleles was associated to oocyte retrieval or oocyte maturation results in their population. Their results reinforce that the variation is not common in Brazilian population, and the association found in the present research is specific to POI. Peluso found a significant association of TT genotype from *BMP15*:c.852 C>T to increased levels of estradiol, a serum abnormality also observed in patients with POI [33].

The Hardy-Weinberg deviation of genotype distribution was observed for this variation on case group POI, due to an increased number of heterozygotes. Deviation from HWE at a marker locus can be due to diverse reasons as population stratification, inbreeding, selection, nonrandom mating, genotyping error, actual association to the disease or trait under study, or a deletion or duplication variation [34]. In fact, in a study of complex diseases in human populations, none of these reasons can be discharged. Therefore, the equilibrium deviation can be a good fact, reinforcing the association of such genotype to the condition or a bad fact, considering that a methodological error could have occurred. Here, reinforcing the idea of association to the condition, we observed no deviation in the control group.

In the literature data, Ma et al. (2015), Tiotiu et al. (2010), Zhang et al. (2007), and Laissue et al. (2006) observed that the frequencies of the *BMP15* c.852C>T genotypes did not vary significantly between the POF and control groups, while Dixit

et al. (2006) showed that the haplotype G-G-C in the three frequent variants c.-9C>G, c.308A>G, and c.852C>T was associated with POI [21, 22, 26, 27, 29].

Concerning *BMP15* c.-9C>G, Leidig et al. (2008) did not find association of with the condition [25]. Ma et al. (2015) observed that the C allele was disproportionally frequent in cases and controls, and Dixit et al. (2006) also observed association of the variant as described previously [21, 29]. Regarding *GDF9*, no difference was observed in the distribution of the *GDF9* variations among patients and controls. Also, in the literature, no association was found between the variants studied with POI.

Voorhuis et al. in 2011 carried out a study with 3616 patients with natural menopause with the objective of studying the association between the 23 single-base variations in five genes, *AMH*, *AMHR2*, *BMP15*, *FOXL2*, and *GDF9* with the timing of menopause. They evaluated rs3810682, rs6521896, rs17249566, rs5961233, and rs3897937 in *BMP15*, of which two of them are common to our study. In *GDF9*, they evaluated six different variations, including rs30177, rs803224, rs11748063, and rs4705974, which are common to the present study. The analysis also showed no significant association of *GDF9* variations to menopause age, but rs6521896 in *BMP15* was associated with later menopause [35].

The divergence of the results of the present study against the results of other populations demonstrates the divergence in the ethnic composition of the Brazilian population, in particular the population of São Paulo's metropolitan area, where

the study was developed. Brazil is a continental country with wide intra-territorial differences. Thus, the application of international models for the attribution of risk factors should be carefully considered. To exemplify, Carvalho-Silva et al. (2001) carried out a genetic study with DNA from Brazilian white population. They found distinct chromosome Y footprints of Italian immigration to southern Brazil, migration of Moroccan Jews to the Amazon region, and Dutch DNA in Brazil northeast. In contrast with the mtDNA data in white Brazilians, which showed that more than 60% of the matrilineages were Amerindian or African, only 2.5% of the Y-chromosome lineages were from Sub-Saharan Africa, and none were Amerindian [36].

One limitation of the present study is the sample size, which despite being larger than that of many POI studies is still small for association analysis. More robust studies could confirm the results observed in the present study. To justify our sample size, we would like to emphasize the rigor of patient selection, excluding genetic alterations common to this group of patients (as *FMR1* premutated alleles and positive anti-TPO). Another limitation was the method of choosing the variations of each gene, which was based on previous association in the literature. It is possible that other variations of the studied genes were associated to the condition, but were not addressed in the present study. Direct sequencing and GWAS studies could evidence different variations associated to POI and are being performed by the research group.

The genetic study and determination of the frequency of gene variants in cases and controls contribute to the understanding of possible risk factors related to the development of premature ovarian insufficiency. To understand these factors is extremely important in a population that has changed their lifestyle, in a population where pregnancies are delayed due to large use of contraceptive methods and the equal insertion of women in universities and in the labor market. Early mapping of risk factors can assist women in reproductive decision-making, such as anticipating motherhood or preserving oocytes or embryos.

Conclusion

The cause of POI has been studied worldwide. The many divergences in results found may be associated with different ethnicities of the populations studied. Variant genotypes from *BMP15* were found at high frequency in women with POI. In view of the important physiological role of this gene in oogenesis, these variants may be associated with the risk of developing POI in Brazilian population.

Compliance with ethical standards The research project was approved by the FMABC's Ethics Committee and all patients read and signed the informed consent form.

References

1. Silva AA, Arent A, Petracco A, Mancebo AC, Bastos AM, Fraga AM, et al. Tratado de Reprodução Assistida. 1ª. Ed. São Paulo: Abril 2010.
2. Goswami D, Conway GS. Premature ovarian failure. Hum Reprod Update. 2005;11(4):391–410.
3. European Society for Human Reproduction and Embryology (ESHRE) Guideline Group on POI, Webber L, Davies M, Anderson R, Bartlett J, Braat D, et al. ESHRE guideline: management of women with premature ovarian insufficiency. Hum Reprod. 2016;31(5):926–37.
4. Vegetti W, Marozzi A, Manfredini E, Testa G, Alagna F, Nicolosi A, et al. Premature ovarian failure. Mol Cell Endocrinol. 2000;161(1–2):53–7.
5. Mattison DR, Evans MI, Schwimmer WB, White BJ, Jensen B, Schulman JD. Familial premature ovarian failure. Am J Hum Genet. 1984;36:1341–8.
6. Alper MM, Garner PR, Seibel MM. Premature ovarian failure. Current concepts. J Reprod Med. 1986;31:699–708.
7. Anasti JN. Premature ovarian failure: an update. Fertil Steril. 1998;70:1–15.
8. Conway GS. Premature ovarian failure. Curr Opin Obstet Gynecol. 1997;9:202–6.
9. Kovanci E, Simpson JL, Amato P, Rohozinski J, Heard MJ, Bishop CE, et al. Oocyte specific G protein coupled receptor-3 (GPR3) gene perturbations are not commonly involved in premature ovarian failure. Fertil Steril. Published online October 20, 2007.
10. Paulini F, Melo EO. The role of oocyte-secreted factors *GDF9* and *BMP15* in follicular development and oogenesis. Reprod Domest Anim. 2011;46:354–61.
11. Trombly DJ, Woodruff TK, Mayo KE. Roles for transforming growth factor beta superfamily proteins in early folliculogenesis. Semin Reprod Med. 2009;27:14–23.
12. Kedem A, Fisch B, Garor R, Ben-Zaken A, Gizunterman T, Felz C, et al. Growth differentiating factor 9 (GDF9) and bone morphogenetic protein 15 both activate development of human primordial follicles in vitro, with seemingly more beneficial effects of GDF9. J Clin Endocrinol Metab. 2011;96:1246–54.
13. Vilodre LC, Moretto M, Kohek MB, Spritzer PM. Premature ovarian failure: present aspects. Arq Bras Endocrinol Metabol. 2007;51(6):920–9.
14. Pennetier S, Uzbekova S, Perreau C, Papillier P, Mermillod P, Dalbès-Tran R. Spatio-temporal expression of the germ cell marker genes *MATER*, *ZAR1*, *GDF9*, *BMP15*, and *VASA* in adult bovine tissues, oocytes, and preimplantation embryos. Biol Reprod. 2004;71:1359–66.
15. Lahiri DK, Numberger JI. A rapid non-enzymatic method for Q2 preparation of HMW DNA from blood for RFLP studies. Nucleic Acids Res. 1991;19:5444.
16. Naslavsky MS, Yamamoto GL, de Almeida TF, Ezquina SAM, Sunaga DY, Pho N, et al. Exomic variants of an elderly cohort of Brazilians in the ABraOM database. Hum Mutat. 2017;38(7):751–63.
17. Galloway SM, McNatty KP, Cambridge LM, Laitinen MP, Juengel JL, Jokiranta TS, et al. Mutations in an oocyte-derived growth factor gene (*BMP15*) cause increased ovulation rate and infertility in a dosage-sensitive manner. Nat Genet. 2000;25(3):279–83.
18. Di Pasquale E, Beck-Peccoz P, Persani L. Hypergonadotropic ovarian failure associated with an inherited mutation of human bone morphogenetic protein-15 (*BMP15*) gene. Am J Hum Genet. 2004;75(1):106–11.
19. Qin Y, Jiao X, Simpson JL, Chen ZJ. Genetics of primary ovarian insufficiency: new developments and opportunities. Hum Reprod Update. 2015;21(6):787–808.

20. Kumar R, Alwani M, Kosta S, Kaur R, Agarwal S. BMP15 and GDF9 gene mutations in premature ovarian failure. *J Reprod Infertil*. 2017;18(1):185–9.
21. Ma L, Chen Y, Mei S, Liu C, Ma X, Li Y, et al. Single nucleotide polymorphisms in premature ovarian failure-associated genes in a Chinese Hui population. *Mol Med Rep*. 2015;12(2):2529–38.
22. Tiotiu D, Alvaro Mercadal B, Imbert R, Verbist J, Demeestere I, De Leener A, et al. Variants of the BMP15 gene in a cohort of patients with premature ovarian failure. *Hum Reprod*. 2010;25(6):1581–7.
23. Wang B, Wen Q, Ni F, Zhou S, Wang J, Cao Y, et al. Analyses of growth differentiation factor 9 (GDF9) and bone morphogenetic protein 15 (BMP15) mutation in Chinese women with premature ovarian failure. *Clin Endocrinol*. 2010;72(1):135–6.
24. Rossetti R, Di Pasquale E, Marozzi A, Bione S, Toniolo D, Grammatico P, et al. BMP15 mutations associated with primary ovarian insufficiency cause a defective production of bioactive protein. *Hum Mutat*. 2009;30(5):804–10.
25. Ledig S, Röpke A, Haeusler G, Hinney B, Wieacker P. BMP15 mutations in XX gonadal dysgenesis and premature ovarian failure. *Am J Obstet Gynecol*. 2008;198(1):84.e1–5.
26. Zhang P, Shi YH, Wang LC, Chen ZJ. Sequence variants in exons of the BMP-15 gene in Chinese patients with premature ovarian failure. *Acta Obstet Gynecol Scand*. 2007;86(5):585–9.
27. Laissue P, Christin-Maitre S, Touraine P, Kuttann F, Ritvos O, Aittomäki K, et al. Mutations and sequence variants in GDF9 and BMP15 in patients with premature ovarian failure. *Eur J Endocrinol*. 2006;154(5):739–44.
28. Dixit H, Rao LK, Padmalatha V, Kanakavalli M, Deenadayal M, Gupta N, et al. Mutational screening of the coding region of growth differentiation factor 9 gene in Indian women with ovarian failure. *Menopause*. 2005;12(6):749–54.
29. Dixit H, Rao LK, Padmalatha VV, Kanakavalli M, Deenadayal M, Gupta N, et al. Missense mutations in the BMP15 gene are associated with ovarian failure. *Hum Genet*. 2006;119(4):408–15.
30. Di Pasquale E, Rossetti R, Marozzi A, Bodega B, Borgato S, Cavallo L, et al. Identification of new variants of human BMP15 gene in a large cohort of women with premature ovarian failure. *J Clin Endocrinol Metab*. 2006 May;91(5):1976–9.
31. Peluso C, Goldman C, Cavalcanti V, Gastaldo G, Trevisan CM, Christofolini DM, et al. Use of bone morphogenetic protein 15 polymorphisms to predict ovarian stimulation outcomes in infertile Brazilian women. *Genet Test Mol Biomarkers*. 2017;21(5):328–33.
32. Duarte CW, Vaughan LK, Beasley TM, Tiwari HK. Multifactorial Inheritance and Complex Diseases. Emery and Rimoin's principles and practice of medical genetics, 6th edn. 2013; chapter 12: 1–15. ISBN: 978–0–12-383834-6.
33. Voorhuis M, Broekmans FJ, Fauser BC, Onland-Moret NC, van der Schouw YT. Genes involved in initial follicle recruitment may be associated with age at menopause. *J Clin Endocrinol Metab*. 2011;96(3):E473–9.
34. Carvalho-Silva DR, Santos FR, Rocha J, Pena SD. The phylogeography of Brazilian Y-chromosome lineages. *Am J Hum Genet*. 2001;68(1):281–6.
35. Chand AL, Ponnampalam AP, Harris SE, Winship IM, Shelling AN. Mutational analysis of BMP15 and GDF9 as candidate genes for premature ovarian failure. *Fertil Steril*. 2006;86(4):1009–12.
36. Takebayashi K, Takakura K, Wang H, Kimura F, Kasahara K, Noda Y. Mutation analysis of the growth differentiation factor-9 and -9B genes in patients with premature ovarian failure and polycystic ovary syndrome. *Fertil Steril*. 2000;74(5):976–9.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.