



Short communication

Benzenesulfonamides incorporating nitrogenous bases show effective inhibition of β -carbonic anhydrases from the pathogenic fungi *Cryptococcus neoformans*, *Candida glabrata* and *Malassezia globosa*

Silvia Bua^a, Sameh M. Osman^b, Zeid AlOthman^b, Claudiu T. Supuran^{a,b,*}, Alessio Nocentini^{a,*}

^a University of Florence, NEUROFARBA Department, Via Ugo Schiff 6, 50019 Sesto Fiorentino, Italy

^b Department of Chemistry, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Keywords:

Carbonic anhydrase
Inhibitor
Metalloenzymes
Adenine
Uracil
Antifungal

ABSTRACT

There is an urgent need for new chemotherapeutic agents to treat human fungal infections due to emerging and spreading globally resistance mechanisms. Among the new targets that have been recently investigated for the development of antifungal drugs there are the metallo-enzymes Carbonic Anhydrases (CAs, EC 4.2.1.1). The inhibition of the β -CAs identified in many pathogenic fungi leads to an impairment of parasite growth and virulence, which in turn leads to a significant anti-infective effect. Based on antifungal nucleoside antibiotics, the inhibition of the β -CAs from the resistance-showing fungi *Candida glabrata* (CgNce103), *Cryptococcus neoformans* (Can2) and *Malasszia globosa* (MgCA) with a series of benzenesulfonamides bearing nitrogenous bases, such as uracil and adenine, is here reported. Many such compounds display low nanomolar (< 100 nM) inhibitory potency against Can2 and CgNce103, whereas the activity of MgCA is considerably less affected (inhibition constants in the range 138.8–5601.5 nM). The β -CAs inhibitory data were compared with those against α -class human ubiquitous isoforms. Interesting selective inhibitory activities for the target fungal CAs over hCA I and II were reported, which make nitrogenous base benzenesulfonamides interesting tools and leads for further investigations in search of new antifungal with innovative mechanisms of action.

1. Introduction

Anti-microbial resistance is one of the biggest threats to global health and a major concern in clinical practice because it impairs the efficacy of common therapies for infectious diseases, among which fungal infections [1–4]. Invasive fungal infections can cause disability and death. Some species of fungi are naturally resistant to treatment with some antifungal medications. Other species can develop resistance due to improper antifungal use, such as too low dosages or treatment courses that are not long enough [5]. Antifungal resistance is a noteworthy problem with *Candida* infections. Some types of *Candida* are increasingly resistant to the first-line and second-line antifungal medications, such as fluconazole and the echinocandins [6]. Treatment failures against *Cryptococcus neoformans* also continue to occur for a variety of reasons including direct antifungal drug resistance. Cryptococcosis has become a relatively common infection worldwide mainly affecting the lungs, but also responsible for meningitis and encephalitis [7]. *Malassezia globosa* is considered as the most probable causative agent for the onset of dandruff. Many drug-resistant fungal strains of *M.*

globosa to azole antifungals have been reported and impair the efficacy of common anti-dandruff cosmetic treatments [8].

A main strategy to overtake the spread of antimicrobial resistance consists in the identification of novel therapeutic targets and thus anti-infectives with alternative mechanisms of action. The metalloenzymes carbonic anhydrases (CAs, EC 4.2.1.1) are essential in the life cycle (pH homeostasis and biosynthetic reactions) as well as in the virulence of many bacterial, fungal and protozoan pathogens [9]. Many data exist in the literature, which show that interference with CA activity in various parasites leads to an impairment of parasite growth and virulence [9]. CA enzymes are classified into seven structurally unrelated groups, α -, β -, γ -, δ -, ζ - η - and θ -CAs [10–13]. In particular, β -CAs isoforms were identified in many pathogenic fungi, among which *C. glabrata* (CgNce103), *C. neoformans* (Can2) and *M. globosa* (MgCA) [9–11]. Several works demonstrated that these CAs play an important role in the CO₂ sensing of the pathogens and in the regulation of sexual development [9,14–16]. For instance, physiological concentrations of CO₂/HCO₃⁻ induce prominent virulence attributes in *C. glabrata* or *C. neoformans* through direct activation of the fungal adenylyl cyclase

* Corresponding authors at: University of Florence, NEUROFARBA Department, Via Ugo Schiff 6, 50019 Sesto Fiorentino, Italy (C.T. Supuran).

E-mail addresses: claudiu.supuran@unifi.it (C.T. Supuran), alessio.nocentini@unifi.it (A. Nocentini).

<https://doi.org/10.1016/j.bioorg.2019.01.030>

Received 1 October 2018; Received in revised form 12 December 2018; Accepted 17 January 2019

Available online 19 January 2019

0045-2068/ © 2019 Elsevier Inc. All rights reserved.

[14,15].

The inhibitory profile against β -CAs of most important classes of CA inhibitors, such as sulfonamides and bioisosteres, inorganic anions, and carboxylate, have been investigated in detail [16–18]. Recently, new chemotypes, such as phenols, dithiocarbamates, and benzoxaboroles were also reported as efficient fungal β -CA inhibitors [19–25]. MgCA inhibition by sulfonamides led to growth defects of the fungus *in vivo*, exhibiting antidandruff effects equivalent to those of the standardazole drug ketoconazole [16].

Nucleoside antibiotics represent a unique class of antimicrobial products derived from nucleosides and nucleotides [26]. They display diverse biological activities including antibacterial, antifungal, antiviral, insecticidal, immunostimulative, immunosuppressive, and anti-tumor activities.

For all these reasons, we investigate here nitrogenous bases-bearing benzenesulfonamides as inhibitors of CgNce103, Can2 and MgCA. The compounds were designed according to the molecular hybridization approach to strengthen the anti-fungal effect of the pathogen CA inhibition with the anti-microbial activity of nitrogenous bases, such as uracil and adenine, derivatives.

2. Chemistry


A benzenesulfonamide scaffold was variably appended with uracil and adenine moieties by means of different length spacer of the ether, amide and its bioisoster triazole types, as previously reported by us [27]. The bioisosteric amide/triazole substitution was pursued due to the anti-fungal activity that triazoles and their derivatives were shown to possess [28]. Uracil and adenine were linked to the main scaffold through the N1 and N9 moieties, respectively, to keep the connection by which nitrogenous bases are incorporated in nucleotides and nucleic acids.

The triazole derivatives (11–12, 15–16) were prepared by copper-catalyzed azide-alkyne cycloadditions (CuAAC), using N1-propargyluracil (8) or the N9-propargyladenine (14) (Schemes 1 and 2) [27].

The amide derivatives (3–7, 22–29) were prepared according to two diverse synthetic pathways depending on the considered nitrogenous base (Schemes 1 and 2) [27].

3. Carbonic anhydrase inhibition

The CA inhibitory activities of compounds 3–7, 11–12, 15–16,



Scheme 1. General synthetic procedure for uracil derivatives 2–12.

22–29, 32, in addition to acetazolamide (AAZ) as standard inhibitor, were measured against MgCA, Can2, and CgNce103 by a stopped flow CO₂ hydrase assay [29]. The displayed inhibitory activities were compared to those against the off-target widely-distributed hCA I and hCA II. The following structure–activity relationship (SAR) were obtained from the data reported in Table 1.

(i) The inhibitory capability of derivatives 3–7, 11–12, 15–16, 22–29, 32 strongly depends on the length and kind of the spacer and the type of nitrogenous base for both the three fungal isozymes and human isoforms. Most of the adenine- and uracil-bearing compounds reported a moderate inhibitory potency against the fungal isoform MgCA, with K_is ranging between 138.8 and 5012.8 nM. Adenine compounds 32, 25 and 29 reported the best inhibitory activity with K_is value of 138.8, 153.9 and 190.2 nM, respectively. These compounds share the longest linkers between the inhibitory scaffold and the amide portion or the adenine core. The free NH₂ confers to 29 better inhibition activity than the protected-amine 25. The length of the linker between main scaffold and nitrogenous base has minor influence on the inhibition in case of uracil compounds. For instance, the uracil derivatives 4 and 6 report very similar K_is (391.2 and 349.3 nM, respectively), although 4 lacks the ethyl spacer present in compound 6. The minor steric hindrance represented by uracil compared to adenine likely justifies this effect.

A remarkable fall of efficacy is observed by switching any substituent on the benzenesulfonamide from the *para* to the *meta* position. Compound 3 is 3.2 time less potent (K_i value 1238.6 nM) than its analog 4. Similar trends are evident from data in Table 1 for the remaining meta-derivatives 7, 11, 15, 22 and 26 which report the worst inhibitory activities, with K_is ranging between 981.8 and 5012.8 nM.

(ii) Can2 is the most inhibited isoform by the sulfonamides herein reported. Compounds 4, 5, 6, 12, 25, 27, 29 and 32 report K_i values even below 100 nM (9.1–87.1 nM). In this case the length of the linker has a major influence on the inhibitory efficacy compared to the type of nitrogenous base and elicits analog trends for both uracil and adenine compounds. In detail, a methylene linker between the amide and benzenesulfonamide ring (5, 24, 28) ascribes to the derivative a worse Can2 inhibition efficacy than the absence of a linker (4, 23, 27), that is in turn less effective than a two-carbon atoms spacer (6, 25, 29).




Table 1
Inhibition data of hCA I, hCA II, and β -CA isoforms MgCA, Can2, and CgNce103, with sulfonamides 3–7, 11–12, 15–16, 22–29, and 32 and the standard inhibitor acetazolamide (AAZ) by a Stopped Flow CO₂ hydrase assay.

Cmpd	K _i ^a (nM)				
	MgCA	Can2	CgNce103	hCA I	hCA II
3	1238.6	309.5	383.1	7966.7	704.3
4	391.2	53.2	83.1	645.5	17.7
5	620.3	87.1	146.6	1565.7	495.4
6	349.3	19.3	46.2	658.9	42.1
7	981.8	182.7	304.4	9840.2	4357.7
11	5012.8	750.7	1010.2	4871	388.1
12	718.9	50.4	149.6	256.2	0.85
15	5601.5	1146.8	1126.4	> 10000	655.3
16	578.4	124.1	124.6	410.9	5.6
22	1861.3	843.9	950.1	> 10000	756.2
23	617.7	143.2	75.3	966.3	30.7
24	862.2	108.6	180.6	4721.4	87
25	153.9	16.8	31.9	1962.6	57.8
26	1474.6	660.2	460.5	> 10000	889.2
27	699.3	49.8	94.6	666.6	8.4
28	908.7	117.5	174.7	977.1	59.2
29	190.2	9.1	56.9	811.8	8.2
32	138.8	27.5	87.3	552.6	78
AAZ	74,000	10	11	250	12

* Mean from 3 different assays, by a stopped flow technique (errors were in the range of \pm 5–10% of the reported values).

(iii) The isoform CgNce103 is intermediately inhibited by the reported sulfonamides with K_is in the range 31.9–1126.4 nM. The general tendency described for Can2 can be applied for this isoform too. The meta-substituted derivatives are again the worst inhibitors

with K_i values in the range 304.4–1126.4 nM. Uracil 6 (K_i of 46.2 nM) and adenine 25 (K_i of 31.90 nM), both appending the ethylene linker, showed the best CgNce103 inhibitory potencies herein reported. Conversely, the introduction of methylene spacer between main scaffold and connection reduces the inhibitory potency of 5 (K_i = 146.6 nM), 24 (K_i = 180.6 nM) and 28 (K_i = 174.7 nM) in comparison to both aliphatic linker-devoid and ethylene bearing derivatives.

Likewise, against MgCA, meta-substituted compounds (3, 7, 11, 15, 22, 26) reported the worst KI compared to para-substituted ones. The bioisosteric substitution of the amide linker with a bulkier triazole one lowers the inhibitory efficacy of both meta- (11 and 15) and para-substituted (12 and 16, K_is of 50.4 and 124.1 nM) derivatives compared to analog amides devoid of the alkyl spacer. As a result, uracil 6 and adenine 25 and 29 stand out as the best Can2 inhibitors herein reported (K_is of 19.3, 16.8, 9.1 nM, respectively). The introduction of p-OH group on the benzenesulfonamide ring positively impacts the inhibitory activity of the compound 7 (K_i value of 182.7 nM) with respect to derivative 3 (K_i of 309.5 nM).

(iv) The data in Table 1 outline interesting target/off-target CAs selectivity profiles. First, MgCA is inhibited in a comparable range with the ubiquitous isoform hCA I, but rather less than the other human isoform hCA II. Conversely, all sulfonamides more effectively inhibit the β -CAs Can2 and CgNce103 than hCA I, and some such compounds even than hCA II, though to a minor extent. In detail, the *m*-substituted uracil compounds (amides) 3 and 7 show selectivity index (SI) of two to twenty for the target Can2/CgNce103 over hCA II. Compounds 5 and 6 inhibit Can2 two- to five-fold better than the human isozyme. Adenines 25 and 32 show

a Can2/hCA II SI of 3. All triazole derivatives more efficiently inhibit hCA II than all β -CAs herein evaluated.

4. Conclusions

A main strategy to overtake the spread of antifungal resistance consists in the identification of novel therapeutic targets and anti-infectives with alternative mechanisms of action. Carbonic anhydrases were shown to be essential in the life cycle (pH homeostasis and biosynthetic reactions) as well as in the virulence of many bacterial, fungal and protozoan pathogens. In particular, the inhibition of the β -CAs identified in many pathogenic fungi leads to an impairment of parasite growth and virulence, which in turn leads to a significant anti-infective effect. Based on antifungal antibiotics incorporating nucleosides, we reported here the inhibition study of β -CAs from the resistance-showing fungi *Candida glabrata* (CgNce103), *Cryptococcus neoformans* (Can2) and *Malassezia globosa* (MgCA) with a series of benzenesulfonamides bearing nitrogenous bases. Many such compounds display excellent low nanomolar inhibitory potency against Can2 and CgNce103, whereas the activity of MgCA is considerably less affected (inhibition constants in the range 138.8–5601.5 nM). The inhibition profiles were shown to depend on the length and positioning of the spacer which connects the two pharmacophores. The β -CAs inhibitory data were compared with those against the off-target hCA I and II. Interesting target/off-target selective efficacies were reported, which make nitrogenous base benzenesulfonamides interesting leads for further studies in search of new antifungal with innovative mechanisms of action.

5. Experimental sections

5.1. Chemistry

The synthesis of sulfonamides **3–7**, **11–12**, **15–16**, **22–29**, and **32** was reported earlier by our group [27].

5.2. Carbonic anhydrase inhibition

An Applied Photophysics stopped-flow instrument has been used for assaying the CA-catalysed CO₂ hydration activity [29]. Phenol red (at a concentration of 0.2 mM) has been used as indicator, working at the absorbance maximum of 557 nm, with 20 mM Hepes (pH 7.5) as buffer, and 20 mM Na₂SO₄ (for maintaining constant the ionic strength), following the initial rates of the CA-catalysed CO₂ hydration reaction for a period of 10–100 s. The CO₂ concentrations ranged from 1.7 to 17 mM for the determination of the kinetic parameters and inhibition constants. For each inhibitor, at least six traces of the initial 5–10% of the reaction have been used for determining the initial velocity. The uncatalyzed rates were determined in the same manner and subtracted from the total observed rates. Stock solutions of inhibitor (0.1 mM) were prepared in distilled-deionised water and dilutions up to 0.01 nM were done thereafter with the assay buffer. Inhibitor and enzyme solutions were preincubated together for 15 min at room temperature prior to assay, in order to allow for the formation of the E–I complex. The inhibition constants were obtained by nonlinear least-squares methods using PRISM 3 and the Cheng-Prusoff equation, as reported earlier, and represent the mean from at least three different determinations [30–33]. All CA isoforms were recombinant ones obtained in-house as reported earlier [34–36].

Acknowledgments

This work was financed in part by a DSFP project to Profs. Z. AlOthman and CT Supuran.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bioorg.2019.01.030>.

References

- <https://www.cdc.gov/fungal/antifungal-resistance.html>.
- M.A. Ghannoum, L.B. Rice, Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance, *Clin. Microbiol. Rev.* 12 (1999) 501–517.
- E. Koksul, H. Tohma, O. Kilic, Y. Alan, A. Aras, I. Gulcin, E. Bursal, Assessment of antimicrobial and antioxidant activities of *Nepeta trachonitica*-Analysis of its phenolic compounds using HPLC-MS/MS, *Scientia Pharmaceutica* 15 (2017) 85.
- U.M. Kocyyigit, Y. Budak, M.B. Gurdere, F. Erturk, B. Yencilek, P. Taslimi, I. Gulcin, M. Ceylan, Synthesis of chalcone-imide derivatives and investigation of their anticancer and antimicrobial activities, carbonic anhydrase and acetylcholinesterase enzymes inhibition profiles, *Arch. Physiol. Biochem.* 124 (2018) 61–68.
- N.P. Wiederhold, Antifungal resistance: current trends and future strategies to combat, *Infect. Drug. Resist.* 10 (2017) 249–259.
- S.G. Whaley, E.L. Berkow, J.M. Rybak, A.T. Nishimoto, K.S. Barker, P.D. Rogers, Azole antifungal resistance in *Candida albicans* and emerging non-*albicans* *Candida* species, *Front Microbiol.* 7 (2016) 2173.
- J.R. Perfect, G.M. Cox, Drug resistance in *Cryptococcus neoformans*, *Drug. Resist. Updat.* 4 (1999) 259–269.
- F.D. Rojas, L. Sosa Mde, M.S. Fernández, M.E. Cattana, S.B. Córdoba, G.E. Giusiano, Antifungal susceptibility of *Malassezia furfur*, *Malassezia sympodialis*, and *Malassezia globosa* to azole drugs and amphotericin B evaluated using a broth microdilution method, *Med. Mycol.* 52 (2014) 641–646.
- C. Capasso, C.T. Supuran, Bacterial, fungal and protozoan carbonic anhydrases as drug targets, *Expert Opin. Ther. Targets.* 19 (2015) 1689–1704.
- C.T. Supuran, Carbonic anhydrases: novel therapeutic applications for inhibitors and activators, *Nat. Rev. Drug Discov.* 7 (2008) 168–181.
- M. Boztas, Y. Cetinkaya, M. Topal, I. Gulcin, A. Menzek, E. Sahin, M. Tanc, C.T. Supuran, Synthesis and carbonic anhydrase isoenzymes I, II, IX, and XII inhibitory effects of dimethoxy-bromophenol derivatives incorporating cyclopropane moieties, *J. Med. Chem.* 58 (2015) 640–650.
- A. Yildirim, U. Atmaca, A. Keskin, M. Topal, M. Celik, I. Gulcin, C.T. Supuran, N-Acylsulfonamides strongly inhibit human carbonic anhydrase isoenzymes I and II, *Bioorg. Med. Chem.* 23 (2015) 2598–2605.
- M. Huseynova, T. Ajdar, M. Vagif, F. Mahizar, A. Gulnar, G. Onur, S. Bahattin, Y. Afsun, S. Baturhan, O. Ali, R. Ozkaya, I. Gulcin, Synthesis, characterization, crystal structure, electrochemical studies and biological evaluation of metal complexes with thiosemicarbazone of glyoxylic acid, *Polyhedron* 155 (2018) 25–33.
- F. Cottier, W. Leewattanasakul, L.R. Kemp, M. Murphy, C.T. Supuran, O. Kurzai, F.A. Mühlshlegel, Carbonic anhydrase regulation and CO₂ sensing in the fungal pathogen *Candida glabrata* involves a novel Rca1p ortholog, *Bioorg. Med. Chem.* 21 (2013) 1549–1554.
- C. Schlicker, R.A. Hall, D. Vullo, S. Middelhaufe, M. Gertz, C.T. Supuran, F.A. Mühlshlegel, C. Steegborn, Structure and Inhibition of the CO₂-sensing carbonic anhydrase Can2 from the pathogenic fungus *Cryptococcus neoformans*, *J. Mol. Biol.* 385 (2009) 1207–1220.
- K.S. Hewitson, D. Vullo, A. Scozzafava, A. Mastrolorenzo, C.T. Supuran, Molecular cloning, characterization, and inhibition studies of a beta-carbonic anhydrase from *Malassezia globosa*, a potential antidandruff target, *J. Med. Chem.* 55 (2012) 3513–3520.
- S. Del Prete, D. Vullo, S.M. Osman, Z. AlOthman, C. Capasso, C.T. Supuran, Anion inhibition studies of the dandruff-producing fungus *Malassezia globosa* β -carbonic anhydrase MgCA, *Bioorg. Med. Chem. Lett.* 25 (2015) 5194–5198.
- C.T. Supuran, Advances in structure-based drug discovery of carbonic anhydrase inhibitors, *Expert Opin. Drug Discov.* 12 (2017) 61–88.
- D. Vullo, S. Del Prete, A. Nocentini, S.M. Osman, Z. AlOthman, C. Capasso, M. Bozdog, F. Carta, P. Gratteri, C.T. Supuran, Dithiocarbamates effectively inhibit the β -carbonic anhydrase from the dandruff-producing fungus *Malassezia globosa*, *Bioorg. Med. Chem.* 25 (2017) 1260–1265.
- A. Nocentini, R. Cadoni, P. Dumy, C.T. Supuran, J.Y. Winum, Carbonic anhydrases from *Trypanosoma cruzi* and *Leishmania donovani* chagasi are inhibited by benzoxaboroles, *J. Enzyme Inhib. Med. Chem.* 33 (2018) 286–289.
- Y. Entezari Heravi, S. Bua, A. Nocentini, S. Del Prete, A.A. Saboury, H. Sereshti, C. Capasso, P. Gratteri, C.T. Supuran, Inhibition of *Malassezia globosa* carbonic anhydrase with phenols, *Bioorg. Med. Chem.* 25 (2017) 2577–2582.
- A. Nocentini, D. Vullo, S. Del Prete, S.M. Osman, F.A.S. Alasmary, Z. AlOthman, C. Capasso, F. Carta, P. Gratteri, C.T. Supuran, Inhibition of the β -carbonic anhydrase from the dandruff-producing fungus *Malassezia globosa* with monothiocarbamates, *J. Enzyme Inhib. Med. Chem.* 32 (2017) 1064–1070.
- A. Nocentini, R. Cadoni, S. Del Prete, C. Capasso, P. Dumy, P. Gratteri, C.T. Supuran, J.Y. Winum, Benzoxaboroles as efficient inhibitors of the β -carbonic anhydrases from pathogenic fungi: activity and modeling study, *ACS Med. Chem. Lett.* 8 (2017) 1194–1198.
- A. Nocentini, S. Bua, S. Del Prete, Y.E. Heravi, A.A. Saboury, A. Karioti, A.R. Bilia, C. Capasso, P. Gratteri, C.T. Supuran, Natural polyphenols selectively inhibit β -carbonic anhydrase from the dandruff-producing fungus *Malassezia globosa*: activity and modeling studies, *ChemMedChem.* 13 (2018) 816–823.
- M.A. Mohamed, A.A. Abdel-Aziz, H.M. Sakr, A.S. El-Azab, S. Bua, C.T. Supuran,

- Synthesis and human/bacterial carbonic anhydrase inhibition with a series of sulfonamides incorporating phthalimido moieties, *Bioorg. Med. Chem.* 25 (2017) 2524–2529.
- [26] G. Niu, J. Zheng, H. Tan, Biosynthesis and combinatorial biosynthesis of antifungal nucleoside antibiotics, *Sci. China Life Sci.* 60 (2017) 939–947.
- [27] A. Nocentini, S. Bua, C.L. Lomelino, R. McKenna, M. Menicatti, G. Bartolucci, B. Tenci, Mannelli L Di Cesare, C. Ghelardini, P. Gratteri, C.T. Supuran, Discovery of new sulfonamide carbonic anhydrase IX inhibitors incorporating nitrogenous bases, *ACS Med. Chem. Lett.* 8 (2017) 1314–1319.
- [28] L.R. Peyton, S. Gallagher, M. Hashemzadeh, Triazole antifungals: a review, *Drugs Today (Barc)*. 51 (2015) 705–718.
- [29] R.G. Khalifah, The carbon dioxide hydration activity of carbonic anhydrase, *J. Biol. Chem.* 246 (1971) 2561–2573.
- [30] S. Bua, E. Berrino, S. Del Prete, V.S. Murthy, V. Vijayakumar, Y. Tamboli, C. Capasso, E. Cerbai, A. Mugelli, F. Carta, C.T. Supuran, Synthesis of novel benzenesulfamide derivatives with inhibitory activity against human cytosolic carbonic anhydrase I and II and *Vibrio cholerae* α - and β -class enzymes, *J. Enzyme. Inhib. Med. Chem.* 33 (2018) 1125–1136.
- [31] M. Ferraroni, R. Gaspari, A. Scozzafava, A. Cavalli, C.T. Supuran, Dioxygen, an unexpected carbonic anhydrase ligand, *J. Enzyme. Inhib. Med. Chem.* 33 (2018) 999–1005.
- [32] J. Leitans, A. Kazaks, A. Balode, J. Ivanova, R. Zalubovskis, C.T. Supuran, K. Tars, Efficient expression and crystallization system of cancer-associated carbonic anhydrase isoform IX, *J. Med. Chem.* 58 (2015) 9004–9009.
- [33] A. Nocentini, C.T. Supuran, Carbonic anhydrase inhibitors as antitumor/antimetastatic agents: a patent review (2008–2018), *Expert. Opin. Ther. Pat.* (2018), <https://doi.org/10.1080/13543776.2018.1508453>.
- [34] A. Nocentini, M. Ceruso, S. Bua, C.L. Lomelino, J.T. Andring, R. McKenna, C. Lanzi, S. Sgambellone, R. Pecori, R. Matucci, L. Filippi, P. Gratteri, F. Carta, E. Masini, S. Selleri, C.T. Supuran, Discovery of β -adrenergic receptors blocker-carbonic anhydrase inhibitor hybrids for multitargeted antiglaucoma therapy, *J. Med. Chem.* 61 (2018) 5380–5394.
- [35] R. Kumar, S. Bua, S. Ram, S. Del Prete, C. Capasso, C.T. Supuran, Sharma PK Benzenesulfonamide bearing imidazothiadiazole and thiazolotriazole scaffolds as potent tumor associated human carbonic anhydrase IX and XII inhibitors, *Bioorg. Med. Chem.* 25 (2017) 1286–1293.
- [36] S. Bua, M. Bozdog, S. Del Prete, F. Carta, W.A. Donald, C. Capasso, C.T. Supuran, Mono- and di-thiocarbamate inhibition studies of the δ -carbonic anhydrase TweCA δ from the marine diatom *Thalassiosira weissflogii*, *J. Enzyme. Inhib. Med. Chem.* 33 (2018) 707–713.