



Substituted cinnamic anhydrides act as selective inhibitors of acetylcholinesterase

Josephine M. Gießel, Immo Serbian, Anne Loesche, René Csuk*

Martin-Luther-University Halle-Wittenberg, Organic Chemistry, Kurt-Mothes Str. 2, D-06120 Halle (Saale), Germany

ARTICLE INFO

Keywords:

Cinnamic anhydrides
Acetylcholinesterase
Butyrylcholinesterase
Inhibitors

ABSTRACT

Cinnamic anhydrides have been shown to be more than reactive reagents, but they also act as inhibitors of the enzyme acetylcholinesterase (AChE). Thus, out of a set of 33 synthesised derivatives, several of them were mixed type inhibitors for AChE (from electric eel). Thus, (*E*)-3-(2,4-dimethoxyphenyl)acrylic anhydride (**2c**) showed $K_i = 8.30 \pm 0.94 \mu\text{M}$ and $K_i' = 9.54 \pm 0.38 \mu\text{M}$, and for (*E*)-3-(3-chlorophenyl)acrylic anhydride (**2u**) $K_i = 8.23 \pm 0.93 \mu\text{M}$ and $K_i' = 13.07 \pm 0.46 \mu\text{M}$ were measured. While being not cytotoxic to many human cell lines, these compounds showed an unprecedented and noteworthy inhibitory effect for AChE but not for butyrylcholinesterase (BChE).

1. Introduction

Cinnamic acid moieties are part of a large number of natural products. Since many of them can be found in our daily nutrition [1–3] an investigation of their biological effects is of major interest. While some of these compounds have been investigated in detail [4–7], many derivatives have not been a subject of biological investigations at all. The effects of cholinesterase inhibitors in combating Alzheimer's disease (AD) remain interesting but still not completely understood. Acetyl- and butyrylcholinesterase are enzymes which influence the cognitive function and memory, thus making these enzymes of special interest in humans being affected by AD [8–10]. Approximately one percent of the world's population suffers from some kind of dementia; as a consequence, 50 million people are affected by AD [11]. Recent work from Kim et al. [12] showed derivatives of cinnamic acid to hold promising cholinesterase (ChE) inhibitory effects. Hence, we were looking for unexplored derivatives close to cinnamic acids, and interestingly cinnamic anhydrides revealed to be a class of compounds having never before tested for their ability to act as inhibitors of ChE (Fig. 1). Interestingly, quite recently another anhydride, 2,3-dimethylmaleic anhydride has been found to act as a strong inhibitor on an acetylcholinesterase from insects [13].

Usually, carboxylic anhydrides are seen as reactive species to synthesize carboxylic acids derivatives that cannot be obtained directly from the carboxylic acid [14,15]. Furthermore, due to this reactivity, it is generally assumed that they are easily hydrolysed, hence resulting in a short lifetime in the presence of water. This holds true for many

anhydrides, however cinnamic anhydrides are remarkable stable in aqueous solutions [16], and they should be investigated in more detail. As a consequence, anhydrides did not seem to be very appealing as potential drugs, and investigations of them are scarcely found in literature. Thus, a mixed anhydride between a triterpene and cinnamic acid [17] has been investigated as novel anti-malarial agent, and cytotoxic properties have been reported for plant extracts [18,19].

2. Results and discussion

2.1. Chemistry

Knoevenagel-Doebner-condensations [20,21] are well-known to access carboxylic acids from malonic acid and an appropriate aldehyde. To investigate the influence of the electronic profile of the aromatic system on reactivity and biological activity, a variety of differently substituted benzaldehydes holding both electron-withdrawing and -donating substituents in various substitution patterns has been included in our investigations (Table 1).

Substituted benzaldehydes were used to prepare 13 commercially not available cinnamic acids applying Knoevenagel-Doebner reactions. These compounds are characterized by the presence of the olefinic protons in their ^1H NMR spectra (as exemplified for **1a**: $\delta = 7.74$ and 6.33 ppm); in the ^{13}C NMR spectra the olefinic carbons were detected at $\delta = 147.1$ and 115.0 ppm, respectively. The yields in these condensation reactions differed between moderate to very good, a fact that can be explained by the different electronic properties due to the differences

* Corresponding author.

E-mail address: rene.csuk@chemie.uni-halle.de (R. Csuk).

<https://doi.org/10.1016/j.bioorg.2019.103058>

Received 11 March 2019; Received in revised form 1 June 2019; Accepted 7 June 2019

Available online 08 June 2019

0045-2068/ © 2019 Elsevier Inc. All rights reserved.




Fig. 1. Structure of cinnamic anhydrides and of 2,3-dimethylmaleic anhydride.

Table 1

Substitution pattern for cinnamic acids (1) and anhydrides (2).

	R ¹	R ²	R ³	R ⁴
a	H	OMe	OMe	H
b	H	OMe	OMe	H
c	OMe	H	OMe	H
d	H	OMe	H	OMe
e	H	OMe	OMe	OMe
f	OH	OMe	H	H
g	H	OH	OMe	H
h	H	OMe	OH	H
i	H	OMe	OAc	H
j	H	Me	Me	H
k	NO ₂	H	H	H
l	H	NO ₂	H	H
m	H	H	N(Me) ₂	H
n	H	H	H	H
o	H	OH	OH	H
p	H	OMe	OH	OMe
q	OMe	H	H	H
r	H	OMe	H	H
s	H	H	OMe	H
t	Cl	H	H	H
u	H	Cl	H	H
v	Cl/H	H/Cl	H	H
w	H	F	H	H
x	H	H	F	H

in the substitution pattern.

Symmetrical and non-symmetrical anhydrides were prepared according to Scheme 1. For the synthesis, a combination of cinnamic acids from previous reactions and commercially available compounds was used. Again, reaction yields were moderate to very good, with compounds 1f–1h being the exceptions, reacting to polymeric structures and not forming anhydrides.

2.2. Biological evaluation

2.2.1. Inhibition of ChE

The inhibitory activity of the compounds for AChE and BChE was determined by Ellman's assays [22]. To support these experiments, some extra molecular docking calculations with the enzyme AChE (from electric eel) were performed using Autodock4 [23]. The results from the biological assays and calculations from the docking of cinnamic acid derivatives are compiled in Table 2. Interestingly, while some compounds proved to be good inhibitors for AChE, none of them was a good inhibitor for BChE.

The results from the biological assays are in excellent agreement with the results from the molecular modelling calculations. Thus, the cinnamic acid finds its way inside the enzyme pocket of AChE (Fig. 2), but shows only show weak interactions with the enzyme and cannot

Table 2

Results from the biological screening (Ellman's assay; %-inhibition at a concentration of 10 μ M) of compounds 1a–1m (compounds not mentioned in the table were insoluble under the conditions of the assay) with AChE (from *electricus*) and BChE (from *equine serum*); the inhibition [22] constants K_i and K_i' are given in μ M and galantamine hydrobromide (GH) was used as a standard; the values are averaged from experiments performed in triplicate, and binding energies are reported in kcal/mol for AChE (PDB:1C2O).


AChE			BChE
Compounds	Inhibition [%]	Binding energy [kcal/mol]	Inhibition [%]
GH	87.70	−7.94	53.00
1a	33.70	−5.30	12.3
1b	7.22	−5.24	1.91
1c	25.91	−5.88	18.34
1d	12.65	−5.67	0.00
1e	9.75	−4.84	2.72
1g	6.89	−5.22	1.58
1h	12.32	−5.29	2.09
1i	12.09	−5.67	3.08
1j	11.85	−6.29	0.79
1k	8.52	−6.33	2.11
1l	10.60	−6.35	0.73
1m	13.61	−5.87	1.28

bind to the active site of the enzyme. In Fig. 2, the active site is located at the lower right with the corresponding amino acids His447, Glu334 and Ser203.

As depicted in Fig. 2, the anhydrides were also docked into the enzyme pocket of AChE. Although these compounds are larger, it is obvious that they fit even better than the parent acids into the pocket thus blocking the active site. The results from the biological assays showed cinnamic anhydrides 2a–2x (Table 3) to be able of binding to the enzyme, and revealed some of them as good inhibitors. These anhydrides were only weak inhibitors for butyrylcholinesterase. Molecular modelling calculations (PDB:6EMI) revealed that these compounds have less interaction with the amino acid residues of the active site (Fig. 3).

The assays were carried out in aqueous solution, and thus it might be a valid assumption that the cinnamic anhydrides will be hydrolysed in this environment. Furthermore, one would assume that the products of hydrolysis induce the inhibitory effect. However, cinnamic acid anhydrides are rather stable under the conditions of the assays, and - as checked by TLC - even after a prolonged period of 72 h less than 10 percent of the anhydride was hydrolyzed into their corresponding acids. Therefore, the anhydride but not the cinnamic acid acts as inhibitor for AChE. The assumption that cinnamic anhydrides act as irreversible inhibitors of the enzyme could also not be confirmed since measurements of the enzyme kinetic showed these compounds as mixed type inhibitors for AChE. Thus, in this series, compounds 2c, 2d, 2j, 2t, 2u and 2v showed good to moderate inhibition of AChE. These biological activities are not entirely unexpected because these cinnamic anhydrides can be considered as isosteric derivatives of curcumin. Of the latter, inhibitory activity is known with respect to AChE [24–28].

Extra screening of cinnamic acids 1a–1m and of anhydrides 2a–2x showed them being not cytotoxic for a variety of human tumor cells lines [cut off: EC₅₀ > 30 μ M; cell lines: FaDu (hypopharyngeal



Scheme 1. . Synthesis of cinnamic acid derivatives and the appropriate anhydrides: (i): piperidine, pyridine, reflux 120 °C, 4 h; (ii): (1) oxalyl chloride, CH₂Cl₂, 0 °C → 25 °C, 2 h (2) pyridine, toluene, 25 °C, overnight.




Fig. 2. Compounds 1a-1m (left) and 2a-2x (right) inside the AChE active site pocket.

carcinoma), A2780 (ovarian carcinoma), HT29 (colorectal carcinoma), MCF7 (breast carcinoma), SW1736 (thyroid carcinoma), A375 (malignant melanoma), A549 (epithel carcinoma)] and to non-malignant NIH 3T3 (mouse fibroblasts) cells. This finding is a prerequisite for developing anhydrides of improved properties.


3. Conclusion

As a result, cinnamic anhydrides have been shown to represent a class of interesting and promising compounds with unforeseen properties. In this study, 33 derivatives were prepared and screened for their biological activity. While these compounds exhibited no cytotoxic effect neither on malignant human tumor cells nor on non-malignant mouse fibroblasts, they proved to be moderate to good inhibitors of the enzyme acetylcholinesterase while being only negligible inhibitors for butyrylcholinesterase. As the cinnamic anhydrides are of remarkable stability in aqueous solution, derivatives of this kind might be of interest to develop AChE inhibitors.

4. Experimental

4.1. General

The reagents were bought from commercial suppliers and used without further purification. The solvents were dried according to usual procedures. Melting points were determined on Büchi Melting Point M-565 or LEICA hot stage microscope and are uncorrected, NMR spectra were recorded on a Varian spectrometer Unity 500 (δ given in ppm, J in



Macherey-Nagel ALUGRAM® Xtra SIL G/UV₂₅₄ pre-coated silica gel 60 F254 plates were used for thin layer chromatography (detection with cerium molybdate spray reagent and UV absorption). IR spectra were recorded on a Perkin-Elmer FT-IR spectrometer Spectrum 1000 and wave numbers are expressed in cm^{-1} . The absorption spectra were measured on Perkin Elmer Lambda14 spectrometer. Microanalyses were performed with a Foss-Heraeus Vario EL (CHNS) instrument.

4.2. Molecular docking

The crystal structures of the *ee*AcChE (PDB = 1C2O) *h*BChE-7 (PDB = 6EMI) were retrieved from the protein databank ([rcsb.org](https://www.rcsb.org)). The enzyme was prepared according to usual procedures. Polar hydrogen atoms were added, water molecules removed, and Gasteiger charges were added. The ligand minimisation and preparation was performed with the MMFF94 force field in Datawarrior. Openbabel was used to create the pdbqt files for Autodock. Calculations were done with Autodock4 [23] Lamarckian genetic algorithm, and the analysis of the docking poses was done with MGLTools 1.5.6.; figures were created with PyMOL.

4.3. Biological screening

The Ellman's assay have been performed as previously described [29,30].

4.4. Syntheses

4.4.1. General procedure A

To a solution of the substituted benzaldehyde (1 eq.) and malonate (1.2 eq.) in pyridine (2.5 mL/mmol) at 120 °C piperidine (0.04 eq.) was added, and the reaction mixture was heated under reflux for 4 h [31]. The solvents were removed *in vacuo* as an azeotrope with toluene (3 × 70 mL). Water (100 mL) was added, and the aqueous phase was extracted with ethyl acetate (3 × 100 mL), brine (1 × 100 mL), dried (MgSO_4) and concentrated. Compounds **1a** and **1c–1m** were prepared following procedure A.

4.4.2. (2E)-3-(3,4-Dimethoxyphenyl)-2-propenoic acid (**1a**)

Recrystallization from ethanol gave **1a** (8.28 g, 80%) as an off-white solid; R_F = 0.05 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 178 °C (lit.: [32] 181–183 °C); IR (KBr): ν = 3440m, 2939m, 2839m, 1683s, 1625s, 1597s, 1516s, 1464m, 1426m, 1340m, 1298m, 1264s, 1210m, 1168m, 1142s, 1025m cm^{-1} ; UV-vis (CHCl_3): λ_{max} (log ϵ) = 238 (4.05), 324 (4.20) nm; ^1H NMR (400 MHz, CDCl_3): δ = 7.74 (d, J = 15.9 Hz, 1H, 3-H), 7.14 (dd, J = 8.3, 1.6 Hz, 1H, 6'-H), 7.08 (s, 1H, 5'-H), 6.88 (d, J = 8.3 Hz, 1H, 2'-H), 6.33 (d, J = 15.9 Hz, 1H, 2-H), 3.92 (s, 6H, 7'-H + 8'-H) ppm; ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ = 172.4 (C-1), 151.7 (C-4'), 149.4 (C-3'), 147.1 (C-3), 127.2 (C-1'), 123.3 (C-6'), 115.0 (C-2), 111.2 (C-2'), 109.9 (C-5'), 56.1 (C-8'), 56.0 (C-7') ppm; MS (ESI, MeOH): m/z (%) = 207.0 ($[\text{M}-\text{H}]^-$, 11), 415.1 ($[\text{2M}-\text{H}]^-$, 100), 436.9 ($[\text{2M}-2\text{H}+\text{Na}]^-$, 13); analysis calcd for $\text{C}_{11}\text{H}_{12}\text{O}_4$ (208.21): C 63.45, H 5.81; found C 63.11, H 6.18.

4.4.3. 3-(3,4-Dimethoxyphenyl)propanoic acid (**1b**)

A solution of **1a** (1.0 g, 4.8 mmol) in THF (20 mL) was hydrogenated at 75 psi pressure in the presence of Pd/C (10%, 0.116 g) for 12 h. Usual work-up gave **1b** (0.83 g, 82%) as a colorless solid; R_F = 0.09 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 98 °C (lit.: [33] 98–99 °C); IR (ATR): ν = 2935w, 1697s, 1591w, 1516m, 1463m, 1430m, 1343w, 1306w, 1236s, 1146s, 1027s, 840m, 809s, 768m, 567m cm^{-1} ; UV-vis (MeOH): λ_{max} (log ϵ) = 202 (4.42), 229 (3.75), 280 (3.32) nm; ^1H NMR (400 MHz, CDCl_3): δ = 6.80 (dd, J = 8.4, 3.2 Hz, 1H, 6'-H), 6.76 (s, 1H, 5'-H), 6.73 (d, J = 5.5 Hz, 1H, 2'-H), 3.87 (s, 3H, 8'-H), 3.86 (s, 3H, 7'-H), 2.91 (td, J = 7.8, 5.2 Hz, 2H, 3-H), 2.64 (td, J = 7.7, 5.1 Hz, 2H, 2-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ = 177.9 (C-1), 149.1 (C-4'),

147.8 (C-3'), 132.9 (C-1'), 120.3 (C-5'), 115.5 (C-2'), 111.8 (C-6'), 56.1 (C-8'), 56.0 (C-7'), 35.8 (C-3), 30.4 (C-2) ppm; MS (ESI, MeOH): m/z (%) = 209.1 ($[\text{M}-\text{H}]^-$, 100), 441 ($[\text{2M}-2\text{H}+\text{Na}]^-$, 71); analysis calcd for $\text{C}_{11}\text{H}_{14}\text{O}_4$ (210.23): C 62.85, H 6.71; found C 62.65, H 6.92.

4.4.4. (E)-3-(2,4-Dimethoxyphenyl)acrylic acid (**1c**)

Recrystallization from ethanol gave **1c** (1.33 g, 64%) as an off-white solid; R_F = 0.25 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 190 °C (lit.: [34] 187–189 °C); IR (KBr): ν = 3448m, 3004w, 2944w, 2838w, 2586w, 1681s, 1599s, 1505s, 1459m, 1418m, 1329s, 1313s, 1298m, 1268s, 1210s, 1160s, 1117s, 1043m, 1028m, 987m, 829m cm^{-1} ; UV-vis (CHCl_3): λ_{max} (log ϵ) = 254 (3.47), 316 (3.84), 354 (3.97) nm; ^1H NMR (400 MHz, CDCl_3): δ = 8.01 (d, J = 16.1 Hz, 1H, 3-H), 7.47 (d, J = 8.6 Hz, 1H, 3'-H), 6.52 (dd, J = 8.6, 2.3 Hz, 1H, 6'-H), 6.46 (d, J = 2.3 Hz, 1H, 5'-H), 6.45 (d, J = 16.0 Hz, 1H, 2-H), 3.88 (s, 3H, 7'-H), 3.85 (s, 3H, 8'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ = 172.6 (C-1), 163.3 (C-4'), 160.3 (C-2'), 142.5 (C-3), 131.0 (C-3'), 116.5 (C-1'), 115.0 (C-2), 105.5 (C-6'), 98.6 (C-5'), 55.7 (C-7' + C-8') ppm; MS (ESI, MeOH): m/z (%) = 191.1 ($[\text{M}+\text{H}-\text{H}_2\text{O}]^+$, 100), 208.9 ($[\text{M}+\text{H}]^+$, 81), 223.0 ($[\text{M}+\text{H}+\text{MeOH}-\text{H}_2\text{O}]^+$, 25), 231.1 ($[\text{M}+\text{Na}]^+$, 8); analysis calcd for $\text{C}_{11}\text{H}_{12}\text{O}_4$ (208.21): C 63.45, H 5.81; found: C 62.35, H 6.13.

4.4.5. (E)-3-(3,5-Dimethoxyphenyl)acrylic acid (**1d**)

Compound **1d** was obtained by recrystallization from ethanol as a white solid (1.16 g, 60%); R_F = 0.08 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 173 °C (lit.: [35] 174 °C); IR (ATR): ν = 2835w, 1682s, 1632m, 1593s, 1469m, 1433m, 1356w, 1320m, 1285s, 1206s, 1163s, 1057s, 927m, 852m, 837s, 808m, 670m, 645m, 606m, 540m, 472m cm^{-1} ; UV-vis (MeOH): λ_{max} (log ϵ) = 226 (4.39), 284 (4.25) nm; ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ = 7.51 (d, J = 16.0 Hz, 1H, 3-H), 6.86 (d, J = 2.2 Hz, 2H, 2'-H), 6.56 (d, J = 16.1 Hz, 1H, 2-H), 6.53 (d, J = 2.2 Hz, 1H, 4'-H), 3.77 (s, 6H, 5'-H) ppm; ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$): δ = 167.6 (C-1), 160.7 (C-3'), 143.9 (C-3), 136.2 (C-1'), 119.9 (C-2), 106.0 (C-2'), 102.4 (C-4'), 55.4 (C-5') ppm; MS (ESI, ASAP): m/z (%) = 207.1 ($[\text{M}-\text{H}]^-$, 100); analysis calcd for $\text{C}_{11}\text{H}_{12}\text{O}_4$ (208.21): C 63.45, H 5.81; found C 62.29, H 6.11.

4.4.6. (E)-3-(3,4,5-Trimethoxyphenyl)acrylic acid (**1e**)

Compound **1e** was obtained by recrystallization from ethanol as an off-white solid (4.36 g, 36%); R_F = 0.04 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 125 °C (lit.: [36] 124–125 °C); IR (ATR): ν = 3004w, 2837w, 2653w, 1684w, 1625m, 1583m, 1502m, 1470w, 1454m, 1417m, 1393m, 1321m, 1283m, 1202m, 1189m, 1156m, 1117s, 997s, 982s, 826m, 739m, 731m, 620m, 609m, 527m, 513m cm^{-1} ; UV-vis (MeOH): λ_{max} (log ϵ) = 231 (4.30), 302 (4.25) nm; ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ = 7.52 (d, J = 15.9 Hz, 1H, 3-H), 7.02 (s, 2H, 2'-H), 6.53 (d, J = 15.9 Hz, 1H, 2-H), 3.81 (s, 6H, 5'-H), 3.68 (s, 3H, 6'-H) ppm; ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ = 167.7 (C-1), 153.1 (C-3'), 144.1 (C-3), 139.3 (C-4'), 129.9 (C-1'), 118.5 (C-2), 105.8 (C-2'), 60.1 (C-6'), 56.0 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 221.1 ($[\text{M}+\text{H}-\text{H}_2\text{O}]^+$, 56), 239.0 ($[\text{M}+\text{H}]^+$, 100), 255.9 ($[\text{M}+\text{NH}_4]^+$, 10), 261.1 ($[\text{M}+\text{Na}]^+$, 37), 276.9 ($[\text{M}+\text{K}]^+$, 5), 292.7 ($[\text{M}+\text{Na}+\text{MeOH}]^+$, 10); analysis calcd for $\text{C}_{12}\text{H}_{14}\text{O}_7$ (238.24): C 60.50, H 5.92; found: C 62.39, H 6.18.

4.4.7. (E)-3-(3-Hydroxy-4-methoxyphenyl)acrylic acid (**1g**)

Compound **1g** was obtained by recrystallization from ethanol as an off-white solid (5.63 g, 58%); R_F = 0.02 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 232 °C (lit.: [37] 230–232 °C); IR (ATR): ν = 1612m, 1509s, 1442m, 1261s, 1134s, 1023m, 936s, 857s, 815s, 760s, 570s, 504s cm^{-1} ; UV-vis (MeOH): λ_{max} (log ϵ) = 219 (4.19), 242 (4.11), 296 (4.20), 321 (4.23) nm; ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ = 7.44 (d, J = 15.9 Hz, 1H, 3-H), 7.08 (dd, J = 8.2, 2.2 Hz, 1H, 6'-H), 7.06 (d, J = 2.1 Hz, 1H, 2'-H), 6.94 (d, J = 8.2 Hz, 1H, 5'-H), 6.23 (d, J = 15.9 Hz, 1H, 2-H), 3.80 (s, 3H, 7'-H) ppm; ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$): δ = 167.7 (C-1), 149.8 (C-4'), 146.6 (C-3'), 144.1 (C-3), 127.1 (C-1'), 120.9 (C-6'), 116.3 (C-2), 114.1 (C-2'), 112.0 (C-5'), 55.6

(C-7') ppm; MS (ESI, MeOH): m/z (%) = 193.0 ([M-H]⁻, 100), 386.8 ([2M-H]⁻, 16); analysis calcd for C₁₀H₁₀O₄ (194.19): C 61.85, H 5.19; found: C 62.03, H 5.22.

4.4.8. (E)-3-(4-Hydroxy-3-methoxyphenyl)acrylic acid (**1h**)

Compound **1h** was obtained by recrystallization from ethanol as a white solid (5.61 g, 57%); R_F = 0.06 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 170 °C (lit.: [37] 168–169 °C); IR (ATR): ν = 3435w, 1662m, 1590m, 1511m, 1465m, 1431m, 1323w, 1265s, 1163s, 1112m, 1033s, 942s, 851s, 803s, 751m, 685m, 598s, 572s, 519s cm⁻¹; UV-vis (MeOH): λ_{max} (log ϵ) = 218 (3.89), 321 (4.00) nm; ¹H NMR (500 MHz, DMSO-*d*₆): δ = 7.48 (d, J = 15.9 Hz, 1H, 3-H), 7.27 (d, J = 1.9 Hz, 1H, 2'-H), 7.08 (dd, J = 8.2, 1.9 Hz, 1H, 6'-H), 6.78 (d, J = 8.1 Hz, 1H, 5'-H), 6.35 (d, J = 15.9 Hz, 1H, 2-H), 3.81 (s, 3H, 7'-H) ppm; ¹³C NMR (125 MHz, DMSO-*d*₆): δ = 167.9 (C-1), 149.1 (C-4'), 147.9 (C-3'), 144.5 (C-3), 125.7 (C-1'), 122.8 (C-6'), 115.6 (C-2), 115.5 (C-5'), 111.2 (C-2'), 55.7 (C-7') ppm; MS (ESI, ASAP): m/z (%) = 178.1 ([M-H-CH₃]⁻, 100), 193.1 ([M-H]⁻, 98); analysis calcd for C₁₀H₁₀O₄ (194.19): C 61.85, H 5.19; found C 61.75, H 6.19.

4.4.9. (E)-3-(4-Acetoxy-3-methoxyphenyl)acrylic acid (**1i**)

Compound **1i** was obtained by recrystallization from ethanol as a yellowish solid (2.52 g, 62%); R_F = 0.02 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 195 °C; IR (ATR): ν = 2944w, 1760m, 1682s, 1632m, 1601m, 1506m, 1469w, 1421m, 1371w, 1316m, 1260s, 1222s, 1198s, 1154s, 1120s, 1033m, 985s, 913m, 855s, 837m, 691m, 648m, 599m, 527m cm⁻¹; UV-vis (MeOH): λ_{max} (log ϵ) = 277 (3.83) nm; ¹H NMR (500 MHz, DMSO-*d*₆): δ = 7.57 (d, J = 16.0 Hz, 1H, 3-H), 7.48 (d, J = 1.8 Hz, 1H, 2'-H), 7.26 (dd, J = 8.2, 1.8 Hz, 1H, 6'-H), 7.12 (d, J = 8.1 Hz, 1H, 5'-H), 6.58 (d, J = 16.0 Hz, 1H, 2-H), 3.82 (s, 3H, 7'-H), 2.26 (s, 3H, 9'-H) ppm; ¹³C NMR (125 MHz, DMSO-*d*₆): δ = 168.4 (C-8'), 167.5 (C-1), 151.1 (C-3'), 143.3 (C-3), 140.8 (C-4'), 133.2 (C-1'), 123.2 (C-5'), 121.3 (C-6'), 119.6 (C-2), 111.8 (C-2'), 56.0 (C-7'), 20.4 (C-9') ppm; MS (ESI, ASAP): m/z (%) = 235.1 ([M-H]⁻, 100); analysis calcd for C₁₂H₁₂O₅ (236.22): C 61.02, H 5.12; found: C 61.59, H 5.31.

4.4.10. (E)-3-(3,4-Dimethylphenyl)acrylic acid (**1j**)

Compound **1j** was obtained by recrystallization from ethanol as a white solid (1.69 g, 57%); R_F = 0.19 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 174 °C; IR (ATR): ν = 2919w, 1672s, 1621s, 1421m, 1318s, 1285s, 1219s, 956s, 925s, 866s, 815s, 688s, 582m, 546s cm⁻¹; UV-vis (MeOH): λ_{max} (log ϵ) = 223 (4.00), 286 (4.14) nm; ¹H NMR (500 MHz, DMSO-*d*₆): δ = 7.56 (d, J = 16.0 Hz, 1H, 3-H), 7.45 (d, J = 1.9 Hz, 1H, 5'-H), 7.38 (dd, J = 7.7, 1.9 Hz, 1H, 6'-H), 7.17 (d, J = 7.7 Hz, 1H, 2'-H), 6.43 (d, J = 16.0 Hz, 1H, 2-H), 2.23 (s, 6H, 7'-H + 8'-H) ppm; ¹³C NMR (125 MHz, DMSO-*d*₆): δ = 167.7 (C-1), 144.1 (C-3), 139.0 (C-4'), 136.8 (C-3'), 131.8 (C-1'), 130.0 (C-6'), 129.1 (C-2'), 125.8 (C-5'), 117.8 (C-2), 19.3 (C-7'), 19.2 (C-8') ppm; MS (ESI, ASAP): m/z (%) = 175.1 ([M-H]⁻, 100); analysis calcd for C₁₁H₁₂O₂ (176.21): C 74.98, H 6.86; found: C 74.75, H 7.00.

4.4.11. (E)-3-(2-Nitrophenyl)acrylic acid (**1k**)

After recrystallization from ethanol compound **1k** was obtained as an off-white solid (0.89 g, 46%); R_F = 0.07 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 247 °C; IR (KBr): ν = 1694m, 1519m, 1419m, 1280s, 1202s, 981m, 865m, 789s, 754s, 701s, 594m, 542m, 486m cm⁻¹; UV-vis (EtOH): λ_{max} (log ϵ) = 286 (3.73) nm; ¹H NMR (400 MHz, CDCl₃): δ = 8.07 (dd, J = 8.1, 1.2 Hz, 1H, 6'-H), 7.94 (dd, J = 7.8, 1.2 Hz, 1H, 3'-H), 7.86 (d, J = 15.9 Hz, 1H, 3-H), 7.78 (td, J = 7.7, 1.5 Hz, 1H, 4'-H), 7.69–7.65 (m, 1H, 5'-H), 6.54 (d, J = 15.8 Hz, 1H, 2-H) ppm; ¹³C NMR (100 MHz, DMSO-*d*₆): δ = 166.8 (C-1), 148.3 (C-2'), 138.7 (C-3), 133.8 (C-4'), 130.8 (C-5'), 129.3 (C-1'), 129.2 (C-3'), 124.6 (C-6'), 123.8 (C-2) ppm; MS (ESI, MeOH): m/z (%) = 191.9 ([M-H]⁻, 41), 237.8 ([M + HCO₂]⁻, 11), 259.9 ([M-H + NaHCO₂]⁻, 6), 384.9 ([2 M-H]⁻, 100), 406.9 ([2M-2H + Na]⁻, 5); analysis calcd for C₉H₇NO₂ (193.16): C 55.96, H 3.65, N 7.25; found: C 55.71, H 3.79, N

7.07.

4.4.12. (E)-3-(3-Nitrophenyl)acrylic acid (**1l**)

After recrystallization from ethanol compound **1l** was obtained as an off-white solid (1.20 g, 62%); R_F = 0.02 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 204 °C; IR (KBr): ν = 2539w, 1683s, 1630m, 1521s, 1485w, 1434w, 1418m, 1360s, 1319s, 1306s, 1285s, 1223m, 1177m, 1108w, 985m, 922s, 869m, 825m, 805s, 747m, 716s, 667s, 596m, 567m, 544m, 486m cm⁻¹; UV-vis (EtOH): λ_{max} (log ϵ) = 295 (3.91) nm; ¹H NMR (400 MHz, DMSO-*d*₆): δ = 12.60 (s, 1H, OH), 8.51 (t, J = 2.0 Hz, 1H, 4'-H), 8.23 (ddd, J = 8.2, 2.3, 0.9 Hz, 1H, 6'-H), 8.18 (dt, J = 7.7, 1.2 Hz, 1H, 5'-H), 7.72 (d, J = 16.2 Hz, 1H, 3-H), 7.71 (d, J = 7.9 Hz, 1H, 2'-H), 6.74 (d, J = 16.1 Hz, 1H, 2-H) ppm; ¹³C NMR (100 MHz, DMSO-*d*₆): δ = 167.1 (C-1), 148.3 (C-3'), 141.4 (C-3), 136.1 (C-1'), 134.0 (C-5'), 130.3 (C-2'), 124.4 (C-6'), 122.8 (C-4'), 122.3 (C-2) ppm; MS (ESI, MeOH): m/z (%) = 192.0 ([M-H]⁻, 100), 237.9 ([M + HCO₂]⁻, 3), 260.0 ([M-H + NaHCO₂]⁻, 2), 384.8 ([2M-H]⁻, 44), 407.0 ([2M-2H + Na]⁻, 14); analysis calcd for C₉H₇NO₂ (193.16): C 55.96, H 3.65, N 7.25; found: C 55.71, H 3.91, N 6.96.

4.4.13. (E)-3-(4-Dimethylamino)phenyl)acrylic acid (**1m**)

Compound **1m** was obtained by recrystallization from ethanol as a yellowish solid (3.86 g, 40%); R_F = 0.08 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 212 °C (lit.: [38] 210–212 °C); IR (ATR): ν = 2803w, 1665m, 1590m, 1523m, 1434m, 1366m, 1308m, 1260m, 1227m, 1163m, 987m, 940m, 858m, 812s, 675m, 513s, 479m cm⁻¹; UV-vis (MeOH): λ_{max} (log ϵ) = 242 (3.80), 358 (4.27) nm; ¹H NMR (500 MHz, DMSO-*d*₆): δ = 7.49 (d, J = 2.5 Hz, 2H, 2'-H), 7.46 (d, J = 15.8 Hz, 1H, 3-H), 6.70 (d, J = 8.9 Hz, 2H, 3'-H), 6.20 (d, J = 15.8 Hz, 1H, 2-H), 2.96 (s, 6H, 5'-H) ppm; ¹³C NMR (100 MHz, DMSO-*d*₆): δ = 168.1 (C-1), 151.6 (C-4'), 144.6 (C-3), 129.7 (C-2'), 121.5 (C-1'), 112.9 (C-2), 111.8 (C-3'), 39.7 (C-5') ppm; MS (ESI, ASAP): m/z (%) = 190.2 ([M-H]⁻, 100); analysis calcd for C₁₁H₁₃NO₂ (191.23): C 69.09, H 6.85, N 7.32; found: C 68.88, H 7.02, N 7.15.

4.4.14. General procedure B

Under argon atmosphere, to a solution of the acid (1 eq.) in dry DCM (10 mL/mmol) at 0 °C oxalyl chloride (3 eq.) and DMF (2 drops) were added. After stirring for 1 h at 0 °C followed by 1 h at room temperature the solvent was removed *in vacuo*, and the residue was washed with dry DCM (10 mL/mmol). The residue was re-dissolved in toluene (5 mL/mmol), pyridine (2 eq.) and the corresponding acid (1 eq.) was added. The mixture was stirred over night at room temperature. Usual aqueous work-up gave a residue that was washed with ethanol.

4.4.15. (E)-3-(3,4-Dimethoxyphenyl)acrylic anhydride (**2a**)

Compound **2a** was obtained as a white solid (0.53 g, 55%); R_F = 0.42 (silica gel, chloroform); m.p. 177 °C; IR (ATR): ν = 2931w, 2835w, 1754m, 1707s, 1625s, 1598m, 1581w, 1510s, 1460m, 1438w, 1424m, 1354w, 1311w, 1266s, 1240s, 1214s, 1162m, 1144s, 1075s, 1021s, 1001s, 986s, 972m, 943s, 868s, 810s, 773m, 731m, 707m, 602m, 570m, 548m cm⁻¹; UV-vis (CHCl₃): λ_{max} (log ϵ) = 227 (3.89), 250 (3.93), 347 (4.26) nm; ¹H NMR (500 MHz, CDCl₃): δ = 7.80 (d, J = 15.8 Hz, 2H, 3-H), 7.17 (dd, J = 8.3, 1.9 Hz, 2H, 6'-H), 7.09 (d, J = 1.9 Hz, 2H, 5'-H), 6.90 (d, J = 8.3 Hz, 2H, 2'-H), 6.40 (d, J = 15.8 Hz, 2H, 2-H), 3.94 (s, 6H, 7'-H), 3.93 (s, 6H, 8'-H) ppm; ¹³C NMR (125 MHz, CDCl₃): δ = 163.1 (C-1), 152.2 (C-4'), 149.5 (C-3'), 148.7 (C-3), 127.0 (C-1'), 123.7 (C-6'), 114.6 (C-2), 111.3 (C-2'), 110.1 (C-5'), 56.2 (C-8'), 56.1 (C-7') ppm; MS (ESI, MeOH): m/z (%) = 421.5 ([M + Na]⁺, 100), 437.5 ([M + K]⁺, 25), 453.6 ([M + Na + MeOH]⁺); analysis calcd for C₂₂H₂₂O₇ (398.41): C 66.32, H 5.57; found: C 66.18, H 6.73.

4.4.16. 3-(3,4-Dimethoxyphenyl)propanoic anhydride (**2b**)

Compound **2b** was obtained as an off-white solid (0.09 g, 46%);

$R_F = 0.04$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 178 °C; IR (ATR): $\nu = 2935w, 2838w, 1697s, 1591w, 1516m, 1463m, 1430m, 1409w, 1346w, 1306m, 1251m, 1236s, 1213m, 1146s, 1026m, 840m, 809m, 768m\text{ cm}^{-1}$; UV-vis (MeOH): λ_{max} (log ϵ) = 299 (3.80), 279 (3.57) nm; ^1H NMR (500 MHz, CDCl_3): $\delta = 6.80$ (dd, $J = 8.3, 5.0$ Hz, 2H, 5'-H + 5''-H), 6.77–6.72 (m, 4H, 2'-H + 2''-H + 6'-H + 6''-H), 3.88 (s, 3H, 7'-H), 3.87 (s, 3H, 7''-H), 3.86 (s, 3H, 8'-H), 3.85 (s, 3H, 8''-H), 2.91 (td, $J = 7.5, 1.2$ Hz, 4H, 2-H + 2''-H), 2.73 (t, $J = 7.6$ Hz, 2H, 3-H), 2.67 (t, $J = 7.7$ Hz, 2H, 3''-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): $\delta = 168.8$ (C-1), 149.2 (C-3'), 149.1 (C-3''), 147.9 (C-4'), 147.8 (C-4''), 132.9 (C-1'), 132.4 (C-1''), 120.3 (C-6'), 120.2 (C-6''), 111.9 (C-2' + C-2''), 111.6 (C-5' + C-5''), 56.1 (C-7' + C-7''), 56.0 (C-8' + C-8''), 37.3 (C-2), 35.7 (C-2''), 30.4 (C-3), 30.0 (C-3'') ppm; MS (ESI, MeOH): m/z (%) = 425.1 ($[\text{M} + \text{Na}]^+$, 100), 826.9 ($[\text{2M} + \text{Na}]^+$, 8); analysis calcd for $\text{C}_{22}\text{H}_{26}\text{O}_7$ (402.44): C 65.66, H 6.51; found: C 65.41, H 6.72.

4.4.17. (*E*)-3-(2,4-Dimethoxyphenyl)acrylic anhydride (2c)

Compound **2c** was obtained as a white solid (0.85 g, 89%); $R_F = 0.35$ (silica gel, chloroform); m.p. 105–108 °C; IR (KBr): $\nu = 3448w, 2946w, 2843w, 1754m, 1711m, 1602s, 1566m, 1505m, 1457m, 1439w, 1418m, 1330m, 1296s, 1268m, 1215s, 1164s, 1118m, 1066s, 985m, 830m\text{ cm}^{-1}$; UV-vis (CHCl_3): λ_{max} (log ϵ) = 265 (3.81), 385 (4.35) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 8.05$ (d, $J = 16.0$ Hz, 2H, 3-H), 7.48 (d, $J = 8.6$ Hz, 2H, 3'-H), 6.54 (d, $J = 15.9$ Hz, 2H, 2-H), 6.53 (dd, $J = 8.6, 2.3$ Hz, 2H, 6'-H), 6.46 (d, $J = 2.3$ Hz, 2H, 5'-H), 3.89 (s, 6H, 7'-H), 3.86 (s, 6H, 8'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 164.1$ (C-1), 163.7 (C-4'), 160.6 (C-2'), 144.0 (C-3), 131.5 (C-3'), 116.4 (C-1'), 114.9 (C-2), 105.7 (C-6'), 98.6 (C-5'), 55.7 (C-7'), 55.6 (C-8') ppm; MS (ESI, MeOH): m/z (%) = 191.1 ($[\text{M}-\text{C}_{11}\text{H}_{11}\text{O}_4]^+$, 38), 209.0 ($[\text{M}-\text{C}_{11}\text{H}_{11}\text{O}_4 + \text{H}_2\text{O}]^+$, 3), 223.0 ($[\text{M}-\text{C}_{11}\text{H}_{11}\text{O}_4 + \text{MeOH}]^+$, 22), 421.0 ($[\text{M} + \text{Na}]^+$, 49), 818.9 ($[\text{2M} + \text{Na}]^+$, 100); analysis calcd for $\text{C}_{22}\text{H}_{22}\text{O}_7$ (398.41): C 66.32, H 5.57; found: C 66.11, H 5.69.

4.4.18. (*E*)-3-(3,5-Dimethoxyphenyl)acrylic anhydride (2d)

Compound **2d** was obtained as an off-white solid (0.31 g, 81%); $R_F = 0.21$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 140 °C; IR (ATR): $\nu = 1778m, 1706m, 1630m, 1593s, 1471w, 1431m, 1356w, 1302w, 1284w, 1208s, 1158s, 1085s, 1065s, 1055s, 977m, 952s, 925m, 850m, 837s, 818m\text{ cm}^{-1}$; UV-vis (CHCl_3): λ_{max} (log ϵ) = 206 (4.45), 231 (4.29), 309 (4.34) nm; ^1H NMR (500 MHz, CDCl_3): $\delta = 7.77$ (d, $J = 15.9$ Hz, 2H, 3-H), 6.71 (d, $J = 2.2$ Hz, 4H, 2'-H), 6.55 (t, $J = 2.2$ Hz, 2H, 4'-H), 6.49 (d, $J = 15.9$ Hz, 2H, 2-H), 3.83 (s, 12H, 5'-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): $\delta = 162.5$ (C-1), 161.3 (C-3'), 148.8 (C-3), 135.7 (C-1'), 117.4 (C-2), 106.6 (C-2'), 103.7 (C-4'), 55.7 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 421.1 ($[\text{M} + \text{Na}]^+$, 44), 818.9 ($[\text{2M} + \text{Na}]^+$, 100); analysis calcd for $\text{C}_{22}\text{H}_{22}\text{O}_7$ (398.41): C 66.32, H 5.57; found: C 66.15, H 5.80.

4.4.19. (*E*)-3-(3,4,5-Trimethoxyphenyl)acrylic anhydride (2e)

Compound **2e** was obtained as a white solid (0.30 g, 77%); $R_F = 0.06$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 174 °C (lit.: [39] 190–192); IR (ATR): $\nu = 1802w, 1757w, 1698w, 1623w, 1582m, 1505m, 1455w, 1422m, 1342w, 1323w, 1267m, 1251m, 1159w, 1125s, 1040s, 1008s, 987m, 959m, 840m\text{ cm}^{-1}$; UV-vis (CHCl_3): λ_{max} (log ϵ) = 236 (4.23), 335 (4.31) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 7.77$ (d, $J = 15.8$ Hz, 2H, 3-H), 6.80 (s, 4H, 2'-H), 6.43 (d, $J = 15.8$ Hz, 2H, 2-H), 3.91 (s, 12H, 5'-H + 6'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 162.7$ (C-1), 153.7 (C-3'), 148.8 (C-3), 141.3 (C-4'), 129.3 (C-1'), 116.0 (C-2), 106.0 (C-2'), 61.2 (C-6'), 56.4 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 481.1 ($[\text{M} + \text{Na}]^+$, 44), 938.9 ($[\text{2M} + \text{Na}]^+$, 100); analysis calcd for $\text{C}_{24}\text{H}_{26}\text{O}_9$ (458.46): C 62.88, H 5.72; found: C 62.75, H 5.90.

4.4.20. (*E*)-3-(4-Acetoxy-3-methoxyphenyl)acrylic anhydride (2i)

Compound **2i** was obtained as a white solid (0.29 g, 76%); $R_F = 0.06$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 122–124 °C

(lit.: [40] 178–180 °C); IR (ATR): $\nu = 1763s, 1702w, 1627m, 1586w, 1461w, 1419w, 1371w, 1305w, 1257m, 1222m, 1169w, 1123w, 1054s, 1034m, 1001m, 966m\text{ cm}^{-1}$; UV-vis (CHCl_3): λ_{max} (log ϵ) = 206 (4.27), 297 (4.20) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 7.81$ (d, $J = 15.9$ Hz, 2H, 3-H), 7.19 (dd, $J = 8.1, 1.9$ Hz, 2H, 6'-H), 7.14 (d, $J = 1.9$ Hz, 2H, 2'-H), 7.10 (d, $J = 8.1$ Hz, 2H, 5'-H), 6.47 (d, $J = 15.9$ Hz, 2H, 2-H), 3.89 (s, 6H, 7'-H), 2.33 (s, 6H, 9'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 168.8$ (C-8'), 162.4 (C-1), 151.8 (C-3'), 148.1 (C-3), 142.5 (C-4'), 132.8 (C-1'), 123.7 (C-5'), 122.0 (C-6'), 117.0 (C-2), 111.8 (C-2'), 55.2 (C-7'), 20.8 (C-9') ppm; MS (ESI, MeOH): m/z (%) = 477.1 ($[\text{M} + \text{Na}]^+$, 55), 930.9 ($[\text{2M} + \text{Na}]^+$, 100); analysis calcd for $\text{C}_{24}\text{H}_{22}\text{O}_9$ (454.43): C 63.43, H 4.88; found: C 63.21, H 5.02.

4.4.21. (*E*)-3-(3,4-Dimethylphenyl)acrylic anhydride (2j)

Compound **2j** was obtained as a white solid (0.23 g, 60%); $R_F = 0.58$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 195 °C; IR (ATR): $\nu = 2919w, 1751s, 1704s, 1626m, 1605w, 1505w, 1450w, 1268w, 1234w, 1206w, 1121s, 1021w, 987s, 944s, 929s, 890m, 865m, 826s, 813s, 567m\text{ cm}^{-1}$; UV-vis (CHCl_3): λ_{max} (log ϵ) = 206 (4.44), 232 (4.36), 310 (4.64) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 7.80$ (d, $J = 15.9$ Hz, 2H, 3-H), 7.35 (d, $J = 2.0$ Hz, 2H, 2'-H), 7.32 (dd, $J = 7.8, 2.0$ Hz, 2H, 6'-H), 7.18 (d, $J = 7.8$ Hz, 2H, 5'-H), 6.47 (d, $J = 15.9$ Hz, 2H, 2-H), 2.31 (s, 6H, 7'-H), 2.30 (s, 6H, 8'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 163.0$ (C-1), 149.0 (C-3), 140.8 (C-3'), 137.5 (C-4'), 131.7 (C-1'), 130.5 (C-5'), 129.9 (C-2'), 126.4 (C-6'), 115.7 (C-2), 20.1 (C-7'), 19.9 (C-8') ppm; MS (ESI, MeOH): m/z (%) = 357.1 ($[\text{M} + \text{Na}]^+$, 74), 691.0 ($[\text{2M} + \text{Na}]^+$, 100); analysis calcd for $\text{C}_{22}\text{H}_{22}\text{O}_3$ (344.41): C 79.02, H 6.63; found: C 78.83, H 6.82.

4.4.22. (*E*)-3-(2-Nitrophenyl)acrylic anhydride (2k)

Compound **2k** was obtained as a yellowish solid (0.23 g, 60%); $R_F = 0.31$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 184 °C; IR (ATR): $\nu = 1756m, 1706m, 1624w, 1570w, 1516m, 1341m, 1288w, 1263w, 1205w, 1148m, 1121s, 979m, 935s, 891m, 867m, 789s, 751s, 712m, 681m, 536m, 503m, 487m\text{ cm}^{-1}$; UV-vis (CHCl_3): λ_{max} (log ϵ) = 252 (4.29) nm; ^1H NMR (500 MHz, CDCl_3): $\delta = 8.34$ (d, $J = 15.7$ Hz, 1H, 3-H), 8.12 (dd, $J = 8.1, 1.0$ Hz, 1H, 6'-H), 7.72–7.68 (m, 2H, 4'-H + 3'-H), 7.62 (ddd, $J = 8.6, 6.8, 2.1$ Hz, 1H, 5'-H), 6.46 (d, $J = 15.8$ Hz, 1H, 2-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): $\delta = 161.0$ (C-1), 148.4 (C-2'), 144.3 (C-3), 133.9 (C-4'), 131.3 (C-5'), 130.1 (C-1'), 129.5 (C-3'), 125.3 (C-6'), 121.7 (C-2) ppm; MS (ESI, MeOH): m/z (%) = 391.0 ($[\text{M} + \text{Na}]^+$, 100), 758.8 ($[\text{2M} + \text{Na}]^+$, 79); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{N}_2\text{O}_7$ (368.30): C 58.70, H 3.28, N 7.61; found: C 58.55, H 3.51, N 7.34.

4.4.23. (*E*)-3-(3-Nitrophenyl)acrylic anhydride (2l)

Compound **2l** was obtained as an off-white solid (0.19 g, 64%); $R_F = 0.07$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 174 °C; IR (ATR): $\nu = 3077w, 1787m, 1714w, 1637w, 1523s, 1481w, 1443w, 1354s, 1268w, 1227w, 1204m, 1091s, 974s, 938m, 804m, 741m, 710m, 660m\text{ cm}^{-1}$; UV-vis (MeOH): λ_{max} (log ϵ) = 205 (3.98), 262 (3.95) nm; ^1H NMR (500 MHz, CDCl_3): $\delta = 8.46$ (t, $J = 2.0$ Hz, 2H, 4'-H), 8.31 (ddd, $J = 8.2, 2.3, 1.0$ Hz, 2H, 6'-H), 7.91 (d, $J = 16.0$ Hz, 2H, 3-H), 7.91–7.88 (m, 2H, 5'-H), 7.65 (t, $J = 8.0$ Hz, 2H, 2'-H), 6.67 (d, $J = 16.0$ Hz, 2H, 2-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): $\delta = 161.4$ (C-1), 149.0 (C-3'), 146.0 (C-3), 135.4 (C-1'), 134.2 (C-5'), 130.4 (C-2'), 125.7 (C-6'), 123.0 (C-4'), 119.8 (C-2) ppm; MS (ESI, MeOH): m/z (%) = 390.9 ($[\text{M} + \text{Na}]^+$, 58), 758.7 ($[\text{2M} + \text{Na}]^+$, 100); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{N}_2\text{O}_7$ (368.30): C 58.70, H 3.28, N 7.61; found: C 58.61, H 3.43, N 7.47.

4.4.24. (*E*)-3-(4-(Dimethylamino)phenyl)acrylic anhydride (2m)

Compound **2m** was obtained as a yellowish solid (0.18 g, 48%); $R_F = 0.1$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 145 °C; IR (ATR): $\nu = 2816w, 1668m, 1594m, 1523m, 1435m, 1368m, 1309m, 1261m, 1228m, 1186s, 988m, 941m, 814s, 676m, 514m, 481m\text{ cm}^{-1}$; UV-vis

(MeOH): $\lambda_{\max}$ (log ϵ) = 246 (2.49), 349 (3.02) nm; ^1H NMR (500 MHz, CDCl_3): δ = 7.71 (d, J = 15.8 Hz, 2H, 3-H), 7.44 (d, J = 8.8 Hz, 4H, 2'-H), 6.68 (d, J = 8.9 Hz, 4H, 3'-H), 6.22 (d, J = 15.8 Hz, 2H, 2-H), 3.03 (s, 12H, 5'-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ = 171.6 (C-1), 152.2 (C-4'), 147.6 (C-3), 130.3 (C-2'), 122.1 (C-1'), 112.0 (C-3'), 111.2 (C-2), 40.3 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 387.1 ([M + Na] $^+$, 15), 751.0 ([2M + Na] $^+$, 16); analysis calcd for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_3$ (364.44): C 72.51, H 6.64, N 7.69; found: C 72.39, H 6.87, N 7.51.

4.4.25. Cinnamic anhydride (2n)

Compound **2n** was obtained as a white solid (0.61 g, 65%); R_F = 0.12 (silica gel, chloroform); m.p. 136 °C (lit.: [41] 136–137 °C); IR (ATR): ν = 1764m, 1699s, 1630m, 1495w, 1450m, 1304w, 1269w, 1227m, 1200m, 1068s, 958s, 857m, 750s, 689s, 674s, 551s, 510m, 476s cm^{-1} ; UV-vis (CHCl₃): $\lambda_{\max}$ (log ϵ) = 227 (4.14), 298 (4.50) nm; ^1H NMR (400 MHz, CDCl_3): δ = 7.86 (d, J = 15.9 Hz, 2H, 3-H), 7.60–7.57 (m, 4H, 3'-H), 7.46–7.41 (m, 6H, 2'-H + 4'-H), 6.54 (d, J = 15.9 Hz, 2H, 2-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ = 162.6 (C-1), 148.8 (C-3), 133.9 (C-1'), 131.4 (C-4'), 129.2 (C-2'), 128.7 (C-3'), 116.9 (C-2) ppm; MS (ESI, ASAP): m/z (%) = 301.3 ([M + Na] $^+$, 100), 333.4 ([M + Na + MeOH] $^+$, 96); analysis calcd for $\text{C}_{18}\text{H}_{14}\text{O}_3$ (278.31): C 77.68, H 5.07; found: C 77.46, H 5.18.

4.4.26. (E)-3-(3,4-Dihydroxyphenyl)acrylic anhydride (2o)

Compound **2o** was obtained as an off-white solid (0.51 g, 53%); R_F = 0.62 (silica gel, chloroform); m.p. 178 °C; ^1H NMR (400 MHz, CD_3OD): δ = 7.56 (d, J = 15.9 Hz, 2H, 3-H), 7.06 (d, J = 2.1 Hz, 2H, 5'-H), 6.96 (dd, J = 8.2, 2.0 Hz, 2H, 6'-H), 6.80 (d, J = 8.2 Hz, 2H, 2'-H), 6.24 (d, J = 15.9 Hz, 2H, 2-H) ppm; ^{13}C NMR (100 MHz, CD_3OD): δ = 171.0 (C-1), 149.5 (C-4'), 147.0 (C-3), 146.8 (C-3'), 127.8 (C-1'), 122.8 (C-6'), 116.5 (C-2'), 115.5 (C-2), 115.1 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 365.1 ([M + Na] $^+$, 21), 706.9 ([2M + Na] $^+$, 26); analysis calcd for $\text{C}_{18}\text{H}_{14}\text{O}_7$ (342.30): C 63.16, H 4.12; found: C 62.89, H 4.36.

4.4.27. (E)-3-(4-Hydroxy-3,5-dimethoxyphenyl)acrylic anhydride (2p)

Compound **2p** was obtained as an off-white solid (0.52 g, 52%); R_F = 0.56 (silica gel, chloroform); m.p. 187 °C; IR (ATR): ν = 2837w, 1659w, 1620w, 1594w, 1516m, 1460m, 1423m, 1335m, 1296m, 1268m, 1214m, 1156m, 1119s, 973m, 909m, 833m, 816m, 629s, 551m cm^{-1} ; UV-vis (MeOH): $\lambda_{\max}$ (log ϵ) = 237 (4.40), 323 (4.46) nm; ^1H NMR (500 MHz, CDCl_3): δ = 7.70 (d, J = 15.8 Hz, 2H, 3-H), 6.80 (s, 4H, 2'-H), 6.32 (d, J = 15.8 Hz, 2H, 2-H), 3.94 (s, 6H, 5'-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ = 171.5 (C-1), 147.5 (C-3'), 147.4 (C-3), 137.8 (C-4'), 125.7 (C-1'), 114.9 (C-2), 105.6 (C-2'), 56.5 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 453.0 ([M + Na] $^+$, 35); analysis calcd for $\text{C}_{22}\text{H}_{22}\text{O}_9$ (430.41): C 61.39, H 5.15; found: C 61.04, H 5.31.

4.4.28. (E)-3-(2-Methoxyphenyl)acrylic anhydride (2q)

Compound **2q** was obtained as a white solid (0.18 g, 44%); R_F = 0.26 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 101 °C (lit.: [40] 104–106 °C); IR (ATR): ν = 2962w, 2838w, 1765m, 1710m, 1623m, 1599m, 1488m, 1464m, 1438m, 1324w, 1299w, 12630m, 1245m, 1198w, 1163m, 1126s, 1104s, 1046m, 1028s, 988m, 871m, 784m, 753s, 518m cm^{-1} ; UV-vis (CHCl₃): $\lambda_{\max}$ (log ϵ) = 232 (3.87), 288 (4.05), 339 (3.97) nm; ^1H NMR (400 MHz, CDCl_3): δ = 8.15 (d, J = 16.1 Hz, 2H, 3-H), 7.55 (dd, J = 7.7, 1.5 Hz, 2H, 6'-H), 7.43–7.38 (m, 2H, 4'-H), 7.02–6.93 (m, 4H, 5'-H + 3'-H), 6.65 (d, J = 16.1 Hz, 2H, 2-H), 3.89 (s, 6H, 7'-H), 3.86 (s, 6H, 8'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ = 163.5 (C-1), 158.9 (C-2'), 144.2 (C-3), 132.6 (C-4'), 129.8 (C-6'), 123.0 (C-1'), 121.0 (C-5'), 117.6 (C-2), 111.4 (C-3'), 55.7 (C-7') ppm; MS (ESI, MeOH): m/z (%) = 361.1 ([M + Na] $^+$, 100), 698.9 ([2M + Na] $^+$, 92); analysis calcd for $\text{C}_{20}\text{H}_{18}\text{O}_5$ (338.36): C 71.00, H 5.36; found: C 70.73, H 5.47.

4.4.29. (E)-3-(3-Methoxyphenyl)acrylic anhydride (2r)

Compound **2r** was obtained as a white solid (0.317 g, 84%); R_F = 0.28 (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 71 °C; IR (ATR): ν = 2842w, 1759s, 1694m, 1628m, 1609w, 1579m, 1491m, 1462m, 1434m, 1319w, 1272m, 1247m, 1158m, 1122s, 1046m, 978s, 935s, 873m, 849s, 775s, 720m, 671m, 577m, 552m, 510m cm^{-1} ; UV-vis (MeOH): $\lambda_{\max}$ (log ϵ) = 216 (3.56), 285 (3.65) nm; ^1H NMR (500 MHz, CDCl_3): δ = 7.82 (d, J = 15.9 Hz, 2H, 3-H), 7.34 (t, J = 7.9 Hz, 2H, 5'-H), 7.17 (d, J = 7.7 Hz, 2H, 6'-H), 7.09–7.08 (m, 2H, 2'-H), 7.00 (ddd, J = 8.3, 2.6, 0.7 Hz, 2H, 4'-H), 6.51 (d, J = 15.9 Hz, 2H, 2-H), 3.85 (s, 6H, 7'-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ = 162.6 (C-1), 160.2 (C-3'), 148.7 (C-3), 135.2 (C-1'), 130.2 (C-5'), 121.4 (C-6'), 117.3 (C-4'), 117.2 (C-2), 113.5 (C-2'), 55.5 (C-7') ppm; MS (ESI, MeOH): m/z (%) = 361.0 ([M + Na] $^+$, 100), 698.8 ([2M + Na] $^+$, 77); analysis calcd for $\text{C}_{20}\text{H}_{18}\text{O}_5$ (338.36): C 71.00, H 5.36; found: C 70.81, H 5.51.

4.4.30. (E)-3-(4-Methoxyphenyl)acrylic anhydride (2s)

Compound **2s** was obtained as a white solid (0.18 g, 46%); R_F = 0.7 (silica gel, chloroform); m.p. 103 °C (lit.: [40] 104–105 °C); IR (ATR): ν = 1754w, 1717m, 1624w, 1597m, 1571w, 1509m, 1455w, 1439w, 1423w, 1304w, 1289w, 1255m, 1176w, 1150w, 1061s, 1022m, 972s, 923m, 824s, 804m, 535m, 518m cm^{-1} ; UV-vis (CHCl₃): $\lambda_{\max}$ (log ϵ) = 206 (4.19), 235 (3.73), 331 (4.13) nm; ^1H NMR (400 MHz, CDCl_3): δ = 7.80 (d, J = 15.9 Hz, 2H, 3-H), 7.53 (d, J = 8.8 Hz, 4H, 2'-H), 6.94 (d, J = 8.8 Hz, 4H, 3'-H), 6.39 (d, J = 15.8 Hz, 2H, 2-H), 3.86 (s, 6H, 5'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ = 163.2 (C-1), 162.3 (C-4'), 148.4 (C-3), 130.6 (C-2'), 126.7 (C-1'), 114.7 (C-3'), 114.4 (C-2), 55.6 (C-5') ppm; MS (ESI, MeOH): m/z (%) = 361.1 ([M + Na] $^+$, 68), 698.9 ([2M + Na] $^+$, 100); analysis calcd for $\text{C}_{20}\text{H}_{18}\text{O}_5$ (338.36): C 71.00, H 5.36; found: C 70.72, H 5.52.

4.4.31. (E)-3-(2-Chlorophenyl)acrylic anhydride (2t)

Compound **2t** was obtained as white crystals (0.324 g, 92%); R_F = 0.88 (silica gel, chloroform); m.p. 156–157 °C; IR (ATR): ν = 1762m, 1709m, 1626w, 1588w, 1471w, 1442w, 1324w, 1259w, 1203w, 1138s, 1117s, 1051m, 1034m, 981m, 934s, 878m, 754s, 734m, 701m, 573m cm^{-1} ; UV-vis (CHCl₃): $\lambda_{\max}$ (log ϵ) = 206 (4.02), 228 (3.38), 290 (4.09) nm; ^1H NMR (500 MHz, CDCl_3): δ = 8.29 (d, J = 16.0 Hz, 2H, 3-H), 7.68 (dd, J = 7.7, 1.7 Hz, 2H, 6'-H), 7.46 (dd, J = 7.9, 1.2 Hz, 2H, 4'-H), 7.38 (td, J = 7.7, 1.7 Hz, 2H, 5'-H), 7.34–7.31 (m, 2H, 3'-H), 6.54 (d, J = 16.0 Hz, 2H, 2-H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ = 161.9 (C-1), 144.4 (C-3), 135.7 (C-2'), 132.1 (C-1'), 132.0 (C-5'), 130.5 (C-4'), 128.1 (C-6'), 127.4 (C-3'), 119.4 (C-2) ppm; MS (ESI, ASAP): m/z (%) = 369.3 ([M + Na] $^+$, 52), 385.3 ([M + K] $^+$, 3), 387.2 ([M + Na + H₂O] $^+$, 7), 401.3 ([M + Na + MeOH] $^+$, 100); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{Cl}_2\text{O}_3$ (347.19): C 62.27, H 3.48, Cl 20.42; found: C 62.01, H 3.64.

4.4.32. (E)-3-(3-Chlorophenyl)acrylic anhydride (2u)

Compound **2u** was obtained as a white solid (0.20 g, 52%); R_F = 0.98 (silica gel, chloroform); m.p. 117 °C; IR (ATR): ν = 1759m, 1709m, 1630w, 1565w, 1271w, 1204w, 1130s, 986m, 939s, 862m, 785m, 721m, 687m, 666s, 564m, 545m cm^{-1} ; UV-vis (CHCl₃): $\lambda_{\max}$ (log ϵ) = 228 (4.14), 291 (4.35) nm; ^1H NMR (400 MHz, CDCl_3): δ = 7.78 (d, J = 15.9 Hz, 2H, 3-H), 7.58–7.56 (m, 2H, 6'-H), 7.47–7.43 (m, 2H, 2'-H), 7.41 (t, J = 1.7 Hz, 2H, 5'-H), 7.40–7.35 (m, 2H, 4'-H), 6.52 (d, J = 15.9 Hz, 2H, 2-H); ^{13}C NMR (100 MHz, CDCl_3): δ = 162.0 (C-1), 147.2 (C-3), 135.6 (C-1'), 135.3 (C-3'), 131.3 (C-5'), 130.5 (C-4'), 128.4 (C-6'), 126.9 (C-2'), 118.2 (C-2) ppm; MS (ESI, ASAP): m/z (%) = 369.3 ([M + Na] $^+$, 61), 385.3 ([M + K] $^+$, 13), 387.2 ([M + Na + H₂O] $^+$, 14), 401.3 ([M + Na + MeOH] $^+$, 100); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{Cl}_2\text{O}_3$ (347.19): C 62.27, H 3.48; found: C 61.97, H 3.65.

4.4.33. (E)-3-(2-Chlorophenyl)acrylic-(E)-3-(3-Chlorophenyl)acrylic anhydride (2v)

Compound **2v** was obtained as an off-white solid (0.17 g, 44%);

$R_F = 0.98$ (silica gel, chloroform); m.p. 117 °C; IR (ATR): $\nu = 1758m$, 1708m, 1626w, 1589w, 1565w, 1472w, 1442w, 1290w, 1203w, 1136s, 1052m, 1038m, 986m, 935s, 866m, 788m, 757s, 733m, 702m, 686m, 672m, 570m cm^{-1} ; UV-vis (CHCl_3): λ_{max} ($\log \epsilon$) = 228 (3.99), 290 (4.22) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 8.28$ (dd, $J = 16.0$, 6.7 Hz, 1H, 3-H), 7.79 (dd, $J = 15.9$, 3.9 Hz, 1H, 3'-H), 7.68 (dt, $J = 7.6$, 2.1 Hz, 1H, 6''-H), 7.58–7.55 (m, 1H, 6''-H), 7.48–7.46 (m, 1H, 4''-H), 7.46–7.44 (m, 1H, 2'''-H), 7.44–7.42 (m, 1H, 5''-H), 7.41 (t, $J = 1.6$ Hz, 1H, 5''-H), 7.39–7.36 (m, 1H, 4''-H), 7.35–7.32 (m, 1H, 3''-H), 6.53 (dd, $J = 16.0$, 3.5 Hz, 2H, 2-H + 2'-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 162.0$ (C-1'), 161.9 (C-1), 147.1 (C-3'), 144.5 (C-3), 135.6 (C-1''), 135.5 (C-2'), 135.3 (C-1'''), 135.2 (C-3''), 132.2 (C-5''), 131.3 (C-5'''), 130.6 (C-4''), 130.5 (C-4'''), 128.4 (C-6''), 128.1 (C-6'''), 127.4 (C-3''), 126.9 (C-2''), 119.3 (C-2), 118.3 (C-2') ppm; MS (ESI, ASAP): m/z (%) = 369.3 ($[\text{M} + \text{Na}]^+$, 60), 385.3 ($[\text{M} + \text{K}]^+$, 7), 387.2 ($[\text{M} + \text{Na} + \text{H}_2\text{O}]^+$, 8), 401.3 ($[\text{M} + \text{Na} + \text{MeOH}]^+$, 100); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{Cl}_2\text{O}_3$ (347.19): C 62.27, H 3.48; found: C 62.11, H 3.69.

4.4.34. (E)-3-(3-Fluorophenyl)acrylic anhydride (2w)

Compound **2w** was obtained as a white solid (0.37 g, 99%); $R_F = 0.48$ (silica gel, *n*-hexane/ethyl acetate, 4:1); m.p. 192 °C; IR (ATR): $\nu = 1771m$, 1705m, 1634m, 1585m, 1486w, 1448w, 1286w, 1270w, 1238w, 1204m, 1170w, 1149w, 1065s, 985m, 964s, 878m, 863m, 784s, 716m, 665s, 562m, 486m cm^{-1} ; UV-vis (CHCl_3): λ_{max} ($\log \epsilon$) = 206 (4.19), 226 (3.71), 291 (4.22) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 7.81$ (d, $J = 15.9$ Hz, 2H, 3-H), 7.44–7.38 (m, 2H, 5'-H), 7.35 (dt, $J = 7.7$, 1.3 Hz, 2H, 6'-H), 7.30–7.26 (m, 2H, 2'-H), 7.18–7.12 (m, 2H, 4'-H), 6.52 (d, $J = 15.9$ Hz, 2H, 2-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 164.4$ (C-3'), 162.0 (C-1), 147.4 (C-3), 136.0 (C-1'), 130.9 (C-5'), 124.8 (C-6'), 118.5 (C-4'), 118.2 (C-2), 114.9 (C-2') ppm; MS (ESI, MeOH): m/z (%) = 337.1 ($[\text{M} + \text{Na}]^+$, 100), 650.8 ($[\text{2M} + \text{Na}]^+$, 48); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{F}_2\text{O}_3$ (314.29): C 68.79, H 3.85; found: C 68.51, H 4.02.

4.4.35. (E)-3-(4-Fluorophenyl)acrylic anhydride (2x)

Compound **2x** was obtained as white needles (0.18 g, 49%); $R_F = 0.89$ (silica gel, chloroform); m.p. 128 °C (lit.: [40] 125–126 °C); IR (ATR): $\nu = 3078w$, 1756m, 1700m, 1627m, 1595m, 1508s, 1412w, 1320w, 1301w, 1272w, 1227s, 1205w, 1163m, 1126m, 989s, 932m, 862m, 827s, 816s, 784m, 731m, 684m, 551m, 535m, 510s, 468m cm^{-1} ; UV-vis (CHCl_3): λ_{max} ($\log \epsilon$) = 206 (4.32), 228 (4.31), 303 (4.63) nm; ^1H NMR (400 MHz, CDCl_3): $\delta = 7.82$ (d, $J = 15.9$ Hz, 2H, 3-H), 7.61–7.55 (m, 4H, 2'-H), 7.16–7.09 (m, 4H, 3'-H), 6.45 (d, $J = 15.9$ Hz, 2H, 2-H) ppm; ^{13}C NMR (100 MHz, CDCl_3): $\delta = 165.9$ (C-4'), 162.5 (C-1), 147.5 (C-3), 130.7 (C-2'), 130.2 (C-1'), 116.6 (C-3'), 116.4 (C-2) ppm; MS (ESI, MeOH): m/z (%) = 337.1 ($[\text{M} + \text{Na}]^+$, 100), 650.8 ($[\text{2M} + \text{Na}]^+$, 44); analysis calcd for $\text{C}_{18}\text{H}_{12}\text{F}_2\text{O}_3$ (314.29): C 68.79, H 3.85; found: C 68.51, H 3.13.

Acknowledgments

Many thanks are due to Dr. D. Ströhl and his team for the NMR spectra and to Dr. R. Kluge for the MS-measurements. IR- and UV-vis spectra and optical rotations were recorded by Ms V. Simon. Cytotoxic screening (SRB assays) was performed by S. Hoenke.

References

- [1] G. Angeloni, L. Guerrini, P. Masella, M. Innocenti, M. Bellumori, A. Parenti, Characterization and comparison of cold brew and cold drip coffee extraction methods, *J. Sci. Food Agric.* 99 (1) (2019) 391–399.
- [2] S. Lafay, A. Gil-Izquierdo, Bioavailability of phenolic acids, *Phytochem. Rev.* 7 (2) (2008) 301–311.
- [3] V.L. Singleton, P. Esau, Phenolic substances in grapes and wine, and their significance, *Adv. Food Res. Suppl.* 1 (1969) 1–261.
- [4] A. Lupini, A. Sorgona, M.P. Princi, F. Sunseri, M.R. Abenavoli, Morphological and

- physiological effects of trans-cinnamic acid and its hydroxylated derivatives on maize root types, *Plant Growth Regul.* 78 (2) (2016) 263–273.
- [5] E.L.D. Niero, G.M. Machado-Santelli, Cinnamic acid induces apoptotic cell death and cytoskeleton disruption in human melanoma cells, *J. Exp. Clin. Cancer Res.* 32 (2013).
- [6] D. Raffa, B. Maggio, M.V. Raimondi, M.G. Cusimano, G. Amico, A. Carollo, P.G. Conaldi, R.L. Bai, E. Hamel, G. Daidone, 2-Cinnamamido, 2-(3-phenylpropionylamido), and 2-(3-phenylpropanamido)benzamides: Synthesis, antiproliferative activity, and mechanism of action, *Eur. J. Med. Chem.* 65 (2013) 427–435.
- [7] T. Ernawati, Y. Anita, P.D. Lotulung, M. Hanafi, Synthesis of Methyl 2-cinnamamido-3-hydroxy propanoate having activity against P388 leukemia cells, *J. Appl. Pharm. Sci.* 4 (3) (2014) 92–95.
- [8] N. Öztas, R. Kaya, A. Maras, E. Sahin, I. Gülcin, S. Göksu, Synthesis and characterization of novel bromophenols: Determination of their anticholinergic, antidiabetic and antioxidant activities, *Bioorg. Chem.* 87 (2019) 91–102.
- [9] C. Bayrak, P. Taslimi, H.S. Karaman, I. Gulcin, A. Menzek, The first synthesis, carbonic anhydrase inhibition and anticholinergic activities of some bromophenol derivatives with S including natural products, *Bioorg. Chem.* 85 (2019) 128–139.
- [10] S. Burmaoglu, A.O. Yilmaz, M.F. Polat, R. Kaya, I. Gulcin, O. Algul, Synthesis and biological evaluation of novel tris-chalcones as potent carbonic anhydrase, acetylcholinesterase, butyrylcholinesterase and α -glycosidase inhibitors, *Bioorg. Chem.* 85 (2019) 191–197.
- [11] M.J. Prince, A. Comas-Herrera, M. Knapp, M.M. Guerchet, M. Karagiannidou, World Alzheimer Report 2016 - Improving healthcare for people living with dementia: Coverage, quality and costs now and in the future, Alzheimer's Disease International, London, 2016.
- [12] J.H. Kim, N.P. Thao, Y.K. Han, Y.S. Lee, B.T.T. Luyen, H.V. Oanh, Y.H. Kim, S.Y. Yang, The insight of in vitro and in silico studies on cholinesterase inhibitors from the roots of *Cimicifuga dahurica* (Turcz.), Maxim, *J. Enzym. Inhib. Med. Ch.* 33 (1) (2018) 1174–1180.
- [13] K.D. Singh, T.B. Devi, N.I. Singh, H.D. Chanu, S. Sougrakpam, B.S. Nameirakpam, Y. Rajashekar, R.K. Labala, D. Sahoo, Biochemical efficacy, molecular docking and inhibitory effect of 2, 3-dimethylmaleic anhydride on insect acetylcholinesterase, *Sci. Rep.* 7 (1) (2017) 12483.
- [14] J.K. Makrandi, V. Kumari, A convenient synthesis of 2-styrylchromones by modified Baker-Venkataraman transformation using phase-transfer catalysis, *Synthetic Commun.* 19 (11–12) (1989) 1919–1922.
- [15] D.T. Khong, Z.M.A. Judeh, Short synthesis of phenylpropanoid glycosides calceolarioside A and syringalide B, *Synlett* 29 (8) (2018) 1079–1083.
- [16] S.-F. Li, K.A. Connors, Kinetics of trans-cinnamic anhydride reactions catalyzed by pyridine, 4-dimethylaminopyridine, and N-methylimidazole, *J. Pharm. Sci.* 70 (3) (1981) 235–238.
- [17] J.D. Habila, F.O. Shode, G.I. Ndukwe, J.O. Amupitan, A.J. Nok, Novel antimalarial agent (Cinnamic 3 β -hydroxyolean-12-en-28-carboxylic anhydride): synthesis, characterization and in vivo studies, *Afr. J. Pharm. Pharmacol.* 5 (24) (2011) 2667–2675.
- [18] B. Liu, L. Jian, G. Chen, X. Song, C. Han, J. Wang, Chemical constituents and in vitro anticancer cytotoxic activities of *Polalthia plagiocnema*, *Chem. Nat. Compd.* 49 (6) (2014) 1172–1174.
- [19] E.R.S. Nkanwen, M.D. Awouafack, J.J.K. Bankeu, H.K. Wabo, S.A.A. Mustafa, M.S. Ali, M. Lamshoeft, M.I. Choudhary, M. Spittler, P. Tane, Constituents from the stem bark of *Cinnamomum zeylanicum* welw. (Lauraceae) and their inhibitory activity toward *Plasmodium falciparum* enoyl-ACP reductase enzyme, *Rec. Nat. Prod.* 7 (4) (2013) 296–301.
- [20] E. Knoevenagel, Condensation von Malonsäure mit aromatischen Aldehyden durch ammoniak und Amine, *Ber.Dtsch. Chem. Ges.* 31 (1898) 2596–2619.
- [21] O. Döbner, Synthese der Muconsäure aus Glyoxal und Malonsäure, *Ber. Dtsch. Chem. Ges.* 35 (1902) 1147–1148.
- [22] G.L. Ellman, K.D. Courtney, V. Andres, R.M. Featherstone, A New and Rapid Colorimetric Determination of Acetylcholinesterase Activity, *Biochem. Pharmacol.* 7 (2) (1961) 88–95.
- [23] G.M. Morris, R. Huey, W. Lindstrom, M.F. Sanner, R.K. Belew, D.S. Goodsell, A.J. Olson, AutoDock4 and AutoDockTools4: automated docking with selective receptor flexibility, *J. Comput. Chem.* 30 (16) (2009) 2785–2791.
- [24] M.A. Abbasi, M. Ilyas, R. Aziz ur, A. Sonia, D. Shahwar, M.A. Raza, K.M. Khan, M. Ashraf, I. Afzal, N. Ambreen, Curcumin and its derivatives: moderate inhibitors of acetylcholinesterase, butyrylcholinesterase and trypsin, *Sci. Iran., Trans. C* 19 (2) (2012) 1580–1583.
- [25] T. Ahmed, A.-H. Gilani, Inhibitory effect of curcuminoids on acetylcholinesterase activity and attenuation of scopolamine-induced amnesia may explain medicinal use of turmeric in Alzheimer's disease, *Pharmacol., Biochem. Behav.* 91 (4) (2009) 554–559.
- [26] Z. Kalaycioglu, I. Gazioglu, F.B. Erim, Comparison of antioxidant, anticholinesterase, and antidiabetic activities of three curcuminoids isolated from *Curcuma longa* L., *Nat. Prod. Res.* 31 (24) (2017) 2914–2917.
- [27] K. Saravanan, C. Kalaiarasi, P. Kumaradhas, Understanding the conformational flexibility and electrostatic properties of curcumin in the active site of rhAChE via molecular docking, molecular dynamics, and charge density analysis, *J. Biomol. Struct. Dyn.* 35 (16) (2017) 3627–3647.
- [28] V. Tello-Franco, M.C. Lozada-Garcia, M. Soriano-Garcia, Experimental and Computational Studies on the Inhibition of Acetylcholinesterase by Curcumin and Some of its Derivatives, *Curr. Comput.-Aided Drug Des.* 9 (2) (2013) 289–298.
- [29] L. Heller, M. Kahnt, A. Loesche, P. Grabandt, S. Schwarz, W. Brandt, R. Csuk, Amino derivatives of platanic acid act as selective and potent inhibitors of butyrylcholinesterase, *Eur. J. Med. Chem.* 126 (2017) 652–668.
- [30] A. Loesche, A. Köwitsch, S.D. Lucas, Z. Al-Halabi, W. Sippl, A. Al-Harrasi, R. Csuk,

- Ursolic and oleanolic acid derivatives with cholinesterase inhibiting potential, *Bioorg. Chem.* 85 (2019) 23–32.
- [31] S. Liu, W.X. Wei, Y.B. Li, X. Liu, X.J. Cao, K.C. Lei, M. Zhou, Design, synthesis, biological evaluation and molecular docking studies of phenylpropanoid derivatives as potent anti-hepatitis B virus agents, *Eur. J. Med. Chem.* 95 (2015) 473–482.
- [32] D. Suresh, J.S. Kumar, Sandhu, Bismuth(III) Chloride-Mediated, Efficient, Solvent-Free, MWI-Enhanced Doebner Condensation for the Synthesis of (E)-Cinnamic Acids, *Synthetic Commun.* 40 (13) (2010) 1915–1919.
- [33] K. Kindler, K. Lührs, Studien über den Mechanismus chemischer Reaktionen. 22. (über Hydrierungen und über spezifische Hydrierungen mittels gebundenen Wasserstoffs 6), *Liebigs Ann. Chem.* 685 (Jun) (1965) 36–48.
- [34] J.R. Peterson, M.E. Russell, I.B. Surjasmita, Synthesis and Experimental Ionization Energies of Certain (E)-3-Arylpropenoic Acids and Their Methyl-Esters, *J. Chem. Eng. Data* 33 (4) (1988) 534–537.
- [35] Y. Asahina, T. Matsuzaki, Synthesis of 3,5-dihydroxycinnamic acid, *Yakugaku Zasshi* 509 (1924) 527–539.
- [36] F. Benington, R.C. Morin, J.L.C. Clark, Notes - Mescaline Anaologs. VI. Mescaline Homologs, *J. Org. Chem.* 21 (12) (1956) 1545–1546.
- [37] S. Prachayasittikul, S. Suphamong, A. Worachartcheewan, R. Lawung, S. Ruchirawat, V. Prachayasittikul, Bioactive Metabolites from *Spilanthes acmella* Murr, *Molecules* 14 (2) (2009) 850–867.
- [38] H. Khabazzadeh, R.M. Nejad, H. Sheibani, K. Saidi, Solvent free microwave assisted Doebner reaction: A green method for the preparation of cinnamic acids, *Catal. Commun.* 8 (9) (2007) 1411–1413.
- [39] R. Joly, R. Bucourt, E. Toromanoff, Esterifying 18-hydroxyyohimbanes, 1960, US2926167.
- [40] N. Armesto, M. Ferrero, S. Fernandez, V. Gotor, Novel enzymatic synthesis of 4-O-cinnamoyl quinic and shikimic acid derivatives, *J. Org. Chem.* 68 (14) (2003) 5784–5787.
- [41] J.J. Kim, Y.D. Park, W.S. Lee, S.D. Cho, Y.J. Yoon, Facile synthesis of carboxylic anhydrides using 4,5-dichloro-2-[(4-nitrophenyl)sulfonyl]pyridazin-3(2H)-one, *Synthesis* (2003) 1517–1520.