



# Exploiting the 7-methylimidazo[1,5-*a*]pyrazin-8(7H)-one scaffold for the development of novel chemical inhibitors for Bromodomain and Extraterminal Domain (BET) family

Yifei Yang<sup>a</sup>, Pan Chen<sup>a</sup>, Leilei Zhao<sup>b</sup>, Fangqing Zhang<sup>b</sup>, Bing Zhang<sup>b</sup>, Changliang Xu<sup>c,\*</sup>,  
Huibin Zhang<sup>b,\*</sup>, Jinpei Zhou<sup>a,\*</sup>

<sup>a</sup> Department of Medicinal Chemistry, China Pharmaceutical University, Nanjing 210009, PR China

<sup>b</sup> Center of Drug Discovery, Jiangsu Key Laboratory of Drug Discovery for Metabolic Disease, China Pharmaceutical University, Nanjing 210009, PR China

<sup>c</sup> Jiangsu Collaborative Innovation Center of Traditional Chinese Medicine Prevention and Treatment of Tumor, Nanjing University of Chinese Medicine, Nanjing, Jiangsu 210023, PR China

## ARTICLE INFO

### Keywords:

BRD4 inhibitors  
Molecular docking  
Anticancer  
BET family

## ABSTRACT

The bromodomain and extraterminal (BET) family of proteins play a crucial role in promoting gene expression of critical oncogenes. Novel BET bromodomain inhibitors with excellent potency, drug metabolism and pharmacokinetics (DMPK) properties were in strong need for development. We reported a series of potential BET inhibitors through incorporation of imidazole into pyridine scaffold. Among them, a novel BET inhibitor with 7-methylimidazo[1,5-*a*]pyrazin-8(7H)-one core, compound **28**, was considered to be the most promising for in-depth study. Compound **28** exhibited excellent BRD4-inhibitory activity with IC<sub>50</sub> value of **33 nM** and anti-proliferation potency with IC<sub>50</sub> value of **110 nM** in HL-60 (human promyelocytic leukemia) cancer cell lines. Western Blot indicated that compound **28** can effectively trigger apoptosis in BxPc3 cells by modulating the intrinsic apoptotic pathway. In conclusion, these results suggested that compound **28** has merely potential for leukemia treatment.

## 1. Introduction

The acetylation of lysine residues were proposed to play crucial roles in post-translational modifications of histones [1]. Ever since 1992, when the bromodomains comprised of 110-amino acid were first identified as 'readers' of *N*-acetyl-lysine modifications in nucleosomal chromatin, there are a lot of researches focused on the mechanism of bromodomains in epigenetics [2–4]. The human genome codes 46 proteins containing 61 BRDs [5], including the bromodomain and extra-terminal (BET) protein family, which contains four members: BRD2, BRD3, BRD4 and BRDT [6]. These proteins comprise of conservative tandem bromodomains that recognize acetylated lysine residues on histone tails and recruit transcriptional factors to promote gene expression [3,7,8]. For example, BRD4 recruits positive transcription elongation factor (P-TEFb) to acetylated chromatin and plays a role in activating the cyclin dependent kinase (CDK9) subunit,

whereby it acts as a vital transcription moderator [9].

BET proteins were confirmed as a promising therapeutic target for competitive inhibitors displacing bromodomains from chromatin to influence changes in gene expression. Therefore great interests and attention were attracted for the development of novel BET inhibitors for cancer treatment [10–12]. Several BET inhibitors with different scaffolds have been reported since the first small molecule BRD4 inhibitor JQ1 was disclosed [13]. There are some inhibitors undergoing clinical trials for the treatment of different diseases, such as OTX015 (Merck & Co. Phase I/II) [14], I-BET762 (GlaxoSmithKline. Phase II) [15], BI-2536 (Boehringer Ingelheim. Phase II) [16], ABBV-075 (AbbVie. Phase I) [17–19] (Fig. 1). However, the major indications of these inhibitors were hematologic malignancies, such as multiple myeloma (MM) [20].

Recent study also revealed the potency of BET inhibitors in solid tumor, such as pancreatic cancer [21]. Pancreatic ductal adenocarcinoma (PDA) is one of the most aggressive cancers with a median

*Abbreviations:* BET, Bromodomain and Extraterminal Domain; DMPK, drug metabolism and pharmacokinetics; P-TEFb, positive transcription elongation factor; CDK9, cyclin dependent kinase; PDA, Pancreatic ductal adenocarcinoma; AlphaScreen, Amplified Luminescent Proximity Homogeneous; HEXIM1, hexamethylene bisacetamide inducible protein 1; IHC, immunohistochemical; WB, Western Blot

\* Corresponding authors.

*E-mail addresses:* [cl.xu81@gmail.com](mailto:cl.xu81@gmail.com) (C. Xu), [zhanghb80@163.com](mailto:zhanghb80@163.com) (H. Zhang), [jpzhou668@163.com](mailto:jpzhou668@163.com) (J. Zhou).

<https://doi.org/10.1016/j.bioorg.2019.103044>

Received 1 May 2019; Received in revised form 31 May 2019; Accepted 4 June 2019

Available online 11 June 2019

0045-2068/ © 2019 Elsevier Inc. All rights reserved.




Fig. 1. Structures of known BET bromodomain inhibitors.


survival time of 6 months, with less than 5% survival 5 years after diagnosis [22]. PDA surpassed only by lung cancer, will become the second leading cause of cancer-related deaths in the United States by 2030 [23]. According to the important role that BET proteins play in the progression and evolution of pancreatic cancer, we designed and synthesized a series of imidazolopyridone derivatives containing the moiety 7-methylimidazo [1, 5-a]pyrazin-8(7H)-one via scaffold hopping as BET inhibitors. Then their activities in AlphaScreen binding assays and effects on cell viability were evaluated in vitro. Cell cycle analysis was performed to characterize the molecular mechanisms underlying the cytotoxic effect of compound 28 in BxPC-3 cell.

### 1.1. Synthesis


Target compounds (15–17, 25–37) were synthesized as outlined in Schemes 1 and 2. All compounds were purified by chromatography and purity was checked by HPLC before biological evaluation (purity was > 98%). The structures were confirmed by <sup>1</sup>H NMR, <sup>13</sup>C NMR

spectrum, and mass spectra. Some representative spectrometry was shown in Supporting Information. As depicted in Scheme 1, commercially available cytosine 8 reacted with bromine in acetic acid to give 6-amino-5-bromo-1H-pyrimidin-2-one (9). Cyclization of compound 9 with chloroacetaldehyde in dimethyl formamide was utilized to obtain intermediate 10. The key intermediate 11 was prepared by methylation of intermediate 10 in the presence of dimethylsulfate. Then Suzuki-Miyaura cross coupling reaction was used to get compound 12, followed by reacting with different phenol in the presence of potassium carbonate. Iron-catalyzed reduction of the nitro group of 13 in the presence of ammonium chloride followed by functionalization of the resulting aniline 14 provided facile access to sulfonamides 15–17.

As shown in Scheme 2, 5-bromopyrazin-2(1H)-one (19) was got via 5-bromopyrazin-2-amine (18) reacting with sodium nitrite in concentrated sulfuric acid. Methylation reaction was utilized to prepare 5-bromo-1-methylpyrazin-2(1H)-one (20) followed by cyclization with tosylmethyl isocyanide to give 5-bromo-7-methylimidazo[1,5-a]pyrazin-8(7H)-one (21). The bond between imidazolopyridone and biaryl



Scheme 1. Reagents and conditions: (i) Br<sub>2</sub>, CH<sub>3</sub>COOH, 70 °C, 85%; (ii) chloroacetaldehyde, DMF, 100 °C, 65%; (iii) Dimethylsulfate, Cs<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>CN, 60 °C, 49%; (iv) Pd(PPh<sub>3</sub>)<sub>4</sub>, Na<sub>2</sub>CO<sub>3</sub>, toluol/water/MeOH, 110 °C, 69%; (v) phenol, K<sub>2</sub>CO<sub>3</sub>, DMF, 80 °C, 84%; (vi) Fe, NH<sub>4</sub>Cl, H<sub>2</sub>O/EtOH, 100 °C, 83%; (vii) RSO<sub>2</sub>Cl, NET<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, then 2 N NaOH, 1,4-dioxane, 70 °C, 58–87%.



**Scheme 2.** Reagents and conditions: (i) NaNO<sub>2</sub>, conc. H<sub>2</sub>SO<sub>4</sub>, 0 °C to 45 °C, 91%; (ii) dimethyl sulfate, Cs<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>CN, 60 °C, 81%; (iii) Tosylmethyl isocyanide, NaH, THF, 0 °C to rt; (iv) Pd(PPh<sub>3</sub>)<sub>4</sub>, Na<sub>2</sub>CO<sub>3</sub>, toluol/water/MeOH, 110 °C, 68%; (v) phenol, K<sub>2</sub>CO<sub>3</sub>, DMF, 80 °C, 95%; (vi) Fe, NH<sub>4</sub>Cl, H<sub>2</sub>O/EtOH, 100 °C, 83%; (vii) RSO<sub>2</sub>Cl, NEt<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, then 2N NaOH, 1,4-dioxane, 70 °C, 58–87%.

ether was formed through Suzuki-Miyaura cross coupling reaction. Then an analogous approach depicted in Scheme 1 provided access to the target compounds (25–37).

## 2. Results and discussion

### 2.1. Design strategy

Firstly, to further improve the potency of pyridone-based BET bromodomain inhibitors such as Fragment 9 [17], imidazole was introduced to afford target compounds (15–17) with 5,6-dihydroimidazo[1,2-c]pyrimidin scaffold. Biological activities were evaluated using an Amplified Luminescent Proximity Homogeneous (AlphaScreen) binding assay and three complementary cellular assays. The AlphaScreen binding assay was used to evaluate the efficacy of compounds for a construct containing BRD4(BD1) and BRD4(BD2). Growth of cancer cell lines are dependent on BET proteins [24], the effects of compounds on cancer cell proliferation were measured using the lymphoblast-like cell line (Raji), promyelocytic cell line (HL-60) and human pancreatic cancer cell line (BxPC-3). However, potency of BET bromodomain inhibitors with 5,6-dihydroimidazo[1,2-c]pyrimidin scaffold such as 17 was relatively poor (Table 1). To further improve the biochemical efficiency, compound 17 was docked into BRD4 (BD1) and superimposed with ABBV-075 via Glide docking (PDB id: 3P50). Docking studies revealed that although the carbonyl moieties of these cores were situated at an ideal distance away from the Asn433 NH<sub>2</sub> group (2.95 Å), there was not an efficient interaction built between the Asn433 amide carbonyl and

moieties of compound 17 (Fig. 2). However, 7-methylimidazo[1,5-a]pyrazin-8(7H)-one-based compounds were designed and evaluated in order to provide a bidentate contact and improve the binding potency. Compared to compound 17, novel imidazolopyridone of 28 improved both biochemical and cellular activity by 8–20-fold (Table 1).

In order to elaborate the specific interaction between the imidazolopyridone core and BET bromodomain, we performed docking studies for compound 28 and ABBV-075. Docking studies indicated that the ligand-protein interactions of two molecules were similar. The 7-methylimidazo[1,5-a]pyrazin-8(7H)-one acted as a KAc mimic, with the methyl group occupying the small hydrophobic pocket. The pyridone carbonyl formed a water mediated hydrogen bond with the NH<sub>2</sub> of TRY97, and accepted a hydrogen bond from the conserved Asn residue (ASN140) with an ideal 2.83 Å distance measured. In addition, the phenyl ether moiety of compound 28 offers valuable interactions by occupying a hydrophobic part of the WPF shelf and ethylsulfonamide moiety is directed into the ZA channel. The key hydrogen bond with Asn140 and the additional water mediated interactions would be important for improving affinity. Gratifyingly, the second imidazolopyridone core exhibited more remarkable binding effects than the first motif, as evaluated by an improvement in IC<sub>50</sub> from 270 to 33 nM (28 vs 17) (see Fig. 3.).


The 7-methylimidazo[1,5-a]pyrazin-8(7H)-one core exhibited effective and potent inhibitory activity of BET protein, we evaluated the SAR at other sites of the compounds for further improving potency (Table 2). Compound 28 with the BRD4(BD1) IC<sub>50</sub> of 33 nM was selected as the lead compound. Different substituents (compound 26–37)

**Table 1**  
Biochemical and Cellular Potency of Compounds 15, 16, 17 and 28.

Compound ID	15	16	17	28
BRD4(BD1) Alpha Screen IC <sub>50</sub> (μM) <sup>a</sup>	0.57 ± 0.022	0.43 ± 0.012	0.27 ± 0.013	0.033 ± 0.005
BRD4(BD2) Alpha Screen IC <sub>50</sub> (μM)	0.48 ± 0.031	0.39 ± 0.024	0.31 ± 0.011	0.025 ± 0.009
HL-60 Proliferation IC <sub>50</sub> (μM) <sup>b</sup>	4.3 ± 0.025	3.9 ± 0.019	2.2 ± 0.028	0.11 ± 0.012
Raji Proliferation IC <sub>50</sub> (μM) <sup>b</sup>	3.8 ± 0.021	5.1 ± 0.016	1.8 ± 0.024	NT
BxPC-3 Proliferation IC <sub>50</sub> (μM) <sup>b</sup>	> 10	> 10	5.6	NT

<sup>a</sup> BRD4 AlphaScreen IC<sub>50</sub> values are reported as the mean ± SD value of three independent determinations.

<sup>b</sup> IC<sub>50</sub> values presented are the mean derived from three measurements.



**Fig. 2.** Molecular modeling of compound **17** with BRD4(1) (PDB id: **3P50**). (A) Docking conformation of compound **17** in BRD4(1) (PDB id: **3P50**). (B) Docking conformation of **ABBV-075** in BRD4(1) (PDB id: **3P50**). (C) Superimposition docking conformation of compound **17** (Blue) and **ABBV-075** (Green) in BRD4(1) (PDB id: **3P50**). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

were introduced into the phenyl group which was located at the WPF shelf to further improve the binding affinity. Firstly, pyridines (compound **26**) was incorporated here to reduce lipophilicity and improve the developability profile, though substantial potency was significantly reduced compared to the phenyl ethers. The mono fluoro substituted compound **27** exhibited moderate improvement. The biochemical activities of compound **36** and **37** with mono bromo substituent were also limited. These results clearly indicated the importance of the moderate groups located at WPF shelf for the affinity. In addition, ethyl sulfonamide was preferred over methyl sulfonamide based upon its modestly improved potency. With these encouraging results in hand, the anti-proliferation activities were further assessed in several tumor cell lines (Table 2).


## 2.2. Anti-proliferation in cancer cell lines

Furthermore, we assessed the effects of compounds on cancer cell proliferation in the lymphoblast-like cell line (Raji), promyelocytic cell line (HL-60) and human pancreatic cancer cell line (BxPC-3). As our previous reports, BET inhibitors show more robust anti-proliferation activities in hematologic malignancies than solid tumor [25]. In this work, all compounds exhibited preferred effects in HL-60 cell lines. It is intriguing that compounds **27** and **28** exhibited satisfying anti-proliferation in solid cancer cell, especially in human pancreatic cancer cell line (BxPC-3). Among them, compound **28** showed anti-proliferation Inh% of 43.2 in BxPC-3 cell line with 2  $\mu\text{M}$  (Table 2). In addition, cell cycle analysis was performed to characterize the molecular mechanisms underlying the cytotoxic effect of compound **28** in BxPC-3 cell. BxPC-3

cell was cultured with DMSO control or varied concentration of compound **28** (2, 4, 8  $\mu\text{M}$ ) respectively. As shown in Fig. 4A, compound **28** treatment resulted in a remarkable increase in G1 cells at 48 h, which was consistent with previous report [20]. These results indicated that compounds **27** and **28** were demonstrated promising effects in the therapy of pancreatic cancer.


## 2.3. Induce apoptosis via the mitochondrial pathways in pancreatic cancer cells

To further clarify the mechanism of compound **28** triggering apoptosis in pancreatic cancer cells, we examined the expression pattern of BRD4 and several key proteins in the intrinsic apoptotic pathway. The results showed that both compound **28** and **JQ1** up-regulated BRD4 protein, perhaps related to enrichment occurred in signaling regulation. Compound **28** downregulated c-Myc at 24 h, consistent with **JQ1**. Meanwhile, compound **28** downregulated BCL-xL in BxPc3 pancreatic cancer cell lines to trigger apoptosis by modulating the intrinsic apoptotic pathway. Early apoptotic events such as caspase activation (caspase 3 and caspase 9) and PARP cleavage were already apparent at the low concentration of compound **28** (Fig. 5). In addition, the result exhibited that the hexamethylene bisacetamide inducible protein 1 (HEXIM1), well-characterized PD markers reported by Lin et al. was significantly up-regulated [26]. Taken together, these results indicated that compound **28** can effectively trigger apoptosis in BxPC3 cells by modulating the intrinsic apoptotic pathway.



**Fig. 3.** Molecular modeling of compound **28** with BRD4(1) (PDB id: **3P50**). (A) Docking conformation of compound **28** in BRD4(1) (PDB id: **3P50**). (B) Superimposition docking conformation of compound **28** (Yellow) and **AbbV-075** (Green) in BRD4(1) (PDB id: **3P50**). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)


**Table 2**  
Biochemical and cellular activities of compound 25–37.




ID	R <sub>1</sub>	R <sub>2</sub>	BRD4/BD1(IC <sub>50</sub> nM) <sup>a</sup>	BRD4/BD2 (IC <sub>50</sub> nM)	HL60 <sup>b</sup> IC <sub>50</sub> (μM)	Raji Inh%(2 μM)	BxPC3 Inh% (2 μM)
25	2,4-di-F-phenyl	Me	49	76	0.25	40.8	20.3
26	3-pyridyl	Et	61	73	7.7	42.1	2.2
27	m-F-phenyl	Et	33	27	0.2	49.3	46.1
28	2,4-di-F-phenyl	Et	33	25	0.11	46.23	43.2
29	3-pyridyl	Me	120	280	6.1	31.4	9.4
30	3,4,5-tri-F-phenyl	Me	90	520	1.5	45.8	7.9
31	o-F-phenyl	Me	46	NT	0.2	42.9	27.5
32	m-F-phenyl	Me	52	NT	0.64	45.2	27.8
33	2-Me-5-F-phenyl	Me	37	110	0.38	49.9	27.9
34	2-F-5-Me-phenyl	Me	74	NT	1.00	49.7	22.4
35	p-isopropylphenyl	Me	100	84	0.39	27.1	30.4
36	p-Br-phenyl	Et	72	NT	0.27	41.3	25.8
37	m-Br-phenyl	Et	52	NT	0.55	38.7	24.3
JQ1			87	79	0.87	47.9	31.1
ABBV-075			39	47	0.12	51.2	42.7

<sup>a</sup> BRD4 AlphaScreen IC<sub>50</sub> values are reported as the mean ± SD value of three independent determinations.

<sup>b</sup> IC<sub>50</sub> values presented are the mean derived from three measurements.



**Fig. 4.** Cellular effects of compound 28 in BxPC-3 cell lines. (A) Cell cycle analysis was performed on BxPC-3 cell cultured with DMSO or varied concentration of compound 28 (2, 4, 8 μM) for 48 h. (B) Data was demonstrated by graph.



**Fig. 5.** Induce apoptosis via the mitochondrial pathways in pancreatic cancer cells. BxPc3 cells were treated with compound 28 for 24 h, and then the expression of BRD4, c-Myc, PARP, Bcl-xl, caspase 9, caspase 3 and HEXIM1 were examined by western blot. “+” means 5 μM, “++” means 10 μM, “+++” means 15 μM.

### 3. Conclusion

We have designed and synthesized a series of BET inhibitors with 7-methylimidazo[1,5-a]pyrazin-8(7H)-one scaffold. Optimization of bidentate contact afforded compounds with high BRD4 potency and excellent physicochemical properties. Compound 28 exhibited excellent BRD4-inhibitory activity with IC<sub>50</sub> value of 33 nM and anti-proliferation potency with IC<sub>50</sub> value of 110 nM in HL-60 cell lines. Western Blot (WB) indicated that compound 28 can effectively trigger apoptosis in BxPc3 cells by modulating the intrinsic apoptotic pathway. All of these results demonstrated that compound 28 exhibited excellent preclinical development properties and was progressed into further investigation.

### 4. Experimental section

#### 4.1. Chemistry. General methods

All chemical reagents were commercially purchased and used without further purification. Column chromatography was performed

on silica gel 60 (200–300 mesh). Thin-layer chromatography (TLC): silica gel 60F254 plates (250 nm; Qingdao Ocean Chemical Company, Qingdao, China) were used to monitor the reactions. Capillary tubes were used for the measurement of melting points. HPLC analysis (UV detector, wavelength: 272 nm) was employed to evaluate purity of the target compounds and all final compounds were purified to  $\geq 95\%$  purity.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra: Bruker ACF-300Q apparatus (300 MHz for  $^1\text{H}$  NMR and 75 MHz for  $^{13}\text{C}$  NMR), in DMSO- $d_6$  or  $\text{CDCl}_3$  unless otherwise indicated. The data of the chemical shifts were showed in  $\delta$  values (ppm) and the coupling constants ( $J$ ) in Hz. Mass spectrometry (MS): Hewlett-Packard 1100 LC/MSD spectrometer; elemental analyses: CHNO-Rapid instrument.

#### 4.1.1. 4-Amino-5-bromopyrimidin-2(1H)-one (9)

To a solution of 4-aminopyrimidin-2(1H)-one (2 g, 18 mmol) in acetic acid (10 mL) was added bromine (3.6 g, 22.5 mmol). The reaction mixture was stirred at 70 °C for 8 h. The reaction mixture was diluted with  $\text{H}_2\text{O}$  (30 mL) and the light yellow precipitate was filtered and washed with water. The solid was then dried under reduced pressure to give the title compound (2.91 g, 84.8% yield).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  10.86 (s, 1H), 7.75 (s, 1H), 6.85 (s, 1H).

#### 4.1.2. 8-Bromoimidazo[1,2-c]pyrimidin-5(6H)-one (10)

To a solution of **9** (1 g, 5.26 mmol) in DMF (10 mL) was added 40% chloroacetaldehyde (1.24 g, 6.32 mmol) at rt. The reaction mixture was stirred at 100 °C for 10 h. The reaction mixture was diluted with  $\text{H}_2\text{O}$  (20 mL) and the white precipitate was filtered and washed with water. The solid was then dried under reduced pressure to give the title compound (0.73 g, 64.8% yield).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  11.92 (s, 1H), 7.89 (d,  $J = 1.0$  Hz, 1H), 7.65 (s, 1H), 7.42 (d,  $J = 1.0$  Hz, 1H).

#### 4.1.3. 8-Bromo-6-methylimidazo[1,2-c]pyrimidin-5(6H)-one (11)

A solution of **10** (0.25 g, 1.17 mmol) and  $\text{K}_2\text{CO}_3$  (0.32 g, 2.34 mmol) in acetonitrile (20 mL) was stirred at ambient temperature. To this solution was added dimethylsulfate (0.17 g, 0.14 mmol) and the reaction mixture was stirred at 70 °C for 4 h. After cooling to the room temperature, the mixture was filtered through a celite pad, and the filtrate was concentrated under reduced pressure. The crude compound was purified by flash chromatography (silica gel, PE/EA = 10:1–5:1) to afford the title compound (0.13 g, 48.8% yield).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  7.92 (d,  $J = 12.7$  Hz, 2H), 7.42 (s, 1H), 3.50 (s, 3H).

#### 4.1.4. 8-(2-Fluoro-5-nitrophenyl)-6-methylimidazo[1,2-c]pyrimidin-5(6H)-one (12)

A mixture of compound **11** (0.3 g, 1.32 mmol), (2-fluoro-5-nitrophenyl)boronic acid (0.495 g, 1.97 mmol), Pd(PPh $_3$ ) $_4$  (0.152 g, 0.133 mmol), and  $\text{Na}_2\text{CO}_3$  (0.418 g, 3.95 mmol) in toluol (3 mL), water (3 mL) and EtOH (1 mL) and methanol (1.5 mL) was stirred at 110 °C under nitrogen atmosphere for 24 h. Then the mixture was diluted with  $\text{H}_2\text{O}$  and EtOAc, and the insoluble material was filtered through Celite. The organic layer was separated and purified by flash chromatography (silica gel, PE/EA = 5:1–2:1) to afford the title compound (0.26 g, 68.57% yield) as a white solid.  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  8.85 (d,  $J = 2.9$  Hz, 1H), 8.38 (d,  $J = 3.1$  Hz, 1H), 7.96 (s, 2H), 7.69 (d,  $J = 9.3$  Hz, 1H), 7.47 (d,  $J = 1.4$  Hz, 1H), 3.61 (s, 3H).

#### 4.1.5. 8-(2-(2,6-Dimethylphenoxy)-5-nitrophenyl)-6-methylimidazo[1,2-c]pyrimidin-5(6H)-one (13a)

A mixture of **12** (0.14 g, 0.49 mmol), 2,6-dimethylphenol (0.12 g, 0.97 mmol) and  $\text{K}_2\text{CO}_3$  (0.13 g, 0.97 mmol) in DMF (4 mL) was stirred at 80 °C for 12 h. Then the mixture was diluted with  $\text{H}_2\text{O}$  and the white precipitate was filtered and washed with water. The solid was then dried under reduced pressure to give the title compound (0.16 g, 84.4% yield).  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  8.63 (d,  $J = 2.8$  Hz, 1H), 8.21 (dd,  $J = 9.1, 2.9$  Hz, 1H), 7.96–7.84 (m, 2H), 7.45 (d,  $J = 1.5$  Hz, 1H), 7.25–7.10 (m, 3H), 6.55 (d,  $J = 9.1$  Hz, 1H), 3.62 (s, 3H), 2.07 (s, 6H).

#### 4.1.6. 8-(5-Amino-2-(2,6-dimethylphenoxy)phenyl)-6-methylimidazo[1,2-c]pyrimidin-5(6H)-one (14a)

To a solution of **13a** (0.1 g, 0.26 mmol) and  $\text{NH}_4\text{Cl}$  (0.054 g, 1.02 mmol) in 50% EtOH was added Iron powder (0.057 g, 1.02 mmol) at ambient temperature. Then the mixture was stirred at 100 °C for 2 h. Then the mixture was diluted with  $\text{H}_2\text{O}$  and EtOAc, and the insoluble material was filtered through Celite. The organic layer was separated and purified by flash chromatography (silica gel, PE/EA = 8:1–2:1) to afford the title compound (0.077 g, 83.41%). The crude product was further reacted directly for next step without further purification.

#### 4.1.7. N-(4-(2,6-Dimethylphenoxy)-3-(6-methyl-5-oxo-5,6-dihydroimidazo[1,2-c]pyrimidin-8-yl)phenyl)methanesulfonamide (15)

To a solution of compound **14a** (0.12 g, 0.332 mmol) and triethylamine (0.13 g, 1.33 mmol) in dichloromethane (6.0 mL) was added dropwise methanesulfonyl chloride (0.13 g, 0.99 mmol) at ice bath. The reaction mixture was stirred for 5 h at ambient temperature. Then the solution was concentrated under reduced pressure. The residue was diluted with a mixture of 1,4-dioxane (6 mL) and 2 M aqueous sodium hydroxide (3 mL) and heated for 30 min at 70 °C. Then the mixture was diluted with  $\text{H}_2\text{O}$  and EtOAc, brought to pH 7 with 1 M HCl. The organic layer was separated and purified by flash chromatography (silica gel, PE/EA = 3:1–1:1) to afford the title compound (0.12 g, 81%) as a white solid. m.p.: 126–128 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.53 (s, 1H), 7.86 (s, 1H), 7.71 (s, 1H), 7.55 (s, 1H), 7.42 (s, 1H), 7.09 (d,  $J = 8.4$  Hz, 4H), 6.33 (d,  $J = 8.8$  Hz, 1H), 3.61 (s, 3H), 3.00 (s, 3H), 2.06 (s, 6H).  $^{13}\text{C}$  NMR (75 MHz, DMSO)  $\delta$  152.43, 150.83, 146.38, 145.12, 133.68, 132.89, 131.85, 131.19, 129.48, 125.99, 125.71, 123.25, 122.27, 113.13, 107.93, 60.21, 40.86, 40.59, 40.31, 40.03, 39.75, 39.47, 39.20, 36.97, 21.20, 16.41, 14.53. MS (ESI,  $m/z$ ): 439.4[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{22}\text{H}_{22}\text{N}_4\text{O}_4\text{S}$ : C, 60.26; H, 5.06; N, 12.78. Found: C, 60.40; H, 5.11; N, 12.58.

#### 4.1.8. N-(4-(2,6-Dimethylphenoxy)-3-(6-methyl-5-oxo-5,6-dihydroimidazo[1,2-c]pyrimidin-8-yl)phenyl)methanesulfonamide (16)

Compound **16** was prepared in a similar procedure of compound **15**. A white solid was provided. Yield: 84.9%. m.p.: 130–132 °C;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.07 (s, 1H), 7.82 (dd,  $J = 12.3, 2.1$  Hz, 2H), 7.56–7.43 (m, 2H), 7.26 (s, 1H), 7.08 (s, 3H), 6.82 (dd,  $J = 8.8, 2.7$  Hz, 1H), 6.22 (d,  $J = 8.8$  Hz, 1H), 3.67 (s, 3H), 3.15 (q,  $J = 7.4$  Hz, 2H), 2.03 (s, 6H), 1.41 (t,  $J = 7.4$  Hz, 3H).  $^{13}\text{C}$  NMR (75 MHz, DMSO)  $\delta$  185.60, 177.60, 152.24, 150.86, 146.38, 145.12, 133.67, 132.89, 131.84, 131.18, 129.49, 125.66, 122.88, 122.27, 113.15, 113.09, 107.91, 105.33, 90.13, 54.72, 45.47, 40.87, 40.59, 40.32, 40.04, 39.76, 39.48, 39.21, 36.98, 16.42, 8.49. MS (ESI,  $m/z$ ): 453.3[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{23}\text{H}_{24}\text{N}_4\text{O}_4\text{S}$ : C, 61.05; H, 5.35; N, 12.38. Found: C, 61.18; H, 5.41; N, 12.63.

#### 4.1.9. N-(4-(2,4-Difluorophenoxy)-3-(6-methyl-5-oxo-5,6-dihydroimidazo[1,2-c]pyrimidin-8-yl)phenyl)methanesulfonamide (17)

Compound **17** was prepared in a similar procedure of compound **15**. A white solid was provided. Yield: 83.2%; m.p.: 129–131 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.85 (s, 1H), 7.83 (s, 1H), 7.69 (s, 1H), 7.52 (d,  $J = 2.1$  Hz, 1H), 7.37 (d,  $J = 0.9$  Hz, 2H), 7.20 (dd,  $J = 15.3, 8.8$  Hz, 2H), 7.04 (d,  $J = 7.9$  Hz, 1H), 6.90 (d,  $J = 8.7$  Hz, 1H), 3.55 (s, 3H), 3.12 (dd,  $J = 14.4, 7.1$  Hz, 2H), 1.23 (t,  $J = 7.2$  Hz, 3H).  $^{13}\text{C}$  NMR (75 MHz, DMSO- $d_6$ )  $\delta$  199.63, 188.26, 147.35, 145.11, 144.54, 134.25, 133.97, 133.14, 127.32, 124.55, 123.21, 122.76, 122.64, 120.04, 117.18, 112.94, 112.27, 109.68, 106.38, 105.97, 96.93, 78.57, 56.16, 45.61, 44.89, 40.80, 40.60, 40.49, 40.33, 40.22, 40.05, 39.93, 39.77, 39.59, 39.49, 39.34, 37.43, 36.84, 31.43, 19.45, 11.77, 8.45. MS (ESI,  $m/z$ ): 461.3[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{21}\text{H}_{18}\text{F}_2\text{N}_4\text{O}_4\text{S}$ : C, 54.78; H, 3.94; N, 12.17. Found: C, 54.18; H, 4.11; N, 12.23.

The synthesis methods of **19–21** would not be shown here because their preparation has been presented in detail in our previous study [25].

## 4.1.10. 5-Bromopyrazin-2(1H)-one (19)

Yellow solid; Yield: 55.7%;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  12.24 (s, 1H), 8.10 (s, 1H), 7.92 (s, 1H).

## 4.1.11. 5-Bromo-1-methylpyrazin-2(1H)-one (20)

Yellow solid; Yield: 83.4%;  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.97 (s, 1H), 7.32 (s, 1H), 3.52 (s, 3H).

## 4.1.12. 5-Bromo-7-methylimidazo[1,5-a]pyrazin-8(7H)-one (21)

Yellow solid; Yield: 72.5%;  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.08 (d,  $J = 3.3$  Hz, 2H), 6.62 (s, 1H), 3.48 (s, 3H).

## 4.1.13. 5-(2-Fluoro-5-nitrophenyl)-7-methylimidazo[1,5-a]pyrazin-8(7H)-one (22)

The synthesis method of **22** was similar to the preparation of **12**. White solid; Yield: 68.3%;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  8.57 (d,  $J = 5.7$  Hz, 1H), 8.51 (dd,  $J = 8.6, 3.9$  Hz, 1H), 8.15 (s, 1H), 7.91 (s, 1H), 7.75 (t,  $J = 9.1$  Hz, 1H), 7.30 (s, 1H), 3.42 (s, 3H).

## 4.1.14. 5-(2-(2,4-Difluorophenoxy)-5-nitrophenyl)-7-methylimidazo[1,5-a]pyrazin-8(7H)-one (23a)

The synthesis method of **23a** was similar to the preparation of **13a**. White solid; Yield: 95.2%; The crude product was further reacted directly for next step without further purification.

## 4.1.15. 5-(5-Amino-2-(2,4-difluorophenoxy)phenyl)-7-methylimidazo[1,5-a]pyrazin-8(7H)-one (24a)

The synthesis method of **24a** was similar to the preparation of **14a**. White solid; Yield: 83.4%;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  7.83 (s, 1H), 7.76 (s, 1H), 7.28–7.21 (m, 1H), 7.04–6.98 (m, 1H), 6.94–6.89 (m, 2H), 6.81 (d,  $J = 9.5$  Hz, 1H), 6.76–6.71 (m, 2H), 5.21 (s, 2H), 3.35 (s, 3H).

The synthesis method of **25–37** was similar to the preparation of **15–17**.

## 4.1.16. N-(4-(2,4-Difluorophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)phenyl)methanesulfonamide (25)

m.p. > 220–222 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.85 (s, 1H), 7.98 (s, 1H), 7.82 (s, 1H), 7.47–7.27 (m, 4H), 7.10 (s, 2H), 6.94 (d,  $J = 8.7$  Hz, 1H), 3.40 (s, 3H), 3.05 (s, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.89, 152.40, 139.16, 138.96, 134.57, 133.04, 129.89, 124.66, 124.49, 123.64, 123.50, 123.20, 122.09, 120.77, 117.56, 113.28, 112.72, 112.42, 106.09, 40.85, 40.57, 40.29, 40.02, 39.74, 39.46, 39.19, 34.61. MS (ESI,  $m/z$ ): 447.0[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{16}\text{F}_2\text{N}_4\text{O}_4\text{S}$ : C, 53.81; H, 3.61; N, 12.55. Found: C, 53.97; H, 3.51; N, 12.42.

## 4.1.17. N-(3-(7-Methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)-4-(pyridin-3-yloxy)phenyl)ethanesulfonamide (26)

m.p.: 128–130 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.98 (s, 1H), 8.29 (dd,  $J = 4.6, 1.2$  Hz, 1H), 8.24 (d,  $J = 2.7$  Hz, 1H), 8.03 (s, 1H), 7.78 (s, 1H), 7.42–7.29 (m, 4H), 7.12 (d,  $J = 8.5$  Hz, 1H), 7.04 (s, 1H), 3.35 (s, 3H), 3.18 (q,  $J = 7.3$  Hz, 2H), 1.24 (t,  $J = 7.3$  Hz, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.81, 153.11, 151.10, 145.23, 140.95, 135.40, 133.13, 129.91, 125.94, 124.94, 123.78, 123.16, 122.39, 122.15, 120.56, 113.17, 45.75, 40.86, 40.58, 40.31, 40.03, 39.75, 39.47, 39.19, 34.58, 8.53. MS (ESI,  $m/z$ ): 426.1[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{19}\text{N}_5\text{O}_4\text{S}$ : C, 56.46; H, 4.50; N, 16.46. Found: C, 56.58; H, 4.91; N, 16.53.

## 4.1.18. N-(4-(3-Fluorophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)phenyl)ethanesulfonamide (27)

m.p. > 220–222 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.95 (s, 1H), 7.99 (s, 1H), 7.77 (s, 1H), 7.42–7.37 (m, 2H), 7.29 (dd,  $J = 15.2, 8.2$  Hz, 1H), 7.17–7.11 (m, 1H), 7.00 (s, 1H), 6.93–6.85 (m, 1H), 6.84–6.71 (m, 2H), 3.34 (s, 3H), 3.18 (q,  $J = 7.3$  Hz, 2H), 1.24 (t,  $J = 7.3$  Hz, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.81, 150.96,

135.40, 133.12, 131.67, 131.54, 129.87, 123.72, 123.17, 122.55, 122.11, 120.98, 114.58, 113.20, 110.97, 110.69, 106.34, 106.01, 45.76, 40.86, 40.58, 40.30, 40.02, 39.74, 39.47, 39.19, 34.56, 8.52. MS (ESI,  $m/z$ ): 443.0[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{21}\text{H}_{19}\text{FN}_4\text{O}_4\text{S}$ : C, 57.01; H, 4.33; N, 12.66. Found: C, 56.84; H, 4.25; N, 12.89.

## 4.1.19. N-(4-(2,4-Difluorophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)phenyl)ethanesulfonamide (28)

m.p. > 220–222 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.89 (s, 1H), 7.94 (s, 1H), 7.80 (s, 1H), 7.45–7.24 (m, 4H), 7.08 (s, 2H), 6.93 (d,  $J = 8.8$  Hz, 1H), 3.40 (s, 3H), 3.15 (q,  $J = 7.2$  Hz, 2H), 1.23 (t,  $J = 7.3$  Hz, 3H);  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.88, 152.18, 134.61, 132.95, 129.89, 124.38, 124.17, 123.99, 123.63, 123.48, 123.21, 122.06, 120.78, 117.61, 113.26, 112.33, 106.45, 106.09, 105.79, 45.73, 40.87, 40.59, 40.31, 40.03, 39.75, 39.48, 39.20, 34.62. **8.51.MS** (ESI,  $m/z$ ): 461.0[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{21}\text{H}_{18}\text{F}_2\text{N}_4\text{O}_4\text{S}$ : C, 54.78; H, 3.94; N, 12.17. Found: C, 53.58; H, 3.91; N, 12.13.

## 4.1.20. N-(3-(7-Methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)-4-(pyridin-3-yloxy)phenyl)methanesulfonamide (29)

m.p.: 115–117 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.91 (s, 1H), 8.29 (dd,  $J = 4.4$  Hz, 1.2 Hz, 1H), 8.25 (d,  $J = 2.6$  Hz, 1H), 8.04 (s, 1H), 7.78 (s, 1H), 7.44–7.29 (m, 4H), 7.13 (d,  $J = 8.4$  Hz, 1H), 7.05 (s, 1H), 3.35 (s, 3H), 3.07 (s, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.80, 153.17, 151.29, 145.18, 140.89, 135.45, 133.08, 129.92, 125.84, 124.87, 124.34, 124.30, 123.21, 122.47, 122.18, 120.66, 113.15, 40.96, 40.68, 40.41, 40.13, 39.92, 39.85, 39.57, 39.29, 34.54. MS (ESI,  $m/z$ ): 412.0[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{19}\text{H}_{17}\text{N}_5\text{O}_4\text{S}$ : C, 55.47; H, 4.16; N, 17.02. Found: C, 55.52; H, 4.21; N, 17.38.

## 4.1.21. N-(3-(7-Methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)-4-(3,4,5-trifluorophenoxy)phenyl)methanesulfonamide (30)

m.p. > 220–222 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.93 (s, 1H), 8.06 (s, 1H), 7.80 (s, 1H), 7.42–7.36 (m, 2H), 7.18 (d,  $J = 9.2$  Hz, 1H), 7.10 (dd,  $J = 9.0, 6.1$  Hz, 2H), 7.04 (s, 1H), 3.37 (s, 3H), 3.07 (s, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.84, 150.99, 135.62, 133.19, 129.94, 124.13, 123.18, 122.35, 122.19, 120.59, 112.92, 104.67, 104.36, 40.81, 40.53, 40.25, 39.97, 39.81, 39.70, 39.42, 39.14, 34.61. MS (ESI,  $m/z$ ): 487.0[M+Na] $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{15}\text{F}_3\text{N}_4\text{O}_4\text{S}$ : C, 51.72; H, 3.26; N, 12.06. Found: C, 52.02; H, 3.18; N, 12.21.

## 4.1.22. N-(4-(2-Fluorophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)phenyl)methanesulfonamide (31)

m.p. > 220–222 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.93 (s, 1H), 8.06 (s, 1H), 7.80 (s, 1H), 7.42–7.36 (m, 2H), 7.18 (d,  $J = 9.2$  Hz, 1H), 7.10 (dd,  $J = 9.0, 6.1$  Hz, 2H), 7.04 (s, 1H), 3.37 (s, 3H), 3.07 (s, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.87, 152.16, 142.68, 142.52, 134.61, 133.01, 129.88, 125.90, 124.56, 124.46, 123.17, 122.27, 122.08, 121.02, 118.08, 117.67, 117.43, 113.31, 40.81, 40.53, 40.26, 39.98, 39.78, 39.70, 39.42, 39.14, 34.62. MS (ESI,  $m/z$ ): 487.0[M+Na] $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{17}\text{F}_1\text{N}_4\text{O}_4\text{S}$ : C, 56.07; H, 4.00; N, 13.08. Found: C, 56.05; H, 4.11; N, 13.12.

## 4.1.23. N-(4-(3-Fluorophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-5-yl)phenyl)methanesulfonamide (32)

m.p. > 220–222 °C;  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-}d_6$ )  $\delta$  9.88 (s, 1H), 8.00 (s, 1H), 7.78 (s, 1H), 7.46–7.36 (m, 2H), 7.30 (dd,  $J = 15.4, 8.0$  Hz, 1H), 7.16 (d, 1H), 7.01 (s, 1H), 6.95–6.85 (m, 1H), 6.85–6.72 (m, 2H), 3.35 (s, 3H), 3.07 (s, 3H).  $^{13}\text{C NMR}$  (75 MHz,  $\text{DMSO-}d_6$ )  $\delta$  154.81, 151.14, 135.45, 133.08, 131.63, 131.50, 129.89, 124.25, 123.22, 122.64, 122.13, 121.10, 114.54, 113.18, 110.91, 110.63, 106.26, 105.93, 40.96, 40.69, 40.41, 40.13, 39.93, 39.85, 39.57, 39.29, 34.52. MS (ESI,  $m/z$ ): 429.0[M+H] $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{17}\text{F}_1\text{N}_4\text{O}_4\text{S}$ : C, 56.07; H, 4.00; N, 13.08. Found: C, 55.98; H, 4.13; N, 12.95.

#### 4.1.24. *N*-(4-(5-Fluoro-2-methylphenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-5-yl)phenyl)methanesulfonamide (**33**)

m.p. > 220–222 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.87 (s, 1H), 8.01 (s, 1H), 7.79 (s, 1H), 7.44–7.36 (m, 2H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.11–7.03 (m, 2H), 6.86–6.76 (m, 1H), 6.73–6.66 (m, 1H), 3.35 (s, 3H), 3.07 (s, 3H), 1.91 (s, 3H). <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>) δ 154.84, 151.18, 135.13, 133.25, 132.56, 132.43, 129.88, 124.37, 124.30, 123.14, 122.23, 121.91, 120.34, 113.25, 110.70, 110.42, 105.41, 105.08, 40.86, 40.58, 40.30, 40.03, 39.83, 39.75, 39.47, 39.19, 34.57, 15.21. MS (ESI, *m/z*): 457.1[M+H]<sup>+</sup>; Anal. calcd. for C<sub>21</sub>H<sub>19</sub>F<sub>1</sub>N<sub>4</sub>O<sub>4</sub>S: C, 51.07; H, 4.33; N, 12.66. Found: C, 51.08; H, 4.23; N, 12.34.

#### 4.1.25. *N*-(4-(2-Fluoro-5-methylphenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-5-yl)phenyl)methanesulfonamide (**34**)

m.p. > 220–222 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.84 (s, 1H), 7.95 (s, 1H), 7.79 (s, 1H), 7.42–7.33 (m, 2H), 7.20–7.10 (m, 1H), 7.07 (s, 1H), 7.02–6.90 (m, 3H), 3.38 (s, 3H), 3.06 (s, 3H), 2.19 (s, 3H). <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>) δ 154.85, 152.27, 142.12, 135.38, 134.59, 132.87, 129.85, 126.38, 126.29, 124.69, 124.53, 123.23, 122.30, 122.03, 120.98, 118.31, 117.06, 116.83, 113.23, 40.97, 40.68, 40.41, 40.13, 39.85, 39.57, 39.29, 34.53, 20.57. MS (ESI, *m/z*): 443.0 [M+H]<sup>+</sup>; Anal. calcd. for C<sub>21</sub>H<sub>19</sub>F<sub>1</sub>N<sub>4</sub>O<sub>4</sub>S: C, 51.07; H, 4.33; N, 12.66. Found: C, 51.12; H, 4.31; N, 12.54.

#### 4.1.26. *N*-(4-(4-Isopropylphenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-5-yl)phenyl)met hanesulfonamide (**35**)

m.p.: 108–110 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.83 (s, 1H), 7.99 (s, 1H), 7.78 (s, 1H), 7.36 (d, *J* = 9.4 Hz, 2H), 7.15 (d, *J* = 7.7 Hz, 2H), 7.01 (d, *J* = 5.5 Hz, 2H), 6.87 (d, *J* = 7.7 Hz, 2H), 3.36 (s, 3H), 3.05 (s, 3H), 2.88–2.76 (m, 1H), 1.14 (d, *J* = 6.6 Hz, 6H). <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>) δ 154.83, 154.16, 152.56, 144.43, 134.45, 133.16, 129.76, 128.03, 124.60, 123.12, 121.99, 121.70, 119.63, 119.00, 113.74, 40.76, 40.48, 40.20, 39.93, 39.65, 39.37, 39.09, 34.57, 33.17, 24.39. MS (ESI, *m/z*): 453.1[M+H]<sup>+</sup>; Anal. calcd. for C<sub>23</sub>H<sub>24</sub>N<sub>4</sub>O<sub>4</sub>S: C, 61.05; H, 5.35; N, 12.38. Found: C, 61.07; H, 5.41; N, 12.61.

#### 4.1.27. *N*-(4-(4-Bromophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-5-yl)phenyl)ethan esulfonamide (**36**)

m.p. > 220–222 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.95 (s, 1H), 7.98 (s, 1H), 7.79 (s, 1H), 7.42 (dd, *J* = 20.5, 8.1 Hz, 4H), 7.11 (d, *J* = 9.2 Hz, 1H), 7.00 (s, 1H), 6.89 (d, *J* = 8.9 Hz, 2H), 3.36 (s, 3H), 3.17 (d, *J* = 7.3 Hz, 2H), 1.24 (t, *J* = 7.3 Hz, 3H); MS (ESI, *m/z*): 500.90 [M+H]<sup>+</sup>. Anal. calcd. for C<sub>21</sub>H<sub>19</sub>BrN<sub>4</sub>O<sub>4</sub>S: C, 50.11; H, 3.80; N, 11.13. Found: C, 50.05; H, 3.41; N, 11.34.

#### 4.1.28. *N*-(4-(3-Bromophenoxy)-3-(7-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-5-yl)phenyl)ethan esulfonamide (**37**)

m.p.: 108–110 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.95 (s, 1H), 8.00 (s, 1H), 7.77 (s, 1H), 7.39 (s, 2H), 7.20 (d, *J* = 29.3 Hz, 4H), 6.97 (d, *J* = 23.3 Hz, 2H), 3.35 (s, 3H), 3.18 (s, 2H), 1.23 (s, 3H); <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>) δ 156.77, 154.30, 150.51, 134.77, 132.67, 131.53, 129.36, 126.56, 123.19, 122.62, 122.03, 121.81, 121.66, 121.08, 120.19, 117.32, 112.72, 45.23, 34.09, 8.03; MS (ESI, *m/z*): 500.90 [M+H]<sup>+</sup>. Anal. calcd. for C<sub>21</sub>H<sub>19</sub>BrN<sub>4</sub>O<sub>4</sub>S: C, 50.11; H, 3.80; N, 11.13. Found: C, 50.15; H, 3.79; N, 11.22.

## 4.2. Docking studies

These docking small molecules were processed with Ligand preparation protocol in Maestro 10.2. Then all ligands were minimized with OPLS 2005 force field [27]. Glide generation protocol of Maestro 10.2 was performed to process the BRD4 (BD1) protein (PDB id: 3P5O), which was downloaded from PDB (<http://www.rcsb.org/pdb>). Seven molecules of water around the Kac-pocket were reserved (id: HOH18, HOH27, HOH33, HOH256, HOH267, HOH268, HOH358), and the bond orders were assigned [25]. After that, a glide box using the

centroid of the bound ligand and a maximum size of 12 Å was generated following hydrogen atoms modified. The optimal docking conformations were saved.

## 4.3. Biological evaluation

### 4.3.1. AlphaScreen bromodomain binding assay

The AlphaScreen bromodomain binding assay was performed by Shanghai ChemPartner Co. Firstly 1 × assay buffer (modified HEPES Buffer) was prepared for this experiment. Small molecules were transferred to assay plates with Echo. DMSO's final concentration is 0.1% following protein solution prepared in 1 × assay buffer (5 nM). Then, to generate the substrate solution, peptide (H4) was added in 1 × assay buffer. 5 μL of protein solution or 5 μL of 1 × assay buffer was transferred to assay plates. After that, these assay plates were incubated at room temperature for 15 min. To each well 5 μL of substrate solution was added for reaction. Next, these assay plates were incubated at room temperature for one hour. In subdued light environment, 15 μL acceptor and donor solution was added and incubated for another one hour at room temperature. The Alpha mode of EnSpire was performed to show the endpoint. Finally, the data was imported into Excel to obtain inhibition values using equation (1). Equation (1): Inh % = (Max – Signal)/(Max – Min) × 100. Meanwhile, max signal was provided from the action of Enzyme and Substrate and min signal was acquired from the Substrate. GraphPad Prism 5 was used to get IC<sub>50</sub> values using equation (2). Equation (2): Y = Bottom + (Top – Bottom)/(1 + 10<sup>(LogIC<sub>50</sub> – X) × Hill Slope</sup>). Y is %inhibition and X is compound concentration.

### 4.3.2. Cell proliferation assay

The anti-proliferation activities of these compounds in cancer cell proliferation lines, lymphoblast-like cell line(Raji), promyelocytic cell line (HL-60) and human pancreatic cancer cell line(BxPC-3) were evaluated using MTT colorimetric assay and the procedure and conditions were described in our previous work [25,27]. Raji and HL-60 cell line from Key laboratory of SATCM for Empirical Formulae Evaluation and Achievements Transformation were grown in RPMI 1640 supplemented with 16% fetal bovine serum (FBS). BxPC-3 cell line from Nanjing Tech University were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS. HL-60 DNA content analysis was performed by flow cytometry (BD, LSR Fortessa TM, San Diego, CA, USA). The percentage of cells in different cell cycle phases was determined with modfit LT 2.0 software (BD Biosciences, Pharmingen, San Diego, CA, USA). Cell cycle analysis was performed following standard protocols.

### 4.3.3. Western Blot analysis

Cell lysates were prepared in cell lysis buffer (RIPA Lysis Buffer, Beyotime, P0013B) with protease inhibitor cocktail (Roche, 5892970001). 30 μg of total protein was resolved on a 12% SDS polyacrylamide gel and probed with anti-cMyc (Abcam Ab32072), anti-BRD4 (Cell Signaling #13440), anti-HEXIM1 (Abcam, Ab-25388), anti-PARP (Cell Signaling #9542), anti-Bcl-xl (Abcam Ab-32370), anti-caspase3 (Cell Signaling # 9665), anti-caspase9 (Cell Signaling # 9508), GAPDH loading control Cell Signaling #5174. Western blotting was performed following standard protocols.

## Acknowledgments

This study was supported by the Natural Science Foundation of Jiangsu Province (No. BK 20141349) and the China National Key HiTech Innovation Project for the R&D of Novel Drugs (No. 2013ZX09301303-002).

## Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bioorg.2019.103044>.

## References

- [1] T. Kouzarides, *Cell* 128 (2007) 693–705.
- [2] S.R. Haynes, C. Dollard, F. Winston, S. Beck, J. Trowsdale, I.B. Dawid, *Nucleic Acids Res.* 20 (1992) 2603.
- [3] C. Dhalluin, J.E. Carlson, L. Zeng, C. He, A.K. Aggarwal, M.M. Zhou, *Nature* 399 (1999) 491–496.
- [4] G.V. Denis, *Front Biosci.* 6 (2001) D1065–D1068.
- [5] P. Filippakopoulos, S. Picaud, M. Mangos, T. Keates, J.P. Lambert, D. Barsyte-Lovejoy, I. Felletar, R. Volkmer, S. Muller, T. Pawson, A.C. Gingras, C.H. Arrowsmith, S. Knapp, *Cell* 149 (2012) 214–231.
- [6] Z. Sun, H. Zhang, Z. Chen, Y. Xie, H. Jiang, L. Chen, H. Ding, Y. Zhang, H. Jiang, M. Zheng, C. Luo, *Bioorg. Med. Chem. Lett.* 27 (2017) 2003–2009.
- [7] S. Rahman, M.E. Sowa, M. Ottinger, J.A. Smith, Y. Shi, J.W. Harper, P.M. Howley, *Mol. Cell Biol.* 31 (2011) 2641–2652.
- [8] G. LeRoy, B. Rickards, S.J. Flint, *Mol. Cell* 30 (2008) 51–60.
- [9] Z. Yang, J.H. Yik, R. Chen, N. He, M.K. Jang, K. Ozato, Q. Zhou, *Mol. Cell* 19 (2005) 535–545.
- [10] S. Picaud, D. Da Costa, A. Thanasopoulou, P. Filippakopoulos, P.V. Fish, M. Philpott, O. Fedorov, P. Brennan, M.E. Bunnage, D.R. Owen, J.E. Bradner, P. Taniere, B. O'Sullivan, S. Muller, J. Schwaller, T. Stankovic, S. Knapp, *Cancer Res.* 73 (2013) 3336–3346.
- [11] R. Gosmini, V.L. Nguyen, J. Toum, C. Simon, J.M. Brusq, G. Krysa, O. Mirguet, A.M. Riou-Eymard, E.V. Boursier, L. Trottet, P. Bamborough, H. Clark, C.W. Chung, L. Cutler, E.H. Demont, R. Kaur, A.J. Lewis, M.B. Schilling, P.E. Soden, S. Taylor, A.L. Walker, M.D. Walker, R.K. Prinjha, E. Nicodeme, *J. Med. Chem.* 57 (2014) 8111–8131.
- [12] I.A. Asangani, V.L. Dommeti, X. Wang, R. Malik, M. Cieslik, R. Yang, J. Escara-Wilke, K. Wilder-Romans, S. Dhanireddy, C. Engelke, M.K. Iyer, X. Jing, Y.M. Wu, X. Cao, Z.S. Qin, S. Wang, F.Y. Feng, A.M. Chinnaiyan, *Nature* 510 (2014) 278–282.
- [13] P. Filippakopoulos, J. Qi, S. Picaud, Y. Shen, W.B. Smith, O. Fedorov, E.M. Morse, T. Keates, T.T. Hickman, I. Felletar, M. Philpott, S. Munro, M.R. McKeown, Y. Wang, A.L. Christie, N. West, M.J. Cameron, B. Schwartz, T.D. Heightman, N. La Thangue, C.A. French, O. Wiest, A.L. Kung, S. Knapp, J.E. Bradner, *Nature* 468 (2010) 1067–1073.
- [14] S. Amorim, A. Stathis, M. Gleeson, S. Iyengar, V. Magarotto, X. Leleu, F. Morschhauser, L. Karlin, F. Broussais, K. Rezaei, P. Herait, C. Kahatt, F. Lokiec, G. Salles, T. Facon, A. Palumbo, D. Cunningham, E. Zucca, C. Thieblemont, *Lancet Haematol.* 3 (2016) e196–e204.
- [15] O. Mirguet, R. Gosmini, J. Toum, C.A. Clement, M. Barnathan, J.M. Brusq, J.E. Mordaunt, R.M. Grimes, M. Crowe, O. Pineau, M. Ajakane, A. Daugan, P. Jeffrey, L. Cutler, A.C. Haynes, N.N. Smithers, C.W. Chung, P. Bamborough, L.J. Uings, A. Lewis, J. Witherington, N. Parr, R.K. Prinjha, E. Nicodeme, *J. Med. Chem.* 56 (2013) 7501–7515.
- [16] L. Chen, J.L. Yap, M. Yoshioka, M.E. Lanning, R.N. Fountain, M. Raj, J.A. Scheenstra, J.W. Strovel, S. Fletcher, *ACS Med. Chem. Lett.* 6 (2015) 764–769.
- [17] K.F. McDaniel, L. Wang, T. Soltwedel, S.D. Fidanze, L.A. Hasvold, D. Liu, R.A. Mantei, *J.K. Pratt* 60 (2017) 8369–8384.
- [18] M.H. Bui, X. Lin, D.H. Albert, L. Li, L.T. Lam, E.J. Faivre, S.E. Warder, X. Huang, D. Wilcox, C.K. Donawho, G.S. Sheppard, L. Wang, S. Fidanze, J.K. Pratt, D. Liu, L. Hasvold, T. Uziel, X. Lu, F. Kohlhapp, G. Fang, S.W. Elmore, S.H. Rosenberg, K.F. McDaniel, W.M. Kati, Y. Shen, *Cancer Res.* 77 (2017) 2976–2989.
- [19] E.J. Faivre, D. Wilcox, X. Lin, P. Hessler, M. Torrent, W. He, T. Uziel, D.H. Albert, K. McDaniel, W. Kati, Y. Shen, *Mol. Cancer Res.* 15 (2017) 35–44.
- [20] K.T. Siu, J. Ramachandran, A.J. Yee, H. Eda, L. Santo, C. Panaroni, J.A. Mertz, R.J. Sims Iii, M.R. Cooper, N. Raj, *Leukemia* 31 (2017) 1760–1769.
- [21] P.L. Garcia, A.L. Miller, K.M. Kreitzburg, L.N. Council, T.L. Gamblin, J.D. Christein, M.J. Heslin, J.P. Arnoletti, J.H. Richardson, D. Chen, C.A. Hanna, S.L. Cramer, E.S. Yang, J. Qi, J.E. Bradner, K.J. Yoon, *Oncogene* 35 (2016) 833–845.
- [22] R.L. Siegel, K.D. Miller, A. Jemal, *CA Cancer J. Clin.* 66 (2016) 7–30.
- [23] L. Rahib, B.D. Smith, R. Aizenberg, A.B. Rosenzweig, J.M. Fleshman, L.M. Matrisian, *Cancer Res.* 74 (2014) 2913–2921.
- [24] X. Zhang, H.C. Lee, F. Shirazi, V. Baladandayuthapani, H. Lin, I. Kuitase, H. Wang, R.J. Jones, Z. Berkova, R.K. Singh, J. Lu, Y. Qian, K. Raina, K.G. Coleman, C.M. Crews (2018).
- [25] L. Zhao, Y. Yang, Y. Guo, L. Yang, J. Zhang, J. Zhou, H. Zhang, *Bioorg. Med. Chem.* 25 (2017) 2482–2490.
- [26] X. Lin, X. Huang, T. Uziel, P. Hessler, D.H. Albert, L.A. Roberts-Rapp, K.F. McDaniel, W.M. Kati, Y. Shen, *Mol. Cancer Ther.* 16 (2017) 388–396.
- [27] Y. Yang, L. Zhao, B. Xu, L. Yang, J. Zhang, H. Zhang, J. Zhou, *Bioorg. Chem.* 68 (2016) 236–244.