



## Aqueous *Coriandrum sativum* L. extract promotes neuroprotection against motor changes and oxidative damage in rat progeny after maternal exposure to methylmercury



Keuri Eleutério Rodrigues<sup>a,b</sup>, Fábio Rodrigues de Oliveira<sup>a,c</sup>, Benilson Ramos Cassunde Barbosa<sup>b</sup>, Ricardo S. Oliveira Paraense<sup>b</sup>, Cahy Manoel Bannwart<sup>b</sup>, Bruno Gonçalves Pinheiro<sup>d</sup>, Anderson de Santana Botelho<sup>e</sup>, Nilton Akio Muto<sup>f</sup>, Cristine Bastos do Amarante<sup>e</sup>, Moises Hamoy<sup>g</sup>, Barbarella de Matos Macchi<sup>b</sup>, Cristiane do Socorro Ferraz Maia<sup>d</sup>, Alejandro Ferraz do Prado<sup>h</sup>, José Luiz Martins do Nascimento<sup>a,b,i,j,\*</sup>

<sup>a</sup> Neuroscience and Cellular Biology Post Graduation Program, Biological Sciences Institute, Federal University of Para, Belem, PA, Brazil

<sup>b</sup> Molecular and Cellular Neurochemistry Laboratory, Biological Sciences Institute, Federal University of Para, Belem, PA, Brazil

<sup>c</sup> Bromatology and Quality Control Laboratory, Health and Biological Sciences Department, Federal University of Amapa (UNIFAP), Macapa, Ap, Brazil

<sup>d</sup> Behavioral and Inflammatory Pharmacology Laboratory, Health Sciences Institute, Pharmacy College, Federal University of Para, Belem, PA, Brazil

<sup>e</sup> Chemical Analysis Laboratory, Museum Paraense Emílio Goeldi, Belem, PA, Brazil

<sup>f</sup> Amazonian Bioactive Compounds Valorization Center, Federal University of Para, Belem, PA, Brazil

<sup>g</sup> Natural Products' Toxicology and Pharmacology Laboratory, Biological Sciences Institute, Federal University of Para, Belem, PA, Brazil

<sup>h</sup> Structural Biology Laboratory, Biological Sciences Institute, Federal University of Para, Belem, PA, Brazil

<sup>i</sup> National Institute of Science and Technology in Neuroimmunomodulation (INCT-NIM), Rio de Janeiro, RJ, Brazil

<sup>j</sup> Pharmaceutical Sciences Post Graduation Program, Health and Biological Sciences Department, Federal University of Amapa (UNIFAP), Macapa, Ap, Brazil

### ARTICLE INFO

#### Keywords:

*Coriandrum sativum*  
Methylmercury  
Rat offspring  
Behavioral parameters  
Oxidative stress

### ABSTRACT

This study aimed to investigate the effects of *Coriandrum sativum* aqueous extract (CSAE) on the rat progeny of mothers exposed to methylmercury (MeHg). The presence of bioactive compounds and CSAE's antioxidant capacity been evaluated, and the offspring were assessed for their total mercury levels, motor behavioral parameters and oxidative stress in the cerebellum. The analysis of the bioactive compounds revealed significant amounts of polyphenols, flavonoids, and anthocyanins, as well as a variety of minerals. A DPPH test showed the CSAE had important antioxidant activity. The MeHg + CSAE group performed significantly better spontaneous locomotor activity, palmar grip strength, balance, and motor coordination in behavioral tests compared the MeHg group, as well as in the parameters of oxidative stress, with similar results to those of the control group. The MeHg + CSAE group also had significantly reduced mercury levels in comparison to the MeHg group. Based on the behavioral tests, which detected large locomotor, balance, and coordination improvements, as well as a reduction in oxidative stress, we conclude that CSAE had positive functional results in the offspring of rats exposed to MeHg.

### 1. Introduction

Mercury is an extremely toxic heavy metal widely distributed in the

environment, and in its organic form of methylmercury (MeHg), capable of biomagnification (Bisi et al., 2012; Boudou and Ribeyre, 1997; Coelho et al., 2013). It is important to note that 80%–90% of human

**Abbreviations:** ATP, Adenosine triphosphate; BDNF, derived neurotrophic factor; Ca, Calcium; CAT, Catalase; CNS, Central Nervous System; CSAE, *Coriandrum sativum* aqueous extract; Cu, Copper; DNA, deoxyribonucleic acid; DPPH, 2,2 diphenyl-1-picrylhydrazyl; DTNB, 5,5'-dithiobis-2-nitrobenzoic acid; Fe, Iron; H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide; H<sub>2</sub>SO<sub>4</sub>, Sulfuric acid; HClO<sub>4</sub>, Perchloric Acid; HNO<sub>3</sub>, Nitric acid; KMnO<sub>4</sub>, Potassium permanganate; MeHg, methylmercury; Mg, Magnesium; Mn, Manganese; NBT, Nitroblue Tetrazolium; NO, Nitric oxide; O<sub>2</sub><sup>-</sup>, Reactive oxygen; PBS, Phosphate buffered saline; ROS, Reactive oxygen species; SnCl<sub>2</sub>, Tin chloride; SOD, Superoxide dismutase; TBARS, Thiobarbituric acid reactive substances; TNB, 5-thio-2-nitrobenzoic acid; Zn, Zinc

\* Corresponding author. Jose Luiz M. do Nascimento. Molecular and Cellular Neurochemistry Laboratory, Biological Sciences Institute, Federal University of Para, Belem-Pa, Brazil, Rua Augusto Corrêa, 01, Belém, PA, 66075-110, Brazil.

E-mail address: [jlmn@ufpa.br](mailto:jlmn@ufpa.br) (J.L.M. do Nascimento).

<https://doi.org/10.1016/j.fct.2019.110755>

Received 3 May 2019; Received in revised form 3 August 2019; Accepted 8 August 2019

Available online 10 August 2019

0278-6915/ © 2019 Published by Elsevier Ltd.

exposure occurs through the ingestion of contaminated fish or water, which, due to the bioaccumulation phenomenon, can result in high organic mercury levels in human populations (Buchanan et al., 2015; Liu et al., 2018; McDowell et al., 2004; Pinheiro et al., 2000; Vieira et al., 2015). Most (95%) ingested MeHg is readily absorbed in the gastrointestinal tract and assimilated, leading to protein complexes containing thiol groups with high mobility through bodily tissues and barriers such as the placental and blood-brain barriers (Amonpatumrat et al., 2008; Aschner and Clarkson, 1989; Balthasar et al., 2017; Roos et al., 2010). Methylmercury accumulates mostly in the central nervous system (CNS), with the cerebellum being one of the most affected areas (Cheng et al., 2015; Heimfarth et al., 2018; Mancini et al., 2009; Ruskiewicz et al., 2016). This leads to neurotoxicity, especially after fetal and neonatal exposure during the early stages of neural development, which are characterized by an immature blood-brain barrier and the rapid growth, proliferation, and differentiation of brain tissue (Ballester et al., 2018; Murcia et al., 2016; Sakamoto et al., 2017).

Methylmercury-induced toxicity contributes to epigenetic alterations that change the expression of important enzymes such as glutathione peroxidase and tyrosine hydroxylase and growth factors such as brain-derived neurotrophic factor (Go et al., 2018; Onishchenko et al., 2008; Usuki et al., 2011). In addition, MeHg leads to alterations in glutamatergic transmission (Deng et al., 2014; Feng et al., 2014; Gutierrez et al., 2018; Liu et al., 2016; Xu et al., 2012), calcium homeostasis (Shao et al., 2015; Shen et al., 2016), and oxidative stress (Al-Osaimi et al., 2018; Dos Santos et al., 2018; Fujimura and Usuki, 2018). All these effects can lead to neurological dysfunction and severe motor disorders (Fujimura et al., 2012; Tavares et al., 2005; Yamamoto et al., 2019; Zimmermann et al., 2014).

Research on various floras to prevent the negative effects of xenobiotic poisoning has significantly increased in the last decade (Fujimura et al., 2012; Kopec et al., 2016; Mzid et al., 2017; Ola-Mudathir and Suru, 2015; Tavares et al., 2005; Yamamoto et al., 2019; Zimmermann et al., 2014). Plant extracts have been extensively investigated in both *in vivo* and *in vitro* models, yielding several interesting positive results (Akhter et al., 2013; Dewanjee et al., 2017; Kujawska et al., 2016; Nazima et al., 2016; Olaleye et al., 2010).

*Coriandrum sativum* L. (Umbelliferae), a species native to Europe and Asia and known popularly as coriander, is commonly used as a condiment in the northern and northeastern regions of Brazil (Al-Mofleh et al., 2006; Bhuiyan et al., 2009; Cavalcante et al., 2016; Melo et al., 2003; Msaada et al., 2014). The species has numerous reported biological activities (Aissaoui et al., 2008, 2011; Bogavac et al., 2015; Hosseinzadeh et al., 2016; Jabeen et al., 2009; Kazempour et al., 2015; Paarakh et al., 2016), including effects on the CNS such as anxiolytic activity (Emamghoreishi and Heidari-Hamedani, 2006; Emamghoreishi et al., 2005; Mahendra and Bisht, 2011; Pathan et al., 2011) and both *in vitro* and *in vivo* antioxidant activity (Anaiegoudari et al., 2016; Chithra and Leelamma, 1999; Harsha and Anilakumar, 2014; Hwang et al., 2014; Kozłowska et al., 2016; Pellegrini et al., 2018; Sreelatha and Inbavalli, 2012; Wangenstein et al., 2004). These activities may be related to coriander's reported chemical composition, which has shown high levels of total phenolic compounds and flavonoids that can attenuate oxidative damage (Justesen and Knuthsen, 2001; Melo et al., 2003, 2005; Pereira and Tavano, 2014).

Previous reports have shown the capacity of aqueous *C. sativum* extracts to remove inorganic mercury and MeHg from plants (Karunasagar et al., 2005). Considering this species can decrease mercury levels, and that high systemic levels of this metal can cause oxidative damage and induce motor dysfunction, we hypothesized that a *C. sativum* aqueous extract (CSAE) would reduce MeHg levels, and consequently, reduce lipid peroxidation and improve motor function. Therefore, this study aimed to evaluate the effects of CSAE on the oxidative damage and behavioral and motor parameters in progeny rats after maternal exposure to MeHg.

## 2. Materials and methods

### 2.1. Preparation of the *Coriandrum sativum* aqueous extract

*Coriandrum sativum* leaves and stalks were obtained in São Francisco do Pará-Pa (01°10'03" S, 47°47'45" W). The specimen was identified by Dr. Antônio Elielson Rocha, and a voucher specimen was deposited in the Museum Paraense Emílio Goeldi herbarium (MG: 228890). After collection, the fresh material was washed under running water, rinsed in distilled water, then frozen for later lyophilization in a vacuum chamber at  $-20^{\circ}\text{C}$  for 24 h. After lyophilization, the material was pulverized in a Willye-type mill (Star FT 50/1; 620 rpm) coupled to 60-mesh sieves, weighed, vacuum packed in polyethylene bags, and stored at  $-20^{\circ}\text{C}$  until use. The extract was prepared by adding 13.5 g of ground material to 300 mL deionized water and homogenizing at room temperature for 1 h for a final concentration of 45 mg/mL (Kansal et al., 2011).

### 2.2. Mineral and phytochemical content of the *Coriandrum sativum* extract

#### 2.2.1. Mineral content determination

The mineral content was determined as described by Amarante et al. (2010). After transferring 500 mg of coriander dry mass to a digester tube, a  $\text{HNO}_3\text{:HClO}_4$  solution (3:1) was added for cold and hot digestions performed in triplicate. The digests were then quantitated in an atomic absorption spectrophotometer (Thermo, ICE3000). The calcium (Ca), magnesium (Mg), copper (Cu), iron (Fe), zinc (Zn), and manganese (Mn) results were compared to their respective standard curves to determine their concentrations, which are expressed in mg/L.

#### 2.2.2. Phytochemical composition of the *Coriandrum sativum* extract

To identify any secondary metabolites with antioxidant activity in the CSAE, the total phenolic (Singleton and Rossi JR, 1965), flavonoid (Quettier-Deleu et al., 2000), and anthocyanin (Askar and Treptow, 1993) contents were determined. The tests were performed in triplicate, and the results are expressed as  $\text{mg}\cdot 100\text{ g}^{-1}$  dry matter.

### 2.3. Antioxidant capacity of the *Coriandrum sativum* extract

The antioxidant capacity of the CSAE was assessed using the DPPH (2,2-diphenyl-1-picryl-hydrazyl) method (Frankel and Meyer, 2000; Vicentino and Menezes, 2007). Standard solutions were prepared with the extract and ascorbic acid at concentrations of 0.176–21  $\mu\text{g}/\text{mL}$ , which were then incubated with a 21  $\mu\text{g}/\text{mL}$  solution of DPPH at room temperature in the dark for 30 min. The reaction product was measured in a spectrophotometer at a wavelength of 515 nm (BIO-RAD Model 450 Microplate Reader). The EC50 calculation is the concentration of extract that provides a 50% response (Sridhar and Charles, 2019). This experiment was performed in triplicate.

### 2.4. Animals

Fifty six primiparous female rats (*Rattus norvegicus*, Wistar strain) weighing  $\cong 250\text{ g}$  and in the first trimester of gestation were obtained from the Federal University of Pará breeding facilities and kept in individuals cages. The environment had a controlled room temperature ( $22 \pm 2^{\circ}\text{C}$ ) and a light-dark cycle of 12 h. Water and food were provided *ad libitum* (periodically measured for assessment). The study followed the NIH Guide for the Care and Use of Laboratory Animals, and the Ethics Committee on Animal Use in Research at the Federal University of Pará, Brazil (CEPAE-UFPA 243–14) approved the experimental protocols.

### 2.5. Experimental design

After 7 days of acclimatization, pregnant rats were divided into 4

groups composed of 14 animals each ( $n = 14$ ): Control; CSAE; MeHg; and MeHg + CSAE.

The control and CSAE groups received filtered water, while the MeHg and MeHg + CSAE groups received mercury at a concentration of 40  $\mu\text{g}/\text{mL}$  diluted in drinking water (Andersen and Andersen, 1993; Farina et al., 2005). The CSAE and MeHg + CSAE groups received the CSAE treatments by gavage with CSAE extract (360 mg/kg day), from 14th of pregnancy to 14th day of lactation (corresponding 21 days of treatment) (Gandhi et al., 2014). The control and MeHg groups were treated by gavage with filtered water over the same test period. After birth, 8 animals per mother were retained to normalize the number of animals per group. For the behavioral assays in pups at 30 days post-natal, a single male was randomly selected from each litter for a total of 14 animals per group. This was done to reduce gender-related assay distortions. For the biochemical assessments, a pool from each mother's litter was used, making a total of 14 pools per group (EPA, 1998).

## 2.6. Behavioral assays

When the progeny were 30-days old, they were subjected to behavioral tests after acclimatizing in a test room with attenuated light and sound for 1 h. All behavioral tests (open-field, grip, beam-walking, and rotarod) were performed between 10:00 a.m. and 6:00 p.m.

### 2.6.1. Open-field test

Open-field tests were performed to evaluate spontaneous locomotor activity (von Horsten et al., 1998). The apparatus consisted of an acrylic arena ( $100 \times 100 \times 40$  cm) divided into 25 quadrants ( $20 \times 20$  cm each), 9 central and 16 peripheral. Each animal was placed at the center of the apparatus and assessed for their exploratory behavior for 5 min. The number of crossings between quadrants, the total number of quadrants, and the distance crossed (measured in meters) were determined for each animal (Fernandes et al., 2018a, 2018b). The tests were recorded via a camera coupled to a computer, and the images were analyzed with Any-Maze® software (Stoelinf Co., Wood Dale, IL, USA).

### 2.6.2. Grip-strength test

This method assesses muscular strength and takes advantage of the natural behavior of rats to cling to horizontal surfaces when supported by the tail. For the test, the animals were allowed to grasp a mounted dynamometer bar while being firmly held by the tail and slowly pulled back by an inspector. The peak pull force was recorded and expressed in grams (g). The tests were performed in triplicate, and the results represent the mean of the 3 values (Takeshita et al., 2017).

### 2.6.3. Rotarod test

A motor coordination assay was performed with a rotating cylinder test on an apparatus consisting of a 20 cm high, 8 cm diameter transversal rod that rotated at variable speeds. The animals were tested on the rotating bar at speeds of 16, 20, 25, 28, and 37 rpm. The latency time until a fall occurred (recorded for a maximum 120 s) and the number of falls were recorded for each animal (da Silva et al., 2018; Slamberova et al., 2006).

### 2.6.4. Beam-walking test

Motor coordination and balance were also assessed by a beam-walking test (Stanley et al., 2005). The apparatus consisted of two supports holding a 1 m wooden beam suspended 50 cm off the floor. The serial beams had 2 types of cross sections: square (12 and 5 mm) and circular (17 and 11 mm). Each apparatus had a platform with a safe box ( $20 \text{ cm}^2$ ) at one end. The animals were placed at the starting point and had 60 s to cross the beam to reach the safe box, with 2 attempts per beam. We counted the number of times that at least one of the hind legs slipped. If two legs slipped at the same time, it was counted as two slips (Carter et al., 2001).

## 2.7. Tissue samples

When they were 30-days old, the progeny rats were anesthetized with ketamine hydrochloride (50 mg/kg) and xylazine hydrochloride (5 mg/kg) before perfusing with cold phosphate buffered saline (PBS; pH 7.4). Approximately 1 g of hair was collected and stored in a polypropylene bag until the mercury testing. The cerebellum was dissected and homogenized in PBS (1:1 w/v) with an Ultra-Turrax homogenizer (IKA-Werke GmbH & Co.). The samples were centrifuged for 10 min at 5000 rpm at 4 °C, and the supernatant was collected and frozen at  $-80$  °C until the biochemical analysis.

## 2.8. Biochemical analysis

### 2.8.1. Oxidative damage assessment

The thiobarbituric acid-reactive substances (TBARS) were determined as described by Ohkawa et al. (1978). This test indicates the presence of products derived from lipid peroxidation, mainly malonaldehyde. Proteins were recovered with 15% trichloroacetic acid, and the supernatant was incubated with 10 mM thiobarbituric acid and heated at 80 °C for 45 min. The resulting lightly pink-colored product was placed in a 96-well microplate for spectrophotometer quantification at 535 nm (BIO-RAD Model 450 Microplate Reader). The results are expressed in nmol TBARS/mg protein.

In high amounts, nitric oxide (NO) can lead to oxidative stress. Therefore, the samples were incubated with Griess reagent (Moncada, 1992; Nims et al., 1995), placed in a 96-well microplate, and measured in a spectrophotometer at 550 nm (BIO-RAD Model 450 Microplate Reader). The amount of NO is expressed as  $\mu\text{mol}/\text{mg}$  protein based on a standard curve for sodium nitrite.

The superoxide anion was quantified with a nitroblue tetrazolium (NBT) assay. This test detects NBT oxidation by intracellular reactive oxygen ( $\text{O}_2^-$ ), which generates formazan crystals that can be quantified by spectrophotometry using a 620 nm filter (BIO-RAD Model 450 Microplate Reader). The results are expressed as absorbance intensity/mg protein (Baehner et al., 1976; Meerovich et al., 2014).

### 2.8.2. Evaluating total glutathione

Total glutathione levels (Foyer and Shigeoka, 2011; Tietze, 1969) were determined by the detection of recycled enzyme after a glutathione reductase treatment in which glutathione's sulfhydryl group reacts with 1 mM 5,5'-dithiobis-(2-nitrobenzoic acid) to form 5-thio-2-nitrobenzoic acid, which has a yellow color. The absorbance was immediately measured by spectrophotometry, once per minute for 5 min, at the 412 nm wavelength (BIO-RAD Model 450 Microplate Reader). Results are expressed in pmol/ $\mu\text{g}$  protein.

### 2.8.3. Enzymatic antioxidant activities

Superoxide dismutase (SOD) was estimated through a reaction of the samples with hypoxanthine and NBT (Maier and Chan, 2002). The absorbance was measured at the 470 nm wavelength (BIO-RAD Model 450 Microplate Reader), and the results are expressed as units of SOD activity per milligram protein.

Catalase enzyme activity was measured after reacting a sample with hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) to form formaldehyde, which is revealed by treating with 34.2 mM Purpald® reagent (4-amino-3-hydrazino-5-mercaptop-1, 2, 4-triazole). The results are expressed as  $\mu\text{M}$  formaldehyde/min/mg protein, determined by comparing the absorbance at 540 nm (BIO-RAD Model 450 Microplate Reader) with a standard curve for 4.25  $\mu\text{M}$  formaldehyde (Johansson and Borg, 1988; Wheeler et al., 1990).

## 2.9. Protein quantification

The Bradford (1976) method was utilized to quantify the protein level. A standard curve was generated with solutions containing known

concentrations of bovine serum albumin, and the results are expressed in mg/mL.

### 2.10. Total mercury dosage (total Hg)

The procedure to extract total mercury was based on a modified method by Costa et al. (2017). Atomic fluorescence spectrometry was used to detect the amount of Hg in 100 mg of animal hair digested in  $\text{HNO}_3/\text{H}_2\text{SO}_4$  (1:1) and fixed in 5% (m/v) potassium permanganate ( $\text{KMnO}_4$ ). The mercury sample was reduced with 20% (m/v) tin chloride ( $\text{SnCl}_2$ ) and quantified with a Quick Trace M-7500 Mercury Analyzer (Teledyne CETAC Technologies), which has a detection limit of  $1 \text{ ng g}^{-1}$  (Costa et al., 2017).

### 2.11. Statistical analyses

The statistical analyses were performed with GraphPad Prism® 6.0. software (GraphPad Software, San Diego, CA, EUA). All results are expressed as the mean  $\pm$  standard deviation (SD). Data normality was assessed with the Shapiro-Wilk test, after which, a two-way ANOVA followed by a Tukey post-test was applied. Each experimental group comprised a total of 14 animals, and  $p$  values  $< 0.05$  were considered statistically significant.

## 3. Results

### 3.1. Minerals and phytochemicals in the *Coriandrum sativum* extract

A wide array of minerals was present in the CSAE (Table 1), with Ca ( $11,024.5 \pm 944 \text{ mg/L}$ ) and Mg ( $3041 \pm 121 \text{ mg/L}$ ) being the most abundant, followed by Fe ( $110.7 \pm 9.18 \text{ mg/L}$ ), Zn ( $72.4 \pm 1.21 \text{ mg/L}$ ), Mn ( $34.5 \pm 0.174 \text{ mg/L}$ ), and Cu ( $13.6 \pm 0.173 \text{ mg/L}$ ). The phytochemical analysis revealed that polyphenols ( $14 \pm 2.42 \text{ mg/g}$ ) were the most abundant secondary metabolites in the CSAE, followed by anthocyanins ( $9.8 \pm 0.174 \text{ mg/g}$ ) and flavonoids ( $5.1 \pm 0.174 \text{ mg/g}$ ).

### 3.2. Antioxidant and free radical scavenging activities of the *Coriandrum sativum* extract

The CSAE inhibited the DPPH radical in a concentration-dependent manner. The  $\text{EC}_{50}$  was  $13.4 \pm 2.3 \mu\text{g/mL}$ , a capacity to donate hydrogen similar to that of ascorbic acid ( $\text{EC}_{50}$   $14.6 \pm 3.8 \mu\text{g/mL}$ ; Fig. 1).

### 3.3. Total mercury in the progeny's hair

The total mercury in the progeny's hair was assessed after 21 days of treating the progenitor with MeHg (MeHg and MeHg + CSAE groups; Fig. 2). Compared to the control group ( $0.019 \pm 0.004 \mu\text{g/g}$ ), the




Fig. 1. Logarithmic curve of the effect of concentration on the DPPH inhibition capacity of the *Coriandrum sativum* aqueous extract (CSAE) compared to that of ascorbic acid in  $\mu\text{g/mL}$ . Data are shown as the mean  $\pm$  SD of 3 independent experiments;  $p$  values  $< 0.05$  were considered statistically significant. Abbreviation: DPPH, 2,2-diphenyl-1-picrylhydrazil; AA, ascorbic acid.




Fig. 2. Mercury levels ( $\mu\text{g/L}$ ) in the progeny's hair after intoxicated the progenitors with methylmercury (MeHg) and treating with a *Coriandrum sativum* aqueous extract (CSAE). Data are shown as the mean  $\pm$  SD ( $n = 14/\text{group}$ ). \*,  $p < 0.05$  compared to the control group; #,  $p < 0.05$  compared to the MeHg-treated group (two-way ANOVA and Tukey test).

MeHg group had significantly higher mercury levels ( $0.693 \pm 0.148 \mu\text{g/g}$ ;  $p < 0.05$ ), while the CSAE group ( $0.020 \pm 0.010 \mu\text{g/g}$ ) had levels similar to those of the control group. The MeHg + CSAE group ( $0.068 \pm 0.032 \mu\text{g/g}$ ) had higher mercury levels than the control group, but compared to the MeHg group, they had significantly less total mercury in their hair.

### 3.4. Treatment with CSAE ameliorated the motor deficits induced by methylmercury


To assess the effects of mercury on spontaneous locomotor activity and the effectiveness of the CSAE treatment, the progeny were subjected to an open-field test. The total and central distances covered were reduced in the MeHg group, with means of  $14.35 \pm 2.948 \text{ m}$  and  $0.38 \pm 0.093 \text{ m}$ , respectively, versus  $19.43 \pm 4.349 \text{ m}$  ( $p < 0.05$ ) and  $2.03 \pm 0.676 \text{ m}$  ( $p < 0.05$ ) for the control group. The distances for the CSAE ( $21.92 \pm 3.680 \text{ m}$ ;  $1.84 \pm 0.732 \text{ m}$ ) and MeHg + CSAE ( $20.47 \pm 2.731 \text{ m}$ ;  $1.49 \pm 0.615 \text{ m}$ ) groups did not significantly differ from the control group. Importantly, when the MeHg and MeHg + CSAE groups were compared, the intoxicated animals treated with CSAE performed better than the untreated intoxicated animals, reaching distances similar to those of the control group (Fig. 3A and B). It is noteworthy that the reduction in central distance traveled qualifies the spontaneous locomotion as anxiogenic.

Cerebellar damage is accompanied by reduced muscle strength. Hence, we utilized a grip strength test. The MeHg group

Table 1

Minerals (mg/mL) and phytochemical constituents (mg/g) of the *Coriandrum sativum* aqueous extract. Data are shown as the mean  $\pm$  SD of 3 independent experiments.

Components	Values
<b>Minerals</b>	
Calcium (Ca)	$11,024.5 \pm 944$
Magnesium (Mg)	$3041.2 \pm 121$
Copper (Cu)	$13.6 \pm 0.173$
Iron (Fe)	$110.7 \pm 9.18$
Zinc (Zn)	$72.48 \pm 1.21$
Manganese (Mn)	$34.6 \pm 0.174$
<b>Secondary metabolites</b>	
Total polyphenols	$14.01 \pm 2.42$
Total flavonoids	$5.1 \pm 0.174$
Anthocyanins	$9.8 \pm 0.174$



**Fig. 4.** Methylmercury (MeHg) intoxication and *Coriandrum sativum* aqueous extract (CSAE) treatment effects on paw grip strength. Data are shown as the mean  $\pm$  SD of 3 test sequences with 14 animals per group. \*,  $p < 0.05$  compared to the control group; #,  $p < 0.05$  compared to the MeHg group (two-way ANOVA and Tukey test).

(106  $\pm$  11.32 g) showed a reduced grip strength relative to the control group (189  $\pm$  43.81 g;  $p < 0.05$ ), whereas the CSAE (207  $\pm$  50.26 g) and MeHg + CSAE (196  $\pm$  56.59 g) groups did not significantly differ. Comparing the MeHg and MeHg + CSAE groups, the positive effects of the CSAE treatment on the intoxicated animals was shown by their increased muscle strength (Fig. 4).

A rotarod test was utilized to assess motor coordination and balance, with the latency and number of slips evaluated at rotational speeds of 16, 20, 25, 28, and 37 rpm. The latency times for the MeHg group (83.3  $\pm$  49.4, 81.7  $\pm$  31.5, 53.3  $\pm$  22.6, 43  $\pm$  23.0, and 16.6  $\pm$  9.2 s, respectively) were shorter in comparison to the control group (126.6  $\pm$  30.4, 157.9  $\pm$  19.1, 153.6  $\pm$  18.6, 145.0  $\pm$  21.0, and 127.6  $\pm$  48.0 s, respectively), while those for the CSAE group did not differ from the control group. In comparison to the control group, the MeHg + CSAE group (159.1  $\pm$  16.1, 154.3  $\pm$  16.0, 142.5  $\pm$  20.8, 137.1  $\pm$  26.4, and 123.9  $\pm$  18.8 s, respectively) had no significant differences, whereas in comparison with the MeHg group, they had significantly longer latency times on all rotational speeds (Fig. 5A).

When the number of falls was assessed (Fig. 5B), the MeHg group (2.5  $\pm$  1.4, 5.1  $\pm$  1.4, 7.5  $\pm$  3.4, 10.2  $\pm$  1.3, and 9.5  $\pm$  3.7) had significantly more than the control group (1.2  $\pm$  1.0, 1.7  $\pm$  1.3, 2.7  $\pm$  1.3, 3.9  $\pm$  2.2, and 4.0  $\pm$  2.2;  $p < 0.05$ ) at all rotational speeds. In contrast, the numbers of falls for the CSAE (1.3  $\pm$  1.0, 1.0  $\pm$  1.4, 3.2  $\pm$  1.7, 3.5  $\pm$  2.2, and 3.2  $\pm$  2.2) and MeHg + CSAE (1.2  $\pm$  0.8, 1.4  $\pm$  1.2, 2.4  $\pm$  2.2, 3.5  $\pm$  1.7, and 3.4  $\pm$  1.7) groups did not statistically differ from those of the control group. Comparing the MeHg + CSAE and MeHg groups, a significant improvement was seen in the balance quality of the MeHg + CSAE group (Fig. 5B).

The beam walk test was used as a second assay to evaluate balance. Again, the MeHg group (4.85  $\pm$  1.2, 5.2  $\pm$  2.6, 4.6  $\pm$  1.9, and 12.5  $\pm$  4.3) had significantly more slips than the control group

**Fig. 3.** Methylmercury (MeHg) intoxication and *Coriandrum sativum* aqueous extract (CSAE) treatment effects on neonate rats' spontaneous locomotor activity. (A) Total distance; (B) Central distance. Data are the mean  $\pm$  SD of the total number of quadrants crossed after 300 s ( $n = 14$ /group). \*,  $p < 0.05$  compared to the control group; #,  $p < 0.05$  compared to the MeHg group (two-way ANOVA and Tukey test).

(2.5  $\pm$  0.9, 2.7  $\pm$  1.4, 2.1  $\pm$  1.2, and 5.3  $\pm$  2.0;  $p < 0.05$ ), while the CSAE (2.2  $\pm$  1.0, 3.0  $\pm$  1.4, 2.2  $\pm$  1.0, and 4.7  $\pm$  2.5) and MeHg + CSAE (2.3  $\pm$  1.0, 3.3  $\pm$  1.7, 2.4  $\pm$  1.0, and 5.7  $\pm$  2.0) groups showed no significant differences. The MeHg + CSAE group had significantly fewer slips than the MeHg group, suggesting the CSAE treatment improved their balance (Fig. 6).

### 3.5. Treatment with CSAE decreased the oxidative damage in the progeny induced by MeHg

A TBARS assay was performed to assess for possible oxidative damage after the intrauterine and lactation mercury exposure and whether the CSAE treatment ameliorated lipid peroxidation in the neonate animals. Lipid peroxidation was significantly higher in the MeHg group (1.4  $\pm$  0.2 pg/mg protein;  $p < 0.05$ ) than in the control group (0.6  $\pm$  0.1 pg/mg protein), while that of the CSAE (0.5  $\pm$  0.1 pg/mg protein) and MeHg + CSAE (0.7  $\pm$  0.1 pg/mg protein) groups remained similar to the control group value. Compared to the MeHg group, the MeHg + CSAE group showed reduced levels of lipid peroxidation, demonstrating an attenuated oxidative pattern (Fig. 7).

After identifying the ability of CSAE to lower lipid peroxidation levels, we evaluated the nitrite and superoxide anion levels to see if they also varied after the CSAE treatment. The nitrite levels were significantly higher in the MeHg group (8.2  $\pm$  1.1  $\mu$ mol/mg protein;  $p < 0.05$ ) than in the control group (4.1  $\pm$  1.3  $\mu$ mol/mg protein), while the levels in the CSAE (4.5  $\pm$  1.2  $\mu$ mol/mg protein) and MeHg + CSAE (4.6  $\pm$  0.9  $\mu$ mol/mg protein) groups remained similar to the control group. However, there was a significant difference between the MeHg and MeHg + CSAE groups (Fig. 8A).

The superoxide anion levels were also significantly higher in the MeHg group (0.2  $\pm$  0.03 U/mg protein;  $p < 0.05$ ) than in the control group (0.1  $\pm$  0.02 U/mg protein), while the CSAE (0.1  $\pm$  0.02 U/mg protein) and MeHg + CSAE (0.17  $\pm$  0.02 U/mg protein) groups had levels comparable to those in the control group. The MeHg + CSAE group had lower superoxide anion levels than the MeHg group (Fig. 8B).

After assessing the reactive oxygen species (ROS) levels, the antioxidant systems were investigated. A reduction in the total glutathione levels was observed in the MeHg group (8.5  $\pm$  1.5 pmol/ $\mu$ g protein) relative to the control group (13.2  $\pm$  2.6 pmol/ $\mu$ g protein;  $p < 0.05$ ), while the levels in the CSAE (13.3  $\pm$  2.0 pmol/ $\mu$ g protein) and MeHg + CSAE (12.7  $\pm$  2.4 pmol/ $\mu$ g protein) groups were comparable to the control group. The MeHg + CSAE group had significantly higher total glutathione levels than the MeHg group (Fig. 9).

Relative to the control group (14.4  $\pm$  1.3 nmol/min/mg protein), less catalase activity was seen in the MeHg group (6.5  $\pm$  0.9 nmol/min/mg protein;  $p < 0.05$ ). The enzyme activity was preserved in the CSAE (13.9  $\pm$  1.5 nmol/min/mg protein) and MeHg + CSAE (13.7  $\pm$  2.0 nmol/min/mg protein) groups, with levels comparable to those of the control group. Importantly, the MeHg + CSAE group had significantly higher catalase activity than the MeHg group (Fig. 10A).

Superoxide dismutase activity was also significantly lower in the MeHg group (25.2  $\pm$  8.4 U/mg protein;  $p < 0.05$ ) than in the control




Fig. 5. Methylmercury (MeHg) intoxication and *Coriandrum sativum* aqueous extract (CSAE) treatment effects on neonates' balance and motor coordination. (A) Latency; (B) Number of falls. Data are shown as the mean  $\pm$  SD (n = 14/group). \*, p < 0.05 compared to the control group; #, p < 0.05 compared to MeHg group (two-way ANOVA with repeated measures and Tukey test).




Fig. 6. Methylmercury (MeHg) intoxication and *Coriandrum sativum* aqueous extract (CSAE) treatment effects on neonates' balance. Data are shown as the mean  $\pm$  SD (n = 14/group). \*, p < 0.05 compared to the control group; #, p < 0.05 compared to the CSAE-treated group (two-way ANOVA with repeated measures and Tukey test); BWT/Q-12MM, 12 mm wide quadrangular pole; BWT/Q-5MM, 5 mm wide quadrangular pole; BWT/C-17MM, 17 mm diameter circular pole; BWT/C-11MM, 11 mm diameter circular pole.




Fig. 7. Malondialdehyde (MDA) levels in the offspring. Results are expressed in pg/mg protein. Data are shown as the mean  $\pm$  SD (n = 14/group). \*, p < 0.05 compared to the control group; #, p < 0.05 compared to the CSAE-treated group (two-way ANOVA and Tukey test); CSAE, *Coriandrum sativum* aqueous extract; MeHg, methylmercury.

group ( $38.9 \pm 12.0$  U/mg protein), while the CSAE ( $39.0 \pm 10.8$  U/mg protein) and MeHg + CSAE ( $39.3 \pm 8.0$  U/mg protein) groups had enzyme activities similar to that of the control group. Significantly higher superoxide dismutase activity was observed in the MeHg + CSAE group than in the MeHg group (Fig. 10B).

#### 4. Discussion

The results show that the progeny of mothers subjected to MeHg


intoxication and treated with CSAE had functional motor improvements during spontaneous (Fig. 3A and B, and 6) and forced locomotion (Fig. 5A and B) assessments. The progeny also showed increased grip strength (Fig. 4). Both outcomes may be linked to a reduction in total mercury levels in the CSAE-treated group.

The harmful effects of MeHg on motor performance have been described previously. Su and Okita (1976) were pioneers in investigating the effects of a single dose of MeHg on Day 10 of gestation. They observed reduced spontaneous locomotor activity in the offspring, which reverted as the specimens grew older. In another study with a similar animal model, Ghizoni et al. (2018) utilized polyunsaturated fatty acids as a treatment for MeHg intoxication, but the test subjects showed no improvement in their behavioral parameters, had compromised motor coordination, and had an accumulation of MeHg in their cerebellar tissue. We hypothesized that the bioactive compounds with intense antioxidant activity present in *C. sativum* might ameliorate the motor-function damage that results from MeHg poisoning during neurodevelopment. Thus, we investigated several parameters related to motor function: spontaneous and forced ambulation, balance and equilibrium through two paradigms, and grip strength.


The mercury reduction seen in the MeHg + CSAE group might be associated with the extract's antioxidant activity (Fig. 1). The antioxidant properties observed in this study have also been seen in other studies that utilized *C. sativum* extracts (Abbassi et al., 2018; Dias et al., 2011; Harsha and Anilakumar, 2014; Pereira and Tavano, 2014; Sreelatha and Inbavalli, 2012). An aqueous extract was preferred, because it has been shown to contain greater amounts of polyphenols (Melo et al., 2003) with high antioxidant capacities such as caffeic acid and glycitin (Melo et al., 2005). Our results demonstrated that the utilized CSAE had high antioxidant activity, free-radical scavenging capacity, and total phenolic content.

An increase in mercury level promotes oxidative stress, with cell membranes and organelle phospholipids as the primary targets (Nath et al., 1996), which leads to higher ROS production, lipid peroxidation, protein synthesis disturbance, and enzyme disruption, all of which contribute to reduced antioxidant enzyme activity (Abdel-Salam et al., 2018; Olguin et al., 2018; Wilson et al., 2005). Methylmercury also leads to neurotoxicity in humans and in animal models, but to date, there is no efficient treatment available for mercury poisoning (Cao et al., 2011; Nielsen and Andersen, 1991; Schutzmeier et al., 2018).

This is the first time the protective properties of CSAE against mercury-induced neurotoxicity have been demonstrated in an *in vivo* model, where the behavioral parameters improved after treatment, likely due to a decrease in oxidative stress. *Coriandrum sativum* has already been shown to have protective effects against oxidative damage in rat brains after lead poisoning at doses of 250 and 500 mg/kg (Vekaria et al., 2012; Velaga et al., 2014). Other extracts have also been investigated to assess their effects on heavy metal poisoning. A *Polygala paniculata* extract prevented lipid peroxidation in the rat cerebellum



**Fig. 8.** Reactive oxygen species assessment in the offspring. (A) Nitrite levels expressed in  $\mu\text{mol}/\text{mg}$  protein; (B) Superoxide anion intensity expressed in absorbance/ $\text{mg}$  protein. Data are shown as the mean  $\pm$  SD ( $n = 14/\text{group}$ ). \*,  $p < 0.05$  compared to the control group; #,  $p < 0.05$  compared to the CSAE-treated group (two-way ANOVA and Tukey test); CSAE, *Coriandrum sativum* aqueous extract; MeHg, methylmercury.



**Fig. 9.** Total glutathione levels in the offspring. Results are expressed in  $\text{pmol}/\mu\text{g}$  protein. Data are shown as the mean  $\pm$  SD ( $n = 14/\text{group}$ ). \*,  $p < 0.05$  compared to the control group; #,  $p < 0.05$  compared to the CSAE-treated group (two-way ANOVA and Tukey test); CSAE, *Coriandrum sativum* aqueous extract; MeHg, methylmercury.

and ameliorated motor damage in an *in vivo* model similar to the one utilized in this study (Farina et al., 2005).


The groups treated with CSAE showed normal ROS levels in the cerebellum, protection against anion superoxide and nitrite damage, and reduced lipid peroxidation. These effects can be attributed to the

extract's chemical composition (Table 1), which is rich in flavonoids such as quercetin-3-glucuronide, isoquercetin, and rutin (Kunzemann and Herrmann, 1977; Nagata et al., 1999), phenolic compounds, and anthocyanins. It is important to note that these compounds are linked to higher antioxidant enzyme production and protective properties against ROS (Dai et al., 2012; Kozłowska et al., 2016; Rashid et al., 2014).

Yang et al. (2012) stimulated protein expression related to neuroplasticity and antioxidant activity in rat brains by administering an anthocyanin extract that provided neuroprotection and increased functional performance. This result was similar to those obtained in our study, where the offspring showed increased functional performance after treating with CSAE.

We identified several minerals in CSAE that can contribute to both antioxidant enzyme synthesis and redox balance modulation, properties involved in the reduced oxidative patterns and improved behavioral performance seen in the treated offspring. Micronutrients and macronutrients play a role in CNS homeostasis and the synthesis of antioxidant enzymes and protein structures, but they can also interact with heavy metals (Brose et al., 2014; Lau et al., 2019; Liu et al., 1992). Elements such as Cu and Fe are redox-active elements and act as co-factors to enzymes in various metabolic reactions, free-radical scavenging, and the biosynthesis of neurotransmitters (Lebovitz et al., 1996; Salama et al., 2009; Strange et al., 2003). A dysfunction in the balance of these minerals can lead to neural changes and maladies such as Parkinson disease (Ji et al., 2017; Mischley et al., 2017).

Elements such as Mn and Zn are essential for proper enzyme



**Fig. 10.** Antioxidant enzyme activity. (A) Catalase (CAT) and (B) superoxide dismutase (SOD) activities in the progeny of MeHg-poisoned mothers. Results are expressed in  $\text{nmol}$  formaldehyde/ $\text{min}/\text{mg}$  protein and  $\text{U}/\text{mg}$  protein, respectively. Data are shown as the mean  $\pm$  SD ( $n = 14/\text{group}$ ). \*,  $p < 0.05$  compared to the control group; #,  $p < 0.05$  compared to the CSAE-treated group (two-way ANOVA and Tukey test); CSAE, *Coriandrum sativum* aqueous extract; MeHg, methylmercury.

functioning (hydrolases, isomerases, ligases, lyases, reductases, and transferases) in various biological systems such as lipid, protein, and carbohydrate metabolism, ATP production, redox homeostasis (SOD2), DNA synthesis, and cellular signaling, where they act as neuromodulators in protein structures with antioxidant properties such as SOD1 (Adebayo et al., 2016; el-Sewedy et al., 1974; Sharma et al., 2010; Sommer et al., 2018). Zinc also acts in glutamatergic synaptic transmission, inhibiting N-methyl-D-aspartate and  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor activity, thus modulating neuronal excitability and long-term synaptic plasticity (Takeda et al., 2004). The levels of components such as Zn and Cu are altered in the CNS after heavy metal poisoning due to oxidative stress, but the treatment with CSAE was able to restore the Zn, Cu, and Fe levels to their physiological state (Ren et al., 2009; Velaga et al., 2014).

The results presented in this study support the hypothesis that a CSAE can protect an offspring's cerebellum after its progenitor is poisoned with mercury by reducing oxidative stress, thereby reducing damages to motor function. The total glutathione levels and catalase and SOD antioxidant-enzyme activities returned to their basal levels after treatment with CSAE. Further studies are necessary for a better understanding of the protective effects of CSAE against cerebellar mercury-induced toxicity.

#### Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Acknowledgments

The authors acknowledge CNPq and CAPES for their financial support.

#### References

- Abbasi, A., Mahmoudi, H., Zaouali, W., MRabet, Y., Casabianca, H., Hosni, K., 2018. Enzyme-aided release of bioactive compounds from coriander (*Coriandrum sativum* L.) seeds and their residue by-products and evaluation of their antioxidant activity. *J. Food Sci. Technol.* 55, 3065–3076.
- Abdel-Salam, A.M., Al Hemaied, W.A., Afifi, A.A., Othman, A.I., Farrag, A.R.H., Zeitoun, M.M., 2018. Consolidating probiotic with dandelion, coriander and date palm seeds extracts against mercury neurotoxicity and for maintaining normal testosterone levels in male rats. *Toxicol. Rep.* 5, 1069–1077.
- Adebayo, O.L., Adenuga, G.A., Sandhir, R., 2016. Selenium and zinc protect brain mitochondrial antioxidants and electron transport chain enzymes following postnatal protein malnutrition. *Life Sci.* 152, 145–155.
- Aissaoui, A., El-Hilaly, J., Israili, Z.H., Lyoussi, B., 2008. Acute diuretic effect of continuous intravenous infusion of an aqueous extract of *Coriandrum sativum* L. in anesthetized rats. *J. Ethnopharmacol.* 115, 89–95.
- Aissaoui, A., Zizi, S., Israili, Z.H., Lyoussi, B., 2011. Hypoglycemic and hypolipidemic effects of *Coriandrum sativum* L. in *Meriones shawi* rats. *J. Ethnopharmacol.* 137, 652–661.
- Akhter, F., Hashim, A., Khan, M.S., Ahmad, S., Iqbal, D., Srivastava, A.K., Siddiqui, M.H., 2013. Antioxidant,  $\alpha$ -amylase inhibitory and oxidative DNA damage protective property of *Boerhaavia diffusa* (Linn.) root. *South Afr. J. Bot.* 88, 265–272.
- Al-Mofleh, I.A., Alhaider, A.A., Mossa, J.S., Al-Sohaibani, M.O., Rafatullah, S., Qureshi, S., 2006. Protection of gastric mucosal damage by *Coriandrum sativum* L. pretreatment in Wistar albino rats. *Environ. Toxicol. Pharmacol.* 22, 64–69.
- Al-Osaimi, A., El-Ansary, A., Al-Daihan, S., Bhat, R.S., Ben Bacha, A., 2018. Therapeutic and protective potency of bee pollen against neurotoxic effects induced by prenatal exposure of rats to methyl mercury. *J. Mol. Neurosci.* MN 65, 327–335.
- Amarante, C.B., Müller, R.C.S., Dantas, K.G.F., Alves, C.N., Müller, A.H., Palheta, D.C., 2010. Chemical composition and nutritional value of leaves and fruits of aninga (*Montrichardia linifera*, Araceae) for large herbivores. *Acta Amazonica* 40, 729–736.
- Amonpatumrat, S., Sakurai, H., Wiriyasermkul, P., Khunweeraphong, N., Nagamori, S., Tanaka, H., Piyachaturawat, P., Kanai, Y., 2008. L-glutamate enhances methylmercury toxicity by synergistically increasing oxidative stress. *J. Pharmacol. Sci.* 108, 280–289.
- Anaigoudari, A., Hosseini, M., Karami, R., Vafaei, F., Mohammadpour, T., Ghorbani, A., Sadeghnia, H.R., 2016. The effects of different fractions of *Coriandrum sativum* on pentylenetetrazole-induced seizures and brain tissues oxidative damage in rats. *Avicenna J. Phytomed.* 6, 223–235.
- Andersen, H.R., Andersen, O., 1993. Effects of dietary alpha-tocopherol and beta-carotene on lipid peroxidation induced by methyl mercuric chloride in mice. *Pharmacol. Toxicol.* 73, 192–201.
- Aschner, M., Clarkson, T.W., 1989. Methyl mercury uptake across bovine brain capillary endothelial cells in vitro: the role of amino acids. *Pharmacol. Toxicol.* 64, 293–297.
- Askar, A., Treptow, H., 1993. Quality Assurance in Tropical Fruit Processing. Springer-Verlag, Berlin ; New York.
- Baehner, R.L., Boxer, L.A., Davis, J., 1976. The biochemical basis of nitroblue tetrazolium reduction in normal human and chronic granulomatous disease polymorphonuclear leukocytes. *Blood* 48, 309–313.
- Ballester, F., Iniguez, C., Murcia, M., Guxens, M., Basterretxea, M., Rebagliato, M., Vioque, J., Lertxundi, A., Fernandez-Somoano, A., Tardon, A., Sunyer, J., Llop, S., 2018. Prenatal exposure to mercury and longitudinally assessed fetal growth: relation and effect modifiers. *Environ. Res.* 160, 97–106.
- Balthasar, C., Stangl, H., Widhalm, R., Granitzer, S., Hengstschlager, M., Gundacker, C., 2017. Methylmercury uptake into BeWo cells depends on LAT2-4F2hc, a system L amino acid transporter. *Int. J. Mol. Sci.* 18.
- Bhuiyan, M.N.I., Begum, J., Sultana, M., 2009. Chemical composition of leaf and seed essential oil of *Coriandrum sativum* L. from Bangladesh. *Bangladesh J. Pharmacol.* 4, 150–153.
- Bisi, T.L., Lepoint, G., Azevedo, A.F., Dorneles, P.R., Flach, L., Das, K., Lailson-Brito, J., 2012. Trophic relationships and mercury biomagnification in Brazilian tropical coastal food webs. *Ecol. Indicat.* 18, 291–302.
- Bogavac, M., Karaman, M., Janjusevic, L., Sudji, J., Radovanovic, B., Novakovic, Z., Simeunovic, J., Bozin, B., 2015. Alternative treatment of vaginal infections - in vitro antimicrobial and toxic effects of *Coriandrum sativum* L. and *Thymus vulgaris* L. essential oils. *J. Appl. Microbiol.* 119, 697–710.
- Boudou, A., Ribeyre, F., 1997. Aquatic ecotoxicology: from the ecosystem to the cellular and molecular levels. *Environ. Health Perspect.* 105 (Suppl. 1), 21–35.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- Brose, J., La Fontaine, S., Wedd, A.G., Xiao, Z., 2014. Redox sulfur chemistry of the copper chaperone Atox1 is regulated by the enzyme glutaredoxin 1, the reduction potential of the glutathione couple GSSG/2GSH and the availability of Cu(I). *Metalloids: Integr. Biometal Sci.* 6, 793–808.
- Buchanan, S., Anglen, J., Turyk, M., 2015. Methyl mercury exposure in populations at risk: analysis of NHANES 2011–2012. *Environ. Res.* 140, 56–64.
- Cao, Y., Chen, A., Jones, R.L., Radcliffe, J., Dietrich, K.N., Caldwell, K.L., Peddada, S., Rogan, W.J., 2011. Efficacy of succimer chelation of mercury at background exposures in toddlers: a randomized trial. *J. Pediatr.* 158, 480–485 e481.
- Carter, R.J., Morton, J., Dunnett, S.B., 2001. Motor coordination and balance in rodents. *Curr. Protocols Neurosci.* 1–14 (Chapter 8), Unit 8.12.
- Cavalcante, A.R., Junior, J.A.S., Gheyk, H.R., Dias, N.S., Paz, V.P.S., 2016. Produção e composição mineral do coentro em sistema hidroponico de baixo custo. *Irriga* 21, 685–696.
- Cheng, J., Fujimura, M., Bo, D., 2015. Assessing pre/post-weaning neurobehavioral development for perinatal exposure to low doses of methylmercury. *J. Environ. Sci.* 38, 36–41.
- Chithra, V., Leelamma, S., 1999. *Coriandrum sativum* changes the levels of lipid peroxides and activity of antioxidant enzymes in experimental animals. *Indian J. Biochem. Biophys.* 36, 59–61.
- Coelho, J.P., Mieirol, C.L., Pereira, E., Duarte, A.C., Pardal, M.A., 2013. Mercury biomagnification in a contaminated estuary food web: effects of age and trophic position using stable isotope analyses. *Mar. Pollut. Bull.* 69, 110–115.
- Costa, J.M.F.J., Lima, A., Rodrigues, D.J., Khoury, E.D.T., Souza, G.D.S., Silveira, L.C.L., Pinheiro, M., 2017. Emotional and motor symptoms in riverside dwellers exposed to mercury in the Amazon. *Revista brasileira de epidemiologia = Brazil. J. Epidemiol.* 20, 212–224.
- da Silva, F.B.R., Cunha, P.A., Ribera, P.C., Barros, M.A., Cartagenes, S.C., Fernandes, L.M.P., Teixeira, F.B., Fontes-Junior, E.A., Prediger, R.D., Lima, R.R., Maia, C.S.F., 2018. Heavy chronic ethanol exposure from adolescence to adulthood induces cerebellar neuronal loss and motor function damage in female rats. *Front. Behav. Neurosci.* 12, 88.
- Dai, L., Dong, X., Ma, H., 2012. Antioxidative and chelating properties of anthocyanins in *Azolla imbricata* induced by cadmium. *Pol. J. Environ. Stud.* 21, 837–844.
- Deng, Y., Xu, Z., Xu, B., Liu, W., Wei, Y., Li, Y., Feng, S., Yang, T., 2014. Exploring cross-talk between oxidative damage and excitotoxicity and the effects of riluzole in the rat cortex after exposure to methylmercury. *Neurotox. Res.* 26, 40–51.
- Dewanjee, S., Joardar, S., Bhattacharjee, N., Dua, T.K., Das, S., Kalita, J., Manna, P., 2017. Edible leaf extract of *Ipomoea aquatica* Forssk. (Convolvulaceae) attenuates doxorubicin-induced liver injury via inhibiting oxidative impairment, MAPK activation and intrinsic pathway of apoptosis. *Food Chem. Toxicol. : Int. J. Publ. Br. Industr. Biol. Res. Assoc.* 105, 322–336.
- Dias, M.I., Barros, L., Sousa, M.J., Ferreira, I.C., 2011. Comparative study of lipophilic and hydrophilic antioxidants from in vivo and in vitro grown *Coriandrum sativum*. *Plant Foods Hum. Nutr.* 66, 181–186.
- Dos Santos, A.A., Lopez-Granero, C., Farina, M., Rocha, J.B.T., Bowman, A.B., Aschner, M., 2018. Oxidative stress, caspase-3 activation and cleavage of ROCK-1 play an essential role in MeHg-induced cell death in primary astroglial cells. *Food Chem. Toxicol. : Int. J. Publ. Br. Industr. Biol. Res. Assoc.* 113, 328–336.
- el-Sewedy, S.M., Abdel-Tawab, G.A., el-Zoghby, S.M., Zeitoun, R., Mostafa, M.H., Shalaby, S.M., 1974. Studies with tryptophan metabolites in vitro. Effect of zinc, manganese, copper and cobalt ions on kynurenine hydrolase and kynurenine aminotransferase in normal mouse liver. *Biochem. Pharmacol.* 23, 2557–2565.
- Emamghoreishi, M., Heidari-Hamedani, G., 2006. Sedative-hypnotic activity of extracts and essential oil of coriander seeds. *Iran. J. Med. Sci.* 31, 22–27.
- Emamghoreishi, M., Khasaki, M., Aazam, M.F., 2005. *Coriandrum sativum*: evaluation of

- its anxiolytic effect in the elevated plus-maze. *J. Ethnopharmacol.* 96, 365–370.
- Farina, M., Franco, J.L., Ribas, C.M., Meotti, F.C., Missau, F.C., Pizzolatti, M.G., Dafre, A.L., Santos, A.R., 2005. Protective effects of *Polygala paniculata* extract against methylmercury-induced neurotoxicity in mice. *J. Pharm. Pharmacol.* 57, 1503–1508.
- Feng, S., Xu, Z., Liu, W., Li, Y., Deng, Y., Xu, B., 2014. Preventive effects of dextromethorphan on methylmercury-induced glutamate dyshomeostasis and oxidative damage in rat cerebral cortex. *Biol. Trace Elem. Res.* 159, 332–345.
- Fernandes, L.M.P., Cartagenes, S.C., Barros, M.A., Carvalheiro, T., Castro, N.C.F., Schamne, M.G., Lima, R.R., Prediger, R.D., Monteiro, M.C., Fontes-Junior, E.A., Cunha, R.A., Maia, C.S.F., 2018a. Repeated cycles of binge-like ethanol exposure induce immediate and delayed neurobehavioral changes and hippocampal dysfunction in adolescent female rats. *Behav. Brain Res.* 350, 99–108.
- Fernandes, L.M.P., Lopes, K.S., Santana, L.N.S., Fontes-Junior, E.A., Ribeiro, C., Silva, M.C.F., de Oliveira Paraense, R.S., Crespo-Lopez, M.E., Gomes, A.R.Q., Lima, R.R., Monteiro, M.C., Maia, C.S.F., 2018b. Repeated cycles of binge-like ethanol intake in adolescent female rats induce motor function impairment and oxidative damage in motor cortex and liver, but not in blood. *Oxid. Med. Cell. Longev.* 2018, 3467531.
- Foyer, C.H., Shigeoka, S., 2011. Understanding oxidative stress and antioxidant functions to enhance photosynthesis. *Plant Physiol.* 155, 93–100.
- Frankel, N.E., Meyer, A.S., 2000. The problems of using one-dimensional methods to evaluate multifunctional food and biological antioxidants. *J. Sci. Food Agric.* 80, 1925–1941.
- Fujimura, M., Cheng, J., Zhao, W., 2012. Perinatal exposure to low-dose methylmercury induces dysfunction of motor coordination with decreases in synaptophysin expression in the cerebellar granule cells of rats. *Brain Res.* 1464, 1–7.
- Fujimura, M., Usuki, F., 2018. Methylmercury induces oxidative stress and subsequent neural hyperactivity leading to cell death through the p38 MAPK-CREB pathway in differentiated SH-SY5Y cells. *Neurotoxicology* 67, 226–233.
- Gandhi, D.N., Panchal, G.M., Dhull, D.K., 2014. Neurobehavioral toxicity in progeny of rat mothers exposed to methylmercury during gestation. *Annali dell'Istituto superiore di sanità* 50, 28–37.
- Ghizoni, H., Ventura, M., Colle, D., Goncalves, C.L., de Souza, V., Hartwig, J.M., Santos, D.B., Naime, A.A., Cristina de Oliveira Souza, V., Lopes, M.W., Barbosa Jr., F., Brocardo, P.S., Farina, M., 2018. Effects of perinatal exposure to n-3 polyunsaturated fatty acids and methylmercury on cerebellar and behavioral parameters in mice. *Food Chem. Toxicol. : Int. J. Publ. Br. Industr. Biol. Res. Assoc.* 120, 603–615.
- Go, S., Kurita, H., Matsumoto, K., Hatano, M., Inden, M., Hozumi, I., 2018. Methylmercury causes epigenetic suppression of the tyrosine hydroxylase gene in an in vitro neuronal differentiation model. *Biochem. Biophys. Res. Commun.* 502, 435–441.
- Gutiérrez, J., Baraibar, A.M., Albinana, E., Velasco, P., Solis, J.M., Hernandez-Guijo, J.M., 2018. Methylmercury reduces synaptic transmission and neuronal excitability in rat hippocampal slices. *Pflug. Arch. Eur. J. Physiol.* 470, 1221–1230.
- Harsha, S.N., Anilakumar, K.R., 2014. In vitro free radical scavenging and DNA damage protective property of *Coriandrum sativum* L. leaves extract. *J. Food Sci. Technol.* 51, 1533–1539.
- Heimfarth, L., Delgado, J., Mingori, M.R., Moresco, K.S., Pureur, R.P., Gelain, D.P., Moreira, J.C.F., 2018. Delayed neurochemical effects of prenatal exposure to MeHg in the cerebellum of developing rats. *Toxicol. Lett.* 284, 161–169.
- Hosseinzadeh, S., Ghalesefidi, M.J., Azami, M., Mohaghegh, M.A., Hejazi, S.H., Ghomashlooyan, M., 2016. In vitro and in vivo anthelmintic activity of seed extract of *Coriandrum sativum* compared to Niclosamid against *Hymenolepis nana* infection. *J. Parasit. Dis. : Off. Organ Indian Soc. Parasitol.* 40, 1307–1310.
- Hwang, E., Lee, D.G., Park, S.H., Oh, M.S., Kim, S.Y., 2014. Coriander leaf extract exerts antioxidant activity and protects against UVB-induced photoaging of skin by regulation of procollagen type I and MMP-1 expression. *J. Med. Food* 17, 985–995.
- Jabeen, Q., Bashir, S., Lyoussi, B., Gilani, A.H., 2009. Coriander fruit exhibits gut modulatory, blood pressure lowering and diuretic activities. *J. Ethnopharmacol.* 122, 123–130.
- Ji, B., Wang, M., Gao, D., Xing, S., Li, L., Liu, L., Zhao, M., Qi, X., Dai, K., 2017. Combining nanoscale magnetic nimodipine liposomes with magnetic resonance image for Parkinson's disease targeting therapy. *Nanomedicine* 12, 237–253.
- Johansson, L.H., Borg, L.A., 1988. A spectrophotometric method for determination of catalase activity in small tissue samples. *Anal. Biochem.* 174, 331–336.
- Justesen, U., Knuthsen, P., 2001. Composition of flavonoids in fresh herbs and calculation of flavonoid intake by use of herbs in traditional Danish dishes. *Food Chem.* 73, 245–270.
- Kansal, L., Sharma, V., Sharma, A., Lodi, S., Sharma, S.H., 2011. Protective role of *Coriandrum sativum* (Coriander) extracts against lead nitrate induced oxidative stress and tissue damage in the liver and kidney in male mice. *Int. J. Appl. Biol. Pharm. Technol.* 2, 65–83.
- Karunasagar, D., Krishna, M.V., Rao, S.V., Arunachalam, J., 2005. Removal and pre-concentration of inorganic and methyl mercury from aqueous media using a sorbent prepared from the plant *Coriandrum sativum*. *J. Hazard Mater.* 118, 133–139.
- Kazempour, S.F., Langehbiz, S.V., Hosseini, M., Shafei, M.N., Ghorbani, A., Pourganji, M., 2015. The analgesic effects of different extracts of aerial parts of *coriandrum sativum* in mice. *Int. J. Biomed. Sci.* 11, 23–28.
- Kopec, A., Sikora, E., Piatkowska, E., Borczak, B., Czech, T., 2016. Possible protective role of elderberry fruit lyophilizate against selected effects of cadmium and lead intoxication in Wistar rats. *Environ. Sci. Pollut. Res. Int.* 23, 8837–8848.
- Kozłowska, M., Gruczynska, E., Scibisz, I., Rudzinska, M., 2016. Fatty acids and sterols composition, and antioxidant activity of oils extracted from plant seeds. *Food Chem.* 213, 450–456.
- Kujawska, M., Kant, P., Mayoral, I.H., Ignatowicz, E., Sikora, J., Oszmianski, J., Czapski, J., Jodynis-Liebert, J., 2016. Effect of chokeberry juice on N-Nitrosodimethylamine-Induced rat liver carcinogenesis. *J. Environ. Pathol. Toxicol. Oncol. : Off. Organ Int. Soc. Environ. Toxicol. Canc.* 35, 317–331.
- Kunzemann, J., Herrmann, K., 1977. [Isolation and identification of flavon(ol)-O-glycosides in caraway (*Carum carvi* L.), fennel (*Foeniculum vulgare* Mill.), anise (*Pimpinella anisum* L.), and coriander (*Coriandrum sativum* L.) and of flavon-C-glycosides in anise. I. Phenolics of spices (author's transl)]. *Zeitschrift für Lebensmittel-Untersuchung und -Forschung* 164, 194–200.
- Lau, H., Shahar, S., Mohamad, M., Rajab, N.F., Yahya, H.M., Din, N.C., Hamid, H.A., 2019. Relationships between dietary nutrients intake and lipid levels with functional MRI dorsolateral prefrontal cortex activation. *Clin. Interv. Aging* 14, 43–51.
- Lebovitz, R.M., Zhang, H., Vogel, H., Cartwright Jr., J., Dionne, L., Lu, N., Huang, S., Matzuk, M.M., 1996. Neurodegeneration, myocardial injury, and perinatal death in mitochondrial superoxide dismutase-deficient mice. *Proc. Natl. Acad. Sci. U.S.A.* 93, 9782–9787.
- Liu, W., Xu, Z., Yang, T., Deng, Y., Xu, B., Feng, S., 2016. Tea polyphenols protect against methylmercury-induced cell injury in rat primary cultured astrocytes, involvement of oxidative stress and glutamate uptake/metabolism disorders. *Mol. Neurobiol.* 53, 2995–3009.
- Liu, X., Nordberg, G.F., Jin, T., 1992. Increased urinary excretion of zinc and copper by mercuric chloride injection in rats. *Biomaterials: Int. J. Role Metal Ions Biol. Biochem. Med.* 5, 17–22.
- Liu, Y., Buchanan, S., Anderson, H.A., Xiao, Z., Persky, V., Turyk, M.E., 2018. Association of methylmercury intake from seafood consumption and blood mercury level among the Asian and Non-Asian populations in the United States. *Environ. Res.* 160, 212–222.
- Mahendra, P., Bisht, S., 2011. Anti-anxiety activity of *Coriandrum sativum* assessed using different experimental anxiety models. *Indian J. Pharmacol.* 43, 574–577.
- Maier, C.M., Chan, P.H., 2002. Role of superoxide dismutases in oxidative damage and neurodegenerative disorders. *The Neuroscientist : Rev. J. Bring. Neurobiol. Neurol. Psychiatr.* 8, 323–334.
- Mancini, J.D., Autio, D.M., Atchison, W.D., 2009. Continuous exposure to low concentrations of methylmercury impairs cerebellar granule cell migration in organotypic slice culture. *Neurotoxicology* 30, 203–208.
- McDowell, M.A., Dillon, C.F., Osterloh, J., Bolger, P.M., Pellizzari, E., Fernando, R., Montes de Oca, R., Schober, S.E., Sinks, T., Jones, R.L., Mahaffey, K.R., 2004. Hair mercury levels in U.S. children and women of childbearing age: reference range data from NHANES 1999–2000. *Environ. Health Perspect.* 112, 1165–1171.
- Meerovich, I., Muthukrishnan, N., Johnson, G.A., Erazo-Oliveras, A., Pellois, J.P., 2014. Photodamage of lipid bilayers by irradiation of a fluorescently labeled cell-penetrating peptide. *Biochim. Biophys. Acta* 1840, 507–515.
- Melo, E.A., Bion, F.M., Filho, J.M., Guerra, N.B., 2003. In vivo antioxidant effect of aqueous and etheric coriander (*Coriandrum sativum* L.) extracts. *Eur. J. Lipid Sci. Technol.* 105, 483–487.
- Melo, E.A., Filho, J.M., Guerra, N.B., 2005. Characterization of antioxidant compounds in aqueous coriander extract (*Coriandrum sativum* L.). *Lebensm. Wiss. Technol.* 38, 15–19.
- Mischley, L.K., Lau, R.C., Bennett, R.D., 2017. Role of diet and nutritional supplements in Parkinson's disease progression. *Oxid. Med. Cell. Longev.* 2017, 6405278.
- Moncada, S., 1992. The 1991 Ulf von Euler Lecture. The L-arginine: nitric oxide pathway. *Acta Physiol. Scand.* 145, 201–227.
- Msaada, K., Jemia, M.B., Salem, N., Bachrouh, O., Sriti, J., Tammar, S., Bettaieb, I., Jabri, I., Kefi, S., Limam, F., Marzouk, B., 2014. Antioxidant activity of methanolic extracts from three coriander (*Coriandrum sativum* L.) fruit varieties. *Arab. J. Chem.* 10, S3176–S3183.
- Murcia, M., Ballester, F., Enning, A.M., Iniguez, C., Valvi, D., Basterrechea, M., Rebagliato, M., Vioque, J., Maruri, M., Tardon, A., Riano-Galan, I., Vrijheid, M., Llop, S., 2016. Prenatal mercury exposure and birth outcomes. *Environ. Res.* 151, 11–20.
- Mzid, M., Badraoui, R., Khedir, S.B., Sahnoun, Z., Rebai, T., 2017. Protective effect of ethanolic extract of *Urtica urens* L. against the toxicity of imidacloprid on bone remodeling in rats and antioxidant activities. *Biomed. Pharmacother. = Biomedicine & pharmacotherapie* 91, 1022–1041.
- Nagata, H., Takekoshi, S., Takagi, T., Honma, T., Watanabe, K., 1999. Antioxidative action of flavonoids, quercetin and catechin, mediated by the activation of glutathione peroxidase. *Tokai J. Exp. Clin. Med.* 24, 1–11.
- Nath, K.A., Croatt, A.J., Likely, S., Behrens, T.W., Warden, D., 1996. Renal oxidant injury and oxidant response induced by mercury. *Kidney Int.* 50, 1032–1043.
- Nazima, B., Manoharan, V., Miltonprabu, S., 2016. Oxidative stress induced by cadmium in the plasma, erythrocytes and lymphocytes of rats: attenuation by grape seed proanthocyanidins. *Hum. Exp. Toxicol.* 35, 428–447.
- Nielsen, J.B., Andersen, O., 1991. Effect of four thiol-containing chelators on disposition of orally administered mercuric chloride. *Hum. Exp. Toxicol.* 10, 423–430.
- Nims, R.W., Darbyshire, J.F., Saavedra, J.E., Christodoulou, D., Hanbauer, I., Cox, G.W., Grisham, M.B., Laval, F., Cook, J.A., Krishna, M.C., Wink, D.A., 1995. Colorimetric methods for the determination of nitric oxide concentration in neutral aqueous solutions. *Comp. Methods Enzymol.* 7, 48–54.
- Ohkawa, H., Ohishi, N., Yagi, K., 1978. Assay for lipid peroxides in animal tissues by Thiobarbituric acid reaction. *Anal. Biochem.* 95, 351–358.
- Ola-Mudathir, F.K., Suru, S.M., 2015. Onion and garlic extracts as potential antidotes for cadmium-induced biochemical alterations in prostate glands of rats. *Andrologia* 47, 1075–1082.
- Olaleye, M.T., Akinmoladun, A.C., Ogunboye, A.A., Akindahunsi, A.A., 2010. Antioxidant activity and hepatoprotective property of leaf extracts of *Boerhaavia diffusa* Linn against acetaminophen-induced liver damage in rats. *Food Chem. Toxicol. : Int. J. Publ. Br. Industr. Biol. Res. Assoc.* 48, 2200–2205.
- Olguin, N., Muller, M.L., Rodriguez-Farre, E., Sunol, C., 2018. Neurotransmitter amines and antioxidant agents in neuronal protection against methylmercury-induced cytotoxicity in primary cultures of mice cortical neurons. *Neurotoxicology* 69, 278–287.

- Onishchenko, N., Karpova, N., Sabri, F., Castren, E., Ceccatelli, S., 2008. Long-lasting depression-like behavior and epigenetic changes of BDNF gene expression induced by perinatal exposure to methylmercury. *J. Neurochem.* 106, 1378–1387.
- Paarakh, M.P., Kumar, J.S., Rupesh, M., 2016. Analgesic activity of methanol extract of coriandrum sativum linn fruit. *World J. Pharm. Pharm. Sci.* 5, 1456–1464.
- Pathan, A.R., Kothawade, K.A., Logade, M.N., 2011. Anxiolytic and analgesic effect of seeds of Coriandrum sativum Linn. *Int. J. Res. Pharm. Chem.* 1, 1087–1099.
- Pellegriani, M., Ricci, A., Serio, A., Chaves-Lopez, C., Mazzarino, G., D'Amato, S., Lo Sterzo, C., Paparella, A., 2018. Characterization of essential oils obtained from abruzzo autochthonous plants: antioxidant and antimicrobial activities assessment for food application. *Foods* 7.
- Pereira, M.P., Tavano, O.L., 2014. Use of different spices as potential natural antioxidant additives on cooked beans (*Phaseolus vulgaris*). Increase of DPPH radical scavenging activity and total phenolic content. *Plant Foods Hum. Nutr.* 69, 337–343.
- Pinheiro, M.C.N., Junco Nakanishi, J., Teichii Oikawa, T., Guimarães, G., Quaresma, M., Cardoso, B., Amoras, W.W., Harada, M., Magno, C., Vieira, J.L.F., Xavier, M.B., Bacelar, D.R., 2000. Exposição humana ao metilmercúrio em comunidades ribeirinhas da Região do Tapajós, Pará, Brasil. *Rev. Soc. Bras. Med. Trop.* 33, 265–269.
- Quettier-Deleu, C., Gressier, B., Vasseur, J., Dine, T., Brunet, C., Luyckx, M., Cazin, M., Cazin, J.C., Bailleul, F., Trotin, F., 2000. Phenolic compounds and antioxidant activities of buckwheat (*Fagopyrum esculentum* Moench) hulls and flour. *J. Ethnopharmacol.* 72, 35–42.
- Rashid, K., Wachira, F.N., Nyabuga, J.N., Wanyonyi, B., Murilla, G., Isaac, A.O., 2014. Kenyan purple tea anthocyanins ability to cross the blood brain barrier and reinforce brain antioxidant capacity in mice. *Nutr. Neurosci.* 17, 178–185.
- Ren, H., Jia, H., Endo, H., Hayashi, T., 2009. Cadmium detoxification effect of Chinese parsley *Coriandrum sativum* in liver and kidney of rainbow trout *Oncorhynchus mykiss*. *Fish. Sci.* 75, 731–741.
- Roos, D.H., Puntel, R.L., Lugokenski, T.H., Ineu, R.P., Bohrer, D., Burger, M.E., Franco, J.L., Farina, M., Aschner, M., Rocha, J.B., de Vargas Barbosa, N.B., 2010. Complex methylmercury-cysteine alters mercury accumulation in different tissues of mice. *Basic Clin. Pharmacol. Toxicol.* 107, 789–792.
- Ruszkiewicz, J.A., Bowman, A.B., Farina, M., Rocha, J.B.T., Aschner, M., 2016. Sex- and structure-specific differences in antioxidant responses to methylmercury during early development. *Neurotoxicology* 56, 118–126.
- Sakamoto, M., Kakita, A., Domingo, J.L., Yamazaki, H., Oliveira, R.B., Sarrazin, S.L., Eto, K., Murata, K., 2017. Stable and episodic/bolus patterns of methylmercury exposure on mercury accumulation and histopathologic alterations in the nervous system. *Environ. Res.* 152, 446–453.
- Salama, Z.A.E., El-Beltagi, H.S., El-Hariri, D.M., 2009. Effect of Fe deficiency on antioxidant system in leaves of three flax cultivars. *Not. Bot. Horti Agrobot. Cluj-Napoca* 37, 122–128.
- Schutzmeier, P., Focil Baquerizo, A., Castillo-Tandazo, W., Focil, N., Bose-O'Reilly, S., 2018. Efficacy of N,N'-bis-(2-mercaptoethyl) isophthalamide on mercury intoxication: a randomized controlled trial. *Environ. Health: Glob. Access Sci. Source* 17, 15.
- Shao, Y., Yamamoto, M., Figeys, D., Ning, Z., Chan, H.M., 2015. Proteomic analysis of cerebellum in common marmoset exposed to methylmercury. *Toxicol. Sci.: Off. J. Soc. Toxicol.* 146, 43–51.
- Sharma, V., Kansal, L., Sharma, A., 2010. Prophylactic efficacy of Coriandrum sativum (Coriander) on testis of lead-exposed mice. *Biol. Trace Elem. Res.* 136, 337–354.
- Shen, A.N., Cummings, C., Pope, D., Hoffman, D., Newland, M.C., 2016. A bout analysis reveals age-related methylmercury neurotoxicity and nimodipine neuroprotection. *Behav. Brain Res.* 311, 147–159.
- Singleton, V.L., Rossi JR., J.A., 1965. Colorimetry of total phenolics with phosphomolybdenic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* 16, 144–158.
- Slamberova, R., Pomelova, M., Charousova, P., 2006. Postnatal development of rat pups is altered by prenatal methamphetamine exposure. *Progress Neuro-Psychopharmacol. Biol. Psychiatr.* 30, 82–88.
- Sommer, T., Bjerregaard-Andersen, K., Uribe, L., Etzerodt, M., Diezemann, G., Gauss, J., Casella, M., Morth, J.P., 2018. A fundamental catalytic difference between zinc and manganese dependent enzymes revealed in a bacterial isatin hydrolase. *Sci. Rep.* 8, 13104.
- Sreelatha, S., Inbavalli, R., 2012. Antioxidant, antihyperglycemic, and antihyperlipidemic effects of Coriandrum sativum leaf and stem in alloxan-induced diabetic rats. *J. Food Sci.* 77, T119–T123.
- Sridhar, K., Charles, A.L., 2019. In vitro antioxidant activity of Kyoho grape extracts in DPPH and ABTS assays: estimation methods for EC50 using advanced statistical programs. *Food Chem.* 275, 41–49.
- Stanley, J.L., Lincoln, R.J., Brown, T.A., McDonald, L.M., Dawson, G.R., Reynolds, D.S., 2005. The mouse beam walking assay offers improved sensitivity over the mouse rotarod in determining motor coordination deficits induced by benzodiazepines. *J. Psychopharmacol.* 19, 221–227.
- Strange, R.W., Antonyuk, S., Hough, M.A., Doucette, P.A., Rodriguez, J.A., Hart, P.J., Hayward, L.J., Valentine, J.S., Hasnain, S.S., 2003. The structure of holo and metal-deficient wild-type human Cu, Zn superoxide dismutase and its relevance to familial amyotrophic lateral sclerosis. *J. Mol. Biol.* 328, 877–891.
- Su, M.Q., Okita, G.T., 1976. Embryocidal and teratogenic effects of methylmercury in mice. *Toxicol. Appl. Pharmacol.* 38, 207–216.
- Takeda, A., Minami, A., Seki, Y., Oku, N., 2004. Differential effects of zinc on glutamatergic and GABAergic neurotransmitter systems in the hippocampus. *J. Neurosci. Res.* 75, 225–229.
- Takeshita, H., Yamamoto, K., Nozato, S., Inagaki, T., Tsuchimochi, H., Shirai, M., Yamamoto, R., Imaizumi, Y., Hongyo, K., Yokoyama, S., Takeda, M., Oguro, R., Takami, Y., Itoh, N., Takeya, Y., Sugimoto, K., Fukada, S.I., Rakugi, H., 2017. Modified forelimb grip strength test detects aging-associated physiological decline in skeletal muscle function in male mice. *Sci. Rep.* 7, 42323.
- Tavares, L.M., Camara, V.M., Malm, O., Santos, E.C., 2005. Performance on neurological development tests by riverine children with moderate mercury exposure in Amazonia, Brazil. *Cad. Saúde Pública* 21, 1160–1167.
- Tietze, F., 1969. Enzymic method for quantitative determination of nanogram amounts of total and oxidized glutathione: applications to mammalian blood and other tissues. *Anal. Biochem.* 27, 502–522.
- U.S. Environmental Protection Agency (EPA), 1998. Guidelines for neurotoxicity risk assessment. *Fed. Regist. Fed. Regis.* 63, 26926–26954.
- Usuki, F., Yamashita, A., Fujimura, M., 2011. Post-transcriptional defects of antioxidant selenoenzymes cause oxidative stress under methylmercury exposure. *J. Biol. Chem.* 286, 6641–6649.
- Vekaria, R.H., Patel, M.N., Bhalodiya, P.N., Patel, V., Desai, T., Tirgar, P., 2012. Evaluation of neuroprotective effect of Coriandrum sativum linn. Against ischemic-reperfusion insult in brain. *Int. J. Phytotherm.* 3, 186–193.
- Velaga, M.K., Yallapragada, P.R., Williams, D., Rajanna, S., Bettaiya, R., 2014. Hydroalcoholic seed extract of Coriandrum sativum (Coriander) alleviates lead-induced oxidative stress in different regions of rat brain. *Biol. Trace Elem. Res.* 159, 351–363.
- Vicentino, A.R.R., Menezes, F.S., 2007. Atividade antioxidante de tinturas vegetais, vendidas em farmácias com manipulação e indicadas para diversos tipos de doenças pela metodologia do DPPH. *Revista Brasileira de Farmacognosia* 17, 384–387.
- Vieira, H.C., Morgado, F., Soares, A.M., Abreu, S.N., 2015. Fish consumption recommendations to conform to current advice in regard to mercury intake. *Environ. Sci. Pollut. Res. Int.* 22, 9595–9602.
- von Horsten, S., Exton, M.S., Voge, J., Schult, M., Nagel, E., Schmidt, R.E., Westermann, J., Schedlowski, M., 1998. Cyclosporine A affects open field behavior in DA rats. *Pharmacol. Biochem. Behav.* 60, 71–76.
- Wangensteen, H., Samuelsen, A.B., Malterud, K.E., 2004. Antioxidant activity in extracts from coriander. *Food Chem.* 88, 293–297.
- Wheeler, C.R., Salzman, J.A., Elsayed, N.M., Omaye, S.T., Korte Jr., D.W., 1990. Automated assays for superoxide dismutase, catalase, glutathione peroxidase, and glutathione reductase activity. *Anal. Biochem.* 184, 193–199.
- Wilson, D.T., Polunas, M.A., Zhou, R., Halladay, A.K., Lowndes, H.E., Reuhl, K.R., 2005. Methylmercury alters Eph and ephrin expression during neuronal differentiation of P19 embryonal carcinoma cells. *Neurotoxicology* 26, 661–674.
- Xu, B., Xu, Z.F., Deng, Y., Liu, W., Yang, H.B., Wei, Y.G., 2012. Protective effects of MK-801 on methylmercury-induced neuronal injury in rat cerebral cortex: involvement of oxidative stress and glutamate metabolism dysfunction. *Toxicology* 300, 112–120.
- Yamamoto, M., Motomura, E., Yanagisawa, R., Hoang, V.A.T., Mogi, M., Mori, T., Nakamura, M., Takeya, M., Eto, K., 2019. Evaluation of neurobehavioral impairment in methylmercury-treated KK-Ay mice by dynamic weight-bearing test. *J. Appl. Toxicol.: JAT* 39, 221–230.
- Yang, H., Xu, Z., Liu, W., Wei, Y., Deng, Y., Xu, B., 2012. Effect of grape seed proanthocyanidin extracts on methylmercury-induced neurotoxicity in rats. *Biol. Trace Elem. Res.* 147, 156–164.
- Zimmermann, L.T., dos Santos, D.B., Colle, D., dos Santos, A.A., Hort, M.A., Garcia, S.C., Bressan, L.P., Bohrer, D., Farina, M., 2014. Methionine stimulates motor impairment and cerebellar mercury deposition in methylmercury-exposed mice. *J. Toxicol. Environ. Health Part A* 77, 46–56.