



ELSEVIER

Contents lists available at ScienceDirect

Food and Chemical Toxicology

journal homepage: www.elsevier.com/locate/foodchemtox

Evaluation of immunologic and intestinal effects in rats administered an E 171-containing diet, a food grade titanium dioxide (TiO₂)

Lance K. Blevins^a, Robert B. Crawford^a, Anthony Bach^{a,b}, Michael D. Rizzo^{a,c}, Jiajun Zhou^{a,d}, Joseph E. Henriquez^{a,e}, D. M. Isha Olive Khan^{a,e}, Sera Sermet^a, Lora L. Arnold^f, Karen L. Pennington^f, Nathalia P. Souza^f, Samuel M. Cohen^{f,g}, Norbert E. Kaminski^{a,b,e,*}

^a Institute for Integrative Toxicology, Michigan State University, East Lansing, MI, USA

^b Center for Research on Ingredient Safety, Michigan State University, East Lansing, MI, USA

^c Cell and Molecular Biology Program, Michigan State University, East Lansing, MI, USA

^d Department of Microbiology & Molecular Genetics, Michigan State University, East Lansing, MI, USA

^e Department of Pharmacology and Toxicology, Michigan State University, East Lansing, MI, USA

^f University of Nebraska Medical Center, Omaha, NE, USA

^g Havlik-Wall Professor of Oncology, USA

ARTICLE INFO

Keywords:

E 171

Dietary titanium dioxide

Gastrointestinal inflammation

T_{regulatory} cells

Intestinal carcinogenesis

ABSTRACT

The toxicity of dietary E 171, a food grade titanium dioxide was evaluated. A recent study reported rats receiving E 171 in water developed inflammation and aberrant crypt foci (ACF) in the gastrointestinal tract. Here, rats received food containing E 171 (7 or 100 days). The 100-day study included feeding E 171 after dimethylhydrazine (DMH) or vehicle only pretreatment. Food consumption was similar between treatment groups with maximum total cumulative E 171 exposure being 2617 mg/kg in 7 days and 29,400 mg/kg in 100 days. No differences were observed due to E 171 in the percentage of dendritic, CD4⁺ T or T_{reg} cells within Peyer's patches or the periphery, or in cytokine production in plasma, sections of jejunum, and colon in 7- or 100-day E 171 alone fed rats. Differences were observed for IL-17A in colon (400 ppm E 171 + DMH) and IL-12p70 in plasma (40 ppm E 171 + DMH). E 171 had no effect on histopathologic evaluations of small and large intestines, liver, spleen, lungs, or testes, and no effects on ACF, goblet cell numbers, or colonic gland length. Dietary E 171 administration (7- or 100-day), even at high doses, produced no effect on the immune parameters or tissue morphology.

1. Introduction

Additives are routinely incorporated into food products to enhance properties such as texture, taste, and color with potential applications including protection from microbial contamination and oxidation. However, consumer awareness of food additives is currently heightened and as such, there is much interest in the health impacts of such additives. Titanium dioxide (TiO₂) is a naturally occurring metal oxide that has been used as a food additive for more than 50 years. It confers a naturally white opaqueness to substances to which it is added due to its high refractive index. In the US, the Food and Drug Administration allows up to 1% by weight of TiO₂ as a food additive (Code of Federal Regulation, 2018) while in the European Union (EU) there is no established threshold as long as levels are not higher than what is required for the intended purpose (e.g., coloring) (Regulation (EC) No 1333,

2008).

Food grade TiO₂ can be in either the rutile or anatase form or a mixture of both and is primarily larger than 100 nm in particle diameter (Diebold, 2003; Yang et al., 2014; Theissmann et al., 2014). Transmission electron microscopy (TEM) of several lots of E 171, the EU designation for food-grade TiO₂, revealed an average frequency of 17–35% of particles by number falling below 100 nm in diameter (nano-fraction) (Yang et al., 2014). More relevantly, TEM analysis of various chewing gum coatings revealed that on average, 4.2% of TiO₂ particles were of the nano fraction (Dudefoi et al., 2017, 2018). TiO₂ has been extensively studied for potential hazard to human health, and it is generally accepted that inhalation of high doses of nano-sized TiO₂ presents a toxicological risk, especially in an occupational setting (Oberdorster et al., 1994; Warheit et al., 2006; Sager et al., 2008; Baan et al., 2006; Lee et al., 1985; Humans, 2010). In rodent models, orally

* Corresponding author. Michigan State University, East Lansing, MI, 48824, USA.

E-mail address: kamins11@msu.edu (N.E. Kaminski).

<https://doi.org/10.1016/j.fct.2019.110793>

Received 21 June 2019; Received in revised form 22 August 2019; Accepted 24 August 2019

Available online 29 August 2019

0278-6915/ © 2019 Elsevier Ltd. All rights reserved.

ingested TiO₂ was largely innocuous, even when administered in the diet at levels of 50,000 ppm (5.0% of the diet) to rats and mice for up to two years (National Toxicology Program, 1979). Adverse effects were only reported in study animals dosed with high amounts of entirely nano-sized particles (Wang et al., 2007, 2013; Tassinari et al., 2014; Mohamed, 2015; Shukla et al., 2013; Azim et al., 2015; Orazizadeh et al., 2014; Jia et al., 2014; Ruiz et al., 2017). However, while it is generally accepted that > 99% of orally ingested TiO₂ particles are passed in the feces (Kreyling et al., 2017a, 2017b, 2017c), it has been shown that TiO₂ particles can accumulate in intestinal crypts (Brun et al., 2014; Powell et al., 2010), bind *in vitro* and *in vivo* to phagocytic microfold cells (M-cells) (Powell et al., 1996, 2010), and pass through the gut into the *lamina propria* (Brun et al., 2014; Powell et al., 1996, 2010). Further, intestinal dendritic cells can directly internalize TiO₂ particles (Brun et al., 2014; Powell et al., 1996, 2015; Manolova et al., 2008). Many of these studies were performed with TiO₂ particles of a diameter less than 100 nm (Winkler et al., 2018). It has also been reported that TiO₂ particle-uptake can provoke inflammatory responses as TiO₂ nano-particles induced inflammasome activation *in vitro* (Ruiz et al., 2017; Winter et al., 2011). It has been proposed that these effects could potentially provoke or exacerbate gut inflammatory disorders such as inflammatory bowel disease (Ruiz et al., 2017; Hummel et al., 2014).

Given the potential for TiO₂ particles to accumulate in the gastrointestinal tract and interact with the gut immune system, it is critical to examine the effects of relevant oral TiO₂ exposure on gut immune homeostasis. While TiO₂ has been studied extensively, many of these studies were not relevant to the human route of exposure, oral dietary ingestion, instead relying on inhalation, direct injection, oral gavage, or administration in drinking water with purified TiO₂ particles of various size fractions (See (Winkler et al., 2018) for review). Further, it has been shown that TiO₂ particles routinely incorporate surrounding macromolecules onto the surface architecture of their particle structure, and depending on the types of molecules, the physical properties of TiO₂ particles, such as absorption, aggregation, and tissue uptake can be drastically altered (McCracken et al., 2013; Nel et al., 2009; Ruh et al., 2012; Walkey and Chan, 2012; DeLoid et al., 2017; Nagayama et al., 2007; McClements et al., 2016). Food-grade TiO₂, E 171, is consumed by humans already in food preparations, allowing for the incorporation of food macromolecules onto the surfaces of E 171 particles. Further, food passes slowly through the digestive system compared to liquids such as water, which is absorbed primarily in the small intestine (Boland, 2016). Given that TiO₂ particles are insoluble in water and tend to aggregate, delivery of E 171 particles in water may not recapitulate the biology of TiO₂ particles already incorporated into food products.

In addition to possible immunologic effects, recent publications have raised concerns about a possible carcinogenic effect on the intestines (Bettini et al., 2017; Urrutia-Ortega et al., 2016; Proquin et al., 2018). A two-year bioassay of dietary TiO₂ to F344 rats and B6C3F1 mice showed no evidence of proliferative or neoplastic effects on the gastrointestinal tract or neoplastic effects in any tissues (National Toxicology Program, 1979). Recently, studies have focused on shorter-term evaluations of aberrant crypt foci (ACF) and goblet cells as indicators of possible carcinogenic effects. However, these utilized administration of TiO₂ by oral gavage or in drinking water rather than diet, with potential agglomeration of particles occurring, altering their interaction with host tissues (Urrutia-Ortega et al., 2016; Additives and Food, 2016; Additives et al., 2018).

The objectives of the present study were to evaluate the acute (7 days) and long term (100 days) effects of dietary E 171 exposure on the immune system of the gastrointestinal (GI) tract and periphery as well as to evaluate chronic exposure either alone or after pre-administration of a known intestinal genotoxic carcinogen, dimethylhydrazine (DMH). Following the 7- and 100-day feeding periods, rats were euthanized and measurements of inflammatory cytokines and phenotyping of immune

cells in the periphery and GI tract were performed. Peyer's patches, peripheral blood mononuclear cells (PBMC), and spleen cells were analyzed for inflammatory and regulatory T-cell responses directly *ex vivo* or after *in vitro* stimulation (7 days). Histopathology, ACF, and goblet cell evaluations were performed on rats in the 100-day study. All tissues were collected from well-defined areas, and measurements, procedures, and evaluations were performed in a standardized and blinded manner. In addition, evaluations of possible proliferative or tumor enhancing effects were performed focusing on histopathology, assessment of ACF (also evaluated in a blinded manner), and quantification of goblet cells in the distal large intestine.

2. Materials and methods

2.1. Chemicals

Food grade sample E171-E was supplied by the Titanium Dioxide Manufacturers Association based on an assessment and characterization of the different grades of E 171 placed on the market. Chemical analysis, including particle size distribution, was performed and diets (control and E 171 containing) were analyzed for TiO₂ content and homogeneity. It was stored at room temperature. N, N'-dimethylhydrazine dihydrochloride (DMH · 2HCl) (98% purity) was purchased from Sigma Aldrich (St. Louis, MO) and stored at room temperature.

2.2. E171 chemical analysis

Chemical analysis of the particle size and particle size distribution of the respective lot of E 171 used in food preparations containing TiO₂ were performed by 2 independent contract laboratories (Kronos; VENATOR). Analyses were performed using 2 independent techniques, electron microscopy and direct volume/mass based methods. Consistent with published findings, the specific method used to quantify E 171 particle size and distribution yielded noticeably different results (Ortlieb, 2010). For example, analysis using scanning electron microscopy (SEM) showed 36% of TiO₂ particles were < 100 nm in diameter with the average diameter recorded being between 110 and 115 nm. Volume/mass based approaches showed similar average particle diameter (150 nm) but showed only 1–2% or particles falling below 100 nm in diameter. In spite of the differences observed between the two analytical methods in the percentage of particles possessing a diameter below 100 nm, the results obtained by the two independent laboratories were similar with each of the two methodologies, demonstrating good concordance between the two laboratories.

2.3. E 171 food preparation

All diets containing E 171 were prepared by Dyets, Inc. (Bethlehem, PA). E 171 was administered in irradiated and pelleted Certified Purina 5002R33 (LabDiet, Richmond, IN) at a concentration of 40, 400 or 5000 ppm and containing 2% maltose dextrin throughout the 7- and 100- day studies. Two batches of diet were prepared. The first batch of diet was fed throughout the 7-day study and through week 10 of the 100-day study. The second batch of diet was fed during the remainder of the 100-day study. Food and water were provided *ad libitum* to all animals during the quarantine and acclimation period and the 7- and 100-day exposure periods. The homogeneity and concentration of E 171 in the test diets were analyzed by Eurofins Food Chemistry Testing US, Inc. (Madison, WI). Analysis of triplicate samples from top, middle and bottom of the diet preparations confirmed homogeneity (data not presented). The concentrations of E 171 in the first and second batches of diets based on the mean of six determinations per diet ± S.D. were found to be, respectively: 0 ppm dose (22.3 ± 1.2, 23.7 ± 1.8 ppm); 40 ppm dose (59.6 ± 1.1, 61.0 ± 2.6 ppm); 400 ppm dose (384 ± 8, 387 ± 13 ppm) and the 5000 ppm dose (4310 ± 132, 4610 ± 160 ppm).

2.4. Animals

Six-week-old male Wistar Han IGS (CrI:WI (Han)) rats were purchased from Charles River Laboratories (Raleigh, NC) for all studies. Animals were housed in facilities fully accredited by the American Association for Accreditation of Laboratory Animal Care (AAALAC). The level of care provided to the animals is based on federal regulations including the U.S. Government Principles for the Utilization and Care of Vertebrate Animals Used in Testing, Research and Training, USDA implementing regulations, (9CFR), of the Animal Welfare Act and U.S. Public Health Service Policy Assurance for the Humane Care and Use of Laboratory Animals negotiated with the Office of Laboratory Animal Welfare, [OLAW](#). The 7-day study protocol was approved by the Michigan State University Institutional Animal Care & Use Committee (IACUC) and performed at Michigan State University. The 100-day study protocol was approved by the University of Nebraska Medical Center (UNMC) IACUC and performed at UNMC. Rats were housed individually during replicate 7-day studies and 2/cage during the 100-day study in plastic cages with dry corn-cob bedding containing environmental enrichment.

2.5. Study design

2.5.1. 7-Day study

Animals were acclimated for 12 days prior to the start of two replicate feeding studies. At the initiation of each of the replicate studies, rats were randomized into 4 groups of 5 animals each using an in-house weight-based randomization program ([Martin et al., 1984](#)). Total food and water consumption were determined at the end of each study. Body weights were determined at the start and end of the 7-day exposure period at the time of euthanasia.

2.5.2. 100-Day study

Note that all doses are for dimethylhydrazine dihydrochloride not dimethylhydrazine. After acclimation for 7 days, the animals were randomized as described above, into 8 groups of 15 animals each. Animals in groups 1–4 were treated with a single intraperitoneal (i.p.) injection of a sterile dose of 180 mg/kg BW dimethylhydrazine dihydrochloride (DMH · 2HCl) in 1.5% EDTA-0.9% NaCl, pH 6.5. Animals in groups 5–8 were treated with a single sterile dose of 1.5% EDTA-0.9% NaCl, pH 6.5. Seven days after i.p. injection, dietary administration of 0, 40, 400, or 5000 ppm E 171, respectively, was initiated in groups 1–4 and 5–8 and continued for 100 days. The dose of DMH · 2HCl was chosen as the highest dose of DMH · 2HCl that could be administered at pH 6.5 with no signs of toxicity based on 3 pilot studies. During the first 2 pilot studies, rats exhibited signs of extreme toxicity including death, when administered i.p. doses of 90, 130, 180 or 398 mg/kg BW DMH · 2HCl (no pH adjustment) which correspond to DMH doses of 41, 59, 81, and 180 mg/kg BW, respectively. In a third pilot study, there was a statistically significant decrease in body weights for rats administered i.p. doses of 287 and 398 mg/kg BW DMH · 2HCl, pH 6.5. A dose of 287 mg/kg BW DMH · 2HCl, pH 6.5, corresponds to a DMH dose of 130 mg/kg BW. There were no signs of toxicity at i.p. doses of 90, 130 and 180 mg/kg BW DMH · 2HCl, pH 6.5. Animals were checked daily for moribundity and mortality, and detailed clinical observations were performed during study weeks 8 and 13. Body weights were determined weekly beginning on day 0 of the study and just prior to euthanasia. Food consumption was determined weekly beginning with administration of the E 171 supplemented diets. Water consumption was determined during weeks 3, 8 and 13 of the study.

2.6. Blood and tissue collection, processing and analysis

At the end of the 7- and 100-day studies, rats were euthanized by administration of a lethal dose (150 mg/kg) of Fatal-Plus® (Vortech Pharmaceuticals, Dearborn, MI) by intraperitoneal injection. All

samples were blinded prior to analysis except for samples from four control animals in each study that were used to calibrate the flow cytometry analyses, and samples from three control animals in the 100-day study that were used to establish the normal histopathology for this strain of rat and samples used for goblet cell analysis.

2.6.1. 7-Day study

Whole blood was collected by cardiac puncture from euthanized rats using a heparin coated needle and syringe. Red blood cells (RBC) in whole blood were lysed using ACK lysis buffer, washed twice with HBSS and resuspended in RPMI containing 10% FBS (Hyclone, Logan, UT) and 100 U/ml of penicillin and streptomycin (Gibco Life Technologies, Grand Island, NY). Leukocytes were either phenotyped by flow cytometry or PBMCs were isolated by layering the cell suspension onto Lympholyte-Rat (CederLane, Burlington, NC) and centrifuged at $1500 \times g$ for 20 min, washed twice with HBSS and resuspended in RPMI containing 10% FBS (Hyclone, Logan, UT) and 100 U/ml of penicillin and streptomycin (Gibco Life Technologies, Grand Island, NY) and activated using anti-CD3/anti-CD28 antibodies to induce cytokine production *ex vivo* (see below). Plasma was collected from the remaining whole blood by centrifugation at $500 \times g$ and stored at -80°C for cytokine quantification. A 1–2 cm segment of jejunum, approximately 15 cm from the cecum and a 1–2 cm segment of colon, approximately 10 cm from the anus were harvested from each rat for evaluation of immunologic parameters. The tissues were mechanically disrupted in 1 ml of RIPA buffer (0.5% deoxycholate, 0.1% SDS, and 1% Igepal in TBS) containing complete protease inhibitor cocktail (Roche Diagnostics, Mannheim, Germany) for determination of inflammatory cytokines within the gastrointestinal track. Cytokine levels were normalized based on total protein concentrations as determined by a BCA kit (Sigma, St. Louis, MO) following manufacturers protocol. Peyer's patches were removed along the jejunum and ileum and leukocytes were isolated for phenotyping and activated using anti-CD3/CD28 antibodies to induce cytokine production *ex vivo* (see below). Briefly, isolated jejunum and ileum Peyer's patches were washed 4 times with HBSS. After the last wash, 2 ml of Spleen Dissociation Medium (StemCell Technologies, Cambridge, MA) were added to the harvested Peyer's patches and incubated at RT for 30 min. After incubation, EDTA was added to a final concentration of 10 mM. Cell suspensions were filtered through a 70 μm nylon mesh strainer and then layered onto Lympholyte-Rat Cell Separation Media (CederLane, Burlington, NC) and centrifuged at $1500 \times g$ for 20 min. Rat spleens were excised, weighed and cut transversely. One half of the spleen was frozen at -80°C , and the other half was made into a single cell suspension, washed with HBSS and resuspended in RPMI containing 10% FBS and 100 U/ml of penicillin and streptomycin. Splenocytes were either used for phenotyping by flow cytometry or for cytokine quantification after *ex vivo* activation (see below). The whole liver was removed, weighed and frozen at -80°C .

2.6.2. 100-Day study

All blood and tissue samples for determination of immunological parameters by flow cytometry were collected, processed and analyzed as described in Section 2.6.1 above. For evaluation of aberrant crypt foci (ACF) analysis, three 2 cm sections were removed from the colon: 1) one from the distal end, 2) one approximately 7 cm from the distal end (mid-portion), and 3) one at the proximal end. Each segment of colon for ACF analysis was placed separately between 2 strips of dental wax and fixed in 10% buffered formalin prior to being processed for ACF evaluation. For evaluation of histopathology and goblet cells, a 1 cm transverse segment approximately 2 cm from the distal end of the colon was removed and fixed in 10% buffered formalin. The spleen, liver and 1 testis were removed and weighed. Sections from the liver, lung, spleen, duodenum, jejunum, and ileum were placed in 10% buffered formalin. The testis was fixed in modified Davidson's solution. All fixed tissues were paraffin embedded in the Tissue Sciences Facility,

UNMC. Approximately 4–5 μm sections were stained with hematoxylin and eosin and examined histopathologically (Cohen et al., 2007). The second testis was removed and along with sections from the lungs, colon, and liver were frozen at -20°C .

2.7. Anti-rat CD3/CD28 activation

Leukocytes from whole blood, Peyer's patches and spleen were counted on a Coulter Counter Z1 (Beckman Coulter, FL) and adjusted to a cell concentration of 1×10^6 cells/ml. Leukocytes from whole blood and spleen were placed into 24-well tissue culture plates while leukocytes from the Peyer's patches were placed in 96 well flat bottom culture plates (Corning, Corning, NY) that were precoated with anti-rat CD3 (5 $\mu\text{g}/\text{ml}$) (Biolegend, CA). Immediately after addition of cells to the 96-well culture plates, anti-rat CD28 (5 $\mu\text{g}/\text{ml}$) (Biolegend, CA) was added to each well. After a 4-day incubation at 37°C with 5% CO_2 , supernatants were collected and stored at -80°C for quantification of cytokines.

2.8. Flow cytometry

Phenotyping of leukocytes from whole blood, Peyer's patches and spleen was performed by flow cytometry using antibodies directed against rat CD45 (OX-1, Biolegend, San Diego, CA) to identify rat leukocytes. Antibodies directed against rat CD103 (OX-62, Biolegend), MHC-II (RT1B)(OX-6, Biolegend) and CD11b/c (OX-42, Biolegend) were used to quantify dendritic cells (DC); CD4 (W3/25, Biolegend) to quantify T_{helper} cells, CD25 (OX-39, Biolegend) to quantify activated T_{helper} cells. Similarly antibodies directed against rat FoxP3 (FJK-16s, eBioscience) were used to identify total $T_{\text{regulatory}}$ (T_{reg}) (CD4^+ , FoxP3 $^+$) cells and activated T_{reg} (CD4^+ , CD25 $^+$ and FoxP3 $^+$). Flow cytometry data were collected on a Canto II (BD Biosciences, San Jose, CA) and the collected data were analyzed using Flowjo v10.2 (Tree Star Inc., Ashland, OR).

2.9. Quantification of cytokines

Cytokines in plasma, tissue extracts and cell culture supernatants were quantified using the LEGENDplex rat inflammation panel (Biolegend, San Diego, CA) for IL-10, IFN- γ , CXCL1 (KC), MCP-1, TNF- α , GM-CSF, IL-18, IL-12 (p70), IL-17A, IL-33, IL-1 α , IL-1 β and IL-6 per the manufacturer's protocol.

2.10. ACF analysis

Colon sections were removed from 10% buffered formalin one at a time and rinsed briefly in 0.9% saline. Each section was stained for 30 s in 0.5% toluidine blue in 0.5% sodium borate. Excess dye was removed by rinsing in clean 0.9% saline. The section was placed on a glass slide, lumen side up, that had previously been divided on the back-side into 25 sections with dimensions of approximately 4 mm^2 each so that ACF would not be counted more than once. The stained colon tissue section was covered with a few drops of aqueous dry mounting medium and cover slipped. The number of ACF in each 4 mm^2 and the number of aberrant crypts (ABC)/foci was determined by light microscopic examination at 40X. If necessary for confirmation of identification of ABC, the 4 mm^2 section was examined at 400X.

2.11. Goblet cell analysis

Beginning with the lowest numbered animal in each group, hemolysin & eosin stained slides of colon tissue were evaluated for acceptability to assess goblet cells, which included no evidence of dilatation of the lumen (secondary to presence of feces) and no evidence of autolysis. The first five colons from each group that met those criteria were evaluated. Two glands were randomly chosen for which the entire

length of the gland from the colonic luminal surface to the base of the gland could be identified, and the glandular lumen was present the entire length of the gland. As close as possible, glands were chosen that were perpendicular to the intestinal lumen and perpendicular to the underlying connective tissue to avoid tangential cutting artifact. For the glands that were evaluated, the length of the gland from the epithelial luminal surface to the base of the crypt was measured using an ocular micrometer. The number of goblet cells in that gland was then evaluated. Two glands were evaluated for each colon.

2.12. Statistical analysis

Prism 6 (GraphPad, San Diego, CA) software was used to determine statistical significance. Data was first assessed for homogeneity using a Bartlett's test. Homogenous data was then evaluated for significant differences from comparative controls using a one-way ANOVA (Dunnett's multiple comparisons test). For data sets deemed non-homogeneous per the Bartlett's test, values were log transformed to meet the variance of the standard deviation and then tested using a one-way analysis of variance ANOVA (Dunnett's multiple comparisons test). For data sets that still did not pass the variance test (i.e., Bartlett's test), an outlier test was performed (Rout test) followed by a reassessment of variance and then a one-way ANOVA. The only data set for which the outlier test was applied was the 100-day plasma IL-6 cytokine levels. For statistical analysis of the ACF, a value of 0.5 was added to all samples so a log transformation could be performed to replace values of "0" in all data sets (Yamamura, 1999). Statistical analysis was performed using SAS for Windows, version 9.4. Group means for body weights, food and water consumptions, goblet cell analysis, and organ tissue weights were evaluated using ANOVA, followed by Duncan's multiple range test for group-wise comparisons. Histopathology was compared using the 2-tailed, Fisher's Exact test. A p value of < 0.05 was deemed statistically significant.

3. Results

3.1. 7- and 100-day studies: in life parameters

In order to investigate the effects of E 171 delivered in a relevant manner, 6-week-old Wistar Hans rats were fed diets containing E 171 (0, 40, 400 and 5000 ppm) for 7 and 100 days. As shown in Table 1, rats in the study groups had comparable body weights by the end of the 7 and 100 day feeding studies. Rats fed acute and chronic high-content dietary E 171 did not demonstrate significantly different food or water consumption (Tables 2 and 3). It is noteworthy that despite comparable body weight in rats receiving 400 or 5000 ppm dietary E 171, there was a trend toward increased food consumption in rats receiving 5000 ppm E 171. At the 5000 ppm dose, rats consumed approximately 10 more grams of food during the 7 day study compared to rats receiving the control diet (Table 2A). However, the difference in food consumption by weight was not significantly different between any of the treatment groups and this trend was not present in the 100 day study at any time point, including after 7 days (Table 2B).

Based on the weight of food consumed, the amount of E 171 each group received was calculated and is shown in Table 4 for both the 7- and 100-day studies. For the 100-day study, two lots of E 171 containing food were used and E 171 content and exposure is broken down by respective lot used (Table 4B). Rats in the highest E 171 treatment groups consumed a total of approximately 361.3 mg of E 171 per kilogram of body weight per day during the 7- and 100-day studies (Table 4). Rats in the next highest E 171 group consumed approximately 10-fold less E 171 during the 7- and 100-day studies compared to the 5000 ppm E 171 group (Table 4). To this end, the 5000 ppm E 171 group received an average dietary exposure of E 171 per animal that is orders of magnitude higher than the average reported human exposure of 1.4 mg/kg (Winkler et al., 2018). Together these results

Table 1

Body weights of male Wistar rats treated with E 171 for 100 days with or without DMH · 2HCl pretreatment.

Group	Treatment	Body Weight (g)			
		Mean ± S.E. ^a			
		Week 1	Week 5	Week 10	Week 15
1	0 ppm E 171	244.0 ± 3.9	353.0 ± 7.5	416.4 ± 10.1	443.7 ± 11.1
2	40 ppm E 171	237.3 ± 2.6	341.2 ± 5.4	408.1 ± 8.3	439.9 ± 10.1
3	400 ppm E 171	241.4 ± 3.4	355.8 ± 6.9	426.4 ± 9.6	454.3 ± 10.7
4	5000 ppm E 171	240.8 ± 3.3	348.3 ± 6.2	415.4 ± 7.8	446.4 ± 9.1
5	180 mg/kg DMH · 2HCl	224.3 ± 2.9 ^b	330.7 ± 4.9 ^b	393.9 ± 6.7	423.7 ± 7.2
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	226.4 ± 3.2 ^c	337.8 ± 6.1	406.6 ± 8.1	436.8 ± 9.6
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	221.5 ± 3.5 ^d	330.9 ± 6.3 ^d	398.6 ± 8.2 ^d	428.5 ± 8.7 ^d
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	223.3 ± 3.3 ^c	335.6 ± 8.4	405.5 ± 10.9	435.4 ± 12.7

^a N = 15 for all groups and all time points.^b Significantly different compared to 0 ppm E 171 group, p < 0.05.^c Significantly different compared to 40 ppm E 171 group, p < 0.05.^d Significantly different compared to 400 ppm E 171 group, p < 0.05.^e Significantly different compared to 5000 ppm E 171 group, p < 0.05.

demonstrate that high acute and chronic E 171 intake did not alter the in life measures of study animals.

Previous studies have shown that high-dose TiO₂ administration in albino mice resulted in increased liver weight and cellularity due to the infiltration of nano-sized TiO₂ particles into the sinuses of the liver (Shukla et al., 2013). Another study suggested that TiO₂ can induce systemic inflammation (Bettini et al., 2017). As part of a general assessment of acute toxicity due to dietary E 171, liver and spleen weights were also determined. As shown in Table 5, E 171 did not alter the liver

to body weight ratio in any of the 7- or 100-day study groups. Similarly, there was no change in the spleen to body weight ratio in any experimental groups with the exception of the 400 ppm E 171 + DMH treatment group, which was modestly increased compared to 400 ppm E 171. There were no changes in spleen cellularity across any of the treatment groups (Table 5). Collectively, these results suggest that dietary E 171 produced no general signs of overt toxicity even at doses as high as 5000 ppm of E 171 over 100-days.

Table 2

Food consumption by male Wistar rats treated with E 171.

A. 7-day Study				Food Consumption					
Group	Treatment			Mean ± S.E. ^a					
						g/animal/day		g/kg BW/day	
1	0 ppm E 171					21.4 ± 0.6		75.8 ± 1.8	
2	40 ppm E 171					22.1 ± 0.4		78.9 ± 2.0	
3	400 ppm E 171					22.3 ± 0.5		81.6 ± 3.0	
4	5000 ppm E 171					23.0 ± 0.3		83.8 ± 1.8 ^b	

B. 100-day Study													
Group	Treatment			Food Consumption									
				Mean ± S.E. ^c									
						Week 2		Week 5		Week 10		Week 15	
						g/animal/day		g/kg BW/day		g/animal/day		g/kg BW/day	
1	0 ppm E 171	23.9 ± 1.4		86.5 ± 5.2		22.8 ± 0.6		64.7 ± 1.7		22.4 ± 0.6		53.8 ± 1.4	
2	40 ppm E 171	22.9 ± 0.4		85.3 ± 1.6		22.7 ± 0.2		66.7 ± 0.7		21.3 ± 0.4		52.2 ± 0.9	
3	400 ppm E 171	23.1 ± 0.5		83.1 ± 1.8		24.8 ± 0.6 ^d		69.8 ± 1.6 ^d		22.6 ± 0.6		52.9 ± 1.4	
4	5000 ppm E 171	23.3 ± 0.2		84.9 ± 0.8		24.7 ± 0.3 ^d		70.9 ± 0.8 ^d		22.7 ± 0.4		54.7 ± 1.1	
5	180 mg/kg DMH · 2HCl	22.3 ± 0.3		85.5 ± 1.2		22.0 ± 0.4		66.5 ± 1.2		21.5 ± 0.3		54.6 ± 0.9	
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	23.4 ± 0.5		89.1 ± 1.7		23.2 ± 0.5		68.8 ± 1.6		22.6 ± 0.5		55.5 ± 1.3	
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	22.1 ± 0.5		85.8 ± 1.9		22.3 ± 0.4 ^e		67.4 ± 1.3		21.5 ± 0.3		53.9 ± 0.9	
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	23.5 ± 0.8		89.6 ± 3.2		24.0 ± 0.8 ^f		71.6 ± 2.3		22.2 ± 0.8		54.7 ± 1.8	

^a N = 10 for all groups^b Significantly different compared to 0 ppm E 171 group p < 0.05^c N = 8 for all groups and all time points.^d Significantly different compared to 0 ppm E 171 group, p < 0.05.^e Significantly different compared to 400 ppm E 171 group, p < 0.05.^f Significantly different compared to 180 mg/kg DMH · 2HCl group, p < 0.05.

Table 3
Water consumption by male Wistar rats treated with E 171.

A. 7-day Study							
Group	Treatment	Water Consumption Mean ± S.E. ^a					
		g/animal/day	g/kg BW/day				
1	0 ppm E 171	68.1 ± 1.3	241.1 ± 4.7				
2	40 ppm E 171	66.3 ± 1.7	235.9 ± 5.6				
3	400 ppm E 171	68.9 ± 1.3	252.9 ± 8.3				
4	5000 ppm E 171	69.4 ± 1.4	253.3 ± 6.6				
B. 100-day Study							
Group	Treatment	Water Consumption Mean ± S.E. ^b					
		Week 3		Week 8		Week 13	
		g/animal/day	g/kg BW/day	g/animal/day	g/kg BW/day	g/animal/day	g/kg BW/day
1	0 ppm E 171	37.3 ± 1.6	122.1 ± 5.1	35.6 ± 1.0	90.0 ± 2.6	31.2 ± 1.3	71.5 ± 3.0
2	40 ppm E 171	37.9 ± 3.2	127.4 ± 10.8	34.8 ± 1.6	90.0 ± 4.2	35.2 ± 2.7	81.8 ± 6.3
3	400 ppm E 171	34.9 ± 1.0	113.2 ± 3.3	37.9 ± 2.0	94.1 ± 5.0	32.1 ± 1.1	71.8 ± 2.5
4	5000 ppm E 171	38.8 ± 1.4	128.2 ± 4.6	35.9 ± 2.1	91.9 ± 5.4	33.2 ± 1.8	75.8 ± 4.1
5	180 mg/kg DMH · 2HCl	37.1 ± 1.4	128.5 ± 4.7	35.5 ± 1.3	95.4 ± 3.4	31.8 ± 1.2	76.5 ± 2.9
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	35.6 ± 1.6	121.3 ± 5.5	35.1 ± 2.1	91.3 ± 5.3	32.7 ± 2.1	76.6 ± 4.9
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	37.0 ± 1.6	129.1 ± 5.5 ^c	35.6 ± 1.3	94.9 ± 3.4	32.0 ± 1.8	76.3 ± 4.4
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	36.5 ± 1.8	124.2 ± 6.2	37.7 ± 2.5	98.3 ± 6.4	32.6 ± 2.3	76.5 ± 5.3

^a N = 10 for all groups

^b N = 7 for groups 2 and 8, Week 3. For all other groups and all other time points, N = 8.

^c Significantly different compared to 400 ppm E 171 group, p < 0.05.

3.2. Effects of acute and chronic dietary E 171 exposure on CD103⁺ dendritic cell (DC) populations in the periphery and in the gut

While studies have demonstrated that > 99% of TiO₂ particles ingested in food pass through the digestive system within feces, it has been shown that gut DC, which are CD103 positive, can actively take up TiO₂ particles from the gut lumen (Manolova et al., 2008; Winter et al., 2011). TiO₂ particles have also been shown to be reactive, particularly

in inducing cellular reactive oxygen species (Sayes et al., 2006). Further, despite the low levels of TiO₂ retention in the host gut, the possibility of CD103⁺ DC recruitment to the gut and uptake of E 171 over time by DC in Peyer's patches exists. Therefore, the number of CD103⁺ DC in the periphery and in Peyer's patches was quantified.

After 7- and 100- days of dietary exposure to E 171, rats were euthanized and blood, spleens, and Peyer's patches were isolated and CD103⁺ DC evaluated. Peripherally, CD103⁺ DC made up less than 1%

Table 4
Consumption of E 171.

A. 7-day Study					
Group	Treatment	E 171 Consumption ^a			
		mg E 171/rat/day	mg E 171/kg BW/day		
1	0 ppm E 171	0.51	1.81		
2	40 ppm E 171	1.33	4.76		
3	400 ppm E 171	8.58	31.43		
4	5000 ppm E 171	102.5	373.86		
B. 100-day Study					
Group	Treatment	E 171 Consumption - First Batch of Diet ^a Weeks 2–10		E 171 Consumption - Second Batch of Diet ^a Weeks 11–15	
		mg E 171/rat/day	mg E 171/kg BW/day	mg E 171/rat/day	mg E 171/kg BW/day
		1	0 ppm E 171	0.5	1.5
2	40 ppm E 171	1.3	3.9	1.3	3.0
3	400 ppm E 171	9.1	25.5	8.5	19.0
4	5000 ppm E 171	103	294	103	236
5	180 mg/kg DMH · 2HCl	0.5	1.5	0.5	1.1
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	1.4	4.1	1.3	3.1
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	8.6	25.7	8.1	19.2
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	102	300	101	237

N = 10 for all groups

^a Consumption of E 171 was determined based on the concentration of E 171 measured in the diets and on body weights and food consumption.

Table 5
Terminal body and tissue weights of male Wistar rats treated with E 171.

A. 7-day Study									
Group	Treatment	Mean \pm S.E. ^a							
		Body Weight (g)	Spleen Weight (g)	Relative Spleen Weight (mg/g)	Splenocyte Cellularity (106 cells/mg)	Liver Weight (g)	Relative Liver Weight (mg/g)		
1	0 ppm E 171	282.8 \pm 5.5	0.77 \pm 0.04	2.73 \pm 0.15	0.23 \pm 0.04	12.1 \pm 0.2	43.0 \pm 1.0		
2	40 ppm E 171	281.5 \pm 6.3	0.73 \pm 0.04	2.60 \pm 0.14	0.27 \pm 0.03	12.0 \pm 0.4	42.7 \pm 1.3		
3	400 ppm E 171	273.7 \pm 5.9	0.74 \pm 0.04	2.72 \pm 0.17	0.28 \pm 0.06	11.9 \pm 0.3	43.4 \pm 0.8		
4	5000 ppm E 171	275.0 \pm 6.1	0.69 \pm 0.05	2.53 \pm 0.19	0.31 \pm 0.04	11.6 \pm 0.4	42.2 \pm 0.9		
B. 100-day Study									
Group	Treatment	Mean \pm S.E. ^b							
		Body Weight (g)	Spleen Weight (g)	Relative Spleen Weight (mg/g)	Splenocyte Cellularity (106 cells/mg)	Liver Weight (g)	Relative Liver Weight (mg/g)	Testes Weight (g)	Relative Testes Weight (mg/g)
1	0 ppm E 171	451.6 \pm 12.2	0.86 \pm 0.04	1.91 \pm 0.07	0.56 \pm 0.06	14.2 \pm 0.5	31.4 \pm 0.6	3.69 \pm 0.10	8.21 \pm 0.19
2	40 ppm E 171	445.7 \pm 11.1	0.83 \pm 0.04	1.85 \pm 0.07	0.53 \pm 0.06	13.9 \pm 0.4	31.2 \pm 0.4	3.60 \pm 0.09	8.10 \pm 0.17
3	400 ppm E 171	461.3 \pm 11.9	0.90 \pm 0.03	1.95 \pm 0.06	0.51 \pm 0.04	13.9 \pm 0.4	30.3 \pm 0.6	3.80 \pm 0.10	8.31 \pm 0.30
4	5000 ppm E 171	451.0 \pm 10.0	0.94 \pm 0.04	2.08 \pm 0.09	0.45 \pm 0.04	13.8 \pm 0.4	30.6 \pm 0.5	3.64 \pm 0.09	8.13 \pm 0.27
5	180 mg/kg DMH · 2HCl	430.1 \pm 7.7	0.86 \pm 0.03	1.99 \pm 0.06	0.50 \pm 0.06	13.2 \pm 0.5	30.6 \pm 0.7	3.46 \pm 0.08	8.07 \pm 0.20
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	440.1 \pm 9.9	0.87 \pm 0.04	1.97 \pm 0.07	0.48 \pm 0.05	13.7 \pm 0.4	31.1 \pm 0.7	3.74 \pm 0.12	8.51 \pm 0.22
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	425.0 \pm 11.3 ^c	0.90 \pm 0.03	2.13 \pm 0.07 ^c	0.49 \pm 0.03	13.1 \pm 0.4	30.9 \pm 0.9	3.76 \pm 0.10	8.95 \pm 0.36
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	443.2 \pm 13.7	0.85 \pm 0.03	1.92 \pm 0.06	0.39 \pm 0.05	13.4 \pm 0.6	30.2 \pm 0.7	3.69 \pm 0.12	8.39 \pm 0.34

^a N = 10 for all group

^b N = 15 for all data.

^c Significantly different compared to 400 ppm E 171 group, $p < 0.05$.




Fig. 1. Dietary exposure to E 171 did not significantly alter systemic or gastrointestinal tract resident CD103⁺ dendritic cell frequency. The frequency of gut resident and circulating CD103⁺ DC was determined in rats fed dietary E 171 for 7 or 100 consecutive days. Rats in the 100 day study were pretreated with 180 mg/kg of DMH or Vh control by i.p. injection. Following the indicated feeding period, animals were euthanized and PBMC, spleen, and Peyer's patches were collected and processed into single cell suspension. Cells were then stained for CD11b/c, CD103, and MHCII and analyzed by flow cytometry to quantify CD103⁺ DC populations. CD103⁺ DC from the 7 day study are shown in panel A and from the 100 day study are shown in panel B. Data are presented as a percentage of the live cells within the lymphocyte gate. Data were analyzed for statistical significance using a one-way ANOVA with a Dunnett's multiple comparisons test when there were no significant differences in the variance of the standard deviation.

of the total leukocyte population in peripheral blood as well as in spleen (Fig. 1A) and these percentages were not affected by the E 171-containing diet. These results were unsurprising as CD103-expressing DC are typically found in tissue parenchyma such as the lung and the GI tract (Scott et al., 2011). As such, CD103⁺ DC made up a substantially larger percentage of the total leukocyte population in the Peyer's patches from the small intestine, accounting for approximately 1% of the total leukocyte pool (Fig. 1A). Likewise, consumption of dietary E 171 did not alter the percent of CD103⁺ DC Peyer's patches. When we quantified CD103⁺ DC in rats fed E 171 for 100-days, we again did not observe an effect with increasing E 171 consumption in any of the tissues analyzed (Fig. 1B), although the frequency of CD103⁺ DC increased modestly in the peripheral blood and spleen compared to the 7-day study (Fig. 1). Surprisingly, pretreatment with DMH, an inducer of GI tract inflammation, had no effect on CD103⁺ DC content, with or without E 171 consumption (Fig. 1B). Together these results suggest that there was no change in the percentage of CD103⁺ DC in peripheral blood, spleen or Peyer's patches due to acute or chronic dietary E 171 consumption.

3.3. Effect of dietary E 171 on CD4⁺ T_{helper} populations in the periphery and locally in Peyer's patches

Based on a previous report suggesting CD4⁺ T cells were affected in response to E 171 administration in drinking water, CD4⁺ T cells were quantified after dietary administration of E 171 for 7- or 100-days. Specifically, the total percentage of CD4⁺ T_{helper} cells and the percentage of T_{helper} cells expressing CD25, an indicator of T_{helper} activation, were quantified in peripheral blood, spleen, and Peyer's patches (Bettini et al., 2017).

Activated CD4⁺ T_{helper} cells traffic in the peripheral blood from lymphoid organs such as the spleen or draining lymph nodes to areas of inflammation or toward chemokine gradients where they exert their effector functions, typically in the form of inflammatory cytokine release (Hirahara and Nakayama, 2016). We first examined the total CD4⁺ T cell population in PBMC, spleen, and Peyer's patches. As shown in Fig. 2, a robust population of CD4⁺ T cells was detected in all tissues examined from all treatment groups. When the relative proportion of

CD4⁺ T cells was compared to animals exposed to E 171 for 7 or 100 days, we did not observe any E 171 mediated alterations in this population of cells (Fig. 2A and B). Since the potential exists for E 171 to alter activation state, and subsequently effector cytokine release, we next determined if E 171 modulated CD4⁺ T_{helper} activation. As such, PBMC were examined to determine if dietary E 171 altered the activation status of these cells. As shown in Fig. 2C and D, a sizable percentage of activated CD4⁺ T_{helper} cells were not detected in the peripheral blood of rats fed E 171 acutely or chronically. Likewise, a small proportion of activated CD4⁺ T_{helper} cells were detected in the spleen, a major site for T cell recruitment and activation, at 7- or 100-days (Fig. 2C and D). These results were not surprising as in the absence of an immunogenic stimuli, one would not expect a high proportion of activated T_{helper} cells (Schmitt et al., 2014). Furthermore, rats fed E 171 for 100 days exhibited no difference in the percentage of activated CD4⁺ T_{helper} cells in peripheral blood or the spleen, compared to controls, suggesting that E 171 does not induce activation of this population of cells in the periphery (Fig. 2C and D). However, there remains the potential that activated CD4⁺ T_{helper} cells may have been recruited to the gut in response to local immune activation. Counter to this possibility, no significant difference in the percentage of activated CD4⁺ T_{helper} cells was observed in the Peyer's patches of rats fed E 171 containing food compared to rats fed a control diet for either 7- or 100-days (Fig. 2). DMH pretreatment also did not alter the percentage of activated T_{helper} cells in any of the tissues analyzed in the 100-day feeding study, despite the predicted increase in gut inflammation (Yoshimi et al., 1994) (Fig. 2D). Moreover, we made the unexpected observation that DMH and E 171 appeared to modestly reduce T_{helper} cell activation in Peyer's patches contrary to previously reported results (Bettini et al., 2017). Collectively, these data suggest that despite the potential for accumulation of E 171 particles locally in the gut or in gut-resident DC, there was no detectable impact on the percentage of activated CD4⁺ T_{helper} cells either peripherally or locally in the Peyer's patches of rats fed E 171 containing diets for 7- or 100-days.




Fig. 2. Dietary E 171 exposure did not change the frequency of CD4⁺ T_{helper} cells systemically or in intestinal Peyer's patches. The frequency of gut resident and circulating total and activated T_{helper} cells were determined in rats fed dietary E 171 for 7 or 100 days. Rats in the 100 day study were pretreated with 180 mg/kg of DMH or Vh control by i.p. injection. After the feeding period, animals were euthanized and PBMC, spleen, and Peyer's patches were collected and processed into single cell suspensions. Cells were then stained for CD4 and CD25 and analyzed by flow cytometry. Activated T_{helper} cells were identified as being CD4/CD25 double positive. Total T_{helper} from 7 (A) and 100 (B) day animals and activated T_{helper} from 7 (C) and 100 (D) day studies are shown. Data are presented as a percentage of the live cells in the lymphocyte gate staining positive. Data were analyzed for statistical significance using a one-way ANOVA with a Dunnett's multiple comparisons test when there were no significant differences in the variance of the standard deviation.




Fig. 3. Acute or chronic exposure to dietary E 171 did not significantly affect the frequency of circulating or GI-resident $T_{regulatory}$ (T_{reg}) cells. The frequency of gut resident and circulating total and activated T_{reg} cells were determined in rats fed dietary E 171 for a total of 7 or 100 days. Rats in the 100 day study were pretreated with 180 mg/kg of DMH or Vh control by i.p. injection. Following the 7- and 100-day feeding period, animals were euthanized and PBMC, spleen, and Peyer's patches were collected and processed into single cell suspensions. Cells were then surface stained for CD4 and CD25 and subsequently stained intracellularly for FoxP3. Following antibody stains, cells were analyzed via flow cytometry to identify T_{reg} cell populations. Data are presented as a percentage of the live cells falling in the lymphocyte gate. Total T_{reg} from 7 (A) and 100 (B) as well as activated T_{reg} from 7 (C) and 100 (D) day studies are shown. Data were analyzed for statistical significance using a one-way ANOVA with a Dunnett's multiple comparisons test when there were no significant differences in the variance of the standard deviation.

3.4. Effect of dietary E 171 on the percentage of $T_{regulatory}$ (T_{reg}) cells in the periphery and locally in Peyer's patches

We did not observe an impact of dietary E 171 at any of the dose levels on CD103⁺ DC populations or on the total percentage of CD4⁺ T_{helper} or activated T_{helper} cells. However, there remains the potential that E 171 may alter the percentage of T_{reg} cells, a critical mediator of local and systemic inflammation, which could lead to a low level inflammatory response in the absence of increased inflammatory cells (Luu et al., 2017). To test this possibility, the percentage of leukocytes staining positive for CD4, CD25 and FoxP3, the critical transcription factor and marker of T_{reg} cells (Bluestone and Tang, 2005), was quantified in PBMC, spleen, and Peyer's patches from rats fed a control or E 171-containing diet for 7 or 100 days. We first determined if there were gross differences in the total T_{reg} population (CD4⁺ FoxP3⁺) in whole blood, spleen, and Peyer's patches. As shown in Fig. 3A and B, we did not observe any significant alterations of the total T_{reg} population due to E 171 administration in any of the tissues examined. Pretreatment with DMH also did not impact the gross T_{reg} population in the periphery (Fig. 3B).

However, as with CD4⁺ T_{helper} cells, T_{reg} cells express CD25 when they are activated and exert suppressive effector function. As such, one would not expect to find T_{reg} cells in excess either in the periphery or locally within the tissue in the absence of an immune reaction (Luu et al., 2017; Bluestone and Tang, 2005). Indeed, studies have shown that the percentage of T cells comprised by T_{reg} cells steadily increases as inflammation increases (Joosse et al., 2019). Corroborating this notion, less than 1% of PBMC and 2% of splenocytes (Fig. 3C) were identified as being of the T_{reg} population suggesting no evidence of systemic inflammation. Likewise, rats fed the E 171-containing diet for 100 days did not have significant differences in the percentages of peripheral leukocytes staining positive for FoxP3 compared to controls (Fig. 3D).

As with activated CD4⁺ T_{helper} cells, the local distribution of T_{reg} cells can be as critical in mediating biological activity through local

exertion of immune suppressive functions (Bluestone and Tang, 2005). To this end, the percentages of T_{reg} and activated T_{reg} cells in the gut Peyer's patches were determined for both the acute and chronic feeding studies. As observed peripherally, no significant differences were detected in the percentage of total or activated T_{reg} cells in the Peyer's patches of rats fed a control diet compared to rats fed the E 171 containing diets for 7 days (Fig. 3A and C). Likewise, the percentage of total and activated T_{reg} cells in the Peyer's patches of rats receiving dietary E 171 chronically for 100 days was unaffected compared to rats receiving the control diet (Fig. 3B and D). DMH pretreatment also did not alter activation state or tissue accumulation of T_{reg} cells (Fig. 3B and D). Collectively, these results suggest that E 171 consumption does not alter T cell-mediated mechanisms of immune control. Further, these data are in agreement with the results obtained in Fig. 2, where no increase in the percentage of activated CD4⁺ T_{helper} cells in Peyer's patches of animals fed E 171 was observed. As stated previously, there is a direct correlation between the level of inflammation and the number of T_{reg} cells controlling normal immunological homeostasis. In total, the results presented in Figs. 2 and 3 suggest that dietary E 171 is not pro-inflammatory in the context of either promoting inflammatory CD4⁺ T_{helper} cell activation or in reducing the percentage of anti-inflammatory T_{reg} cells.

3.5. Dietary E 171 did not significantly alter inflammatory cytokine levels in plasma, small intestine, or colon

While the safety of E 171 as a food additive has been studied extensively for the past 50+ years, recent studies have suggested that E 171 and other micro-particles can trigger intestinal inflammation (Ruiz et al., 2017; Winter et al., 2011; Hummel et al., 2014), inflammasome activation (Ruiz et al., 2017; Winter et al., 2011), and the presence of TiO₂ particles weakly associates with the development of inflammatory bowel disease (IBD) (Hummel et al., 2014). More recently, a study by Bettini and colleagues suggested that TiO₂ intake resulted in increased inflammatory cytokine profiles strikingly similar to those found in IBD

patients (Bettini et al., 2017). As stated previously, the method of delivery of TiO₂ particles can change the properties of the particles themselves and lead to differential interactions between TiO₂ particles and host tissues (Nel et al., 2009; Ruh et al., 2012; Walkey and Chan, 2012). To this end, we attempted to recapitulate the findings of Bettini et al. using a model of TiO₂ delivery in food containing E 171.

Six-week-old Wistar Han rats were fed a diet consisting of either control chow or chow with increasing concentrations of E 171 for 7 or 100 days. Rats in the 100-day study were pretreated with either DMH or vehicle (Vh) before being fed E 171 containing diets. Following the feeding period rats were euthanized and colon, small intestine, and plasma were collected. Peyer's patches from rats acutely fed E 171 were also collected for *ex vivo* analysis of anti-CD3/CD28 induced T_{helper} cell inflammatory cytokine release. A rat inflammatory cytokine bead array was used to quantify tissue levels of IL-1 α , IL-1 β , IL-6, interferon γ (IFN γ), IL-12p70, IL-17A, IL-18, IL-33, CCL2/MCP-1, CXCL1/KC (IL-8), GM-CSF, and tissue necrosis factor α (TNF α), many of which have been implicated in gut inflammatory disorders such as IBD (Neurath, 2014; Francescone et al., 2015).

First, changes in circulating inflammatory cytokine levels were examined in response to acute exposure to dietary E 171. Two of the cytokines examined were below the limit of detection (TNF α and IL-17) while the others were not altered by dietary E 171 in any of the treatment groups, when compared to control diet (Supplemental Fig. 1), suggesting that acute exposure to dietary E 171 did not alter the levels of circulating inflammatory cytokines.

We also investigated the possibility of local inflammatory cytokine production in the GI tract, specifically in the small intestine and colon where inflammatory cytokine production has been shown to correspond with inflammatory GI disorders (Neurath, 2014; Francescone et al., 2015). Of the inflammatory cytokines that were detected in both tissues, no statistical differences were observed in the concentration of these cytokines between any of the E 171 treatment groups (Supplemental Figs. 2 and 3) suggesting 7 days of dietary E 171 did not induce GI inflammation. Likewise, the anti-inflammatory cytokine, IL-10, which increases in concentration in response to local inflammation (Bluestone and Tang, 2005), was not altered in plasma, intestine, or colon further suggesting that inflammation was not induced by consumption of dietary E 171 even in the 5000 ppm treatment group (Supplemental Figs. 1, 2, and 3). Together these data suggest that dietary E 171 does not induce inflammation peripherally or in the GI tract.

As no significant effect of acute dietary E 171 was observed compared to controls on the levels of inflammatory cytokines in the periphery or GI tract, additional studies were performed to explore the possibility that E 171 might alter the effector cytokine profile of T_{helper} cells in lymphoid tissue or circulation, which may not be manifest without T cell specific stimuli. To address this potential, lymphocytes were isolated from peripheral blood, spleen, and Peyer's patches of rats treated with E 171 and activated *ex vivo* with anti-CD3/anti-CD28 for 4 days to induce T_{helper} cell cytokine production. Following the 4-day culture period, supernatants were collected and assayed for inflammatory cytokine production. As shown in Supplemental Figs. 4, 5, and 6, we did not observe any acute effects of E 171 exposure on any of the induced cytokines produced from *ex vivo* stimulated T_{helper} cells, which is contrary to a previous study (Bettini et al., 2017).

While there were no overt effects of acute E 171 administration, we also examined the inflammatory cytokine profile in rats chronically exposed to E 171 as alterations in immune homeostasis may become evident later. As noted in the 7-day feeding study, we did not observe any significant E 171-mediated alteration of any of the 13 cytokines assayed in any of the tissues examined (Figs. 4–6). Furthermore, DMH pretreatment also did not significantly alter tissue specific cytokine release, with the sole exceptions being elevated IL-12p70 in plasma (Fig. 4) and IL-17A in the colon (Fig. 6). These results were surprising as we would have expected more inflammatory cytokine release in the GI

tract in response to DMH, a known inflammagen (Yoshimi et al., 1994; Hamiza et al., 2012).

These results suggest that E 171, fed acutely or chronically, does not alter inflammatory cytokine production either peripherally or locally in the GI tract. However, it is noteworthy that the increase in IL-12p70 occurred in the 40 ppm E 171 + DMH treatment group (Fig. 4), and there was no indication of an E 171 dose-dependent effect rendering this finding difficult to interpret. Conversely, while increased IL-17A was detected in the colons of DMH treated rats fed 400 ppm E 171 (Fig. 6), this effect was not observed in the 5000 ppm + DMH treatment group. The biological significance of increased IL-17A in the colon at only 400 ppm E 171 + DMH is unclear, especially in light of the fact IL17A levels in this treatment group were not different from background levels (no treatment control) in the colon.

3.6. Effect of 100-day dietary administration of E 171 on histopathology

As prior studies have suggested that chronic E 171 exposure resulted in increased neoplastic lesions (Bettini et al., 2017), we next used histopathological analysis to determine any gross abnormalities associated with E 171 exposure in various tissues. Histologic evaluation of the duodenum, jejunum, ileum, spleen, liver, lung, and testes was normal for nearly all animals. Occasional animals had mononuclear infiltrate in the livers, which is a normal finding in this strain of rats (Thoolen et al., 2010). There were no significant differences in the incidences between groups. In the spleen, a single animal in the 5000 ppm E 171 group showed possible decreased lymphoid tissue based only on a visual proportion of lymphoid to non-lymphoid tissue, but this could be well within the normal range for this strain of rats. The testes were all normal except for one atrophic testis with focal tubular calcification in one rat in the 5000 ppm E 171 group, a common lesion in rats even at this age (Thoolen et al., 2010; Wright et al., 1982; Foley, 2001; James and Heywood, 1979a, 1979b). There were no histologic changes in any of the segments of the small intestine.

While we did not observe any gross pathologies in the E 171 treated animals, rats that were pretreated with DMH displayed several histopathologic abnormalities, with or without E 171 exposure. There were two invasive adenocarcinomas in one animal in the 180 mg/kg DMH · 2HCl + 0 ppm E 171 group, and single adenomas in one animal in the 180 mg/kg DMH · 2HCl + 40 ppm E 171 group and one animal in the 180 mg/kg DMH · 2HCl + 400 ppm E 171 group (Data not shown). There were no other histologic changes in the large intestines of the other animals treated with E 171 with or without pretreatment with DMH · 2HCl. One rat in the 180 mg/kg DMH · 2HCl + 0 ppm E 171 group and one rat in the 400 ppm group had subpleural lymphocytes in the lung, but without any evidence of acute inflammatory changes or hyperplasia. Collectively, these data suggest that in the absence of a known GI carcinogen, that E 171 does not induce gross or histopathologic changes or abnormalities.

3.7. Effect of 100-day dietary E 171 on formation of aberrant crypt foci (ACF) and aberrant crypts (ABC)

While many of the observed histopathological abnormalities occurred in DMH pretreated rats, there exists the possibility that 100 days is not adequate to observe the full transition to cancer. As such, we quantified ACF and ABC as described precursor to GI tract carcinogenesis (Corpet and Tache, 2002; Wargovich et al., 2010). Unfortunately, much of the epithelial surface of the colon samples (proximal, middle and distal) were obscured when observed by light microscopy, therefore we were unable to examine the entire surface of the colon samples (Table 6). This was not observed in a previous study performed in mice (Zhou et al., 2016). In an effort to determine if the epithelial surface was obscured due to mucus, we dehydrated a portion of one section in an ascending series of alcohols (70–100%) (Strugala et al., 2003); however, this did not increase the area of visible




Fig. 4. Chronic dietary E 171 exposure did not alter systemic inflammation as evidenced by inflammatory cytokine accumulation. Rats were pretreated with 180 mg/kg DMH or Vh control by i.p. injection and fed dietary E 171 for a total of 100 days. Following the 100 day feeding period, animals were euthanized and plasma was collected and assayed for inflammatory cytokines and chemokines using the LEGENDplex rat inflammation panel. Data are shown as cytokine/chemokine concentration (pg/ml) for plasma. Data were analyzed for statistical significance using a one-way ANOVA with a Dunnett's multiple comparisons test when there were no significant differences in the variance of the standard deviation. * = $p < 0.05$ indicates significance compared to no E 171 control.

epithelium compared to an undehydrated portion of the same section. It is also possible this obscuring of the surface was the result of autolysis due to delayed fixation, which was a result of the processing of the colonic tissue during necropsy, although this was not apparent by histopathology. The results for the areas of epithelium that were observed show that there is the expected significant increase in ACF/cm² and ABC/cm² in groups 5–8 that were pretreated with 180 mg/kg BW DMH · 2HCl compared to the groups that were not pretreated with DMH (Table 6). E 171 treatment administered after DMH did not result in statistically significant increases in ACF or ABC (Table 6). No change in the number of AFC and ABC were observed due to E 171 exposure alone in the absence of DMH.

3.8. Evaluation of goblet cells

As expressed above, dietary E 171 consumption by rats pretreated with or without DMH did not increase the incidence of aberrant crypts or tumorigenesis. However, 100 days may not be an adequate amount of time for this transition. It has been reported that an early contributing event to intestinal inflammation, morphologic changes, and eventual tumor formation is the loss of mucin secretion by intestinal goblet cells (Van der Sluis et al., 2006). These cells produce mucins, particularly mucin 2, which forms a barrier that maintains intestinal immune homeostasis by segregating commensal bacteria from the intestinal

epithelia (Van der Sluis et al., 2006; Pelaseyed et al., 2014). It has been reported that a loss of goblet cells, mucin secretion, and a lengthening of the colonic gland precede intestinal inflammation and tumor formation (Van der Sluis et al., 2006; Velcich et al., 2002). Furthermore, another study also suggested that these cells could preferentially take up nanoparticles, such as TiO₂, from intestinal M cells, allowing for accumulation (Hansen et al., 2009). As such, we evaluated colonic gland length and enumerated gland goblet cell content as an early indicator of intestinal inflammation as a result of dietary E 171. First, we measured the colonic gland length in defined sections of colon from rats fed E 171 with or without DMH pretreatment. The length of the colonic glands examined was not significantly different in any treatment group, with the exception of a significant lengthening of colonic gland length in DMH pretreated rats receiving no E 171 (Table 7). Next we quantified the number of goblet cells per colonic gland (unit) to determine if dietary E 171 resulted in a loss of these cells. As shown in Table 7, the number of goblet cells/unit was similar in all treatment groups. Our results suggest that dietary E 171, with or without treatment with DMH, for 100 days had no effect on the length of the colonic glands examined or the number of goblet cells/unit.

4. Discussion

Food-grade TiO₂, otherwise referred to as E 171 in the European




Fig. 5. Chronic dietary E 171 exposure did not modulate small intestine inflammatory cytokine accumulation. Rats were pretreated with 180 mg/kg DMH or Vh control by i.p. injection and fed dietary E 171 for a total of 100 days. Following the 100 day feeding period, animals were euthanized and a 1-2 cm segment of jejunum was collected and assayed for inflammatory cytokines and chemokines using the LEGENDplex rat inflammation panel. Data are shown as cytokine/chemokine concentration (pg/mg) for small intestine. Cytokine protein levels were normalized to total protein levels for each sample of small intestine. Data were analyzed for statistical significance using a one-way ANOVA with a Dunnett's multiple comparisons test when there were no significant differences in the variance of the standard deviation.

Union, is a widely used food additive that is ingested daily by human consumption of numerous manufactured food products. Despite the extensive risk assessment performed on E 171 over the course of the past 5 decades, it has undergone renewed scrutiny as recent reports have suggested that E 171 can bind intestinal cells, pass through intestinal lumen, and induce inflammatory cytokine production in the GI tract (Wang et al., 2007; Ruiz et al., 2017; Kreyling et al., 2017c; Brun et al., 2014; Powell et al., 1996, 2010, 2015; Manolova et al., 2008). However, few *in vivo* studies have used E 171 as a dietary component at relevant levels with relevant characteristics, i.e. > 100 nm diameter TiO₂ particle size (Winkler et al., 2018). Here we show that acute and chronic dietary consumption of E 171 at increasing doses (40–5000 ppm) resulted in no significant effects on either peripheral or GI tract immune homeostasis as evidenced by immune cell phenotyping or inflammatory cytokine analysis. The sole exceptions to this were significantly elevated IL-17A levels in the colons of DMH + 400 ppm E 171 treated rats and IL-12p70 from plasma of DMH + 40 ppm E 171 treated rats (Figs. 4 and 6). Furthermore, there were no effects of E 171 on histopathologic evaluations of small and large intestine, liver, spleen, lungs, or testes, as well as no effects on ACF, colonic goblet cell numbers, or colonic gland length, which is in contrast to recent reports (Bettini et al., 2017; Urrutia-Ortega et al., 2016) but is consistent with the lack of preneoplastic or neoplastic changes in rats or mice in a two-

year bioassay at dietary doses as high as 50,000 ppm (5%) (National Toxicology Program, 1979).

Recently, it was reported that acute exposure to TiO₂ preparations, including E 171, by intragastric gavage for 7 days resulted in altered frequency of GI resident DC, T_{helper} cells, and T_{reg} cells (Bettini et al., 2017). Specifically, it was reported that the frequency of CD103⁺ DC increased in the Peyer's patches of rats fed E 171, while the frequencies of T_{helper} and T_{reg} cells decreased (Bettini et al., 2017). By contrast, we observed no changes in the frequency of local gut DC, T_{helper}, or T_{reg} cells in either the 7- or 100-day feeding studies. The authors did not propose a direct mechanism for why GI resident DC populations would expand with TiO₂ treatment, other than to note this could be compensatory to the local inflammation induced by TiO₂ exposure. However, they did not detect increased frequency of inflammatory T_{helper} cells or anti-inflammatory T_{reg} cells. This finding would be counter to the hypothesis of increased local inflammation. Indeed, our own findings suggest no difference in peripheral or local tissue levels of inflammatory cytokines or tissue frequency of T_{helper} cells, even at dietary E 171 levels greater than those evaluated by Bettini et al. (2017). While we did observe significantly elevated levels of IL-17A in the colons and increased circulating IL-12p70 in plasma of DMH + 400 ppm E 171 treated rats in the 100-day study, this is in stark contrast to published findings where IL-17A was significantly increased in as little as 7 days




Fig. 6. Dietary E 171 exposure did not alter colonic inflammation as evidenced by inflammatory cytokine accumulation. Rats were pretreated with 180 mg/kg DMH or Vh control by i.p. injection and fed dietary E 171 for a total of 100 days. Following the 100 day feeding period, animals were euthanized and a 1-2 cm segment of colon was collected and assayed for inflammatory cytokines and chemokines using the LEGENDplex rat inflammation panel. Data are shown as cytokine/chemokine concentration (pg/mg) for colon. Cytokine protein levels were normalized to total protein levels for each sample of colon. Data were analyzed for statistical significance using a one-way ANOVA with a Dunnett's multiple comparisons test when there were no significant differences in the variance of the standard deviation. * = $p < 0.05$ indicates significance compared to respective no E 171 control.

following E 171 exposure alone (Bettini et al., 2017). Further, the same study did not identify significant increases in IL-17A at the tissue level, but rather from peripheral splenocytes (Bettini et al., 2017) highlighting substantial differences. Despite the elevated levels of inflammatory IL-17A in the colon of DMH + 400 ppm E 171 treated rats, it is important to emphasize that this elevation was in comparison to DMH alone, which was lower than background levels of IL-17A (i.e., no treatment control). Moreover, the IL-17A levels in DMH + 400 ppm E 171 treated rats were not different from the no E 171, mock treatment control group. Furthermore, studies have linked increased IL-17A to tumorigenesis in colitis-associated cancer models (Hyun et al., 2012). Our finding that IL-17A levels were not significantly elevated compared to control animals strongly argues against the ability of E 171 to induce colitis or tumorigenesis. Likewise, there were no changes in the percentage of T_{reg} isolated from the Peyer's patches nor were there any histological features indicative of colonic inflammation. Interestingly, DMH did not significantly alter any of the immune parameters that were evaluated, whether in the intestines, spleen or peripheral blood (Yoshimi et al., 1994). However, there was a general trend in that the colon associated inflammatory cytokine levels in the DMH only control treatment group showed lower levels compared to the no treatment control group, including IL-1 β , IL-6, IL-17A, IL-18 and IL-33.

While altered basal inflammation could be responsible for the increases observed in GI dendritic cells, it would not explain the decreased number of gut-associated T_{helper} cells. The aforementioned study ascribed the effect of decreased T_{helper} cell frequency in the Peyer's patches to decreased T_{helper} cell differentiation (Bettini et al., 2017). However, the Peyer's patch is restricted to the types of T lymphocytes that can be recruited and maintained compared to the spleen, which is less selective (Williams and Butcher, 1997). Splenic selectivity could also account for the observed differences between T_{helper} cells from Peyer's patches and spleen stimulated *ex vivo* with a non-specific activator or all T cells with anti-CD3/anti-CD28 stimulation. Yet, this would not explain the differences observed in our study compared to recent reports as we did not observe any difference in T_{helper} frequency in GI-associated tissue or in the spleens of animals exposed to E 171 in either the 7- or 100-day studies.

Another explanation for the observed difference between our study and the one conducted by Bettini and colleagues (Bettini et al., 2017) could potentially be how E 171 was delivered. In the aforementioned study, E 171, which was suspended in water, was administered directly into the stomach of rats by intragastric gavage in a 7-day study. Further, in a 100-day study, they administered E 171 in the drinking water. These modes of exposure are strikingly different from the typical mode

Table 6

Evaluation of E 171 treatment for 100 days with or without pretreatment with DMH · 2HCl on the induction of colonic aberrant crypt foci (ACF) and aberrant crypts (ABC) in male Wistar rats.

Group	Treatment	Total cm2 Evaluated	Mean No. of ACF/cm2, ^a	Mean No. of ABC/cm2, ^a
1	0 ppm E 171	25.4	0.8 ± 0.5	1.9 ± 1.1
2	40 ppm E 171	19.7	0.1 ± 0.1	0.2 ± 0.2
3	400 ppm E 171	27.6	0.9 ± 0.4	2.1 ± 1.1
4	5000 ppm E 171	23.9	0.9 ± 0.5	2.7 ± 1.6
5	180 mg/kg DMH · 2HCl	24.2	5.4 ± 1.2 ^b	17.1 ± 4.1 ^b
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	27.3	5.3 ± 1.3 ^c	14.8 ± 3.9 ^c
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	30.4	7.2 ± 1.3 ^d	19.7 ± 3.6 ^d
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	26.9	10.1 ± 2.6 ^e	28.4 ± 6.6 ^e

^a Results expressed as mean ± S.E. N = 15 for groups 1 and 6–8 and 14 for groups 2–5.

^b Significantly different compared to 0 ppm E 171 group, $p < 0.05$.

^c Significantly different compared to 40 ppm E 171 group, $p < 0.05$.

^d Significantly different compared to 400 ppm E 171 group, $p < 0.05$.

^e Significantly different compared to 5000 ppm E 171 group, $p < 0.05$.

Table 7

Evaluation of colonic goblet cells in male Wistar rats treated with E 171 for 100 days with or without pretreatment with DMH · 2HCl.

Group	Treatment	Mean Length of Gland (units) ^{a,b}	Mean No. of Goblet Cells/Unit ^b
1	0 ppm E 171	4.6 ± 0.1	6.5 ± 0.5
2	40 ppm E 171	5.2 ± 0.2	5.9 ± 0.2
3	400 ppm E 171	4.9 ± 0.3	6.4 ± 0.3
4	5000 ppm E 171	5.3 ± 0.2	6.2 ± 0.3
5	180 mg/kg DMH · 2HCl	5.5 ± 0.3 ^c	6.2 ± 0.4
6	180 mg/kg DMH · 2HCl + 40 ppm E 171	5.0 ± 0.2	5.9 ± 0.4
7	180 mg/kg DMH · 2HCl + 400 ppm E 171	5.3 ± 0.3	6.0 ± 0.4
8	180 mg/kg DMH · 2HCl + 5000 ppm E 171	5.5 ± 0.2	6.2 ± 0.2

^a The length of the gland from the epithelial luminal surface to the base of the crypt was measured using an ocular micrometer.

^b Results expressed as mean ± S.E. N = 5 for all groups except 3 and 8. Only 4 animals in these groups met the criteria for inclusion in goblet cell evaluation.

^c Significantly different compared to 0 ppm E 171 group, $p < 0.05$.

of human exposure in which E 171 is delivered to the GI tract as a constituent of food preparations. As stated previously, the surface structure of TiO₂ molecules is rather malleable to alteration by surrounding macromolecules, resulting in widely different coronas depending on the preparation of TiO₂ (Lundqvist et al., 2008; Cedervall et al., 2007), which could have impacted the way TiO₂ particles interact with the host gut environment in our respective studies.

Equally important, it is well established that TiO₂ is water insoluble (Winkler et al., 2018). In the Bettini study (Bettini et al., 2017), the E 171 used for gavage was delivered in a water suspension (Cho et al., 2013). Delivery of TiO₂ in water can greatly change the nature of the particles themselves, as TiO₂ tends to aggregate and agglomerate in water suspension (Winkler et al., 2018), raising the possibility that delivery of E 171 in an aqueous solution could lead to clumps of TiO₂ passing through to the gut environment. As mentioned above, the aforementioned study administered, E 171 in drinking water, which requires sonication, a process with unknown effects on the form of

TiO₂. Agglomeration and particle settling could also occur over time. This would represent a key difference in how TiO₂ is seen by the host when consumed in food, where more particles would be part of the food matrix resulting in a more uniform profile of single particles as compared to clumps (McCracken et al., 2013). In fact, some studies utilizing differing routes of delivery for TiO₂ nanoparticles have suggested that the aggregation/agglomeration of TiO₂ particles can impact their interface with the host immune system (Fu et al., 2014; Kobayashi et al., 2009).

Despite the reported effects of E 171 on immune homeostasis, the most pertinent finding by Bettini and colleagues was the increase in ACF by E 171 in a model of DMH-induced carcinogenesis, suggesting that E 171 increased intestinal carcinogenesis (Bettini et al., 2017). However, no significant differences between groups administered E 171 at any of the three concentrations evaluated, whether fed alone or after pretreatment with DMH, were observed in ACF or ABC. Although there was a weak trend observed between increasing E 171 exposure and increased ACF/ABC in DMH-treated animals, this was not statistically significant. The failure to reach significance is critical as we utilized 15 rats per group to increase statistical power of the study. Further, evaluation of the effects of E 171 on proliferative lesions of the intestine involved histopathology, ACF numbers and size, large intestinal goblet cell number, and gland length in defined areas of intestine that were evaluated in a blinded manner (except for goblet cell number and gland length), a key factor in performing reproducible scientific experiments (Begley, 2013).

Urrutia-Ortega et al. (2016) reported that they detected decreases in goblet cells in the distal colon, and suggested that the decreased mucin could increase the risk of colon cancer. It was also reported by Talbot et al. (2018) that TiO₂ was trapped by intestinal mucus but no effect on mucin O-glycosylation or short-chain fatty acid synthesis, concluding that TiO₂ did not alter the intestinal barrier protection under healthy conditions. However we did not observe any differences between treatment groups in numbers of goblet cells per gland, the size of the glands, or number of goblet cells per unit length of the glands. The one exception to this was a significant lengthening of colonic glands in rats pretreated with DMH only (Table 7). As with other parameters, we used well-defined tissue samples, methods, and criteria to evaluate the goblet cells. We especially were careful to only evaluate glands from segments of colons that were not dilated because of the presence of feces; such dilation distorts the shape and size of the glands and can affect the number of goblet cells. Importantly, no lesions were observed in rats administered E 171 alone. Thus our data suggests that even under pathologic conditions in the intestine (DMH pretreatment) that dietary E 171 is innocuous in regards to reported effects on intestinal goblet cells and mucosal barrier maintenance.

In rats pretreated with DMH, one rat in the DMH + control group developed two separate invasive adenocarcinomas, one of which could be identified as arising from an adenoma. In each of the DMH + low-dose E 171 and DMH + mid-dose E 171 groups, there was an adenoma. In the DMH + high-dose E 171 group there were no tumors or any abnormalities of the intestine detected histopathologically. No abnormalities in the small or large intestines were observed in rats treated with E 171 without DMH pretreatment.

While DMH is a known, potent, genotoxic intestinal carcinogen in rodents, variability in numbers and types of lesions in the intestine are usually seen in bioassays of DMH and other carcinogens, particularly when a relatively low dose is administered as in our experiments (Perse, 2005; Rosenberg et al., 2009). There is also considerable variability in the number and size of ACF that are induced by DMH, as seen in the present experiment with the inter-individual variability seen within each group. In the rats fed E 171 without pretreatment with DMH, there were few ACF in any group, a typical finding for control rats. We did not observe the increase in ACF reported by Bettini et al. (2017). In their study, there were no groups administered only E 171; they only evaluated groups pre-treated with DMH. In their publication, they did

not mention whether they neutralized the highly acidic DMH·2HCl, but they almost certainly did as the acid without neutralization is highly irritating and highly toxic to the rats when injected ip. Also, they stated that they injected 180 mg/kg of DMH. However, they used DMH·2HCl as did we, and at that level of DMH (which would be 398 mg/kg DMH·2HCl, there is severe toxicity. Thus, it is likely that they used 180 mg DMH·2HCl, which is 81 mg/kg DMH, the dose that we used.

Based on our results it is unlikely that E 171 affects intestinal carcinogenesis in a DMH model of colonic cancer. In agreement with this finding, E 171 showed no evidence of direct carcinogenesis in a National Toxicology Program bioassay in either rats or mice when fed at levels of 2.5 or 5.0% of the diet (National Toxicology Program, 1979), doses considerably higher than those used by Bettini et al. (2017) Urrutia-Ortega et al. (2016), or in the study presented here. Also, it is well known that non-genotoxic substances that increase intestinal tumorigenesis show increased proliferative lesions in short term studies when administered alone, without pretreatment with a genotoxic carcinogen like DMH (Rosenberg et al., 2009; Cooper et al., 2000; Kozoni et al., 2000; Redston, 2001). Furthermore, non-genotoxic substances that enhance intestinal carcinogenesis will increase intestinal tumor incidences when administered alone in a two-year bioassay (National Toxicology Program, 1979). Thus, it is highly unlikely that TiO₂ is an intestinal carcinogen or enhances intestinal carcinogenesis.

Lastly, histopathology of liver, spleen, lungs, and testes did not show evidence of E 171 treatment-related effects. Subpleural lymphocytes in the lung occur occasionally in this strain of rats, and mononuclear foci in the liver commonly occur (Thoolen et al., 2010). The finding of atrophy in 1 testis also is not unusual for this strain of rats, even by 21 weeks of age as in our experiment.

In conclusion, we report the finding that acute and chronic dietary exposure to E 171 in food preparation did not result in significant alteration of gut immune homeostasis as evidenced by altered frequency of intestinal DC, T_{helper}, or T_{reg} cells. Despite changes at a single dose in colonic IL-17A and plasma levels of IL-12p70 in DMH + E 171 treated rats, no other changes in intestinal tissue or blood inflammatory cytokine production was observed. Furthermore, no E 171 effects on intestinal histopathology, ACF, or goblet cells were observed alone or after pretreatment with the genotoxic intestinal carcinogen DMH. Assessment of E 171 delivered via food represents a critical model of human TiO₂ exposure, ingestion through food consumption, and had notably different effects as compared to recent studies utilizing different models of TiO₂ delivery to experimental animals with other modes of administration. Given the varied nature of reports of adverse health effects with TiO₂ in specific contexts, this would suggest that the matrix in which TiO₂ is received when administered orally might be one important determinant, as E 171 administered in food does not appear to affect peripheral or GI tract immune homeostasis or intestinal carcinogenesis.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We gratefully acknowledge the technical expertise of the Tissue Sciences Facility at UNMC in the preparation of H&E (Haemotoxylin and Eosin) slides for histopathology and goblet cell examination, and the assistance of Stephanie Henkel in the preparation of this manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://>

doi.org/10.1016/j.fct.2019.110793.

Financial support

This work was supported by the Center for Research on Ingredient Safety, Michigan State University, the Grocery Manufacturers Association, the Titanium Dioxide Manufacturers Association, the International Association of Color Manufacturers, and the Fred and Pamela Buffett Cancer Center Tissue Sciences Facility Shared Resource, supported by the National Cancer Institute under award number P30 CA036727.

References

- Additives, E. Panel o.F., Food, N.S.a.t., 2016. Re-evaluation of titanium dioxide (E 171) as a food additive. *EFSA J.* 14 (9), e04545.
- Additives, E. Panel o.F., et al., 2018. Evaluation of four new studies on the potential toxicity of titanium dioxide used as a food additive (E 171). *EFSA J.* 16 (7), e05366.
- Azim, S.A., et al., 2015. Amelioration of titanium dioxide nanoparticles-induced liver injury in mice: possible role of some antioxidants. *Exp. Toxicol. Pathol.* 67 (4), 305–314.
- Baan, R., et al., 2006. Carcinogenicity of carbon black, titanium dioxide, and talc. *Lancet Oncol.* 7 (4), 295–296.
- Begley, C.G., 2013. Six red flags for suspect work. *Nature* 497 (7450), 433–434.
- Bettini, S., et al., 2017. Food-grade TiO₂ impairs intestinal and systemic immune homeostasis, initiates preneoplastic lesions and promotes aberrant crypt development in the rat colon. *Sci. Rep.* 7, 40373.
- Bluestone, J.A., Tang, Q., 2005. How do CD4 + CD25 + regulatory T cells control autoimmunity? *Curr. Opin. Immunol.* 17 (6), 638–642.
- Boland, M., 2016. Human digestion—a processing perspective. *J. Sci. Food Agric.* 96 (7), 2275–2283.
- Brun, E., et al., 2014. Titanium dioxide nanoparticle impact and translocation through ex vivo, in vivo and in vitro gut epithelia. *Part. Fibre Toxicol.* 11, 13.
- Cedervall, T., et al., 2007. Understanding the nanoparticle-protein corona using methods to quantify exchange rates and affinities of proteins for nanoparticles. *Proc. Natl. Acad. Sci. U. S. A.* 104 (7), 2050–2055.
- Cho, W.S., et al., 2013. Comparative absorption, distribution, and excretion of titanium dioxide and zinc oxide nanoparticles after repeated oral administration. *Part. Fibre Toxicol.* 10, 9.
- Code of Federal Regulations, 2018. Listing of Color Additives Exempt from Certification. Available from: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/cfrsearch.cfm?fr=73.575>.
- Cooper, H.S., et al., 2000. Dysplasia and cancer in the dextran sulfate sodium mouse colitis model. Relevance to colitis-associated neoplasia in the human: a study of histopathology, B-catenin and p53 expression and the role of inflammation. *Carcinogenesis* 21 (4), 757–768.
- Corpet, D.E., Tache, S., 2002. Most effective colon cancer chemopreventive agents in rats: a systematic review of aberrant crypt foci and tumor data, ranked by potency. *Nutr. Cancer* 43 (1), 1–21.
- DeLoId, G.M., et al., 2017. An integrated methodology for assessing the impact of food matrix and gastrointestinal effects on the biokinetics and cellular toxicity of ingested engineered nanomaterials. *Part. Fibre Toxicol.* 14 (1), 40.
- Diebold, U., 2003. The surface science of titanium dioxide. *Surf. Sci. Rep.* 48 (5–8), 53–229.
- Dudefoi, W., et al., 2017. Criteria to define a more relevant reference sample of titanium dioxide in the context of food: a multiscale approach. *Food Addit. Contam. Part A Chem Anal Control Expo Risk Assess* 34 (5), 653–665.
- Dudefoi, W., et al., 2018. Evaluation of the content of TiO₂ nanoparticles in the coatings of chewing gums. *Food Addit. Contam. Part A Chem Anal Control Expo Risk Assess* 35 (2), 211–221.
- Foley, G.L., 2001. Overview of male reproductive pathology. *Toxicol. Pathol.* 29 (1), 49–63.
- Francescone, R., Hou, V., Grivennikov, S.I., 2015. Cytokines, IBD, and colitis-associated cancer. *Inflamm. Bowel Dis.* 21 (2), 409–418.
- Fu, Y., et al., 2014. Systemic immune effects of titanium dioxide nanoparticles after repeated intratracheal instillation in rat. *Int. J. Mol. Sci.* 15 (4), 6961–6973.
- Hamiza, O.O., et al., 2012. Amelioration of 1,2 Dimethylhydrazine (DMH) induced colon oxidative stress, inflammation and tumor promotion response by tannic acid in Wistar rats. *Asian Pac. J. Cancer Prev. APJCP* 13 (9), 4393–4402.
- Hansen, G.H., et al., 2009. Endocytic trafficking from the small intestinal brush border probed with FM dye. *Am. J. Physiol. Gastrointest. Liver Physiol.* 297 (4), G708–G715.
- Hirahara, K., Nakayama, T., 2016. CD4 + T-cell subsets in inflammatory diseases: beyond the Th1/Th2 paradigm. *Int. Immunol.* 28 (4), 163–171.
- Humans, I.W.G.o.t.E.o.C.R.t., 2010. Carbon black, titanium dioxide, and talc. *IARC Monogr. Eval. Carcinog. Risks Hum.* 93, 1–413.
- Hummel, T.Z., et al., 2014. Exogenous pigment in Peyer patches of children suspected of having IBD. *J. Pediatr. Gastroenterol. Nutr.* 58 (4), 477–480.
- Hyun, Y.S., et al., 2012. Role of IL-17A in the development of colitis-associated cancer. *Carcinogenesis* 33 (4), 931–936.
- James, R.W., Heywood, R., 1979a. Age-related variations in the testes and prostate of beagle dogs. *Toxicology* 12 (3), 273–279.

- James, R.W., Heywood, R., 1979b. Age-related variations in the testes of sprague-dawley rats. *Toxicol. Lett.* 4 (4), 257–261.
- Jia, F., et al., 2014. Effect of pubertal nano-TiO₂ exposure on testosterone synthesis and spermatogenesis in mice. *Arch. Toxicol.* 88 (3), 781–788.
- Joesse, M.E., et al., 2019. Tipping the balance: inhibitory checkpoints in intestinal homeostasis. *Mucosal Immunol.* 12 (1), 21–35.
- Kobayashi, N., et al., 2009. Comparative pulmonary toxicity study of nano-TiO₂ particles of different sizes and agglomerations in rats: different short- and long-term post-instillation results. *Toxicology* 264 (1–2), 110–118.
- Kozoni, V., et al., 2000. The effect of lithocholic acid on proliferation and apoptosis during the early stages of colon carcinogenesis: differential effect on apoptosis in the presence of a colon carcinogen. *Carcinogenesis* 21 (5), 999–1005.
- Kreyling, W.G., et al., 2017a. Quantitative biokinetics of titanium dioxide nanoparticles after intravenous injection in rats: Part 1. *Nanotoxicology* 11 (4), 434–442.
- Kreyling, W.G., et al., 2017b. Quantitative biokinetics of titanium dioxide nanoparticles after intratracheal instillation in rats: Part 3. *Nanotoxicology* 11 (4), 454–464.
- Kreyling, W.G., et al., 2017c. Quantitative biokinetics of titanium dioxide nanoparticles after oral application in rats: Part 2. *Nanotoxicology* 11 (4), 443–453.
- Lee, K.P., Trochimowicz, H.J., Reinhardt, C.F., 1985. Pulmonary response of rats exposed to titanium dioxide (TiO₂) by inhalation for two years. *Toxicol. Appl. Pharmacol.* 79 (2), 179–192.
- Lundqvist, M., et al., 2008. Nanoparticle size and surface properties determine the protein corona with possible implications for biological impacts. *Proc. Natl. Acad. Sci. U. S. A.* 105 (38), 14265–14270.
- Luu, M., Steinhoff, U., Visekruna, A., 2017. Functional heterogeneity of gut-resident regulatory T cells. *Clin. Transl. Immunol.* 6 (9), e156.
- Manolova, V., et al., 2008. Nanoparticles target distinct dendritic cell populations according to their size. *Eur. J. Immunol.* 38 (5), 1404–1413.
- Martin, R.A., et al., 1984. Randomization of animals by computer-program for toxicity studies. *J. Am. Coll. Toxicol.* 3 (1), 1–11.
- McClements, D.J., et al., 2016. The Role of the Food Matrix and Gastrointestinal Tract in the assessment of biological properties of ingested engineered nanomaterials (iENMs): state of the science and knowledge gaps. *NanoImpact* 3–4, 47–57.
- McCracken, C., et al., 2013. Minimal intestinal epithelial cell toxicity in response to short- and long-term food-relevant inorganic nanoparticle exposure. *Chem. Res. Toxicol.* 26 (10), 1514–1525.
- Mohamed, H.R., 2015. Estimation of TiO₂ nanoparticle-induced genotoxicity persistence and possible chronic gastritis-induction in mice. *Food Chem. Toxicol.* 83, 76–83.
- Nagayama, S., et al., 2007. Time-dependent changes in opsonin amount associated on nanoparticles alter their hepatic uptake characteristics. *Int. J. Pharm.* 342 (1–2), 215–221.
- National Toxicology Program, 1979. Bioassay of titanium dioxide for possible carcinogenicity. *Natl. Cancer Inst. Carcinog. Tech. Rep. Ser.* 97, 1–123.
- Nel, A.E., et al., 2009. Understanding biophysicochemical interactions at the nano-bio interface. *Nat. Mater.* 8 (7), 543–557.
- Neurath, M.F., 2014. Cytokines in inflammatory bowel disease. *Nat. Rev. Immunol.* 14 (5), 329–342.
- Oberdorster, G., Ferin, J., Lehnert, B.E., 1994. Correlation between particle size, in vivo particle persistence, and lung injury. *Environ. Health Perspect.* 102 (Suppl. 5), 173–179.
- Orazizadeh, M., et al., 2014. Effect of glycyrrhizic acid on titanium dioxide nanoparticles-induced hepatotoxicity in rats. *Chem. Biol. Interact.* 220, 214–221.
- Ortlieb, M., 2010. White giant or white dwarf?: particle size distribution measurements of TiO₂. *GIT Lab J. Eur.* 14, 42–43.
- Pelaseyed, T., et al., 2014. The mucus and mucins of the goblet cells and enterocytes provide the first defense line of the gastrointestinal tract and interact with the immune system. *Immunol. Rev.* 260 (1), 8–20.
- Perse, M., 2005. The dimethylhydrazine induced colorectal tumours in rat - experimental colorectal carcinogenesis. *Radiol. Oncol.* 39 (1).
- Powell, J.J., et al., 1996. Characterisation of inorganic microparticles in pigment cells of human gut associated lymphoid tissue. *Gut* 38 (3), 390–395.
- Powell, J.J., et al., 2010. Origin and fate of dietary nanoparticles and microparticles in the gastrointestinal tract. *J. Autoimmun.* 34 (3), J226–J233.
- Powell, J.J., et al., 2015. An endogenous nanomineral chaperones luminal antigen and peptidoglycan to intestinal immune cells. *Nat. Nanotechnol.* 10 (4), 361–369.
- Proquin, H., et al., 2018. Gene expression profiling in colon of mice exposed to food additive titanium dioxide (E171). *Food Chem. Toxicol.* 111, 153–165.
- Redston, M., 2001. Carcinogenesis in the GI tract: from morphology to genetics and back again. *Mod. Pathol.* 14 (3), 236–245.
- Regulation (EC) No 1333, 2008. Of the European Parliament and of the Council of 16 December 2008 on Food Additives. 2018 12/20/18. Available from: <https://eur-lex.europa.eu/eli/reg/2008/1333/2014-04-14>.
- Rosenberg, D.W., Giardina, C., Tanaka, T., 2009. Mouse models for the study of colon carcinogenesis. *Carcinogenesis* 30 (2), 183–196.
- Ruh, H., et al., 2012. Identification of serum proteins bound to industrial nanomaterials. *Toxicol. Lett.* 208 (1), 41–50.
- Ruiz, P.A., et al., 2017. Titanium dioxide nanoparticles exacerbate DSS-induced colitis: role of the NLRP3 inflammasome. *Gut* 66 (7), 1216–1224.
- Sager, T.M., Komminen, C., Castranova, V., 2008. Pulmonary response to intratracheal instillation of ultrafine versus fine titanium dioxide: role of particle surface area. *Part. Fibre Toxicol.* 5, 17.
- Sayes, C.M., et al., 2006. Correlating nanoscale titania structure with toxicity: a cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicol. Sci.* 92 (1), 174–185.
- Schmitt, N., Benteibibel, S.E., Ueno, H., 2014. Phenotype and functions of memory Tfh cells in human blood. *Trends Immunol.* 35 (9), 436–442.
- Scott, C.L., Aumeunier, A.M., Mowat, A.M., 2011. Intestinal CD103+ dendritic cells: master regulators of tolerance? *Trends Immunol.* 32 (9), 412–419.
- Shukla, R.K., et al., 2013. TiO₂ nanoparticles induce oxidative DNA damage and apoptosis in human liver cells. *Nanotoxicology* 7 (1), 48–60.
- Strugala, V., et al., 2003. Colonic mucin: methods of measuring mucus thickness. *Proc. Nutr. Soc.* 62 (1), 237–243.
- Talbot, P., et al., 2018. Food-grade TiO₂ is trapped by intestinal mucus in vitro but does not impair mucin O-glycosylation and short-chain fatty acid synthesis in vivo: implications for gut barrier protection. *J. Nanobiotechnol.* 16 (1), 53.
- Tassinari, R., et al., 2014. Oral, short-term exposure to titanium dioxide nanoparticles in Sprague-Dawley rat: focus on reproductive and endocrine systems and spleen. *Nanotoxicology* 8 (6), 654–662.
- Theissmann, R., Kluwig, M., Koch, T., 2014. A reproducible number-based sizing method for pigment-grade titanium dioxide. *Beilstein J. Nanotechnol.* 5, 1815–1822.
- Thoolen, B., et al., 2010. Proliferative and nonproliferative lesions of the rat and mouse hepatobiliary system. *Toxicol. Pathol.* 38 (7 Suppl. 1), 5S–81S.
- Urrutia-Ortega, I.M., et al., 2016. Food-grade titanium dioxide exposure exacerbates tumor formation in colitis associated cancer model. *Food Chem. Toxicol.* 93, 20–31.
- Van der Sluis, M., et al., 2006. Muc2-deficient mice spontaneously develop colitis, indicating that MUC2 is critical for colonic protection. *Gastroenterology* 131 (1), 117–129.
- Velcich, A., et al., 2002. Colorectal cancer in mice genetically deficient in the mucin Muc2. *Science* 295 (5560), 1726–1729.
- Walke, C.D., Chan, W.C., 2012. Understanding and controlling the interaction of nanomaterials with proteins in a physiological environment. *Chem. Soc. Rev.* 41 (7), 2780–2799.
- Wang, J., et al., 2007. Acute toxicity and biodistribution of different sized titanium dioxide particles in mice after oral administration. *Toxicol. Lett.* 168 (2), 176–185.
- Wang, Y., et al., 2013. Susceptibility of young and adult rats to the oral toxicity of titanium dioxide nanoparticles. *Small* 9 (9–10), 1742–1752.
- Wargovich, M.J., Brown, V.R., Morris, J., 2010. Aberrant crypt foci: the case for inclusion as a biomarker for colon cancer. *Cancers* 2 (3), 1705–1716.
- Warheit, D.B., et al., 2006. Pulmonary instillation studies with nanoscale TiO₂ rods and dots in rats: toxicity is not dependent upon particle size and surface area. *Toxicol. Sci.* 91 (1), 227–236.
- Williams, M.B., Butcher, E.C., 1997. Homing of naive and memory T lymphocyte subsets to Peyer's patches, lymph nodes, and spleen. *J. Immunol.* 159 (4), 1746–1752.
- Winkler, H.C., et al., 2018. Critical review of the safety assessment of titanium dioxide additives in food. *J. Nanobiotechnol.* 16 (1), 51.
- Winter, M., et al., 2011. Activation of the inflammasome by amorphous silica and TiO₂ nanoparticles in murine dendritic cells. *Nanotoxicology* 5 (3), 326–340.
- Wright Jr., J.R., et al., 1982. Testicular atrophy in the spontaneously diabetic BB Wistar rat. *Am. J. Pathol.* 108 (1), 72–79.
- Yamamura, K., 1999. Transformation using (x + 0.5) to stabilize the variance of populations. *Res. Popul. Ecol.* 41 (3), 229–234.
- Yang, Y., et al., 2014. Characterization of food-grade titanium dioxide: the presence of nanosized particles. *Environ. Sci. Technol.* 48 (11), 6391–6400.
- Yoshimi, N., et al., 1994. Expression of cytokines, TNF-alpha and IL-1 alpha, in MAM acetate and 1-hydroxyanthraquinone-induced colon carcinogenesis of rats. *Carcinogenesis* 15 (4), 783–785.
- Zhou, L., et al., 2016. Suggestive evidence for the induction of colonic aberrant crypts in mice fed sodium nitrite. *Nutr. Cancer* 68 (1), 105–112.