

Components of variation in female common cuckoo calls

Zhuqing Deng^a, Huw Lloyd^b, Canwei Xia^{a,*}, Anders P. Møller^{a,c}, Wei Liang^d, Yanyun Zhang^{a,*}

^a Ministry of Education Key Laboratory for Biodiversity and Ecological Engineering, College of Life Sciences, Beijing Normal University, Beijing 100875, China

^b School of Science and Environment, Manchester Metropolitan University, Manchester M1 5GD, UK

^c Ecologie Systématique Evolution, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, F-91405 Orsay Cedex, France

^d Ministry of Education Key Laboratory for Ecology of Tropical Islands, College of Life Sciences, Hainan Normal University, Haikou 571158, China

ARTICLE INFO

Keywords:

Acoustic signals
Call consistency
Common cuckoo
Female vocalization
Geographical variation
Vocal activity

ABSTRACT

Investigations on bird vocalizations have largely focused on males. Female vocalizations are widespread in birds but few studies have investigated female vocal characteristics, particularly in non-Passeriformes. In this study, we use new field recordings from China, and calls available from an online sound library to examine temporal patterns, call consistency and geographical variation in vocalizations of female common cuckoos *Cuculus canorus*. The peak in vocal activity (both male and female) was in the morning, which contrasts to what would be predicted if the sole function of the female call was to distract the attention of hosts after parasitizing a host's nest in the afternoon. Both male and female common cuckoos were more vocally active in sunny weather, than rainy weather. We also found larger intra-individual variation in female rather than in male calls, which may benefit female cuckoos by increasing stimulation to host species. Peak frequency of female calls decreased with increased latitude, while differences in female call features were not associated with geographic distance. In summary, our findings that female calls are used in the morning, rather than at peak egg laying, yet are highly variable and show little geographic patterns suggest that the function of these female calls may be more variable and intricate than previously thought. However, because research on female vocalizations is underrepresented, future studies are still needed.

1. Introduction

Studies of bird vocalizations have been critical in shaping our understanding of the relationship between avian signal evolution and species differentiation (Andersson, 1994). Historically, most attention on variation in avian acoustic signals has been directed towards males (Beecher and Brenowitz, 2005), with many bird populations showing regional differences in male vocal characteristics over local and/or wider geographical scales (Irwin, 2000; Boughman, 2002; Kaluthota et al., 2016). However, female vocalizations are also widespread in both temperate and tropical species (Garamszegi et al., 2007; Yang et al., 2011; Odom et al., 2014; Mahr et al., 2016).

Whilst females of some species use their acoustic signals to defend territories, coordinate breeding activities and attract mates (Langmore, 1998), mate attraction across bird species does not appear to be the primary function of female vocalizations (Dabelsteen et al., 1998; Cain and Langmore, 2016; Krieg and Getty, 2016). Female vocalisations vary in their complexity (Langmore, 1998; Cain and Langmore, 2016) but few studies have examined whether female vocal characteristics vary

geographically, particularly for non-Passeriformes (Odom and Benedict, 2018). Given the prevalence of female vocalizations, more research should be conducted to build a comprehensive understanding of the function and evolution of bird vocalizations.

Male common cuckoos *Cuculus canorus* utter loud simple and stereotypic “cuck-ooo” calls during the breeding season. Temporal and frequency variables among different male cuckoo calls are sufficient to provide individual information (Jung et al., 2014; Li et al., 2017; Zebok et al., 2017), which can be used to distinguish between neighbors and strangers (Moskat et al., 2017). Furthermore, there appears to be a high degree of consistency in the number of syllables in each bout produced by individual males (Møller et al., 2016a, b). However, males of different subspecies differ significantly in their calls (Wei et al., 2015), even within the same subspecies, and male calls from populations in the same habitat are more similar to each other than those from populations in different habitats (Fuisz and de Kort, 2007). In comparison, our knowledge of the characteristics and function of female cuckoo calls have, until recently, been largely overlooked. Female cuckoos often give a conspicuous ‘chuckle’ call (Payne, 2005). One

* Corresponding author.

E-mail addresses: dengzhuqing@mail.bnu.edu.cn (Z. Deng), H.Lloyd@mmu.ac.uk (H. Lloyd), xiacanwei@126.com (C. Xia), anders.moller@u-psud.fr (A.P. Møller), liangwei@hainnu.edu.cn (W. Liang), zhangyy@bnu.edu.cn (Y. Zhang).

<https://doi.org/10.1016/j.beproc.2018.10.007>

Received 13 July 2018; Received in revised form 18 September 2018; Accepted 17 October 2018

Available online 26 October 2018

0376-6357/ © 2018 Elsevier B.V. All rights reserved.

recent experimental playback study suggests that the chuckle call primarily serves as a distraction of the host parent species (e.g. reed warbler *Acrocephalus scirpaceus*) since female chuckle calls had the same effect on the attention of host and non-host species, as playback of the calls of sparrowhawk *Accipiter nisus* (York and Davies, 2017). In contrast, male common cuckoo calls had no such effect. Such a function would enable female cuckoos to benefit from reducing their egg rejection rate through distraction of the attention of hosts (York and Davies, 2017). Other aspects of female cuckoo calls remain unquantified in the peer-review literature (Liang, 2017; Kim et al., 2017).

In this study, we use new recordings and those deposited in the online avian acoustic sound library Xeno-Canto (<http://www.xeno-canto.org>), to examine temporal patterns, individual consistency, and geographical variation in the chuckle calls of female common cuckoos. Unlike most birds, which lay egg in the morning, female cuckoos predominantly lay their eggs in the afternoon (Payne, 2005), and the function of female cuckoo calls is to distract the host species after laying (York and Davies, 2017). Therefore, we predicted that peak female cuckoo calls should occur during the afternoon. Since sound signals are more easily distorted in bad weather (e.g. rain or strong winds) than good weather (e.g. sunny) (Lengagne and Slater, 2002), we predicted that cuckoo calls would be more frequent on days with good weather. Theoretically, elaborate signals could increase stimulation of sensory perception (Akre and Johnsen, 2014; Cui et al., 2016), which may reduce habituation in the distraction of the attention cuckoo hosts. Consequently, we predicted that female calls should be more variable than male calls. Besides the potential difference in call characteristics among different subspecies and populations in different habitats, there may also be differences in female call characteristics among different host races within the same population. As female cuckoo calls functionally mimic hawk calls (York and Davies, 2017), different populations may exhibit different call characteristics due to different geographic variation in hawk calls and/or hawk species with different calls occurring within the resident bird community. Thus, our final objective was to determine whether female chuckle calls exhibit geographic variation, but because this could be more complex than those for males, we made no specific prediction about geographic variation in female cuckoo calls.

2. Methods

2.1. Study area and sound recording

Fieldwork was conducted during June 8th to July 28th 2017 in the Liaohe Delta Nature Reserve (41.034°N; 121.725°E), Liaoning Province, Northeast China. This region represents one of the most important estuarine wetlands in China, with the largest area of reed-bed habitat along the coastal region of China, and consequently, extensive nesting habitats for Oriental reed warbler *Acrocephalus orientalis*. Here, the common cuckoo is a summer breeding species, and it predominantly parasitizes Oriental reed warbler nests (Li et al., 2016). Density of cuckoos is high in our study population (Li et al., 2016), where several individuals often occur in close proximity (less than 10 m) to each other but data on territory size is lacking. We used a TASCAM DR-100MKIII recorder (Tascam Co., Japan) and a Sennheiser MKH416 P48 external directional microphone (Sennheiser Co.,

Germany), with a sampling rate of 44.1 kHz and a sampling accuracy of 16 bits, to record cuckoo vocalizations. Further recordings were made using seven passive acoustic recorder SM4 Songmeters (Wildlife Acoustics Inc. USA) placed at seven different locations, separated by a minimal distance of 200 m, to continuously record cuckoo calls from June 8th to July 28th. The minimal distance between these recorders is larger than the effective recording range (100 m for cuckoos call, seen in Huang et al., 2017), in order to avoid the same call being recorded by two recorders. Recorders were attached to telegraph poles at a height of 3 m above ground, and set to record continuously at a sampling rate of 44.1 kHz, and a sampling accuracy of 16 bits. Recorders were checked every 10 days to replace the batteries and memory cards. Using mist nets, we also trapped and banded 20 individual adult common cuckoos (6 females and 14 males) around our recording sites, whilst daily observations also revealed many other unbanded individual cuckoos at our recording sites during data collection.

To examine geographic variation in female cuckoo song, we downloaded all recordings of female common cuckoo from the online sound library Xeno-Canto (<http://www.xeno-canto.org>). For multiple recordings collected on the same day at the same location such that individuals could not be identified, we randomly selected one recording for analysis to avoid pseudo-replication. We also used four recordings that we collected from Liaohe Delta Nature Reserve, and four recordings collected from Beijing Wild Duck Lake National Wetland Park (40.410°N; 115.829°E), which is situated approximately 500 km from the Liaohe Delta Nature Reserve. These eight recordings were made in June 2017 with the same equipment mentioned before.

2.2. Acoustic measurements

All recordings were re-sampled with 8 kHz, and saved as wav files. We used Avisoft-SASLab Pro 5.2 audio analysis software (Avisoft Bioacoustics, Germany) to generate spectrograms with the following settings: Fast Fourier transform length 256 points; Hamming window with a frame size of 100% and an overlap of 50%; frequency resolution 31 Hz; and time resolution of 16 ms. Here the phrase ‘number of calls’ refers to number of syllables in males, and the number of calling bouts in females. Male cuckoo calls consist of a repeated series of ‘cu-coo’ syllables (Møller et al., 2016a; Li et al., 2017) i.e. both ‘cu’ and ‘coo’ elements constitute a single syllable with several successive syllables constituting a bout. The pause between successive male calling bouts was always longer than 2 s in our recordings, which is obviously greater than the time interval between successive ‘cu-coo’ syllables within one calling bout (Fig. 1a). Female cuckoo calls consist of a series of rapidly repeated “kwik-kwik-kwik” notes (York and Davies, 2017), which we named a bout (Fig. 1b). Each “kwik” represented a syllable, corresponding to the terminology used for male calls. For each bout (both male and female), we measured maximum frequency, minimum frequency, peak frequency, duration, and the number of syllables. Peak frequency refers to the frequency associated with the maximum energy. Setting a standard to measure minimum frequency was problematic because energy decreases gradually towards low frequency in female cuckoo calls. Consequently, we did not include minimum frequency in the subsequent analyses.

To describe the temporal patterns of male and female cuckoo vocal activity, we used Kaleidoscope Pro Software (Wildlife Acoustics Inc.




Fig. 1. Spectrogram of male (A), and female (B) common cuckoo calls, recorded in Liaohe Delta Nature Reserve (41.034°N; 121.725°E).

USA) to create recognizers for finding all male and female calls from recordings collected with the SM4 Song meters. Firstly, we entered the following acoustic features of our target sound, either a male syllable or female bout, to Kaleidoscope Pro Software: frequency range from 400 to 1200 Hz for a male syllable, and 600 to 2900 Hz for a female calling bout; duration ranges from 0.3 to 0.5 s for a male syllable, and 1.6 to 4 s for a female calling bout. These acoustic features are slightly larger than actual parameters of male and female cuckoo calls, but this was done to increase the detectability of these calls by the Kaleidoscope Pro Software. Lastly, we manually checked all calls identified by the recognizer based on visual inspection of the spectrograms. In total, we obtained 701,661 male syllables, and 2407 female bouts after manual check.

2.3. Temporal patterns of vocal activity

We used both our field recordings and recordings downloaded from Xeno-Canto to examine temporal patterns of vocal behavior by male and female cuckoos. Using our field recordings from June 8th to July 24th we first calculated the number of calls per hour using absolute time since we lacked data on twilight time. Our prediction was that female cuckoos had a peak call output in the afternoon, based on the primary function of female calls being the distraction of hosts during egg laying by cuckoos. Thus, we assumed that the use of absolute time, rather than twilight time, would have little or no effect on tests of our prediction. We used a generalized linear mixed model (GLMM) to estimate the repeatability of number of calls per hour, based on the function *rpt* in the R package *rptR* (Stoffel et al., 2017). After confirming that most variation occurred among hours, rather than during the same hour on different days (repeatability of number of calls per hour in male 0.763 ± 0.056 mean \pm SE; 95% confidence interval ranged from 0.623 to 0.828; $P < 0.001$; in female = 0.872 ± 0.087 mean \pm SE; 95% confidence interval ranged from 0.609 to 0.910; $P < 0.001$), we pooled the data by calculating the mean number of calls per hour, and we used these data to illustrate temporal patterns of cuckoo vocal activity.

For a second data source, we used the time of cuckoo calls from the recordings downloaded from Xeno-Canto, again using absolute rather than twilight time for the analysis. From this second source, we obtained the time of 359 recordings of male calls, and 36 recordings of female calls and calculated the number of recordings for every two-hour period. Pearson correlation coefficient based on the number of calls (or recordings) in each hour from the Liaohe Delta Nature Reserve recordings and those from Xeno-Canto were used to determine patterns of similarity in vocal activity from the two separate data sources. Finally, we related patterns of vocal activity from the first data source to localized weather conditions. We defined weather condition based on the information from Weather China (www.weather.com.cn), and classified a day as bad i.e. days with rain or strong winds (with wind speed greater than 8 m/s) or good i.e. days without rain and strong winds.

2.4. Individual call consistency

We used a TASCAM DR-100MKIII recorder and a Sennheiser MKH416 P48 external directional microphone to obtain recordings from 18 males in the Liaohe Delta Nature Reserve during June and July 2017; six of these males were banded from our mist-netting efforts, whilst further recordings were made from unbanded males from locations separated by at least 2 km from each other, in an effort to reduce the likelihood of repeatedly sampling the same individual twice. Since female calls were much rarer than male calls in our study area (Fig. 2), we supplemented the dataset with recordings made using the passive acoustic recorder SM4 Songmeters. Since we could not be certain which individual calls belonged to the same individual in the passive acoustic recordings, we defined two bouts recorded by the same recorders within 1 min and with similar amplitude as being from the same




Fig. 2. The number of calls by male and female common cuckoos per day during the 2017 breeding season, summarized from data collected by seven recorders.

individual because we found that it is rare to hear different female calls at the same time in adjacent sites. Using this definition, we obtained 43 recordings, each containing two bouts, from 6 songmeters. These recordings were recorded in different locations, or at the same location on different days. Since common cuckoos are abundant in our study area (Li et al., 2017) we treated these 43 recordings as being derived from 43 different female individuals. To avoid pseudo-replication, we used a smaller data set only including six recordings (corresponding to six individual females) recorded from different locations, randomly selected from the original 43 recordings. We calculated the Pearson correlation coefficient from measurements on two successive bouts, and used this to reflect individual call consistency. We compared the correlation coefficients between males and females in the larger data set ($n = 43$), using the Fisher z-transformation to assess the significance of the difference between two correlation coefficients. We did not calculate the significance of the difference between correlation coefficients between males and females in the smaller data set because of the smaller sample size ($n = 6$).

2.5. Geographical variation

We measured 35 female recordings from Xeno-Canto, and supplemented this dataset with four recordings from Liaohe Delta Nature Reserve, and four recordings from Wild Duck Lake National Wetland Park. Although different equipment was used to collect recordings by Xeno-Canto recorders, we assume that any effects of equipment on recordings should only cause noise in the data set, and there is no reason to expect bias in such effects. We restricted our analyses to one bout in each recording, as most recordings downloaded from Xeno-Canto only contained one bout. Since acoustic measures vary by different orders of magnitude e.g. the frequency of cuckoo syllables ranges in the hundreds of Hz, while the duration of syllables lasts nearly a tenth of a second, acoustic measures were transformed into z-scores, and then used to calculate the Euclidean distance of all pairs of bouts. We used Mantel tests to assess the correlation between Euclidean distance and geographic distance, and Pearson correlation to compared the acoustic measures with latitude, as acoustic characteristics are known to change with latitude in many bird species (Kaluthota et al., 2016; Wei et al., 2017). All statistical analyses were performed using R software, v. 3.4.1 (R Core Team, 2015). Data are presented as mean \pm standard deviation. Differences with P values less than 0.05 were considered significant.

3. Results

3.1. Temporal variation in common cuckoo calls

At the Liaohe Delta Nature Reserve, we found that the number of cuckoo calls per day fluctuated widely, with a sharp decline after July 25th (Fig. 2). The number of male and female cuckoo calls per day was




Fig. 3. Call output (number of syllables summarized from data collected by seven recorders per day \pm SE) by (A) male and (B) female common cuckoos in relation to different weather conditions. Asterisk indicates significant difference at $P < 0.05$ (independent samples t test).




Fig. 4. Daily temporal pattern of common cuckoo call (A) based on recordings from seven recorders in Liaohe Delta Nature Reserve; (B) 359 male recordings and 36 female recordings from the online sound library Xeno-Canto. Error bars are SE.

significantly positively correlated (Pearson correlation, $r = 0.075$, $P < 0.001$, $n = 47$) and both male and female common cuckoos showed higher vocal activity during good weather (Fig. 3). Males on average uttered $17,192 \pm 1241$ (mean \pm SD) syllables summarized from data collected by seven recorders per day in good weather ($n = 25$ days), significantly more than the $11,867 \pm 899$ syllables per day during bad weather ($n = 22$ days; independent samples t test, $t_{45} = 3.392$, $P = 0.001$). Females uttered 61 ± 6 bouts summarized from data collected by seven recorders per day in good weather ($n = 25$ days), significantly larger than 40 ± 7 bouts per day during bad weather ($n = 22$ days; independent samples t-test, $t_{45} = 2.406$, $P = 0.02$). Data from both the Liaohe Delta Nature Reserve (Fig. 4a) and Xeno-Canto (Fig. 4b) revealed that peak call output occurred during the morning rather than in the afternoon, with noticeably little female call activity in the afternoon at the Liaohe Delta Nature Reserve. Number of calls per hour from Liaohe Delta Nature Reserve and Xeno-Canto were strongly positively correlated in both males ($r = 0.769$, $P < 0.001$) and females ($r = 0.655$, $P = 0.001$).

3.2. Individual consistency

Males ($n = 18$) generally showed higher individual call consistency




Fig. 5. Pearson correlation coefficients from measurements of vocalization features in common cuckoos in two successive bouts. Asterisk indicates significant difference at $P < 0.05$ (Z test). n is the number of individuals and error bars are SE.

than females ($n = 43$) (Fig. 5). Pearson correlation coefficients of maximum frequency (male: $r = 0.926$; female: $r = 0.618$), duration (male: $r = 0.488$; female: $r = 0.067$), and number of syllables (male: $r = 0.609$; female: $r = 0.087$) were significantly larger for males than for females (Z test, $Z = 3, 1.98, 2.05$; $P = 0.003, 0.048, 0.040$, respectively). Pearson correlation coefficients of peak frequency were larger in females ($r = 0.523$) than in males ($r = 0.179$), but the difference was not significant (Z test, $Z = 1.32$, $P = 0.187$). Pearson's correlation coefficients for females in the smaller data set (6 recordings made from different locations) showed a similar trend as Pearson's correlation coefficients for females in the larger data set (43 recordings made from different locations, or at the same location on different days): If Pearson correlation coefficients for females in the larger data set was larger (or lower) than Pearson correlation coefficients for males, Pearson's correlation coefficients for females in the smaller data set was also larger (or lower) than in males for each variable (Fig. 5).

3.3. Geographical variation

We found in female calls a maximum frequency of bouts of 2228 ± 40 Hz; peak frequency of bouts of 1905 ± 47 Hz; duration of bouts of 1.979 ± 0.056 s, with 19.53 ± 0.78 syllables. There was no significant linear trend of differences in female calls and geographic distance (Mantel test, $P = 0.256$) (Fig. 6a). Peak frequency decreased with increased latitude (Pearson's correlation coefficients, $r = -0.364$, $P = 0.016$, $n = 43$) (Fig. 6b), while maximum frequency ($r = -0.166$, $P = 0.289$), duration ($r = -0.031$, $P = 0.842$), and number of syllables ($r = -0.094$, $P = 0.547$) showed no significant relationship with




Fig. 6. (A) Lack of significant linear trend in female common cuckoo calls with geographic distance (Mantel test, $P = 0.256$); and (B) significant negative correlation between peak frequency and latitude (Pearson's correlation coefficients, $r = -0.364$, $P = 0.016$, $n = 43$).

latitude.

4. Discussion

4.1. Temporal patterns of female vocal activity

Based on the field recordings from Liaohe Delta Nature Reserve and others deposited on the online sound library Xeno-Canto, we found that vocal activity of female cuckoos peaked in morning rather than in the afternoon. We admit that the call recordings in Xeno-Canto are influenced by human activities (Blackburn et al., 2014) e.g. observers are more likely to spend more time recording birds in good weather and in the morning, which may lead to bias in the interpretation of daily vocal behavior patterns. In addition, to avoid potential bias due to differences in time across the different time zones represented in the Xeno-Canto recordings, we used the regional local time rather than twilight time as this would have little or no effect on tests of our prediction of peak call output in the afternoon. We used automatic recognizers generated from our field recordings to automatically identify potential cuckoo calls, all of which were then subsequently checked manually by visual inspection of the spectrograms. Previous research on Large Hawk Cuckoo (*Hierococcyx sparveroides*) found that only about 50% of all cuckoo calls were correctly identified by automatic recognizers from 96 h of recordings (Huang et al., 2017). Most of the common cuckoo calls that were not detected by the automatic recognizers in this study occurred during the dawn chorus, and were largely masked by the dawn songs of oscine passerines. Thus, female cuckoos may have higher peak of call output in the morning than we observed.

The peak of female call output in the morning is in contrast to what would be predicted if the only function of the female cuckoo call was to distract the attention of hosts after parasitizing a host's nest i.e. peak in call activity should occur during the afternoon (see York and Davies, 2017). Perhaps, female calls may be used to find nests of hosts in the morning. As most birds in the morning lay eggs or sit on eggs, female cuckoos produce calls which cause hosts to leave the nest, and then locate nests of hosts. It is also likely that female common cuckoo calls at our study site have other functions besides distraction hosts (Liang, 2017). For example, females calling in the morning may attract the attention of males (e.g. Langmore, 1998). Alternatively, it may be the case that potential host species at Liaohe Delta Nature Reserve are able to distinguish between female common cuckoo calls from hawk calls, or even use female calls as a predictor of risk of parasitism or increased probability of egg rejection, although further experimental playback research is needed.

4.2. Individual call consistency

Individual consistency as measured by repeatability is important in social behavior, but also as an upper limit for heritability and hence micro-evolution (Bell et al., 2009; Nakagawa and Schielzeth, 2010).

Male common cuckoo call features are consistent within individuals (Jung et al., 2014; Li et al., 2017; Zsebok et al., 2017), which can facilitate neighbor—stranger discrimination in birds (Moskat et al., 2017). For females, York and Davies (2017) reported that the primary function of calls is to distract the attention of host species. Elaborate signals could increase stimulation of sensory perception (Akre and Johnsen, 2014; Cui et al., 2016), which may reduce habituation of distraction of the attention of hosts. In agreement with this prediction, females generally show lower individual consistency in call characteristics than males: specifically, Pearson's correlation coefficients of maximum frequency, duration, and number of syllables in females are significantly lower than those of males. These results were consistent across both the larger (43 recordings recorded in different locations, or at the same location on different days) and the smaller (6 recordings recorded in different locations) data sets thus we believe that our sample sizes were sufficient to evaluate both within- and between-individual differences. Pearson's correlation coefficients of peak frequency in female calls was larger than in males, although the difference was not significant. For many bird species, peak frequency is determined by body size (Fletcher, 2004; Rodriguez et al., 2015) and is consistently higher in male cuckoo calls (Jung et al., 2014; Li et al., 2017; Zsebok et al., 2017). This acoustic feature could potentially be used for monitoring female cuckoos.

4.3. Geographical variation in female cuckoo calls

Geographic variation in bird vocalizations is common, and may affect mate choice, pair bonding, and territory defense in both passerine species such as stonechat *Saxicola torquata* (Mortega et al., 2014) and coal tit *Periparus ater* (Pentzold et al., 2016), and also non-passerine species such as gentoo penguin *Pygoscelis papua* (Lynch and Lynch, 2017) and corncrake *Crex crex* (Budka and Osiejuk, 2017). One of the more common patterns in bird acoustic geographic variation can be described in terms of the 'isolation by distance' model (Podos and Warren, 2007), in which vocalization differences in paired populations increase with the distance between those populations (Irwin, 2000; King et al., 2013). This pattern can be caused by both cultural drift and genetic differences, as there are few chances for culture and genetic exchange among remote populations (Irwin et al., 2008; Stewart and MacDougall-Shackleton, 2008; Ramsay and Otter, 2015). In accordance with this model, differences in calls and geographic distance are known to be closely correlated in male common cuckoos (Wei et al., 2015). However, in our study we found no evidence of isolation by distance, and we found no significant linear trend in differences in female cuckoo calls with geographic distance. For female cuckoos, there may be differences in call characteristics among different races within the same population, besides potential differences in call characteristics among different subspecies and populations in different habitats. If the purpose of female cuckoo calls is to functionally mimic hawk calls (York and Davies, 2017), different populations may have different call features

due to different hawk species with different calls occurring in the local bird community. These factors could lead to complexity in geographic variation of female cuckoo calls, and mask any linear trend between differences in calls and geographic distance as predicted in the isolation by distance model.

Peak frequency is thought to correlate negatively with body size (Fletcher, 2004; Rodriguez et al., 2015) but it is not unusual to find evidence to the contrary to this rule in some oscines (e.g. rufous-collared sparrow *Zonotrichia capensis* Handford and Lougheed, 1991; dark-eyed junco *Junco hyemalis* and serin *Serinus serinus* Cardoso et al., 2008), due to song learning and sexual selection (Patel et al., 2010; Cardoso, 2012). However, this rule is generally supported in suboscines and non-passerines, particularly for species with large body size and low acoustic frequency in vocalizations such as doves (Tubaro and Mahler, 1998), tinamous (Bertelli and Tubaro, 2002), and antbirds (Seddon, 2005). In this study, we found that peak frequency of female cuckoo calls decreased at higher latitude. In cuckoos, both male and female body size is known to increase at higher latitudes (Payne, 2005; Erritzøe et al., 2012), and this may lead to the negative relationship between peak frequency and latitude.

5. Conclusions

In this study, we investigated three aspects (temporal patterns, call consistency, geographical variation) concerning female common cuckoo calls. York and Davies (2017) suggested that female common cuckoos mimic sparrowhawk vocalizations in order to distract the host species after laying. In accordance with this prediction, we found larger intra-individual variation in female calls, which may increase stimulation of the host species. Besides, we found peak frequency of female calls decreased with increased latitude, while differences in female call features were not associated with geographic distance. However, the daily vocal pattern contrasts with what would be predicted according to York and Davies (2017). If the only function of the female common cuckoo call was to distract the attention of hosts after parasitizing a host's nest, peak call output should occur during the afternoon because female common cuckoos predominantly lay their eggs in the afternoon (Payne, 2005). Data from both the Liaohe Delta Nature Reserve and Xeno-Canto revealed that peak call output occurred during the morning rather than in the afternoon. Based on this result, we infer that female common cuckoo calls have other functions besides distraction hosts.

Authors' contributions

YZ, WL designed the experiments; ZD participated in the field work; ZD, CX carried out the analyses; CX drafted the earlier version of the manuscript and HL, AM, ZD revised it. All authors have read and approved the final manuscript.

Funding

This study was supported by the National Key Technology R & D Program of China (No. 2016YFC0503200), National Natural Science Foundation of China (No. 31872243 to YZ, No. 31601868 to CX, and 31472453 to WL) and the State Forestry Administration of China (No. KJHX2016135).

Availability of data and materials

All recordings and data in this study are available from the corresponding author (xiacanwei@126.com) on reasonable request.

Ethical standards

Our research protocol was approved by the Animal Management Committee at the College of Life Sciences, Beijing Normal University

under license number CLS-EAW-2016-017. Bird capture and banding were permitted by the National Bird-banding Center of China under license number H20110042. The experiments comply with the current laws of the country in which they were performed.

Declarations of interest

None.

Acknowledgements

We thank Xiaomeng Zhao, Boning Xue, Qi Wang, Qi Luo, Qingbin Wang, Shuang Guan, Ziqiang Huang, Linyu Jin and Juan Chen for assistance in the field work.

References

- Akre, K.L., Johnsen, S., 2014. Psychophysics and the evolution of behavior. *Trends Ecol. Evol. (Amst.)* 29, 291–300. <https://doi.org/10.1016/j.tree.2014.03.007>.
- Andersson, M., 1994. *Sexual Selection*. Princeton University Press, Princeton.
- Beecher, M.D., Brenowitz, E.A., 2005. Functional aspects of song learning in songbirds. *Trends Ecol. Evol.* 20, 143–149. <https://doi.org/10.1016/j.tree.2005.01.004>.
- Bell, A.M., Hankison, S.J., Laskowski, K.L., 2009. The repeatability of behaviour: a meta-analysis. *Anim. Behav.* 77, 771–783. <https://doi.org/10.1016/j.anbehav.2008.12.022>.
- Bertelli, S., Tubaro, P.L., 2002. Body mass and habitat correlates of song structure in a primitive group of birds. *Biol. J. Linn. Soc. Lond.* 77, 423–430. <https://doi.org/10.1046/j.1095-8312.2002.00112.x>.
- Blackburn, T.M., Su, S., Cassey, P., 2014. A Potential metric of the attractiveness of bird song to humans. *Ethology* 120, 305–312. <https://doi.org/10.1111/eth.12211>.
- Boughman, J.W., 2002. How sensory drive can promote speciation. *Trends Ecol. Evol. (Amst.)* 17, 571–577. [https://doi.org/10.1016/S0169-5347\(02\)02595-8](https://doi.org/10.1016/S0169-5347(02)02595-8).
- Budka, M., Osiejuk, T.S., 2017. Microgeographic call variation in a non-learning species, the Corncrake (*Crex crex*). *J. Ornithol.* 158, 651–658. <https://doi.org/10.1007/s10336-017-1438-7>.
- Cain, K.E., Langmore, N.E., 2016. Female song and aggression show contrasting relationships to reproductive success when habitat quality differs. *Behav. Ecol. Sociobiol. (Print)* 70, 1867–1877. <https://doi.org/10.1007/s00265-016-2192-1>.
- Cardoso, G.C., 2012. Paradoxical calls: the opposite signaling role of sound frequency across bird species. *Behav. Ecol.* 23, 237–241. <https://doi.org/10.1093/beheco/arr200>.
- Cardoso, G.C., Mamede, A.T., Atwell, J.W., Mota, P.G., Ketterson, E.D., Price, T.D., 2008. Song frequency does not reflect differences in body size among males in two oscine species. *Ethology* 114, 1084–1093. <https://doi.org/10.1111/j.1439-0310.2008.01552.x>.
- Cui, J., Song, X., Zhu, B., Fang, G., Tang, Y., Ryan, M.J., 2016. Receiver discriminability drives the evolution of complex sexual signals by sexual selection. *Evolution* 70, 922–927. <https://doi.org/10.1111/evo.12889>.
- Dabelsteen, T., McGregor, P.K., Lampe, H., Langmore, N., Holland, J., 1998. Quiet song in song birds: an overlooked phenomenon. *Bioacoustics* 9, 89–105. <https://doi.org/10.1080/09524622.1998.9753385>.
- Erritzøe, J., Mann, C.F., Brammer, F.P., Fuller, R.A., 2012. *Cuckoos of the World*. Bloomsbury, London.
- Fletcher, N.H., 2004. A simple frequency-scaling rule for animal communication. *J. Acoust. Soc. Am.* 115, 2334–2338. <https://doi.org/10.1121/1.1694997>.
- Fuisz, T.I., de Kort, S.R., 2007. Habitat-dependent call divergence in the common cuckoo: is it a potential signal for assortative mating? *Proc. R Soc. Lond. B* 274, 2093–2097. <https://doi.org/10.1098/rspb.2007.0487>.
- Garamszegi, L.Z., Pavlova, D.Z., Eens, M., Möller, A.P., 2007. The evolution of song in female birds in Europe. *Behav. Ecol.* 18, 86–96. <https://doi.org/10.1093/beheco/arl047>.
- Handford, P., Lougheed, S.C., 1991. Variation in duration and frequency characters in the song of the rufous-collared sparrow, *Zonotrichia capensis*, with respect to habitat, trill dialects and body size. *Condor* 93, 644–658. <https://doi.org/10.2307/1368196>.
- Huang, W., Xu, S., Liang, W., Xia, C., 2017. Daily vocal pattern of large hawk cuckoo (*Hierococcyx sparveroides*). *Chinese J. Zool.* 52, 945–953. <https://doi.org/10.13859/j.cjz.201706003>.
- Irwin, D., 2000. Song variation in an avian ring species. *Evolution* 54, 998–1010. [https://doi.org/10.1554/0014-3820\(2000\)054\[0998:SVIAR\]2.3.CO;2](https://doi.org/10.1554/0014-3820(2000)054[0998:SVIAR]2.3.CO;2).
- Irwin, D.E., Thimban, M.P., Irwin, J.H., 2008. Call divergence is correlated with geographic and genetic distance in greenish warblers (*Phylloscopus trochiloides*): a strong role for stochasticity in signal evolution? *J. Evol. Biol.* 21, 435–448. <https://doi.org/10.1111/j.1420-9101.2007.01499.x>.
- Jung, W.J., Lee, J.W., Yoo, J.C., 2014. "Cu-coo": can you recognize my stepparents? A study of host-specific male call divergence in the common cuckoo. *PLoS One* 9, e90468. <https://doi.org/10.1371/journal.pone.0090468>.
- Kaluthota, C., Brinkman, B.E., Dos Santos, E.B., Rendall, D., 2016. Transcontinental latitudinal variation in song performance and complexity in house wrens (*Troglodytes aedon*). *Proc. R Soc. Lond. B* 283, 20152765. <https://doi.org/10.1098/rspb.2015.2765>.

- Kim, H., Lee, J.W., Yoo, J.C., 2017. Comparing vocal structures of the parasitic and nonparasitic groups in Cuculinae. *Avian Res.* 8, 27. <https://doi.org/10.1186/s40657-017-0084-3>.
- Krieg, C.A., Getty, T., 2016. Not just for males: females use song against male and female rivals in a temperate zone songbird. *Anim. Behav.* 113, 39–47. <https://doi.org/10.1016/j.anbehav.2015.12.019>.
- Langmore, N.E., 1998. Functions of duet and solo songs of female birds. *Trends Ecol. Evol. (Amst.)* 13, 136–140. <https://doi.org/10.1163/156853998793066267>.
- Lengagne, T., Slater, P.J.B., 2002. The effects of rain on acoustic communication: tawny owls have good reason for calling less in wet weather. *Proc. R. Soc. Lond. B* 269, 2121–2125. <https://doi.org/10.1098/rspb.2002.2115>.
- Li, D., Ruan, Y., Wang, Y., Chang, A.K., Wan, D., Zhang, Z., 2016. Egg-spot matching in common cuckoo parasitism of the oriental reed warbler: effects of host nest availability and egg rejection. *Avian Res.* 7, 199–209. <https://doi.org/10.1186/s40657-016-0057-y>.
- Li, Y., Xia, C., Lloyd, H., Li, D., Zhang, Y., 2017. Identification of vocal individuality in male cuckoos using different analytical techniques. *Avian Res.* 8, 21. <https://doi.org/10.1186/s40657-017-0079-0>.
- Liang, W., 2017. Crafty cuckoo calls. *Nat. Ecol. Evol.* 1, 1427–1428. <https://doi.org/10.1038/s41559-017-0321-5>.
- Lynch, M.A., Lynch, H.J., 2017. Variation in the ecstatic display call of the gentoo penguin (*Pygoscelis papua*) across regional geographic scales. *Auk* 134, 894–902. <https://doi.org/10.1016/j.auc.2016.09.006>.
- Møller, A.P., Morelli, F., Mousseau, T.A., Tryjanowski, P., 2016a. The number of syllables in Chernobyl cuckoo calls reliably indicate habitat, soil and radiation levels. *Ecol. Indic.* 66, 592–597. <https://doi.org/10.1016/j.ecolind.2016.02.037>.
- Møller, A.P., Morelli, F., Tryjanowski, P., 2016b. Cuckoo folklore and human well-being: cuckoo calls predict how long farmers live. *Ecol. Indic.* 72, 766–768. <https://doi.org/10.1016/j.ecolind.2016.09.006>.
- Mahr, K., Seifert, C.L., Hoi, H., 2016. Female and male blue tits (*Cyanistes caeruleus*) sing in response to experimental predator exposition. *J. Ornithol.* 157, 907–911. <https://doi.org/10.1007/s10336-016-1345-3>.
- Mortega, K.G., Flinks, H., Helm, B., 2014. Behavioural response of a migratory songbird to geographic variation in song and morphology. *Front. Zool.* 11, 85. <https://doi.org/10.1186/s12983-014-0085-6>.
- Moskat, C., Elek, Z., Ban, M., Geltsch, N., Hauber, M.E., 2017. Can common cuckoos discriminate between neighbours and strangers by their calls? *Anim. Behav.* 126, 253–260. <https://doi.org/10.1016/j.anbehav.2017.02.013>.
- Nakagawa, S., Schielzeth, H., 2010. Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists. *Biol. Rev.* 85, 935–956. <https://doi.org/10.1111/j.1469-185X.2010.00141.x>.
- Odom, K.J., Benedict, L., 2018. A call to document female bird songs: applications for diverse fields. *Auk* 135, 314–325. <https://doi.org/10.1642/AUK-17-183.1>.
- Odom, K.J., Hall, M.L., Riebel, K., Omland, K.E., Langmore, N.E., 2014. Female song is widespread and ancestral in songbirds. *Nat. Commun.* 5. <https://doi.org/10.1038/ncomms4379>. 3379–3379.
- Patel, R., Mulder, R.A., Cardoso, G.C., 2010. What makes vocalisation frequency an unreliable signal of body size in birds? A study on black swans. *Ethology* 116, 554–563. <https://doi.org/10.1111/j.1439-0310.2010.01769.x>.
- Payne, R.B., 2005. *The Cuckoos*. Oxford University Press, Oxford.
- Pentzold, S., Foerschler, M.I., Tietze, D.T., Randler, C., Martens, J., Päckert, M., 2016. Geographic variation in coal tit song across continents and reduced species recognition between central European and Mediterranean populations. *Vertebr. Zool.* 66, 191–199. <https://doi.org/10.15496/publikation-14608>.
- Podos, J., Warren, P.S., 2007. The evolution of geographic variation in birdsong. *Adv. Study Behav.* 37, 403–458. [https://doi.org/10.1016/S0065-3454\(07\)37009-5](https://doi.org/10.1016/S0065-3454(07)37009-5).
- R Core Team, 2015. R: A Language and Environment for Statistical Computing. <http://www.R-project.org>.
- Ramsay, S.M., Otter, K.A., 2015. Geographic variation in white-throated sparrow song may arise through cultural drift. *J. Ornithol.* 156, 763–773. <https://doi.org/10.1007/s10336-015-1183-8>.
- Rodriguez, R.L., Araya-Salas, M., Gray, D.A., Reichert, M.S., Symes, L.B., Wilkins, M.R., Safran, R.J., Höbel, G., 2015. How acoustic signals scale with individual body size: common trends across diverse taxa. *Behav. Ecol.* 26, 168–177. <https://doi.org/10.1093/beheco/aru174>.
- Seddon, N., 2005. Ecological adaptation and species recognition drives vocal evolution in neotropical subsocial birds. *Evolution* 59, 200–215. <https://doi.org/10.1554/04-300>.
- Stewart, K.A., MacDougall-Shackleton, E.A., 2008. Local song elements indicate local genotypes and predict physiological condition in song sparrows *Melospiza melodia*. *Biol. Lett.* 4, 240–242. <https://doi.org/10.1098/rsbl.2008.0010>.
- Stoffel, M.A., Nakagawa, S., Schielzeth, H., 2017. rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* 8, 1639–1644. <https://doi.org/10.1111/2041-210X.12797>.
- Tubaro, P.L., Mahler, B., 1998. Acoustic frequencies and body mass in new world doves. *Condor* 100, 54–61. <https://doi.org/10.2307/1369896>.
- Wei, C., Jia, C., Dong, L., Wang, D., Xia, C., Zhang, Y., Liang, W., 2015. Geographic variation in the calls of the common cuckoo (*Cuculus canorus*): isolation by distance and divergence among subspecies. *J. Ornithol.* 156, 533–542. <https://doi.org/10.1007/s10336-014-1153-6>.
- Wei, C., Price, T.D., Liu, J., Alström, P., Zhang, Y., 2017. The evolutionary origin of variation in song length and frequency in the avian family Cettiidae. *J. Avian Biol.* 48, 1295–1300. <https://doi.org/10.1111/jav.01366>.
- Xing, X., Alström, P., Yang, X., Lei, F., 2013. Recent northward range expansion promotes song evolution in a passerine bird, the light-vented bulbul. *J. Evol. Biol.* 26, 867–877. <https://doi.org/10.1111/jeb.12101>.
- Yang, C., Zhang, Y., Cai, Y., Stokke, B.G., Liang, W., 2011. Female crowing and differential responses to simulated conspecific intrusion in male and female Hainan partridge (*Arborophila ardens*). *Zool. Sci.* 28, 249–253. <https://doi.org/10.2108/zsj.28.249>.
- York, J.E., Davies, N.B., 2017. Female cuckoo calls misdirect host defences towards the wrong enemy. *Nat. Ecol. Evol.* 1, 1520–1525. <https://doi.org/10.1038/s41559-017-0279-3>.
- Zsebok, S., Moskat, C., Ban, M., 2017. Individually distinctive vocalization in common cuckoos (*Cuculus canorus*). *J. Ornithol.* 158, 213–222. <https://doi.org/10.1007/s10336-016-1376-9>.