

Contents available at [ScienceDirect](https://www.sciencedirect.com)Diabetes Research  
and Clinical Practicejournal homepage: [www.elsevier.com/locate/diabres](http://www.elsevier.com/locate/diabres)International  
Diabetes  
Federation

## Review

# Efficacy and safety of cinnamon in type 2 diabetes mellitus and pre-diabetes patients: A meta-analysis and meta-regression



Serawit Deyno<sup>a,b,\*</sup>, Kassahun Eneyew<sup>b</sup>, Sisay Seyfe<sup>b</sup>, Naasson Tuyiringire<sup>a,c</sup>, Emanuel L. Peter<sup>a,d</sup>, Rekik Ashebir Muluye<sup>e</sup>, Casim Umba Tolo<sup>a</sup>, Patrick Engeu Ogwang<sup>a</sup>

<sup>a</sup> Pharm-BioTechnology and Traditional Medicine Center of Excellence (PHARMBIOTRAC), Mbarara University of Science and Technology, P.O. Box 1410, Mbarara, Uganda

<sup>b</sup> Faculty of Medicine, College of Medicine and Health Sciences, Hawassa University, P. O. Box 1560, Hawassa, Ethiopia

<sup>c</sup> School of Nursing and Midwifery, College of Medicine and Health Sciences, University of Rwanda, University Avenue, P.O. Box 56, Butare, Rwanda

<sup>d</sup> Department of Innovation, Technology Transfer & Commercialization, National Institute for Medical Research, Barack Obama Drive, P.O. Box 9653, 2448 Dar Es Salaam, Tanzania

<sup>e</sup> Traditional and Modern Drug Research Directorate, Ethiopian Public Health Institute, Ethiopia

## ARTICLE INFO

## Article history:

Received 24 May 2019

Received in revised form

8 August 2019

Accepted 12 August 2019

Available online 16 August 2019

## Keywords:

Herbal therapy

Blood sugar level

Diabetes mellitus

Efficacy

Systematic review

## ABSTRACT

**Introduction:** Cinnamon has been used as a dietary component and in the management of diabetes mellitus. This study systematically reviewed and synthesized evidence on the efficacy of cinnamon for the treatment of type 2 diabetes mellitus (T2DM) and pre-diabetes patients.

**Methods:** Databases of Web of Sciences, the Cochrane library, PubMed, CINAHL and SCOPUS were searched. Stata version 13 (College Station, Texas 77845 USA) and RevMan var. 5.3 software were used for meta-analysis. Heterogeneity was assessed using Chi-square and  $I^2$  tests.

**Results:** Sixteen randomized controlled studies were included in the meta-analysis. Cinnamon significantly reduced fasting blood glucose (FBG) and homeostatic model assessment for insulin resistance (HOMA-IR) level compared to placebo with weighted mean difference (WMD) of  $-0.545$  (95% CI:  $-0.910, -0.18$ ) mmol/L,  $I^2 = 83.6\%$  and  $-0.714$  ( $-1.388, -0.04$ ),  $I^2 = 84.4\%$  respectively. There was no significant change in weighted mean difference of gly-

**Abbreviations:** ALT, alanine transaminase; AST, aspartate transaminase; DM, diabetes mellitus; FBG, fasting blood glucose; Hb1Ac, glycosylated hemoglobin 1Ac; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; RBG, random blood glucose; T2DM, type 2 diabetes mellitus; TC, total cholesterol; TG, triglycerides; WMD, weighted mean difference

\* Corresponding author at: Pharm-BioTechnology and Traditional Medicine Center of Excellence (PHARMBIOTRAC), Mbarara University of Science and Technology, P.O. Box 1410, Mbarara, Uganda.

E-mail addresses: [dserawit@std.must.ac.ug](mailto:dserawit@std.must.ac.ug) (S. Deyno), [ntuyiringire@std.must.ac.ug](mailto:ntuyiringire@std.must.ac.ug) (N. Tuyiringire), [epeter@std.must.ac.ug](mailto:epeter@std.must.ac.ug) (E.L. Peter), [tolocas@must.ac.ug](mailto:tolocas@must.ac.ug) (C.U. Tolo), [pogwang@must.ac.ug](mailto:pogwang@must.ac.ug) (P.E. Ogwang).

<https://doi.org/10.1016/j.diabres.2019.107815>

0168-8227/© 2019 Elsevier B.V. All rights reserved.

cosylated hemoglobin A1C (HbA1c) % and lipid profiles (mmol/L). Meta-regression did not show any factor significantly affecting the treatment response.

**Conclusion:** Cinnamon reduced FBG and HOMA-IR, level in T2DM and pre-diabetes patients compared to placebo. High heterogeneity observed among included studies warrants further clinical trials after standardization of cinnamon formulation.

© 2019 Elsevier B.V. All rights reserved.

## Contents

1. Introduction	2
2. Methods	3
2.1. Study design	3
2.2. Search strategy	3
2.3. Inclusion and exclusion criteria	3
2.4. Data extraction	3
2.5. Study quality assessment	3
2.6. Statistical analysis	3
3. Results	3
3.1. Characteristics of included studies	3
3.2. Heterogeneity and publication bias in the included studies	5
3.3. Risk of bias in the included study	5
3.4. Effect of interventions	5
3.5. Heterogeneity and publication bias	5
3.6. Safety outcomes	5
4. Discussion	6
4.1. Current findings	6
4.2. Findings of the current study in light of previous meta-analysis	6
4.3. Limitation of the study	6
5. Conclusion	12
Authors' contributions	12
Funding	12
Declaration of Competing Interest	12
References	12

## 1. Introduction

Diabetes is a chronic metabolic disorder characterized by elevated blood glucose level resulting from a defect in insulin secretion, insulin action, or both [1]. Long-term complications of diabetes mellitus (DM) include retinopathy, nephropathy, neuropathy, periodontal disease, sexual dysfunction, and increased risk of cardiovascular diseases [2,3]. Over 90% of all DM cases are type 2 DM (T2DM) (IDF, 2017; [4]. The increase in T2DM is predominantly related to increasing urbanization, population ageing, obesity, ethnicity, family history of diabetes [4], history of gestation diabetes [5], increasing physical inactivity [6], and unhealthy diet [7].

Management of T2DM involves the use of modern anti-diabetic drugs. However, the anti-diabetic drugs are expensive [8], have significant adverse effects [9] and fail to effectively control the glycemia [10]. Herbal medicines become the alternative way for DM management as they have few adverse

effects and readily accessible to the majority of the populations. Cinnamon is among the many herbal medicines used for the treatment of DM. It has two main varieties, *Cinnamomum cassia* (also known as *Cinnamomum aromaticum*) and *Cinnamomum zeylanicum* [11]. *In vitro* and *in vivo* studies have shown that cinnamon is an insulin sensitizer [12]. Clinical trials evaluated the efficacy of cinnamon in glycemic control among people with T2DM.

Previous four meta-analytic studies produced inconsistent results [13,14,15,16]. Since the last meta-analysis, several clinical trials were conducted and published. The findings of these studies have been included in the current meta-analysis and meta-regression to update the relevance of the previous systematic review and meta-analysis and also identify possible explanation for heterogeneity. Hence, this study systematically reviewed and synthesized evidence on the efficacy of cinnamon for the treatment of patients with T2DM and pre-diabetes patients.

## 2. Methods

### 2.1. Study design

This is a meta-analysis based on electronic databases. The meta-analysis is prepared and structured in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analysis guidelines [17]. The protocol of this study was registered (PROSPERO 2018 = CRD42018106474) [18].

### 2.2. Search strategy

PubMed databases were initially searched to outline the study and construct keywords. After construction of search words, further searches were made in databases of Web of Sciences, SCOPUS, CINAHL, and the Cochrane library. The keywords used include [cinnamon OR Cinnamomum] AND [diabetes OR glycem\* OR hyperglyce\* OR hypoglyce\*] AND Clinical trial OR controlled trial OR randomized trial. Then text words contained in the title and abstract were analyzed. A second search using all identified keywords and index terms was undertaken across all included databases. Thirdly, the reference list of all identified studies was searched for additional studies. Unpublished studies were searched in Google and Google Scholar. Ongoing clinical trials were also searched through clinicaltrials.gov.

Two independent reviewers (KE & SS) conducted the title and abstract screening in duplicate to identify eligible articles using predefined criteria with support of SD. Full texts of the eligible articles were obtained and assessed against the inclusion and exclusion criteria. Any disagreements between the reviewers during full-text assessment were resolved by discussion and consensus, when no resolution reached, a third reviewer (SD) was involved in a decision.

### 2.3. Inclusion and exclusion criteria

The predefined inclusion criteria for this meta-analysis were as follows: 1, Randomized Controlled Trials (RCTs) in T2DM patients or pre-diabetes patients 2, Participants of the studies should be aged greater than 18 years and older of either sex 3, Studies with follow up duration of at least four weeks for both primary and secondary outcomes were considered. Since clinical studies have indicated that this is the minimum period required for treatments to produce meaningful changes in glucose control as assessed by HbA1c concentrations [19]. This systematic review excluded non-randomized clinical trials, cross-sectional studies, case series and case reports studies, studies conducted on patients younger than 18 years, and T2DM patients.

### 2.4. Data extraction

Joanna Briggs Institutes Meta-Analysis of Statistics Assessment and Review Instrument (JBI-MASARI) tools were used to extract the data [20]. Data extracted include last name of the first author, year of study, the dosage of the cinnamon and the control (placebo), type of population, duration of the

follow-up, and the primary and secondary outcomes of interest. The primary outcomes were FBG, HbA<sub>1c</sub>, insulin level, LDL, HDL, TC, BMI, HOMA-IR, Alanine aminotransferase (ALT), and Aspartate aminotransferase (AST). In general, specific details about the interventions and populations were extracted.

### 2.5. Study quality assessment

Two independent reviewers (KE and SS) assessed the eligibility of articles for methodological validity before inclusion in the review using standardized Cochrane risk of bias tool for a randomized clinical trial [21]. Disagreements between the two reviewers were resolved through discussion with a principal investigator (SD). The bias was graded as high, low, or unclear. Publication bias was tested using Egger's test [22].

### 2.6. Statistical analysis

RevMan5.3 software (Cochrane Informatics and Knowledge Management Department, London, UK) and Stata version 13.0 (StataCorp, LP, college station, TX) were used for the analysis of the data. All of the variables in this study were continuous, and the inverse of the variance-weighted method was used for pooling the weighted mean differences and its 95% confidence intervals (CI). Clinical and methodological heterogeneity were assessed using Chi-square test and  $I^2$  tests [21]. Because of significant heterogeneity amongst the studies, the random-effects model (REM) was used to estimate the pooled mean difference and 95% CIs using the DerSimonian and Laird method [23]. The presence of publication bias was tested using Egger's test [22]. This meta-analysis was conducted as per the Preferred Reporting Items for Systematic Reviews and Meta-Analysis guidelines [17]. Subgroup analysis and meta-regression were performed to identify the sources of heterogeneity.

## 3. Results

### 3.1. Characteristics of included studies

A total of 572 articles were identified through the electronic database search. After removing duplications and screening the articles based on titles and abstracts, 347 articles remained. Further screening for inclusion and exclusion criteria left 35 for full-text examination. Nineteen were further excluded with reasons. Reasons for exclusion: four of the articles were not conducted on DM patients [24,25,26,27], six articles did not address the study question [28–33], three were review papers [34,35,36], four were meta-analysis [13,14,15,16], one did not report standard deviation [37], one was conducted in type 1 DM [38]. This exclusion left sixteen articles for inclusion in the meta-analysis, [39,40,41,42,43 44,45,46,47,48,49,50,51,52, 53,54]. These sixteen studies enrolled a total of 1098 participants. A PRISMA flowchart is presented to show the screened, excluded and included articles (Fig. 1) and characteristics of included studies were summarized in Table 1.




Fig. 1 – Flow diagram of study selection, screening, inclusion and exclusion.

Table 1 – Characteristics of included studies.

Study	Type of DM	Total participants		Dosage	Follow up duration	Cinnamon botanical source & formulation	Placebo formulation
		Enrolled	Completed				
[39]	Type II	58	58	2 g/d	12 week	C. cassia,	Starch
[40]	Pre-diabetes	173	137	1 g/d	3 month	Commercial spray-dried extract of cinnamon	Wheat flour
[47]	Type II	60	43	1 g/d	3 month	C. cassia, capsules	Wheat flour
[41]	Type II	109	89	1 g/d	3 month	C. cassia, capsules	Usual care
[42]	Type II	35	35	1 g/d	60 days	C. cassia, capsules	Not specified
[52]	Type II	14	14	1.5 g	30 days	Not given, capsules	Maize flour
[51]	Type II	60	60	6 g/d	60 days	C. cassia	Wheat flour
[43]	Pre-diabetic	62	52	1.2 g	4 month	C. cassia, capsules	
[53]	Type II	69	66	14.4 g/d	3 month	C. aromaticum	Not specified
[48]	Type II	79	65	3 g/d	4 month	C. cassia,	cellulose
[44]	Type II	105	102	1 g/d	3 month	Not specified	Starch
[54]	Type II	44	39	3 g/d	8 week	C. zeylanicum	Cellulose
[49]	Type II	44	37	3 g/d	8 week	C. zeylanicum	Wheat flour
[50]	Type II	25	25	1.5 g/d	6 week	C. cassia	Wheat flour
[45]	Pre-diabetes	21	17	12 g/d	12 week	C. cassia, capsules	Cellulose
[46]	Type II	140	138	1 g/d	3 month	C. verum, capsules	Starch

### 3.2. Heterogeneity and publication bias in the included studies

All of the sixteen included studies in the systematic review were randomized controlled trials. However, there were significant variations among the included studies, both clinically and statistically. Clinical variations include; the strength of the dosage used, duration of the treatment, and type of patients included. The strength of the cinnamon used for the treatment ranges from 1 g to 14.4 g. The follow-up period ranges from one month to four months. Three studies were conducted in adults with T2DM patients either on oral anti-hyperglycemic agents or recently diagnosed T2DM. Some of the studies have background oral hypoglycemic as standard treatment while others do not have or have not provided evidence on the background therapy. The heterogeneity in the included studies was witnessed by high  $I^2$  ranging from 49.3% to 86%, all p-value greater than 0.05. Egger's publication bias test were significant for TG (p-value = 0.032) while all other parameters which satisfied conditions for Eggers test did not show significant value.

### 3.3. Risk of bias in the included study

The risk of bias was particularly high in the domain of detection, selection and performance bias due to failure of the included studies to adequately blind participants, personnel and outcome assessment as well as to sufficiently conceal allocation sequence. Low risk of bias were given when specific methods for random sequence generation were described using one of computer-generated random sequence, permuted block method, sealed envelope, random numeric table, numerical code, and RAS software. The risk of bias graph and risk of bias summary were given in Fig. 2 and Fig. 3.

### 3.4. Effect of interventions

Fifteen studies evaluated cinnamon effect on FBG. Cinnamon significantly reduced FBG (mmol/L) and compared to placebo

respectively with weighted mean difference (WMD) of  $-0.545$  (95% CI:  $-0.910, -0.18$ ), high heterogeneity was observed,  $I^2 = 83.6\%$ , Fig. 4. Four studies reported HOMA-IR and meta-analysis revealed significant reduction in WMD ( $-0.714$  (95% CI:  $1.388, -0.040$ ),  $I^2 = 84.4.1\%$ ), Fig. 5. Eight studies evaluated effect on insulin ( $\mu\text{U}/\text{mL}$ ) and found non-significant insulin reduction ( $-0.964$ [95% CI:  $-1.97, -0.042$ ],  $I^2 = 55.4\%$ ), Fig. 6. Other evaluating the effectiveness of cinnamon on biochemical parameters such as HbA<sub>1c</sub> (%), LDL (mmol/L), HDL (mmol/L), TC (mmol/L), TG (mmol/L), and BMI ( $\text{kg}/\text{m}^2$ ) respectively included twelve, twelve, eleven, eleven, twelve, and seven studies. All these parameters did not show significant differences in WMD reduction between cinnamon and placebo treated groups. The reduction in WMD for these parameters were: HbA<sub>1c</sub>% ( $-0.104$  [95% CI:  $-0.138, 0.110$ ],  $I^2 = 69.6\%$ ), LDL (mmol/L) ( $-0.115$  [95%CI:  $-0.270, 0.039$ ],  $I^2 = 86.01\%$ ), HDL (mmol/L) ( $-0.004$  [95%CI:  $-0.059$  to  $0.050$ ],  $I^2 = 81.0\%$ ), TC (mmol/L) ( $-0.198$ [95% CI:  $-0.405,0.010$ ],  $I^2 = 86.4\%$ , TG (mmol/L) ( $-0.100$  [95% CI:  $-0.221,0.022$ ],  $I^2 = 69.0\%$ , and BMI ( $\text{kg}/\text{m}^2$ ) ( $-0.37$ [95% CI:  $-1.12, 0.39$ ],  $I^2 = 57.0\%$ ). The forest plots for these parameters were respectively presented as Figs. 7–12 for HbA<sub>1c</sub>, LDL, HDL, TC, TG, and BMI.

### 3.5. Heterogeneity and publication bias

Considerable heterogeneity was observed in the included studies,  $I^2$  ranging from 55.4% to 86.0%. Meta-regression performed to account for heterogeneity using type of DM, type of cinnamon, the dosage of cinnamon, and baseline values did not show significant values, Table 2.

### 3.6. Safety outcomes

Almost all of the studies included reported that cinnamons were tolerated in individuals taking the medications. Two studies evaluated specific parameters for safety [53,45], reported ALT and AST values. Both parameters did not show significant variation in cinnamon compared to placebo. Liver function test performed using AST ( $\mu\text{kat}/\text{L}$ ) and ALT ( $\mu\text{kat}/\text{L}$ )




Fig. 2 – Risk of bias graph: review authors' judgments about each risk of bias item presented as percentages across all included studies.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Akilen et al, 2010	+	+	+	+	+	-	?
Anderson et al, 2016	+	+	+	+	?	?	?
Belivens et al, 2007	?	?	?	+	+	?	?
Crawford, 2009	+	+	-	+	+	?	?
Hasanzade et al, 2013	+	+	+	+	?	?	?
Khan et al, 2003	?	?	?	?	?	?	?
Khan et al, 2010	?	?	?	?	?	?	?
Liu et al, 2015	+	+	+	+	+	?	?
Lu et al, 2012	?	+	+	+	?	?	?
Mang et al, 2006	?	+	+	+	-	?	?
Mirfeizi et al., 2016	+	+	+	+	?	?	?
Talaei et al, 2017	?	+	+	+	+	?	?
Vafa et al, 2012	?	?	+	+	+	-	?
Vanschoonbeek et al, 2006	?	?	+	+	?	?	?
Wickenberg et al, 2014	+	+	?	+	?	?	?
Zare et al, 2018	+	+	+	+	?	+	?

**Fig. 3 – Risk of bias summary: review authors’ judgments about each risk of bias item for each included study.**

on two studies revealed WMD respectively as 0.27(95% CI: -3.20, 3.74),  $I^2 = 58\%$  and 3.5(95% CI: -3.43, 10.44),  $I^2 = 63\%$ .

## 4. Discussion

### 4.1. Current findings

This study revealed that cinnamon significantly reduced FBG and HOMA-IR level in patients with T2DM or pre-diabetes compared to placebo. However, there is no significant difference in HbA1c level, in various lipid profiles and BMI between cinnamon and placebo-treated groups. Cinnamon is generally reported as well tolerated in all of the included studies.

Herbal medications are used along with modern medicine to attain glycemic control as adjuvant therapy. The ability of cinnamon to control FBG but not HbA1c could be due to short duration of therapy where the follow-up period ranges only from one month to four months [55,56]. HbA1c also correlates well with the prevention of the risk of long-term diabetes complications [57,58]. The significant reduction in FBG and HOMA-IR could indicate that the possible mechanism of action could be due to insulin sensitization effect [12]. Cinnamon could, therefore, be anticipated to do better when combined with anti-diabetic agents with secretagogue activity [59] or insulin itself compared to drugs with insulin sensitizers [60]. Cinnamon has shown an increasing effect on Phosphorylation activity of insulin receptors, decreasing effect on tyrosine phosphatase activity, showing insulin-like properties [61] possibly due to its insulin sensitizer effect. Studies have also shown that cinnamon inhibits glycogen synthase activity [62].

### 4.2. Findings of the current study in light of previous meta-analysis

A total of ten reviews on the efficacy of cinnamon for treatment of DM were obtained, among which four were meta-analysis [13,14,15,16]. The conclusion in the meta-analyses varied possibly due to the difference in dosage strength, frequency of administration, cinnamon formulation, source and type, clinical variability, or background therapy. One study ended up with in-depth narrative review due to heterogeneity amongst the included studies [63].

The first meta-analysis concluded that cinnamon does not appear to improve HbA1c, FBG, or lipid parameters in DM [16] while the second meta-analysis showed a significant decrease in HbA1c and FBG level. The meta-analysis by [15] was inconclusive with non-significant difference in HbA1c, serum insulin, or postprandial glucose where the authors concluded that there is insufficient evidence to support the use of cinnamon for DM. The last meta-analysis by [13] established that cinnamon significantly reduced FBG compared to placebo while [15] failed to provide conclusive evidence. No significant differences were observed on the level of HbA1c in both meta-analyses [15,13]. Costello and colleagues concluded that cinnamon supplements added to standard hypoglycemic medications and other lifestyle therapies had modest effects on FBG and HbA1c [63]. In general, all of the meta-analyses revealed high heterogeneity and inconsistency which could be attributed to variation in both clinical and methodological parameters.

### 4.3. Limitation of the study

High heterogeneity was observed which is not resolved by meta-regression. Majority of included studies did not report safety data which make this study fall short of conclusive tangible safety data. High heterogeneity, small sample size and unavailability of specific safety require cautious interpretation of the findings. Despite this limitation, the current study provides useful evidence for clinical decision making and planning future studies.




Fig. 4 – Forest plot showing reduction in FBG (mmol/L) in cinnamon treated group versus placebo treated group.




Fig. 5 – Forest plot showing homeostatic model assessment for insulin resistance (HOMA-IR) in cinnamon treated versus placebo treated group.




Fig. 6 – Forest plot showing Insulin ( $\mu\text{U/mL}$ ) level in cinnamon treated group versus placebo treated group.




Fig. 7 – Forest plot showing HbA1c level in cinnamon treated versus placebo treated group.




Fig. 8 – Forest plot showing LDL (mmol/L) level in cinnamon treated versus placebo treated group.




Fig. 9 – Forest plot showing HDL (mmol/L) in cinnamon treated group versus placebo treated group.




Fig. 10 – Forest plot showing TC level (mmol/L) in cinnamon treated group versus placebo treated group.




Fig. 11 – Forest plot showing TG (mmol/L) level in cinnamon treated group versus placebo treated group.




Fig. 12 – Forest plot showing BMI (kg/m<sup>2</sup>) in cinnamon treated group versus placebo treated group.

Table 2 – Meta-regression on factors affecting treatment response.

Parameters	Independent variable	Coefficient	95% CI		P-value	
					unadjusted	adjusted
FBG	Type of DM	0.6837343	-3.130722	4.498191	0.651	0.977
	Cinnamon type	-0.3673474	-1.474011	0.7393162	0.552	0.939
	Dose strength	-0.1566368	-0.5677789	0.2545053	0.418	0.832
HBA1c	Follow-up days	-0.0164402	-0.0331336	0.0660139	0.489	0.903
	Type of DM	0.1153504	-0.8404791	1.07118	0.828	1.000
	Cinnamon type	-0.1519301	-0.5070144	0.2031542	0.333	0.724
TC	Dose strength	-0.0926305	-0.2728043	0.0875433	0.241	0.610
	Follow-up days	-0.0050237	-0.0175607	0.0075133	0.386	0.803
	Type of DM	0.0931593	-1.18078	1.367099	0.794	0.998
LDL	Cinnamon type	-0.0310705	-0.3674204	0.3052795	0.852	0.999
	Dose strength	-0.0003277	-0.1371922	0.1365368	0.954	1.000
	Follow-up days	0.0041113	-0.0121693	0.020392	0.534	0.930
HDL	Type of DM	0.1633249	-0.8060875	1.132737	0.590	0.957
	Cinnamon type	-0.0255926	-0.273205	0.2220198	0.801	0.995
	Dose strength	0.0060709	-0.0956677	0.1078094	0.942	1.000
TG	Follow-up days	0.0025773	-0.0091249	0.0142795	0.568	0.945
	Type of DM	-0.1083974	-0.4794793	0.2626845	0.548	0.933
	Cinnamon type	-0.0441675	-0.1170607	0.0287256	0.234	0.547
Insulin	Dose strength	0.0001992	-0.0398358	0.0402342	0.989	1.000
	Follow-up days	0.0001691	-0.0038792	0.0042173	0.924	1.000
	Type of DM	0.1905146	-0.5258414	0.9068705	0.558	0.940
Insulin	Cinnamon type	0.0181225	-0.2076415	0.2438865	0.866	1.000
	Dose strength	-0.0166791	-0.0855962	0.0522381	0.602	0.955
	Follow-up days	0.0013554	-0.0076775	0.0103883	0.721	0.991
	Type of DM	5.941361	-0.2838175	12.16654	0.026	0.063
Insulin	Cinnamon type	0.6095515	-1.795996	3.015099	0.504	0.802
	Dose strength	-0.2005707	-0.7162316	0.3150902	0.231	0.587
	Follow-up days	-0.0651718	-0.1446713	0.0143277	0.077	0.150

## 5. Conclusion

Cinnamon significantly reduced elevated FBG and HOMA-IR compared to placebo. However, there is no significant reduction in HbA1c and lipid profiles levels between cinnamon treated and placebo-treated T2DM patients or pre-diabetes patients. Meta-regression did not provide evidence for high level of heterogeneity. Further research is warranted for standardization of cinnamon formulation, clinical validation of the standardized formulation and further clinical trials.

## Authors' contributions

SD, SS, and KE conceptualized the idea and design of the study. KE and SS screened the studies and extracted the data. SD drafted the manuscript. SD and ELP conducted data analysis. CUT and PEO mentored. NT, RA, ELP, CUP, and PEO revised the manuscript for important intellectual contents. All authors read and approved the final manuscript.

## Funding

This research has no specific grant from funding agencies in the public, commercial, or not-for-profit sectors. However, the support from World Bank project, PHARMBIOTRAC, was crucial.

## Declaration of Competing Interest

The authors have no competing interests.

## REFERENCES

- [1] AMERICAN DIABETES, A. 2009. Diagnosis and classification of diabetes mellitus. *Diabetes care*, 32 Suppl 1, S62-S67.
- [2] Viazzi Francesca, Russo Giuseppina Tiziana, Ceriello Antonio, Fioretto Paola, Giorda Carlo, De Cosmo Salvatore, Pontremoli Roberto. Natural history and risk factors for diabetic kidney disease in patients with T2D: lessons from the AMD-annals. *J Nephrol* 2019;32(4):517-25. <https://doi.org/10.1007/s40620-018-00561-3>.
- [3] Stojanovic M, Cvetanovic G, Andelkovic-Apostolovic M, Stojanovic D, Rancic N. Impact of socio-demographic characteristics and long-term complications on quality of life in patients with diabetes mellitus. *Cent Eur J Public Health* 2018;26:104-10.
- [4] WHO, Global report on diabetes mellitus; Geneva, 2016. (ISBN 978 92 4 156525 7)
- [5] Bellamy L, Casas J-P, Hingorani AD, Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *Lancet* 2009;373:1773-9. [https://doi.org/10.1016/S0140-6736\(09\)60731-5](https://doi.org/10.1016/S0140-6736(09)60731-5).
- [6] Whiting DR, Guariguata L, Weil C, Shaw J. IDF Diabetes Atlas: Global Estimates of the Prevalence of diabetes for 2011 and 2030. *Diabetes Res Clin Pract* 2011;94(3):311-21. <https://doi.org/10.1016/j.diabres.2011.10.029>.
- [7] Ley SH, Hamdy O, Mohan V, Hu FB. Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *Lancet* 2014;383(9933):1999-2007. [https://doi.org/10.1016/S0140-6736\(14\)60613-9](https://doi.org/10.1016/S0140-6736(14)60613-9).
- [8] Smith-Spangler CM, Bhattacharya J, Goldhaber-Fiebert JD. Diabetes, its treatment, and catastrophic medical spending in 35 developing countries. *Diabetes Care* 2012;35:319-26. <https://doi.org/10.2337/dc11-1770>.
- [9] Singh A, Dwivedi S. Study of adverse drug reactions in patients with diabetes attending a tertiary care hospital in New Delhi, India. *Indian J Med Res* 2017;145. [https://doi.org/10.4103/ijmr.IJMR\\_109\\_16](https://doi.org/10.4103/ijmr.IJMR_109_16).
- [10] Alhadramy MS. Diabetes and oral therapies: a review of oral therapies for diabetes mellitus. *J Taibah Univ Med Sci* 2016;11(4). <https://doi.org/10.1016/j.jtumed.2016.02.001>.
- [11] Ranasinghe P, Galappaththya P, Constantine GR, Jayawardena R, Weeratunga HD, Premakumara S, et al. Cinnamomum zeylanicum (Ceylon cinnamon) as a potential pharmaceutical agent for type-2 diabetes mellitus: study protocol for a randomized controlled trial. *Trials* 2017;18:017-2192.
- [12] Qin B, Nagasaki M, Ren M, Bajotto G, Oshida Y, Sato Y. Cinnamon extract (traditional herb) potentiates in vivo insulin-regulated glucose utilization via enhancing insulin signaling in rats. *Diabetes Res Clin Pract* 2003;62:139-48.
- [13] Allen RW, Schwartzman E, Baker WL, Coleman CI, Phung OJ. Cinnamon use in type 2 diabetes: an updated systematic review and meta-analysis. *Ann Fam Med* 2013;11:452-9.
- [14] Akilen R, Tsiami A, Devendra D, Robinson N. Cinnamon in glycaemic control: Systematic review and meta analysis. *Clinical Nutrition* 2012;31:609-15.
- [15] Leach MJ, Kumar S. Cinnamon for diabetes mellitus. *Cochrane Database Syst Rev* 2012:12.
- [16] Baker WL, Gutierrez-Williams G, White CM, Kluger J, Coleman CI. Effect of cinnamon on glucose control and lipid parameters. *Diabetes Care* 2008;31:41-3.
- [17] Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010;8:336-41.
- [18] DEYNO, S., ENEYEW, K., SEYFE, S., TUYIRINGIRE, N., PETER, E., ASHEBIR, R., TOLO, C. U. & OGWANG, P. E. 2018. The efficacy and safety of cinnamon in patients with type 2 diabetes mellitus: systematic review and meta-analysis of randomized clinical trials [Online]. 2018].
- [19] Sidorenkov G, Haaijer-Ruskamp FM, de Zeeuw D, Denig P. A longitudinal study examining adherence to guidelines in diabetes care according to different definitions of adequacy and timeliness. *PLoS ONE* 2011;6(9). <https://doi.org/10.1371/journal.pone.0024278>.
- [20] JBI. Meta-analysis of statistics: assessment and review instrument (JBI mastari). Adelaide: Joanna Briggs Institute; 2006.
- [21] Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- [22] Egger M, Davey SG, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34.
- [23] Dersimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
- [24] Markey O, McClean CM, Medlow P, Davison GW, Trinick TR, Duly E, et al. Effect of cinnamon on gastric emptying, arterial stiffness, postprandial lipemia, glycemia, and appetite responses to high-fat breakfast. *Cardiovasc Diabetol* 2011;10:1475-2840.
- [25] Beejmohun V, Peytavy-Izard M, Mignon C, Muscente-Paque D, Deplanque X, Ripoll C, et al. Acute effect of Ceylon cinnamon extract on postprandial glycemia: alpha-amylase inhibition, starch tolerance test in rats, and randomized crossover clinical trial in healthy volunteers. *BMC Complement Altern Med* 2014;14:1472-6882.

- [26] Borzoei A, Rafrat M, Asghari-Jafarabadi M. Cinnamon improves metabolic factors without detectable effects on adiponectin in women with polycystic ovary syndrome. *Asia Pac J Clin Nutr* 2018;27:556–63.
- [27] Wickenberg J, Lindstedt S, Berntorp K, Nilsson J, Hlebowicz J. Ceylon cinnamon does not affect postprandial plasma glucose or insulin in subjects with impaired glucose tolerance. *Br J Nutr* 2012;107:1845–9.
- [28] Aghasi M, Ghazi-Zahedi S, Koochani F, Siassi F, Nasli-Esfahani E, Keshavarz A, et al. The effects of green cardamom supplementation on blood glucose, lipids profile, oxidative stress, sirtuin-1 and irisin in type 2 diabetic patients: a study protocol for a randomized placebo-controlled clinical trial. *BMC Complement Altern Med* 2018;18:017–2068.
- [29] Azimi P, Ghiasvand R, Feizi A, Hariri M, Abbasi B. Effects of cinnamon, cardamom, saffron, and ginger consumption on markers of glycemic control, lipid profile, oxidative stress, and inflammation in Type 2 diabetes patients. *Rev Diabet Stud* 2014;11:258–66.
- [30] Azimi P, Ghiasvand R, Feizi A, Hosseinzadeh J, Bahreynian M, Hariri M, et al. Effect of cinnamon, cardamom, saffron and ginger consumption on blood pressure and a marker of endothelial function in patients with type 2 diabetes mellitus: a randomized controlled clinical trial. *Blood Press* 2016;25:133–40.
- [31] Bernardo MA, Silva ML, Santos E, Moncada MM, Brito J, Proenca L, et al. Effect of cinnamon tea on postprandial glucose concentration. *J Diabetes Res* 2015;913651:14.
- [32] Crawford P, Thai C, Obholz J, Schievenin J, True M, Shah SA, et al. Assessment of the effect of lifestyle intervention plus water-soluble cinnamon extract on lowering blood glucose in pre-diabetics, a randomized, double-blind, multicenter, placebo controlled trial: study protocol for a randomized controlled trial. *Trials* 2016;17:015–1138.
- [33] Hlebowicz J, Hlebowicz A, Lindstedt S, Bjorgell O, Hoglund P, Holst JJ, et al. Effects of 1 and 3 g cinnamon on gastric emptying, satiety, and postprandial blood glucose, insulin, glucose-dependent insulinotropic polypeptide, glucagon-like peptide 1, and ghrelin concentrations in healthy subjects. *Am J Clin Nutr* 2009;89:815–21.
- [34] Pham AQ, Kourlas H, Pham DQ. Cinnamon supplementation in patients with type 2 diabetes mellitus. *Pharmacotherapy* 2007;27:595–9.
- [35] Dugoua JJ, Seely D, Perri D, Cooley K, Forelli T, Mills E, et al. From type 2 diabetes to antioxidant activity: a systematic review of the safety and efficacy of common and cassia cinnamon bark. *Can J Physiol Pharmacol* 2007;85:837–47.
- [36] Kirkham S, Akilen R, Sharma S, Tsiami A. The potential of cinnamon to reduce blood glucose levels in patients with type 2 diabetes and insulin resistance. *Diabetes Obes Metab* 2009;11:1100–13.
- [37] Sahib AS. Anti-diabetic and antioxidant effect of cinnamon in poorly controlled type-2 diabetic Iraqi patients: A randomized, placebo-controlled clinical trial. *J Intercult Ethnopharmacol* 2016;5:108–13.
- [38] Altschuler JA, Casella SJ, Mackenzie TA, Curtis KM. The effect of cinnamon on A1C among adolescents with type 1 diabetes. *Diabetes Care* 2007;30:813–6.
- [39] Akilen R, Tsiami A, Devendra D, Robinson N. Glycated haemoglobin and blood pressure-lowering effect of cinnamon in multi-ethnic Type 2 diabetic patients in the UK: a randomized, placebo-controlled, double-blind clinical trial. *Diabet Med* 2010;27:1159–67.
- [40] Anderson RA, Zhan Z, Luo R, Guo X, Guo Q, Zhou J, et al. Cinnamon extract lowers glucose, insulin and cholesterol in people with elevated serum glucose. *J Tradit Complement Med* 2015;6:332–6.
- [41] Crawford P. Effectiveness of cinnamon for lowering hemoglobin A1C in patients with type 2 diabetes: a randomized, controlled trial. *J Am Board Fam Med* 2009;22:507–12.
- [42] Hasanzade F, Toliat M, Emami SA, Emamimoghaadam Z. The Effect of Cinnamon on Glucose of Type II Diabetes Patients. *J Traditional Complement Med* 2013;3:171–4.
- [43] Liu Y, Cotillard A, Vatieer C, Bastard J-P, Fellahi S, Stévant M, et al. A Dietary supplement containing cinnamon, chromium and carnosine decreases fasting plasma glucose and increases lean mass in overweight or obese pre-diabetic subjects: a randomized Placebo-Controlled Trial. *PLOS One* 2015;10:e0138646.
- [44] Mirfeizi M, Tourzani ZM, Mirfeizi SZ, Jafarabadi MA, Rezvani HR, Afzali M. Controlling type 2 diabetes mellitus with herbal medicines: a triple-blind randomized clinical trial of efficacy and safety. *J Diabetes* 2015;8(5). <https://doi.org/10.1111/1753-0407.12342>.
- [45] Wickenberg J, Lindstedt S, Nilsson J, Hlebowicz J. Cassia cinnamon does not change the insulin sensitivity or the liver enzymes in subjects with impaired glucose tolerance. *Nutr J* 2014;13:1475–2891.
- [46] Zare R, Nadjarzadeh A, Zarshenas MM, Shams M, Heydari M. Efficacy of cinnamon in patients with type II diabetes mellitus: a randomized controlled clinical trial. *Clin Nutr* 2018;11:30114–6.
- [47] Blevins SM, Leyva MJ, Brown J, Wright J, Scofield RH, Aston CE. Effect of cinnamon on glucose and lipid levels in non insulin-dependent type 2 diabetes. *Diabet. Care* 2007;30:2236–7.
- [48] Mang B, Wolters M, Schmitt B, Kelb K, Lichtinghagen R, Stichtenoth DO, et al. Effects of a cinnamon extract on plasma glucose, HbA<sub>1c</sub>, and serum lipids in diabetes mellitus type 2. *Eur J Clin Invest* 2006;36:340–4.
- [49] Vafa M, Mohammadi F, Shidfar F, Sormaghi MS, Heidari I, Golestan B, et al. Effects of cinnamon consumption on glycemic status, lipid profile and body composition in type 2 diabetic patients. *Int J Prevent Med* 2012;3:531–6.
- [50] Vanschoonbeek K, Thomassen BJ, Senden JM, Wodzig WK, van Loon LJ. Cinnamon supplementation does not improve glycemic control in postmenopausal type 2 diabetes patients. *J Nutr* 2006;136:977–80.
- [51] Khan A, Safdar M, Khan MM, Khattak KN, Anderson RA. Cinnamon improves glucose and lipids of people with type 2 diabetes. *Diabet Care* 2003;26:3215–8.
- [52] Khan Radhia, Khan Zakkia, Shah Safdar Hussain. Cinnamon May Reduce Glucose, Lipid and Cholesterol Level in Type 2 Diabetic Individuals. *Pakistan J Nutr* 2010;9(5):430–3. <https://doi.org/10.3923/pjn.2010.430.433>.
- [53] Lu T, Sheng H, Wu J, Cheng Y, Zhu J, Chen Y. Cinnamon extract improves fasting blood glucose and glycosylated hemoglobin level in Chinese patients with type 2 diabetes. *Nutr Res* 2012;32:408–12.
- [54] Talaei B, Amouzegar A, Sahranavard S, Hedayati M, Mirmiran P, Aziz F. Effects of cinnamon consumption on glycemic indicators, advanced glycation end products, and antioxidant status in Type 2 diabetic patients. *Nutrients* 2017;9:991.
- [55] Rosado J. A study to determine the effects of cinnamon on blood glucose and lipid levels in person's with type-2 diabetes. University of Hawaii at Manoa; 2010.
- [56] Yazdanpanah S, Rabiee M, Tahriri M, Abdolrahim M, Rajab A, Jazayeri HE, et al. Evaluation of glycated albumin (GA) and GA/HbA<sub>1c</sub> ratio for diagnosis of diabetes and glycemic control: A comprehensive review. *Crit Rev Clin Lab Sci* 2017;54:219–32.
- [57] Syed IA, Khan WA. Glycated haemoglobin—a marker and predictor of cardiovascular disease. *J Pak Med Assoc* 2011;61:690–5.

- [58] Silbernagel G, Grammer TB, Winkelmann BR, Boehm BO, Marz W. Glycated hemoglobin predicts all-cause, cardiovascular, and cancer mortality in people without a history of diabetes undergoing coronary angiography. *Diabetes Care* 2011;34:1355–61.
- [59] Mascarello A, Frederico MJ, Castro AJ, Mendes CP, Dutra MF, Woehl VM, et al. Novel sulfonyl(thio)urea derivatives act efficiently both as insulin secretagogues and as insulinomimetic compounds. *Eur J Med Chem* 2014;86:491–501.
- [60] Chen Y, Ma H, Zhu D, Zhao G, Wang L, Fu X, et al. Discovery of novel insulin sensitizers: promising approaches and targets. *PPAR Res* 2017;8360919:4.
- [61] Olefsky JM. Treatment of insulin resistance with peroxisome proliferator-activated receptor gamma agonists. *J Clin Invest* 2000;106:467–72.
- [62] Jarvill-Taylor KJ, Anderson RA, Graves DJ. A hydroxychalcone derived from cinnamon functions as a mimetic for insulin in 3T3-L1 adipocytes. *J Am Coll Nutr* 2001;20:327–36.
- [63] Costello RB, Dwyer JT, Saldanha L, Bailey RL, Merkel J, Wambogo E. Do cinnamon supplements have a role in glycemic control in type 2 diabetes? a narrative review. *J Academy Nutrition Dietetics* 2016;116:1794–802.