



## Early and late effects of maternal experience on hippocampal neurogenesis, microglia, and the circulating cytokine milieu



Rand S. Eid<sup>a,b</sup>, Jessica A. Chaiton<sup>b,c</sup>, Stephanie E. Lieblich<sup>b,c</sup>, Tamara S. Bodnar<sup>d</sup>, Joanne Weinberg<sup>a,d</sup>, Liisa A.M. Galea<sup>a,b,c,\*</sup>

<sup>a</sup> Graduate Program in Neuroscience, University of British Columbia, Vancouver, British Columbia, Canada

<sup>b</sup> Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver, British Columbia, Canada

<sup>c</sup> Department of Psychology, University of British Columbia, Vancouver, British Columbia, Canada

<sup>d</sup> Department of Cellular and Physiological Sciences, University of British Columbia, Vancouver, British Columbia, Canada

### ARTICLE INFO

#### Article history:

Received 8 August 2018

Received in revised form 3 January 2019

Accepted 21 January 2019

Available online 2 February 2019

#### Keywords:

Pregnancy

Postpartum

Motherhood

Doublecortin

Cell proliferation

Iba-1

Proinflammatory cytokines

Anti-inflammatory cytokines

Neurogenesis

Middle age

Ki67

### ABSTRACT

The maternal brain displays considerable plasticity, and motherhood is associated with changes in affective and cognitive function. Motherhood can alter the trajectory of brain aging, including modifications to neuroplasticity and cognition. Here, we investigated the short- and long-term effects of motherhood on hippocampal neurogenesis, microglial density and morphology, and circulating cytokines, domains known to be altered with age and implicated in cognition and mood. Female rats were bred then euthanized during gestation or at various postpartum time points, culminating in middle age, and nulliparous rats served as age-matched controls. Hippocampal neurogenesis was significantly suppressed during gestation and the postpartum period. Interestingly, neurogenesis declined significantly in middle-aged nulliparous rats but increased in primiparous rats across the same period. Transient postpartum adaptations to the neuroimmune environment of the hippocampus were evidenced, as Iba-1-immunoreactive microglia assumed a deramified morphology followed by increased density. Intriguingly, aging-related changes in circulating cytokines were dependent on parity. These adaptations in neurogenic and immune processes may have ramifications for maternal mood and cognition across the peripartum period and beyond.

© 2019 Elsevier Inc. All rights reserved.

### 1. Introduction

Dramatic physiological adaptations occur during pregnancy and the postpartum period to ensure offspring development and survival (Dulac et al., 2014; Hall et al., 2011; Lain and Catalano, 2007; PrabhuDas et al., 2015; Rossant and Cross, 2001). The maternal brain exhibits substantial plasticity, including large-scale volumetric changes (Galea et al., 2000; Hoekzema et al., 2016; Oatridge et al., 2002), alterations in cellular architecture (Leuner and Gould, 2010; Pawluski and Galea, 2006), and hippocampal neurogenesis (Darnaudéry et al., 2007; Leuner et al., 2007; Pawluski and Galea, 2007). Although this capacity for plasticity is likely essential for the onset of a repertoire of maternal behaviors (Bridges, 2015), motherhood is also associated with changes in affective function (Bennett et al., 2004; Darcy et al., 2011; O'Hara, 2009),

hypothalamic-pituitary-adrenal (HPA) axis regulation (Slattery and Neumann, 2008), and cognition (Cuttler et al., 2011; De Groot et al., 2006; Galea et al., 2000; Kinsley et al., 1999; Pawluski et al., 2006). Interestingly, motherhood may improve the aging trajectory in terms of cognition (Colucci et al., 2006; Cui et al., 2014; Gatewood et al., 2005), neuroplasticity (Barha et al., 2015; Barha and Galea, 2011; Galea et al., 2018), and cellular aging (Barha et al., 2016), suggesting that the effects of motherhood on the brain may be long lasting. However, the mechanisms underlying alterations in the aging maternal brain are not known but may include modifications in neurogenic or immune processes, both of which were examined in the present study.

The hippocampus produces new neurons across the lifespan (Altman and Das, 1965; Boldrini et al., 2018; Eriksson et al., 1998), and these neurons play a role in certain aspects of learning and memory (Yau et al., 2015), mood regulation (Sahay and Hen, 2007), and the stress response (Snyder et al., 2011). Importantly, several studies found postpartum reductions in hippocampal cell proliferation (Darnaudéry et al., 2007; Leuner et al., 2007; Pawluski and Galea, 2007), cell survival (Pawluski and Galea, 2007), and the

\* Corresponding author at: Djavad Mowafaghian Centre for Brain Health, University of British Columbia, 2215 Wesbrook Mall, Vancouver, BC V6T 1Z3, Canada. Tel.: +1 (604) 822 6536; fax: +1 (604) 822 6923.

E-mail address: [liisa.galea@ubc.ca](mailto:liisa.galea@ubc.ca) (L.A.M. Galea).

density of immature neurons (Workman et al., 2015). Interestingly, motherhood may have contrasting effects on hippocampal neurogenesis with age, as studies have found increased neurogenesis in middle-aged primiparous and multiparous rats relative to age-matched nulliparous controls (Barha et al., 2015; Galea et al., 2018). This finding signifies that parity can have delayed pro-neurogenic effects; thus, the current study aimed to determine the timeline by which these changes may emerge.

Adaptations to the maternal immune system are well documented and necessary for the establishment and maintenance of pregnancy (Aghaeepour et al., 2017; Mor and Cardenas, 2010). In contrast, and despite the growing recognition of plasticity in the maternal brain, there is a paucity of research on potential neuro-immune adaptations with maternal experience. Few studies to date have examined microglia, the innate immune cells of the brain, in pregnant and postpartum rats (Haim et al., 2017; Posillico and Schwarz, 2016). Microglia alterations were found in several regions of the maternal brain and normalized by postpartum day 21 in all regions except the hippocampus (Haim et al., 2017). This indicates that changes in the neuroimmune environment of the maternal hippocampus may be longer lasting. The hippocampus undergoes considerable plasticity in the peripartum period (Galea et al., 2014), perhaps not surprisingly given its role in cognitive function (Sweatt, 2004) and mood regulation (Campbell and MacQueen, 2004). Neuroimmune processes are implicated in cognition (Lee et al., 2008; Parkhurst et al., 2013; vom Berg et al., 2012), stress (Hodes et al., 2014; Kreisel et al., 2014), and mood (Setiawan et al., 2015), raising the possibility that changes in the neuroimmune environment of the hippocampus may represent a substrate for motherhood-related changes in hippocampal function.

In the nonmaternal brain, immune processes have been implicated in the regulation of adult hippocampal neurogenesis, under basal and inflammatory conditions (reviewed in Sierra et al., 2014). For example, inflammation was first demonstrated to suppress neurogenesis by studies using systemic or intrahippocampal administration of the bacterial endotoxin lipopolysaccharide (Ekdahl et al., 2003; Monje, 2003). Microglia also maintain homeostasis in the healthy adult neurogenic niche via phagocytosis of apoptotic new cells (Sierra et al., 2010). In the maternal brain, 1 study found that alterations in T-cell activity accounted for at least some of the postpartum-associated reductions in neurogenesis (Rolls et al., 2008). To date, however, no studies have concurrently examined adaptations in microglia and neurogenesis in the maternal hippocampus, and therefore, the experiments reported here aimed to fill this gap.

Given the extensive cross-talk between the central nervous system and the immune system (Louveau et al., 2015), motherhood-related adaptations in the immune system can potentially drive plasticity in the brain. Reproductive immunology research has been primarily focused on aspects of immune function that affect fetal development and the success of pregnancy (PrabhuDas et al., 2015). Although many of the maternal immune adaptations normalize in the postpartum period (Groer et al., 2015), there is evidence indicating that maternal experience can leave a lasting footprint on the immune system (Barrat et al., 1997b, 1999, 1997a; Helle et al., 2004). For example, the risk of dying of infectious disease after the age of 65 years was increased in mothers of twins, compared with mothers of singletons (Helle et al., 2004). This effect may be related to reproductive effort and is perhaps indicative of accelerated immunosenescence (Helle et al., 2004). Although the long-term effects of parity on the immune system have received little attention in animal models, delayed senescence in certain aspects of immune function is evidenced in parous relative to nonparous mice (Barrat et al., 1997b,a, 1999). In tandem with neurogenic and neuroimmune markers, our present study aimed to assess whether maternal

experience can alter the circulating cytokine profile at various intervals after parturition, ending well after the reproductive event itself. The circulating cytokine profile is not only informative to the general inflammatory state, but may have ramifications for brain and behavior as peripheral cytokine signals propagate to the brain (Miller et al., 2014; Quan and Banks, 2007). Preclinical cytokine data may also be valuable for comparative purposes, as circulating cytokines levels are accessible biomarkers in clinical populations (Guerreiro et al., 2007).

In this study, we examined the short- and long-term effects of parity on microglia density and morphology and on neurogenesis in the hippocampus. These measures were examined across age and time since parturition, extending into middle age. To gain information about the peripheral inflammatory milieu, we quantified concentrations of various serum cytokines across the same time points. We expected parity to suppress hippocampal neurogenesis in the short term and to increase neurogenesis in middle age. At least in the short term, we expected parity to modify microglial density and morphology in the dentate gyrus. Finally, we expected alterations in the circulating cytokine profile during pregnancy and the early postpartum period and hypothesized that parity would modulate the age-related changes in the circulating cytokine milieu.

## 2. Materials and methods

### 2.1. Animals

Young adult female and male Sprague-Dawley rats were purchased from Charles River Laboratories (Montreal, Canada), weighing 200–250g. All rats arrived at our facility at the same time. Rats were maintained on a 12-hour light/dark cycle (lights on at 07:00 hours), in standard laboratory conditions ( $21 \pm 1$  °C;  $50 \pm 10\%$  humidity) and given *ad libitum* access to water and food (Purina Rat Chow). Female rats were initially pair-housed, and except for the breeding period, all rats were housed in female-only colony rooms. Males were used for breeding purposes only. Nulliparous rats were not housed in the same colony room as primiparous rats during breeding or when primiparous rats had active litters. To minimize potential environmental exposure differences between nulliparous and primiparous groups, primiparous rats were transferred to the nulliparous colony room on the day that their litters were weaned (postpartum day 21). All procedures were performed in accordance with ethical guidelines set by the Canadian Council on Animal Care and approved by the Animal Care Committee at the University of British Columbia.

### 2.2. Breeding procedure and experimental groups

Female rats were bred at approximately 7 months of age. At 18:00 hours daily, each pair of female cage mates was placed with 1 male. Vaginal lavage samples were obtained the following morning, between 08:00 and 09:00 hours, and examined for the presence of sperm cells. The detection of sperm cells indicated gestation day 1 (GD1), at which point the pregnant female was weighed and single-housed. Primigravid or primiparous rats (i.e., pregnant or mothering for the first time;  $n = 30$ ) were randomly assigned to one of the 6 groups ( $n = 5$  each) according to the timeline of euthanasia relative to gestation. This included one gestational group at gestation day 13 (GD13) and 4 postpartum groups: postpartum day 8 (PPD8), postpartum day 30 (PPD30), postpartum day 90 (PPD90), and postpartum day 180 (PPD180). Nulliparous rats (i.e., never pregnant;  $n = 30$ ) were randomly assigned to control groups ( $n = 5$  each) that were age-matched to each of the primiparous groups. Specifically, nulliparous rats at approximately 7, 7.5, 8, 10, and 13 months of age served as control groups for primiparous rats at

GD13, PPD8, PPD30, PPD90, and PPD180, respectively (experimental groups are detailed in Fig. 1). These time points were chosen to capture: mid-gestation (GD13), as a previous study found reductions in the survival of hippocampal neurons produced at this time (Rolls et al., 2008); an early postpartum time point (PPD8) that avoids the acute inflammatory state surrounding parturition (Catalano et al., 2010) and is associated with declines in cell proliferation (Leuner et al., 2007); a post-weaning, late postpartum time point (PPD30) shown to be associated with reduced neurogenesis (Workman et al., 2015); and finally, for a time-course analysis of the effects of parity on the aging trajectory, two further time points were selected leading to middle age (PPD90 and PPD180), as previous studies reported increased neurogenesis in middle-aged rats with previous maternal experience (Barha et al., 2015; Galea et al., 2018). To account for potential effects of social housing, nulliparous controls were single-housed at GD1 of their primiparous counterparts and until experimental endpoint. For all postpartum groups, litters were culled to include between 8 and 10 pups, with approximately 50% males and females. When the original sex ratio or litter size was not sufficient to achieve this, pups were cross-fostered between dams that had given birth the same day. Pups were weaned at PPD21 for all postpartum groups, except for the PPD8 group in which the dams remained with their litter until just before euthanasia. After weaning, primiparous rats remained single-housed until experimental endpoint. One rat from the nulliparous 10-month-old group was eliminated from the study due to a mammary gland tumor.

### 2.3. Perfusion and tissue collection

All perfusions were completed between 9:00 and 11:00 am. The rats were deeply anesthetized with an overdose of sodium pentobarbital (i.p.), and blood was collected via cardiac puncture. Brains were collected immediately after transcardial perfusion with 60 mL of cold 0.9% saline, followed by 120 mL of cold 4% paraformaldehyde. Brains were stored at 4 °C in 4% paraformaldehyde for 24 hours and then transferred into a 30% sucrose solution (in 0.1 M phosphate buffer) until sectioning. In the group euthanized during gestation, the uterus was dissected to confirm pregnancy. To obtain serum, blood samples were allowed to clot overnight at 4 °C and then centrifuged at 10g for 15 minutes, and serum aliquots were stored at –20 °C until processing.

### 2.4. Brain tissue processing and immunohistochemistry

Brains were sliced into 40 µm coronal sections using a Leica SM2000 R Microtome (Richmond Hill, Ontario, Canada). Sections were collected in series of 10 along the rostral-caudal axis of the hippocampus and then stored at –20 °C in a cryoprotectant

consisting of 30% ethylene glycol (Sigma-Aldrich, St. Louis, MO, USA) and 20% glycerol (Sigma-Aldrich) in 0.1 M phosphate buffer (pH 7.4). Sections were thoroughly rinsed (5 × 10 minutes) in phosphate-buffered saline (PBS) before staining to remove the cryoprotective medium. All immunohistochemical procedures were conducted on free-floating brain sections and on a rotator at room temperature unless otherwise noted.

#### 2.4.1. Doublecortin


Doublecortin (DCX) is a microtubule-associated protein expressed in immature neurons for 21 days after production in adult rats (Brown et al., 2003) and thus was used as a marker of adult hippocampal neurogenesis. Tissue was rinsed in 0.1 M PBS (pH 7.4; 5 × 10 minutes) between each of the following procedures. Tissue was treated with 0.3% hydrogen peroxide (H<sub>2</sub>O<sub>2</sub> in dH<sub>2</sub>O) for 30 minutes and then incubated for 24 hours at 4 °C in a primary antibody solution containing 1:1000 goat anti-doublecortin (Santa Cruz Biotechnology, Santa Cruz, CA, USA) in 3% normal rabbit serum and 0.4% Triton-X in 0.1 M PBS. Next, tissue was transferred to a secondary antibody solution consisting of 1:500 rabbit anti-goat (Vector Laboratories, Burlington, ON, Canada) in 0.1 M PBS for 24 hours at 4 °C. Finally, tissue was transferred to an avidin-biotin complex (Elite kit; 1:1000, Vector Laboratories) in PBS for 4 hours and then immunoreactants were visualized with a Nickel-enhanced DAB reaction (Vector Laboratories). Sections were mounted onto glass slides, allowed to dry, then dehydrated in increasing graded ethanol, defatted with xylene, and cover-slipped with Permount (Fisher Scientific).

#### 2.4.2. Ionized calcium-binding adaptor molecule-1

Iba-1 is a calcium-binding protein widely used as a microglial marker (Korzhevskii and Kirik, 2016). Tissue was rinsed in 0.1 M PBS (pH 7.4; 3 × 10 minutes) between each of the following procedures. Tissue was incubated in 0.3% hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>, in dH<sub>2</sub>O) for 25 minutes and then blocked with 10% normal goat serum (NGS) in 0.5% Triton-X in 0.1 M PBS. Tissue was then transferred to a primary antibody solution for 18 hours at 4 °C, consisting of 1:1000 anti-Iba-1 (Wako, Osaka, Japan) in 10% NGS and 0.4% Triton-X in 0.1 M PBS. Next, tissue was incubated in a secondary antibody solution for 1 hour, containing 1:500 biotinylated anti-rabbit (Vector Laboratories) in 2.5% NGS and 0.4% Triton X in PBS. Finally, tissue was transferred to an avidin-biotin complex (Elite kit; 1:50, Vector Laboratories) in 0.4% Triton-X in PBS for 1 hour, and immunoreactivity was visualized with a Nickel-enhanced DAB reaction (Vector Laboratories). Sections were mounted onto glass slides, allowed to dry, then counterstained with cresyl violet, dehydrated in a series of ethanol solutions of increasing concentrations, defatted with xylene, and cover-slipped with Permount (Fisher Scientific).

#### 2.4.3. Ki67

Ki67 is expressed during all active phases of the cell cycle but not during G<sub>0</sub> phase (Scholzen and Gerdes, 2000) and therefore was used as a marker of cell proliferation in the dentate gyrus. Tissue was rinsed with 0.1 M PBS (pH 7.4; 3 × 10 minutes) between each of the following procedures. Tissue was incubated in a primary antibody solution for 48 hours at 4 °C, consisting of 1:200 mouse anti-Ki67 (NCL-L-Ki67-MM1; Leica Biosystems, Newcastle, UK) in 3% normal donkey serum and 0.3% Triton-X in 0.1 M PBS. Next, tissue was incubated for 18 hours at 4 °C in a secondary antibody solution consisting of 1:200 donkey anti-mouse IgG, Alexa Fluor 555 (Molecular Probes, Eugene, Oregon, USA) in 3% normal donkey serum in 0.1 M PBS. Sections were counterstained with DAPI (2.5 minutes; 300 nM; ThermoFisher, Waltham, WA, USA), mounted onto glass slides, and cover-slipped with an anti-fade medium (2.5% polyvinyl alcohol-DABCO).



**Fig. 1.** Experimental groups. Primigravid or primiparous rats ( $n = 5/\text{group}$ ) were euthanized at gestation day 13 (GD13), postpartum day 8 (PPD8), postpartum day 30 (PPD30), postpartum day 90 (PPD90), or postpartum day 180 (PPD180). Nulliparous rats ( $n = 5/\text{group}$ ) were age-matched to their primiparous counterparts and euthanized along the same timeline, depicted as approximate age in months. Arrows indicate euthanasia day. Abbreviations: GD, gestation day; PPD, postpartum day.

## 2.5. Microscopy, cell quantification, and morphological analyses

An investigator blinded to experimental conditions quantified DCX-, Iba-1-, and Ki67-immunoreactive cells and analyzed cell morphology. See Figs. 4B, 6D–F, and 7 for representative photomicrographs.

### 2.5.1. Ionized calcium-binding adaptor molecule-1

Under a 40× objective on a Nikon E600 microscope, an exhaustive quantification of Iba-1-IR cells was completed in 4 hippocampal slices from each animal, as we have done previously (Mahmoud et al., 2016a). This included 2 dorsal and 2 ventral sections, with approximate Bregma coordinates of –3.12, –3.48, 6.00, and –6.36. Iba-1-IR cells were quantified in the dentate gyrus, specifically in the granule cell layer (GCL), the subgranular zone (SGZ), and within an approximately 50 μm band of the molecular layer (ML).

Under baseline conditions, microglia typically display ramified morphology, characterized by highly branched and long processes that continuously survey their environment (Nimmerjahn et al., 2005). Under inflammatory states, microglial processes typically retract, and the cell can take on an amoeboid morphology, in which the cell becomes enlarged and void of processes; thus, microglial morphology is widely used as a proxy measure of functional state (Karperien et al., 2013). Here, Iba-1-IR cell morphology was analyzed using NIS Elements Basic Research software (Nikon) under a Nikon E600 microscope. Using the measure feature, soma size, in addition to cell process length and number (Karperien et al., 2013), was measured live at 40× for every cell within a 23,672.24 μm<sup>2</sup> region of interest (ROI), with 3 ROIs each for the dorsal and ventral hippocampus, using 6 different sections. Furthermore, no more than one ROI was taken from an individual tissue slice, and ROIs were defined in 3 consistent locations in the GCL for both the dorsal and ventral hippocampus within each animal. Cells were defined by the presence of an Iba-1-IR cell body within the ROI, and this definition did not necessitate the presence of cell processes, by that ensuring the inclusion of any cells with amoeboid morphology. The average process length per cell was calculated using the total length and number of processes for each cell and subsequently an average process length was calculated for each animal. Both primary (extending directly from the cell body) and secondary processes were taken into account in the analyses. There were no significant differences between groups in the number of Iba-1-IR cells that fell within the ROIs and were used for morphological analyses (see Table 1).

In an additional analysis of microglial morphology, Iba-1-IR cells were categorized into 1 of 3 morphological phenotypes, adapted from other methods (Haim et al., 2017; Schwarz et al., 2012). Specifically, cells were categorized as (1) ramified microglia (highly

branched, with longer processes), stout microglia (few, shorter processes), or (3) amoeboid microglia (no processes). This was performed under a 40× objective, in 4 hippocampal slices per animal, and involved a classification of all Iba-1-IR cells in the GCL, SGZ, and an approximately 50 μm band of the ML. For each animal, the percentage of cells of each morphological phenotype was calculated.

### 2.5.2. Doublecortin

Under a 100× objective on an Olympus CX22LED brightfield microscope, DCX-IR cells in the GCL were exhaustively counted in every 10th section of the hippocampus along the rostral-caudal axis. Thus, raw counts were multiplied by a factor of 10 to obtain an estimate of the total number of DCX-IR cells in the hippocampus. Previous work indicates that certain conditions or experiences, including reproductive experience (Workman et al., 2015), can alter the rate at which newly produced hippocampal neurons mature (Overstreet-Wadiche et al., 2006), and this can have functional consequences as changes in the maturational timeline can alter the rate of functional integration of these new neurons into existing hippocampal networks. Thus, to examine whether pregnancy and motherhood can alter the rate at which newly produced hippocampal neurons mature, we investigated the maturational stages of DCX-IR cells. Specifically, using the 100× objective on an Olympus CX22LED brightfield microscope, 50 DCX-IR cells (25 dorsal GCL and 25 ventral GCL; each taken from 3 slices) were randomly selected for each animal for maturational staging. Cells were categorized into 1 of 3 maturational stages, based on previously established criteria (Plümpe et al., 2006): (1) proliferative (no process or short process), (2) intermediate (medium process with no branching), or (3) postmitotic (long processes with branching in the GCL and ML). DCX-immunolabeled sections were also used to measure GCL areas in every 10th section of the hippocampus, using images taken at 4x and the ImageJ software. Volume was estimated using these areas and following Cavalieri's principle.

### 2.5.3. Ki67

Under a 100× objective on an Olympus CX22LED microscope equipped with epifluorescence, Ki67-IR cells in the GCL and SGZ of the DG were exhaustively counted in every 10th section of the hippocampus along the rostral-caudal axis. Raw counts were multiplied by a factor of 10 to obtain an estimate of the total number of Ki67-IR cells in the DG.

## 2.6. Serum cytokine and hormone quantification

A multiplex electrochemiluminescence immunoassay kit (V-PLEX Proinflammatory Panel 2, Rat) was purchased from Meso Scale Discovery (Rockville, MD) and used according to manufacturer instructions to measure serum cytokine concentrations. The antibody precoated plates allowed for the simultaneous quantification of the following cytokines: interferon-gamma (IFN-γ), interleukin-1-beta (IL-1β), interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-6 (IL-6), chemokine (C-X-C motif) ligand 1 (CXCL1), interleukin-10 (IL-10), interleukin-13 (IL-13), and tumor necrosis factor-alpha (TNF-α). Samples were run in duplicates, plates were read with a Sector Imager 2400 (Meso Scale Discovery), and data were analyzed using the Discovery Workbench 4.0 software (Meso Scale Discovery). The assays' lower limits of detection (LLOD), which varied between analytes and plates (2 plates total), were as follows (pg/mL): IFN-γ: 0.163–0.266; IL-10: 0.233–0.313; IL-13: 0.78–2.7; IL-1β: 1.48–1.62; IL-4: 0.179–0.298; IL-5: 7.64–9.8; IL-6: 2.48–2.49; CXCL1: 0.085–0.164; and TNF-α: 0.156–0.186. Any values below the LLOD were assigned 0 pg/mL, as we have done previously (Bodnar et al., 2017). All samples were within the detection range for TNF-α,

**Table 1**

Mean number of Iba-1-IR cells used for process length analyses ± standard error of the mean

Group	Number of Iba-1-IR cells analyzed
Nulliparous–7 mo	23.40 ± 1.03
Nulliparous–7.5 mo	21.00 ± 2.04
Nulliparous–8 mo	20.80 ± 2.13
Nulliparous–10 mo	21.20 ± 1.16
Nulliparous–13 mo	22.25 ± 2.32
Primiparous–GD13	23.67 ± 1.67
Primiparous–PPD8	22.00 ± 0.77
Primiparous–PPD30	23.60 ± 1.47
Primiparous–PPD90	20.60 ± 1.03
Primiparous–PPD180	23.20 ± 0.37

Key: GD, gestation day; PPD, postpartum day; mo, months. There were no significant differences between groups.

CXCL1, and IL-10. One sample fell below the LLOD for each of IFN- $\gamma$ , IL-4, and IL-13. For 3 cytokines, a number of samples fell below the LLOD ( $n = 12$  for IL-6,  $n = 17$  for IL-1 $\beta$ , and  $n = 36$  for IL-5). This panel was chosen as it includes a broad range of cytokines, some traditionally considered proinflammatory (IL-1 $\beta$ , IFN- $\gamma$ , TNF- $\alpha$ ), anti-inflammatory (IL-4, IL-10), and pleiotropic (IL-6), in addition to the chemokine CXCL1, which is important for neutrophil recruitment. Therefore, combined, these markers provide a comprehensive view of the inflammatory milieu.

Steroid hormones modulate neuroplasticity (Mahmoud et al., 2016b) and interact with the immune system (Schumacher et al., 2014). Furthermore, circulating concentrations increase dramatically during pregnancy, drop abruptly at parturition, and diminish in aging females. To investigate the potential mediating roles of steroid hormones, we measured 17 $\beta$ -estradiol, the most potent of the estrogens and the most abundant in premenopausal women and younger female rats (Rannevik et al., 1986). Serum 17 $\beta$ -estradiol concentrations were quantified in samples taken at perfusion, using an ultrasensitive estradiol radioimmunoassay (Beckman Coulter, Prague, Czech Republic) and in accordance with manufacturer's instructions, with all samples run in duplicates. The assay sensitivity is 2.2pg/mL, and the average inter- and intra-assay coefficients of variation were <15%.


### 2.7. Statistical analyses

Statistical analyses were performed using Statistica software (Tulsa, OK). All dependent variables were subjected to the Kolmogorov-Smirnov test for normality and Bartlett's test for heterogeneity of variance. TNF- $\alpha$ , IL-1 $\beta$ , IL-5, and IL-6 data were not normally distributed and therefore Box-Cox transformed before analyses. There were few violations to the heterogeneity of variance assumption; however, these were corrected after Box-Cox transformation. Neural measures (DCX- and Ki67-IR cell number, Iba-1-IR density, length and number of processes) and serum measures (17 $\beta$ -estradiol, IFN- $\gamma$ , IL-1 $\beta$ , IL-4, IL-5, IL-6, CXCL1, IL-10, IL-13, and TNF- $\alpha$ ) were each analyzed using factorial analysis of variance (ANOVA), with time (GD13/7 months, PPD8/7.5 months, PPD30/8 months, PPD90/10 months, PPD180/13 months) and reproductive status (primigravid/parous, nulliparous) as the between-subject factors. Post hoc analyses used Fisher's least significant difference. *A priori*, we expected parity to modulate the age-related changes in cytokine levels and density/morphology of Iba-1-IR cells. Any *a priori* comparisons were subjected to a Bonferroni correction. Pearson's correlations were performed on dependent variables of interest, and Fisher's z-transformation of correlation coefficients was used to assess the significance of the difference between correlations in nulliparous and primiparous groups. Serum 17 $\beta$ -estradiol concentrations were used as a covariate in ANOVAs, as circulating concentrations can influence neuroplastic and immune measures (Mahmoud et al., 2016b; Schumacher et al., 2014); ANOVAs are reported with the covariate in instances where there was a significant main effect of the covariate. Finally, as an exploratory approach, and to complement our findings from ANOVA analyses, we ran a principal component analysis (PCA) on the cytokine data, with the purpose of deriving information about the amount of variance accounted for by potential cytokine networks within the data set.

## 3. Results

### 3.1. Serum 17 $\beta$ -estradiol was reduced in the early postpartum and increased in the late postpartum

As expected, 17 $\beta$ -estradiol was reduced in the early postpartum period at PPD8, relative to age-matched nulliparous rats ( $p = 0.015$ ;



**Fig. 2.** Serum concentrations of 17 $\beta$ -estradiol at perfusion in primiparous and nulliparous rats. The x-axis represents time relative to gestation and parturition in primiparous rats and approximate age in months. 17 $\beta$ -estradiol was significantly reduced in the early postpartum period at PPD8 but significantly increased in the late postpartum period at PPD30 and PPD90. \* indicates  $p = 0.015$ , significantly lower than age-matched nulliparous rats; # indicates  $p < 0.03$ , significantly higher than 8-month-old nulliparous rats and primigravid/parous rats at GD13, PPD8, and PPD180; + indicates  $p < 0.03$ , significantly higher than primigravid/parous rats at GD13, PPD8, and PPD180. Data are represented in mean values  $\pm$  standard error of the mean (SEM). Abbreviations: GD, gestation day; PPD, postpartum day.

reproductive status by time interaction:  $F(4, 36) = 4.45$ ,  $p = 0.005$ ; Fig. 2). Furthermore, primiparous rats at PPD30 had higher serum 17 $\beta$ -estradiol concentrations than age-matched nulliparous rats ( $p < 0.008$ ; Fig. 2). Finally, 17 $\beta$ -estradiol concentrations at PPD30 and PPD90 were significantly higher than all other primiparous groups ( $p$ 's < 0.03), but there were no significant differences between any of the nulliparous groups (all  $p$ 's > 0.19). There was also a significant main effect of time ( $p < 0.01$ ) but not of reproductive status ( $p = 0.41$ ).


### 3.2. Parity and age had no significant effect on GCL volume

GCL volume was not significantly affected by age, parity, or age by parity interaction (all  $p$ 's > 0.48); thus, all further analyses were performed on the number, rather than density, of Ki67- and DCX-IR cells.

### 3.3. The number of DCX-IR cells was significantly reduced during pregnancy and the postpartum period, and declined in middle age in nulliparous rats only

The number of DCX-IR cells was significantly reduced in primigravid and primiparous rats relative to age-matched nulliparous controls at GD13, PPD8, and PPD30 ( $p$ 's < 0.001; significant time by reproductive status interaction;  $F(4, 38) = 6.7213$ ,  $p < 0.001$ ; Fig. 3). There were also significant main effects of time and reproductive status (all  $p$ 's < 0.001).

There was a significant age-related decline in the number of DCX-IR cells in nulliparous rats, such that each nulliparous group had a significantly lower number of DCX-IR cells than all previous nulliparous age groups (all  $p$ 's < 0.014), with the exception of a non-significant decline from 10 to 13 months ( $p = 0.11$ ; significant time by reproductive status interaction;  $F(4, 38) = 6.7213$ ,  $p < 0.001$ ; Fig. 3). There was also a significant decline in DCX-IR cell number in primiparous rats at PPD30 relative to GD13 and PPD8 ( $p$ 's < 0.017; Fig. 3). However, the trajectory of age-related change in DCX-IR cell number was significantly altered by parity in middle age; unlike nulliparous rats, primiparous groups had no significant difference in the number of DCX-IR cells from 8 to 10 months of age ( $p = 0.10$ ).




**Fig. 3.** Estimated total number of doublecortin (DCX)-immunoreactive (IR) cells in the granule cell layer of primiparous and nulliparous rats across 7–13 months of age. The x-axis represents time relative to gestation and parturition in primiparous groups and approximate age in months. DCX-IR cell number was significantly reduced in primiparous rats in mid-gestation and until postpartum day 30. Between 8 and 13 months of age, DCX-IR cell number significantly declined in nulliparous rats but significantly increased in primiparous rats. Data are represented in mean values  $\pm$  standard error of the mean (SEM). \*\*\* indicates  $p < 0.001$ , significantly different from age-matched primiparous group. # denotes  $p < 0.001$ , significantly different from 8-month-old nulliparous group. † indicates  $p = 0.044$ , significantly different from primiparous rats at PPD30. Abbreviations: GCL, granule cell layer, DCX-IR, doublecortin-immunoreactive; GD, gestations day, PPD, postpartum day.

Based on previous findings (Barha et al., 2015; Galea et al., 2018), we expected parity to increase neurogenesis in middle age. Indeed, we found a significant increase in DCX-IR cells from 8 to 13 months of age in primiparous rats ( $p = 0.044$ ; 1-tailed).

#### 3.4. Parity and age had no significant effect on the maturational stage of DCX-IR cells

Regardless of age and reproductive status, the percentage of proliferative DCX-IR cells was significantly higher than that of intermediate DCX-IR cells ( $p < 0.001$ ), and the percentage of postmitotic DCX-IR cells was significantly higher than that of intermediate and proliferative DCX-IR cells ( $p$ 's  $< 0.001$ ; significant main effect of DCX maturational stage,  $F(2, 76) = 107.17$ ,  $p < 0.001$ ; Table 2). There were no other significant main effects and no significant interactions (all  $p$ 's  $> 0.14$ ).



**Table 2**

Mean percentage of proliferative, intermediate, and postmitotic doublecortin-immunoreactive cells in the granule cell layer  $\pm$  standard error of the mean

Group	% Proliferative	% Intermediate	% postmitotic
Nulliparous–7 mo	37.60 $\pm$ 1.60	17.20 $\pm$ 2.15	45.20 $\pm$ 2.65
Nulliparous–7.5 mo	26.40 $\pm$ 2.86	18.80 $\pm$ 1.85	54.80 $\pm$ 2.87
Nulliparous–8 mo	35.60 $\pm$ 4.71	20.80 $\pm$ 1.02	46.40 $\pm$ 4.45
Nulliparous–10 mo	29.60 $\pm$ 6.52	21.60 $\pm$ 2.64	48.80 $\pm$ 4.84
Nulliparous–13 mo	25.33 $\pm$ 8.74	20.00 $\pm$ 1.15	54.67 $\pm$ 8.97
Primiparous–GD13	32.00 $\pm$ 6.23	18.40 $\pm$ 2.93	49.60 $\pm$ 5.84
Primiparous–PPD8	28.40 $\pm$ 5.31	16.00 $\pm$ 1.10	56.40 $\pm$ 6.71
Primiparous–PPD30	27.60 $\pm$ 4.26	17.20 $\pm$ 0.80	55.20 $\pm$ 4.88
Primiparous–PPD90	23.60 $\pm$ 3.19	16.80 $\pm$ 3.01	59.60 $\pm$ 4.53
Primiparous–PPD180	28.40 $\pm$ 5.71	16.00 $\pm$ 2.45	56.00 $\pm$ 6.23

Key: GD, gestation day; PPD, postpartum day; DCX-IR, doublecortin-immunoreactive; mo, months.


Parity and age did not significantly affect the maturational stage of DCX-IR cells.

#### 3.5. Ki67-IR cells declined in mid-gestation and the early postpartum period in primiparous rats and with age regardless of reproductive status

Parity significantly reduced the number of Ki67-IR cells in the GCL ( $F(1, 38) = 9.2681$ ,  $p = 0.004$ ; significant main effect of reproductive status; Fig. 4A). Regardless of reproductive status, Ki67-IR cells declined significantly with age ( $F(4, 38) = 16.505$ ,  $p < 0.001$ ; main effect of time; Fig. 4A), in which there was a significant difference between all age groups ( $p$ 's  $< 0.005$ ), with the exception of non-significant differences from 7 to 7.5 months, 8–10 months, and 10–13 months ( $p$ 's  $> 0.05$ ). Although there was no significant age by parity interaction ( $p = 0.13$ ), *a priori* we expected a decline in cell proliferation in the early postpartum period based on previous work (Leuner et al., 2007). Planned comparisons revealed a significant decline in Ki67-IR cells on PPD8 ( $p = 0.013$ ; 1-tailed), in addition to a significant decline on GD13 ( $p = 0.004$ ), relative to age-matched nulliparous controls in both instances (Fig. 4A).

#### 3.6. Density of Iba-1-IR cells increased in the late postpartum period and fluctuated significantly with age in primiparous but not in nulliparous rats

Because we quantified Iba-1-IR cells in 4 sections per animal, the density of IR cells was analyzed, as we have done previously (Mahmoud et al., 2016a). Age significantly affected the density of Iba-1-IR cells in the dentate gyrus, with higher densities at 8 and



**Fig. 4.** (A) Estimated total number of Ki67-immunoreactive (IR) cells in the granule cell layer and subgranular zone of primiparous and nulliparous rats across 7–13 months of age. The x-axis represents time relative to gestation and parturition in primiparous groups and approximate age in months. Ki67-IR cell number was significantly reduced in primiparous rats in mid-gestation and the early postpartum period and declined with age regardless of reproductive status. Data are represented in mean values  $\pm$  standard error of the mean (SEM). \* indicates  $p < 0.014$ , significantly different from age-matched primiparous group. # denotes  $p = 0.004$ , significant main effect of reproductive status. (B) Representative photomicrograph of the dentate gyrus, showing Ki67-IR cells (pink), counterstained with DAPI (blue). Abbreviations: Ki67-IR, Ki67-immunoreactive; GD, gestations day; PPD, postpartum day.

13 months relative to 7.5 and 10 months of age ( $p$ 's < 0.02; significant main effect of time,  $F(4, 39) = 3.43, p = 0.017$ ). We expected alterations in the density of Iba-1-IR cells in the postpartum period due to previous findings (Haim et al., 2017). *A priori* comparisons show that the density of Iba-1-IR cells fluctuated significantly across time in primiparous rats; density was increased in the late postpartum period at PPD30 relative to PPD8 ( $p = 0.002$ ) and in middle-aged rats at PPD180 relative to PPD8 ( $p < 0.003$ ; Fig. 5A). On the other hand, density did not significantly change across age in nulliparous rats (all  $p$ 's > 0.17; Fig. 5A). There were no significant differences between nulliparous and primiparous groups across age (all  $p$ 's > 0.3), except for a trend for higher density at PPD30 relative to age-matched nulliparous controls ( $p = 0.09$ ; Fig. 5A). There was no significant main effect of reproductive status and no significant interaction ( $p$ 's > 0.28).


### 3.7. Average length and number of Iba-1-IR cell processes changed significantly with age and reproductive status

Iba-1-IR cells displayed significantly shorter processes in the early postpartum period at PPD8 relative to nulliparous controls ( $p = 0.005$ ; Fig. 5B) and to primiparous rats at GD13 ( $p < 0.002$ ).


There was a decline in average length of Iba-1-IR cell processes at 10 months of age, relative to 7 months in primiparous rats and to 7, 7.5, and 8 months in nulliparous rats (all  $p$ 's < 0.027; significant time by reproductive status interaction  $F(4, 38) = 4.67, p = 0.004$ ; Fig. 5B). However, there was no further significant change in average process length between 10 and 13 months of age, regardless of reproductive status ( $p$ 's > 0.23). There was also a significant main effect of the covariate serum  $17\beta$ -estradiol ( $p = 0.012$ ), a trend toward a significant main effect of time ( $p = 0.067$ ), but no significant main effect of reproductive status ( $p = 0.42$ ).

The average number of Iba-1-IR cell processes increased significantly with age in nulliparous rats, where significantly more processes were found in 13- compared with 8-month-old rats ( $p < 0.002$ ), but missed significance compared to 7.5-month-old rats ( $p = 0.027$ ; planned comparisons; Fig. 5C). On the other hand, there were no significant differences in the average number of cell processes between primiparous rats across age (all  $p$ 's > 0.06). There were trends toward significant main effects of time ( $p = 0.054$ ) and reproductive status ( $p = 0.056$ ) but no significant time by reproductive status interaction ( $p = 0.11$ ).

There was a trend for primiparity to decrease the soma size of Iba-1-IR cells relative to nulliparity ( $F(1, 38) = 5.26, p = 0.092$ ; main



**Fig. 5.** Density and morphology of Iba-1-immunoreactive cells in the dentate gyrus of primiparous and nulliparous rats. The x-axis represents time relative to gestation and parturition in primiparous rats and approximate age in months. (A) Iba-1-IR cell density was stable across age in nulliparous rats but fluctuated significantly in primiparous rats, with increased density at PPD30 and PPD180, relative to PPD8. \* indicates  $p < 0.003$ , significantly different from primiparous group at PPD8. (B) Average length of Iba-1-immunoreactive cell processes was reduced at PPD8 and decreased with age regardless of reproductive status. \* denotes  $p < 0.006$ , significantly different from age-matched nulliparous controls and from primiparous group at GD13. (C) Average number of Iba-1-immunoreactive cell processes increased significantly with age in nulliparous but not primiparous rats. \* indicates  $p < 0.002$ , significantly different from 8-month-old nulliparous group. (D) There was a trend toward significance for parity to reduce soma area of Iba-1-IR cells. \* indicates  $p = 0.092$ , main effect of reproductive status. Data are represented in mean values  $\pm$  SEM. Abbreviations: GD, gestation day; PPD, postpartum day; Iba-1-IR, Iba-1-immunoreactive.




**Fig. 6.** Percentage of Iba-1-immunoreactive cells of ramified (A), stout (B), and ameboid (C) morphology in the dentate gyrus. The x-axis represents time relative to gestation and parturition in primiparous rats and approximate age in months. (A) The percentage of ramified Iba-1-IR cells was transiently reduced in the early postpartum period. \* indicates  $p < 0.02$ , PPD8 significantly lower than 7.5-month-old nulliparous rats, primigravid rats at GD13, and primiparous rats at PPD30. (B) The percentage of stout Iba-1-IR cells was transiently increased in the early postpartum period. \* indicates  $p < 0.007$ , PPD8 significantly higher than 7.5-month-old nulliparous rats and primigravid rats at GD13. (C) The percentage of ameboid Iba-1-IR cells did not differ between groups. Representative photomicrographs of Iba-1-IR cells of ramified (D), stout (E), and ameboid (F) morphology captured at 60 $\times$ . Data are represented in mean values  $\pm$  SEM. Abbreviations: GD, gestation day; PPD, postpartum day; Iba-1-IR, Iba-1-immunoreactive.


effect of reproductive status; Fig. 5D). Although there was no significant age by parity interaction ( $p = 0.80$ ), the effect of primiparity to reduce soma size appears to be driven by the PPD30, 90, and 180 groups. In addition, there was a significant main effect of time ( $F(4, 38) = 5.28, p < 0.001$ ).

**3.8. The percentage of Iba-1-IR cells of stout morphology increased, and that of ramified morphology decreased, in the early postpartum period**

There was a significant Iba-1 morphology by time by reproductive status interaction ( $F(8, 74) = 2.066, p = 0.05$ ; Fig.6), in



**Fig. 7.** Representative photomicrographs of the granule cell layer in the dentate gyrus. (A) Doublecortin-immunoreactive cells in an 8-month-old nulliparous rat. (B) Doublecortin-immunoreactive cells in an 8-month-old primiparous rat on postpartum day 30. (C) Iba-1-immunoreactive cells in a 7.5-month-old nulliparous rat. (D) Iba-1-immunoreactive cells in a 7.5-month-old primiparous rat on postpartum day 8.



**Fig. 8.** Serum levels of interferon- $\gamma$  (A) and interleukin-10 (B) in primiparous and nulliparous rats. The x-axis represents time relative to gestation and parturition in primiparous groups and approximate age in months. There was a significant aging-related increase in IFN- $\gamma$  (A) and IL-10 (B) in nulliparous but not primiparous rats. \* indicates  $p < 0.004$ , significantly different from all other nulliparous groups. # indicates  $p = 0.023$ , significantly different from 13-month-old primiparous rats. Data are represented in mean values  $\pm$  SEM. Abbreviations: IFN- $\gamma$ , interferon gamma; IL-10, interleukin 10; GD, gestation day; PPD, postpartum day.

which the percentage of ramified Iba-1-IR cells was reduced in primiparous rats at PPD8 relative to age-matched nulliparous controls, primigravid rats at GD13, and primiparous rats at PPD30 (all  $p$ 's  $< 0.02$ ; Fig. 6A). In addition, the percentage of stout Iba-1-IR cells was significantly increased in primiparous rats at PPD8 relative to age-matched nulliparous controls and primigravid rats at GD13 ( $p$ 's  $< 0.007$ ; Fig. 6B). There were no significant differences in the percentage of amoeboid Iba-1-IR cells between any of the groups ( $p$ 's  $> 0.05$ ; Fig. 6C). There was also a significant main effect of Iba-1 morphology ( $p < 0.001$ ), a trend for a significant Iba-1 morphology by time interactions ( $p = 0.087$ ), but no other significant main effects or interactions ( $p$ 's  $> 0.49$ ).

### 3.9. Serum IFN- $\gamma$ and IL-10 showed an age-related increase in nulliparous but not in primiparous rats


There was a significant aging-related increase in IFN- $\gamma$  in nulliparous rats, in which 13-month-old nulliparous rats had significantly higher IFN- $\gamma$  levels than all other nulliparous groups (all  $p$ 's  $< 0.004$ ; planned comparisons; Fig. 8A). No significant differences were found in IFN- $\gamma$  levels between any primiparous

groups (all  $p$ 's  $> 0.04$ , non-significant due to Bonferroni correction). Furthermore, at 13 months, IFN- $\gamma$  levels were higher in nulliparous relative to primiparous rats ( $p = 0.023$ ; Fig. 8A). There was also a significant main effect of time ( $p < 0.001$ ), but not reproductive status ( $p = 0.14$ ), nor an interaction ( $p = 0.325$ ).

Similarly, serum IL-10 increased significantly with age in nulliparous but not primiparous rats; significantly higher levels of IL-10 were detected in 13-month-old nulliparous rats relative to all other nulliparous groups ( $p < 0.002$ ; Fig. 8B), but no significant differences were found between any of the primiparous groups (all  $p$ 's  $> 0.05$ ; planned comparisons; Fig. 8B). There was also a significant main effect of time ( $p < 0.003$ ), but not reproductive status ( $p = 0.11$ ) nor an interaction ( $p = 0.45$ ).

### 3.10. Serum IL-4 was transiently increased in the early postpartum and then persistently reduced by parity

Nulliparous rats had higher levels of IL-4 than primiparous rats, regardless of time point (main effect of reproductive status:  $F(1, 38) = 7.63$ ,  $p < 0.009$ , Fig. 9A). Regardless of



**Fig. 9.** Serum levels of interleukin-4 (A) and interleukin-5 (B) in primiparous and nulliparous rats. The x-axis represents time relative to gestation and parturition in primiparous groups and approximate age in months. (A) Serum IL-4 levels were transiently increased in primiparous rats at PPD8 but persistently suppressed by parity thereafter. # indicates  $p < 0.03$ , trend toward significance relative to GD13 and PPD30. \* indicates  $p < 0.009$ , main effect of primiparity to reduce IL-4 levels. (B) Serum IL-5 was transiently blunted during gestation, and a significant age-related decline in serum IL-5 was found only in nulliparous rats. \* indicates  $p = 0.006$ , significantly higher than primigravid rats at GD13. # indicates  $p < 0.02$ , significantly higher than all nulliparous groups between 8 and 13 months of age. Abbreviations: IL-4, interleukin-4; GD, gestation day; PPD, postpartum day.

reproductive status, there was an age-related increase in serum IL-4, with significantly elevated levels at 13 months compared to all groups at 8 months of age and younger (all  $p$ 's < 0.04; main effect of time:  $F(4, 38) = 4.29, p < 0.006$ ). There was no significant reproductive status by time interaction ( $F(4, 38) = 1.73, p = 0.16$ ), but *a priori*, we expected cytokine levels to be altered in the early postpartum period in primiparous rats. Indeed, primiparous rats showed a trend for a transient increase in serum IL-4 in the early postpartum period, with higher levels at PPD8 relative to GD13 ( $p = 0.029$ ) and PPD30 ( $p = 0.027$ ; *a priori* comparisons, missing significance with correction; Fig. 9A). In age-matched nulliparous control groups, serum IL-4 was not significantly different in 7.5-month-old relative to 7- or 8-month-old rats ( $p$ 's > 0.45).

### 3.11. Serum IL-5 was transiently reduced during gestation and showed an age-related decline in nulliparous but not primiparous rats


Serum IL-5 levels were significantly reduced in gestation (GD13), relative to age-matched nulliparous controls ( $p = 0.006$ ; time by reproductive status interaction,  $F(4, 37) = 3.31, p = 0.020$ ; Fig. 9B). Furthermore, IL-5 levels declined with age in nulliparous animals, as higher levels were detected at 7 months relative to 8, 10, and 13 months (all  $p$ 's < 0.02; Fig. 9B). Although non-significant, IL-5 levels increased in primiparous animals with age, suggesting a reversed pattern of age-related changes in IL-5 compared with nulliparous rats. There were no significant main effects of time or reproductive status (all  $p$ 's > 0.1).

### 3.12. Serum IL-13, IL6, CXCL1, IL-1 $\beta$ , and TNF- $\alpha$ were not significantly altered by parity or age

Regardless of reproductive status, there was a trend toward significance for a main effect of time to affect IL-13 ( $F(4, 37) = 2.46, p = 0.06$ ; Fig. 10A), where IL-13 levels were elevated at 13 months compared with 8 months ( $p < 0.002$ ; planned comparisons). There were no significant main effects of reproductive status or 17 $\beta$ -estradiol (covariate) and no time by reproductive status interaction (all  $p$ 's > 0.4). There were no significant main effects of reproductive status or age or an interaction for serum concentrations of IL-6 (all  $p$ 's > 0.21, Fig. 10B), CXCL1 (all  $p$ 's > 0.40 Fig. 10C), or IL-1 $\beta$  (all  $p$ 's > 0.32, Fig. 10D). There were trends toward significance for a main effect of parity to increase TNF- $\alpha$  ( $p = 0.064$ ) and a main effect of time ( $p = 0.061$ ) for TNF- $\alpha$  to decline with age but no significant reproductive status by time interaction ( $p = 0.72$ , Fig. 10E).

### 3.13. Principal component analysis of serum cytokines

The model generated 4 principal components, which accounted for 82.6% of the variance within the data set, with the first principal component explaining 40.4% of the variance, the second 18.2%, the third 12.6%, and the fourth 11.4%. Interestingly, IFN- $\gamma$ , IL-10, IL-13, and IL-4 loaded heavily onto principal component 1 (PC1; see Table 3). These same cytokines also showed the most robust alterations with parity and age with ANOVA, therefore the principal component analysis ultimately verified our individual ANOVA analyses. Subsequently, we analyzed PC1 scores using ANOVA, which revealed a main effect of time ( $F(4, 37) = 3.78, p = 0.011$ ; Fig. 10F), with higher scores at 13 months of age relative to all other age groups (all  $p$ 's < 0.033). Planned comparisons reveal a more robust age-related increase in PC1 scores in nulliparous rats, with higher



**Fig. 10.** Serum concentrations of interleukin-13 (A), interleukin-6 (B), CXCL1 (C), interleukin- $\beta$  (D), and tumor necrosis factor alpha- $\alpha$  (E) in primiparous and nulliparous rats. The x-axis represents time relative to gestation and parturition in primiparous rats and approximate age in months. (A) There was a trend toward significance for serum IL-13 to increase with age regardless of reproductive status. \* indicates  $p < 0.002$ , relative to 8-month-old rats. Reproductive status and age had no significant effects on serum levels of interleukin-6 (B), CXCL1 (C), and interleukin- $\beta$  (D). (E) There were trends toward significance for parity to increase ( $p = 0.064$ ) and for age to decrease ( $p = 0.061$ ), TNF- $\alpha$  concentrations. (F) Principal component 1 scores in primiparous and nulliparous rats, \* indicates  $p$ 's < 0.007, significantly higher PC1 scores in 13-month-old relative to 7- and 8-month-old nulliparous rats. Data are represented in mean values  $\pm$  SEM. Data are represented in mean values  $\pm$  SEM. Abbreviations: GD, gestation day; PPD, postpartum day; IL-4, interleukin-4; IL-5, interleukin-5; IL-13, interleukin-13; GD, gestation day; PPD, postpartum day; IL-4, interleukin-4; IL-5, interleukin-5; IL-13, interleukin-13; GD, gestation day; PPD, postpartum day; IL-4, interleukin-4; IL-5, interleukin-5.

**Table 3**  
Principal component analysis loading table

Cytokine	PC1	PC2	PC3	PC4
IFN- $\gamma$	<b>0.87</b>	-0.28	0.01	-0.04
IL-10	<b>0.91</b>	-0.15	-0.12	0.027
IL-13	<b>0.87</b>	-0.18	0.27	0.12
IL-1 $\beta$	-0.06	0.25	<b>0.78</b>	-0.46
IL-4	<b>0.89</b>	-0.17	-0.05	0.03
IL-5	-0.24	0.10	0.30	<b>0.87</b>
IL-6	0.46	<b>0.77</b>	-0.11	0.091
CXCL1	0.41	0.46	0.42	0.11
TNF- $\alpha$	0.25	<b>0.77</b>	-0.39	-0.096


Key: PC, principal component; IFN- $\gamma$ , interferon-gamma; IL-10, interleukin-10; IL-13, interleukin-13; IL-1 $\beta$ , interleukin-1-beta; IL-4, interleukin-4; IL-5, interleukin-5; IL-6, interleukin-6; CXCL1, chemokine (C-X-C motif) ligand 1; TNF- $\alpha$ , tumor necrosis factor alpha.  
High loadings are indicated in bold.

scores in 13- relative to 7- and 8-month-old rats ( $p$ 's < 0.007). In contrast, there were no significant differences in PC1 scores between any primiparous groups ( $p$ 's > 0.038; non-significant due to Bonferroni correction). There was no significant main effect of reproductive status, nor an interaction ( $p$ 's > 0.11). Interestingly, the

pattern observed here is akin to the age-related increase in IL-10, IFN- $\gamma$ , and IL-4 in nulliparous rats obtained with individual ANOVA analyses. Furthermore, IL-6 and TNF- $\alpha$  loaded heavily onto PC2, and IL-1 $\beta$  and IL-5 loaded heavily onto PC3 and PC4, respectively (Table 1).

### 3.14. Reproductive status influenced the direction and magnitude of correlations between microglia and neurogenesis measures

Previous studies indicate that microglia play a role in adult hippocampal neurogenesis under baseline and inflammatory conditions (Ekdahl et al., 2003; Monje, 2003; Sierra et al., 2010). However, the role of microglia in motherhood-associated neurogenic changes has not been studied; thus, to indirectly examine this, we correlated our measures of neurogenesis and microglia. Increased average length of processes was significantly correlated with a higher number of Ki67-IR cells in nulliparous ( $r = 0.58$ ,  $p = 0.005$ ; Fig. 11A) but not primiparous rats ( $r = 0.089$ ,  $p = 0.67$ ), and these correlations were significantly different ( $z = 1.81$ ,  $p = 0.035$ ). Similarly, increased average length of processes was significantly correlated with a higher number of DCX-IR cells in nulliparous ( $r = 0.62$ ,  $p = 0.002$ ; Fig. 11B) but not primiparous rats ( $r = 0.11$ ,  $p = 0.60$ ;



**Fig. 11.** Correlations between dependent variables of interest. Increased average length of Iba-1-immunoreactive (IR) cell processes was associated with higher Ki67 (A) and doublecortin (DCX; B) expression in nulliparous rats only. Ki67-IR cell number was significantly and positively correlated with DCX-IR cell number in nulliparous rats only (C). Elevated serum concentrations of IL-10 (D), and IL-4 (E) were significantly associated with shorter average Iba-1-IR cell processes in primiparous rats only. In nulliparous rats only, increased Ki67-IR (F) and DCX-IR (G) cell number was significantly associated with lower serum IL-4.

Fig. 11B), and these correlations were significantly different ( $z = 1.96, p = 0.025$ ). Interestingly, increased Ki67-IR cell number was significantly associated with more DCX-IR cells in nulliparous ( $r = 0.86, p < 0.001$ ; Fig. 11C) but not in primiparous rats ( $r = 0.30, p = 0.15$ ; Fig. 11C), and the difference between the correlations was significant ( $z = 3.04, p = 0.001$ ).

### 3.15. Reproductive status influenced the direction and magnitude of the correlations between circulating cytokine concentrations and neural measures

Peripheral cytokine signals propagate to the brain and can influence neuroimmune function (Miller et al., 2014; Quan and Banks, 2007); thus, we examined correlations between serum cytokine levels and microglial measures to assess whether central and peripheral inflammatory indicators are associated in parous and nonparous rats. Interestingly, in primiparous rats, the average length of Iba-1-IR cell processes was negatively correlated with serum IL-10 ( $r = -0.56, p = 0.005$ ; Fig. 11D) and IL-4 ( $r = -0.6, p = 0.002$ ; Fig. 11E). On the other hand, length of cell processes was not significantly correlated with IL-10 or IL-4 in nulliparous rats (all  $p$ 's  $> 0.36$ ). These correlations were significantly different between reproductive status groups for IL-10 ( $z = 2.13, p = 0.017$ ) but not for IL-4 ( $p = 0.054$ ). Regardless of reproductive status, no other significant correlations were found between length of Iba-1-IR cell processes and all other measured cytokines (all  $p$ 's  $> 0.056$ ; all trends toward significance appear in primiparous groups only). Furthermore, there were no significant correlations between any of the cytokines and Iba-1-IR cell density, regardless of reproductive status (all  $p$ 's  $> 0.23$ ).

In nulliparous rats, increased serum concentrations of IL-4 were associated with fewer DCX-IR and Ki67-IR cells (Ki67:  $-0.57, p = 0.006$ ; Fig. 11F; and DCX:  $r = -0.70, p < 0.001$ ; Fig. 11G). There were no significant correlations between any cytokines and Ki67- or DCX-IR cells in primiparous rats (all  $p$ 's  $> 0.19$ ). These correlations were significantly different between reproductive status groups in both instances (DCX:  $z = -2.58, p < 0.005$ ; Ki67:  $z = -1.82, p = 0.034$ ).

### 3.16. Serum 17 $\beta$ -estradiol concentrations were not significantly correlated with hippocampal cell proliferation, Iba-1 measures, or serum cytokine concentrations

Serum 17 $\beta$ -estradiol concentrations were not significantly associated with Ki67 expression in primiparous or nulliparous rats ( $p$ 's  $> 0.1$ ). Regardless of reproductive status, there were also no significant correlations between 17 $\beta$ -estradiol concentrations and any measure of Iba-1-IR cells (density, cell body size, average length and number of processes; all  $p$ 's  $> 0.1$ ) or any of the cytokine concentrations (all  $p$ 's  $> 0.17$ ).

## 4. Discussion

Here, we report short- and long-term effects of maternal experience on hippocampal neurogenesis, microglial density and morphology in the dentate gyrus, and circulating cytokine levels, culminating 6 months after parturition. We found that adult hippocampal neurogenesis was suppressed in mid-gestation and up to 1 month postpartum. Interestingly, the aging trajectory of neurogenesis was modulated by reproductive experience, as neurogenesis levels declined from 7 to 13 months of age in nulliparous rats but showed a slight increase in primiparous rats across the same period. Hippocampal cell proliferation was suppressed in mid-gestation and the early postpartum period in

primiparous rats, but normalized thereafter, as an age-related decline in cell proliferation was observed regardless of previous parity. Furthermore, microglia in the dentate gyrus displayed a more activated morphology in the early postpartum period, followed by a transient increase in microglial density in the later postpartum period and overall smaller microglia soma size in primiparous compared with nulliparous rats. We found alterations in circulating cytokine levels during pregnancy and the early postpartum period, and more intriguingly, we show that the age-related changes in circulating cytokine levels were dependent on parity. We also observed that reproductive status shifted the associations between microglia and neurogenesis, with average length of Iba-1-IR cell processes being positively associated with neurogenesis in nulliparous but not in primiparous rats. Furthermore, parity modulated the correlations between serum cytokines and microglial morphology, and between serum cytokines and neurogenesis levels. These correlations suggest that the relationships between immune processes and neurogenesis may be modified with parity. Finally, 17 $\beta$ -estradiol concentrations were reduced in the early postpartum period (PPD8) and in mid-gestation (GD13), in line with previous findings (Rosenblatt et al., 1988). Beyond the expected pregnancy and postpartum changes in circulating 17 $\beta$ -estradiol, we found that concentrations were not significant moderators of most variables, including hippocampal cell proliferation, microglial density and morphology, and cytokine concentrations. Collectively, our data suggest that maternal experience has transient and delayed effects on hippocampal neurogenesis, microglia, and the peripheral inflammatory milieu.

### 4.1. Adult hippocampal neurogenesis was suppressed during gestation and the postpartum period

We report that adult hippocampal neurogenesis, measured via the expression of DCX, was suppressed beginning in mid-gestation in primigravid rats. Few studies to date have examined the survival of new cells in the maternal hippocampus during gestation (Pawluski et al., 2010; Rolls et al., 2008). Our findings are, however, consistent with a study that found suppressed neurogenesis in pregnant mice during mid- and late-gestation (Rolls et al., 2008). In contrast, expression of the polysialylated neuronal cell adhesion molecule (PSA-NCAM) was increased in the dentate gyrus of pregnant rats at GD18 (Banasr et al., 2001), indicating a potential increase in neurogenesis, as PSA-NCAM is expressed on newly generated and migrating neurons (Rutishauser, 2008). However, because PSA-NCAM is also expressed on neurons undergoing other forms of plasticity (Rutishauser, 2008), its expression provides limited and non-specific information regarding neurogenesis levels. Another study in rats found that the survival of cells produced on GD1 was not significantly altered when examined across gestation (Pawluski et al., 2010), partially contrasting with our current findings. Importantly, however, DCX is expressed in immature neurons between 2 hours and 21 days after production (Brown et al., 2003). Therefore, our current data provide information on the population of cells produced as early as 8 days before impregnation and as late as the day of euthanasia (GD13). Therefore, inconsistencies between the 2 studies are not surprising, considering that the cell populations examined were produced under different conditions. We also observe a concurrent reduction in cell proliferation on GD13 indicating that this may underlie the decline in immature neurons at this time. Although no prior studies have examined cell proliferation in mid-gestation, cell proliferation was not altered on GD1 (Pawluski et al., 2010), GD7 (Shingo et al., 2003), or GD21 (Furuta and Bridges, 2005). Therefore, a more

detailed time-course analysis of hippocampal plasticity during pregnancy is warranted.

Consistent with a prior study (Workman et al., 2015), we also found reduced DCX expression in the postpartum period, evident until PPD30. This finding is also in keeping with past work showing reduced survival in new cells labeled on PPD2 and examined 21 days later (Pawluski and Galea, 2007) and in new cells labeled in mid-gestation (GD11–12) and examined 14 days later, in the early postpartum period (Rolls et al., 2008). We found reductions in hippocampal cell proliferation in the early postpartum period, which normalized by PPD30, in line with previous data (Darnaudéry et al., 2007; Leuner et al., 2007; Pawluski and Galea, 2007; Rolls et al., 2008). Thus, the suppression in immature neurons found at PPD8 likely resulted from a reduction in both cell proliferation and survival, whereas the suppression at PPD30 is likely due to decreased cell survival rather than proliferation. Although the functional significance is not known, suppressed neurogenesis in the maternal brain may be mechanistically associated with the enhanced susceptibility to mood disorders in the peripartum period (Hendrick et al., 1998). Furthermore, separate lines of evidence indicate that adult neurogenesis is involved in hippocampal regulation of the HPA axis at least in males (Schloesser et al., 2009; Snyder et al., 2011), and that the HPA axis undergoes substantial adaptations during pregnancy and the postpartum (De Weerth and Buitelaar, 2005; Lightman et al., 2001; Slattery and Neumann, 2008). Therefore, reductions in neurogenesis in the maternal hippocampus could influence HPA axis function. Suppressed neurogenesis may also be linked to deficits in hippocampus-dependent learning and memory reported in late pregnancy and the early postpartum period (reviewed in (Workman et al., 2012)).

#### 4.2. Maternal experience altered the trajectory of age-related changes in hippocampal neurogenesis

Between 8 and 13 months of age, immature neurons in the dentate gyrus declined significantly in nulliparous rats but showed a slight increase in primiparous rats. We observe an age-related decline in cell proliferation (Ki67-IR cells) regardless of reproductive status, suggesting that the differential effects in immature neurons (DCX-IR cells) are driven by differences in cell survival. Previous work indicates that hippocampal neurogenesis steadily declines with age, with the most substantial decline occurring between adulthood and middle age in female rats (Driscoll et al., 2006; Kuhn et al., 1996; Nacher et al., 2003), consistent with our current data from nulliparous rats. Thus, the increase in neurogenesis levels in middle-aged primiparous rats suggests that reproductive experience can modify the trajectory of age-related alterations in neurogenesis. Alternatively, it may also be reasonable to interpret this finding as merely a normalization of neurogenesis to nulliparous levels. However, 2 previous reports indicate higher neurogenesis levels in primiparous and multiparous relative to nulliparous middle-aged rats (Barha et al., 2015; Galea et al., 2018), and as such, an altered aging trajectory is conceivable. It is possible that a more robust difference in neurogenesis levels would arise only after multiple reproductive experiences or later into middle age. There is emerging evidence from human and rodent studies suggesting that motherhood can alter the course of age-related cognitive decline (Beeri et al., 2009; Colucci et al., 2006; Cui et al., 2014; Gatewood et al., 2005). For example, reproductive experience mitigated the age-related decline in spatial memory in rats (Gatewood et al., 2005) and mice (Cui et al., 2014). Other studies indicate that parity is associated with cognitive impairment in the aging female (Beeri et al., 2009; Colucci et al., 2006). These

inconsistencies may be reconciled by more complex interactions with genetic factors that have been associated with pathological cognitive aging (Corbo et al., 2007; Cui et al., 2014). Although speculative, the modest increase in hippocampal neurogenesis in middle-aged primiparous rats may be associated with enhanced hippocampus-dependent cognition that is seen at that time.

Interestingly, we observed differences in the relationship between levels of cell proliferation and immature neurons depending on reproductive status, such that a significant positive correlation between the 2 measures was only seen in nulliparous rats. In addition, increased IL-4 concentrations were associated with reduced proliferation and immature neurons in the hippocampus of nulliparous but not primiparous rats. Previous work points to a role of IL-4 in the regulation of cell proliferation under conditions of neurodegeneration (Bhattarai et al., 2016), therefore our findings suggest that this role may be altered by parity. These relationships should be further investigated, as they appear when reproductive status groups are collapsed across age but nonetheless indicate that parity may modulate the effects of immune signaling on hippocampal neurogenesis.

#### 4.3. Microglia assumed a de-ramified morphology in the early postpartum period

Microglia display a predominantly ramified morphology under basal conditions, and de-ramification is thought to be indicative of increased classical activation under inflammatory conditions (Luo and Chen, 2012). Here, we show that microglia in the dentate gyrus exhibited significantly shortened processes at 8 days postpartum, suggesting an increase in microglial activation in the early postpartum period. This de-ramification was a transient morphological modification as the average length of processes was not significantly different from nulliparous controls by PPD30. We further show a small but significant shift in the percentages of microglia assuming different morphological states at PPD8. Specifically, we observe a reduction in ramified morphology and an increase in stout morphology, which in tandem with our data on length of processes indicates a shift toward classical microglial activation in the dentate gyrus in the early postpartum period. However, we interpret these findings with caution, as the information that can be deduced from morphological phenotype about functional states is limited (Boche et al., 2013).  $17\beta$ -estradiol concentrations significantly moderated the average length of microglial processes, suggesting that the observed de-ramification at PPD8 might be moderated by postpartum reductions in estradiol. However, as this measure of estradiol concentrations provides information about a small window of time around perfusion, a complete picture of the role of estradiol cannot be extrapolated from the present study. Future studies should also consider the role of other hormones, including progesterone and corticosterone, in motherhood-associated changes in microglia.

To our knowledge, only 2 studies to date have examined microglia in the maternal brain (Haim et al., 2017; Posillico and Schwarz, 2016), with findings partially consistent with our current data. Haim et al. (2017) reported a decrease in the number of microglia with a ramified morphology on PPD8, in several regions including the dorsal hippocampus. This is in line with our current report of microglial deramification and a shift in percentages of morphological states in the dentate gyrus on the same postpartum day (PPD8). In the present study, we found an increase in the density of microglia in the dorsal and ventral dentate gyri at PPD30, but no alteration in density during gestation or the early postpartum period. Although no other studies have examined microglia in the maternal brain as late as 30 days postpartum, our finding that microglial density in the hippocampus remains unchanged during

gestation and the early postpartum period contrasts previous reports (Haim et al., 2017; Posillico and Schwarz, 2016). These previous studies found reduced microglial density in several brain regions from late gestation to the early postpartum (GD20 and PPD1, 8, and 21: Haim et al., 2017; PPD0: Posillico and Schwarz, 2016). Furthermore, Haim et al. (2017) found that microglial density normalized to nulliparous control levels by PPD21 in all regions examined except the dorsal hippocampus. It appears, however, that the inconsistencies in findings may be accounted for by differences in microglial densities within subregions of the dentate gyrus or by methodological differences related to density measurement. For example, Haim et al. (2017) examined Iba-1 density within the dorsal dentate gyrus only, whereas we included samples from both the dorsal and ventral dentate gyri. In addition, as we were primarily interested in the neurogenic niche, we quantified Iba-1-IR cells within the GCL, the SGZ, and a thin band of the ML, whereas Haim et al did not specify subregions within the dentate gyrus. Finally, Haim et al. (2017) used optical density, whereas density here was defined as the number of cells per volume of dentate gyrus. Interestingly, the increase in microglial density that we find at PPD30 coincides with a return to normalized microglial morphology. We speculate that this may represent a resolution from the proinflammatory state at PPD8. Overall, our novel data provide an important addition to the literature indicating that pregnancy-related immune adaptations are not limited to the periphery but also exist in the brain. More specifically, our data indicate the existence of a proinflammatory hippocampal environment in the early postpartum period. Importantly, increased microglial activation may be central to the pathophysiology of depression (Kreisel et al., 2014; Miller and Raison, 2015; Setiawan et al., 2015), thus it is conceivable that similar processes are implicated in postpartum depression. Our current findings which point to an increase in microglial activation in the early postpartum may represent a neural mechanism of enhanced susceptibility to mood disorders at that time, however additional work would be required to directly test this.

The effects of parity on microglial morphology were not limited to the early postpartum period. Specifically, although the number of microglial cell processes increased significantly with age in nulliparous rats, this effect was prevented by parity. Furthermore, parity reduced microglial soma size, an effect which appears to emerge at PPD30 onward. The functional significance of these long-lasting alterations cannot be determined from the present study, but as increased microglial soma size is indicative of classical activation, it is possible that parity may dampen microglial activation in the aging brain. To gain better insight into the functional significance of these changes, future studies should investigate how previous parity may impact microglial structure and function in the aging brain in response to an immune challenge.

Interestingly, we found that reproductive status affected the associations between microglia and neurogenesis in the hippocampus. Specifically, increased cell proliferation and immature neurons were significantly associated with more ramified microglial morphology in nulliparous rats only. These observations suggest that the neuro-immune regulation of adult hippocampal neurogenesis is affected by reproductive status. Importantly, in addition to phagocytic activity during development and disease, microglia are important players in the regulation of adult hippocampal neurogenesis, where they phagocytose apoptotic new cells while maintaining ramified morphology and a non-inflammatory environment (Sierra et al., 2010). Thus, future research should directly examine the possibilities of microglial phagocytosis in the regulation of neurogenesis during pregnancy and the postpartum period.

#### 4.4. Maternal experience modifies the age-related changes in circulating cytokine levels

In addition to expected cytokine alterations during pregnancy and the early postpartum period (Holtan et al., 2015; Shimaoka et al., 2000), we observed both persistent and delayed effects of reproductive experience on the circulating cytokine profile. Specifically, after an initial increase at PPD8, IL-4 levels were persistently blunted in primiparous rats relative to nulliparous controls. Furthermore, in nulliparous but not primiparous rats, IFN- $\gamma$  and IL-10 increased and IL-5 declined significantly with age. Adaptations to the immune systems during pregnancy and the early postpartum period are well established (PrabhuDas et al., 2015). On the other hand, little attention has been paid to potential long-term effects of motherhood on the immune system. To our knowledge, only a few studies have examined the effect of parity on immune systems in aged female mice and 1 in aged rats. Together, these studies suggested that parity may delay certain indicators of immune senescence, as they relate to alterations in cytokine production *in vitro* from activated spleen cells (Barrat et al., 1997a), and the distribution of immune cell populations in the spleen (Barrat et al., 1997b) and bone marrow (Barrat et al., 1999). Another study from our laboratory found a trend for increased serum levels of IL-6 in primiparous compared with nulliparous rats at 15 months of age, in rats that had previous surgery (Galea et al., 2018). Here, we report that a single reproductive experience alters serum cytokine levels when examined in middle age, up to 6 months after the reproductive event.

Importantly, peripheral cytokines can affect brain function, as they can access the central nervous system through various mechanisms, including active and passive transport, and the activation of cytokine receptors on afferent nerve fibers (reviewed in Miller et al., 2014; Quan and Banks, 2007). Thus, the effects of parity to modify age-related changes in peripheral cytokines may have ramifications for the aging brain in general, and more specifically the hippocampus, as it contains one of the highest densities of proinflammatory cytokine receptors in the brain (reviewed in Loftis et al., 2010). Inflammation is a core characteristic of the aging process, and the proinflammatory cytokine IFN- $\gamma$  increases with age (Oxenkrug, 2011; Rodríguez et al., 2007). Thus, our current data suggest that parity may prevent or delay at least certain aspects of aging-related inflammation. Interestingly, from PPD30 onward, we observed a sustained suppression in serum IL-4 in primiparous rats relative to nulliparous controls. Traditionally considered an anti-inflammatory cytokine (Hart et al., 1989), elevated IL-4 in nulliparous rats may be suggestive of a compensatory response to attenuate a proinflammatory state that is indicated by elevated IFN- $\gamma$ . However, IL-4 is pleiotropic (Milner et al., 2010) and as such can also have proinflammatory properties. For example, sustained exposure to elevated levels of IL-4 was associated with increased inflammation (Milner et al., 2010). The same study found prolonged IL-4 exposure to be associated specifically with elevated levels of IL-10 and IFN- $\gamma$ , but not IL-6 and TNF- $\alpha$ . This is in line with our current data in which IL-4, IL-10, and IFN- $\gamma$  were concurrently elevated in middle-aged nulliparous rats. Thus, the cytokine profile in nulliparous groups may be alternatively driven by an increase in IL-4. We investigated correlations between peripheral cytokines and microglial morphology as a plethora of work suggests that systemic inflammation can profoundly affect microglial activation (reviewed in Hoogland et al., 2015). Interestingly, we found that higher concentrations of IL-10 and IL-4 were associated with more de-ramified microglial morphology in primiparous but not nulliparous rats. The mechanisms and consequences of these altered relationships between peripheral cytokines and microglial morphology are not known, and these findings should be interpreted with caution, as the correlations were detected when age was not included as a

factor. In the future, it is also important to examine whether parity may have similar effects on age-related changes in brain cytokines, particularly in the hippocampus. Although the functional consequences of the observed differences in cytokine profiles cannot be determined from our current finding, we demonstrate here that the trajectory of immune senescence is altered by parity.

## 5. Conclusions

In summary, we report that maternal experience suppressed hippocampal neurogenesis (proliferation and immature neurons) during gestation and the postpartum period, but unlike what was observed in nulliparous rats, there was no age-related decline in neurogenesis (immature neurons) in middle-aged primiparous rats. Maternal experience also resulted in transient microglial de-ramification in the dentate gyrus, suggesting the existence of a proinflammatory hippocampal environment in the early postpartum period. In addition to short-term cytokine alterations, maternal experience modified the trajectory of age-related changes in circulating cytokine levels. Furthermore, there were significant associations between neurogenesis and microglia or IL-4 in nulliparous but not in primiparous rats, indicating different enduring relationships between immune characteristics and neurogenesis with maternal experience. These findings should encourage future work aimed at delineating the functional consequences for behavior and immune function across the peripartum period and beyond, especially in relation to maternal mood and cognition. Importantly, our data provide support for the notion that female reproductive history should be regarded as an important determinant of aging-related changes in physiology.

## Disclosure

The authors declare that there are no conflicts of interest associated with this work.

## Acknowledgements

The authors thank Dr. Timothy Kieffer and Travis Webber for generously providing access to their Sector Imager and Arianne Albert for her assistance with principal component analyses.

This work was supported by a grant from the Canadian Institute of Health Research to LAMG (PJT148662) and a Four-Year Doctoral Fellowship from the University of British Columbia to RSE.

## References

- Aghaepour, N., Ganio, E.A., McIlwain, D., Tsai, A.S., Tingle, M., Van Gassen, S., Gaudilliere, D.K., Baca, Q., McNeil, L., Okada, R., Ghaemi, M.S., Furman, D., Wong, R.J., Winn, V.D., Druzin, M.L., El-Sayed, Y.Y., Quaintance, C., Gibbs, R., Darmstadt, G.L., Shaw, G.M., Stevenson, D.K., Tibshirani, R., Nolan, G.P., Lewis, D.B., Angst, M.S., Gaudilliere, B., 2017. An immune clock of human pregnancy. *Sci. Immunol.* 2. pii:eaan2946.
- Altman, J., Das, G.D., 1965. Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J. Comp. Neurol.* 124, 319–335.
- Banasr, M., Hery, M., Brezun, J.M., Daszuta, A., 2001. Serotonin mediates oestrogen stimulation of cell proliferation in the adult dentate gyrus. *Eur. J. Neurosci.* 14, 1417–1424.
- Barha, C.K., Galea, L.A.M., 2011. Motherhood alters the cellular response to estrogens in the hippocampus later in life. *Neurobiol. Aging* 32, 2091–2095.
- Barha, C.K., Hanna, C.W., Salvante, K.G., Wilson, S.L., Robinson, W.P., Altman, R.M., Nepomnaschy, P.A., 2016. Number of children and telomere length in women: a prospective, longitudinal evaluation. *PLoS One* 11, e0146424.
- Barha, C.K., Lieblich, S.E., Chow, C., Galea, L.A.M., 2015. Multiparity-induced enhancement of hippocampal neurogenesis and spatial memory depends on ovarian hormone status in middle age. *Neurobiol. Aging* 36, 2391–2405.
- Barrat, F., Lesourd, B., Boulouis, H.J., Thibault, D., Vincent-Naulleau, S., Gjata, B., Louise, A., Neway, T., Pilet, C., 1997a. Sex and parity modulate cytokine production during murine ageing. *Clin. Exp. Immunol.* 109, 562–568.
- Barrat, F., Lesourd, B.M., Louise, A., Boulouis, H.J., Vincent-Naulleau, S., Thibault, D., Sanaa, M., Neway, T., Pilet, C.H., 1997b. Surface antigen expression in spleen cells of C57B1/6 mice during ageing: influence of sex and parity. *Clin. Exp. Immunol.* 107, 593–600.
- Barrat, F.S., Lesourd, B.M., Louise, A.S., Boulouis, H.J., Thibault, D.J., Neway, T., Pilet, C.A., 1999. Pregnancies modulate B lymphopoiesis and myelopoiesis during murine ageing. *Immunology* 98, 604–611.
- Beeri, M.S., Rapp, M., Schmeidler, J., Reichenberg, A., Purohit, D.P., Perl, D.P., Grossman, H.T., Prohovnik, I., Haroutunian, V., Silverman, J.M., 2009. Number of children is associated with neuropathology of Alzheimer's disease in women. *Neurobiol. Aging* 30, 1184–1191.
- Bennett, H.A., Einarson, A., Taddio, A., Koren, G., Einarson, T.R., 2004. Prevalence of depression during pregnancy: systematic review. *Obstet. Gynecol.* 103, 698–709.
- Bhattarai, P., Thomas, A.K., Cosacak, M.I., Papadimitriou, C., Mashkaryan, V., Froc, C., Reinhardt, S., Kurth, T., Dahl, A., Zhang, Y., Kizil, C., 2016. IL4/STAT6 signaling activates neural stem cell proliferation and neurogenesis upon amyloid- $\beta$ 42 aggregation in adult zebrafish brain. *Cell. Rep.* 17, 941–948.
- Boche, D., Perry, V.H., Nicoll, J.A.R., 2013. Review: activation patterns of microglia and their identification in the human brain. *Neuropathol. Appl. Neurobiol.* 39, 3–18.
- Bodnar, T.S., Taves, M.D., Lavigne, K.M., Woodward, T.S., Soma, K.K., Weinberg, J., 2017. Differential activation of endocrine-immune networks by arthritis challenge: insights from colony-specific responses. *Sci. Rep.* 7, 1–14.
- Boldrini, M., Fulmore, C.A., Tartt, A.N., Simeon, L.R., Pavlova, I., Poposka, V., Rosoklija, G.B., Stankov, A., Arango, V., Dwork, A.J., Hen, R., Mann, J.J., 2018. Human hippocampal neurogenesis persists throughout aging. *Cell. Stem. Cell.* 22, 589–599.
- Bridges, R.S., 2015. Neuroendocrine regulation of maternal behavior. *Front. Neuroendocrinol.* 36, 178–196.
- Brown, J.P., Couillard-Després, S., Cooper-Kuhn, C.M., Winkler, J., Aigner, L., Kuhn, H.G., 2003. Transient expression of doublecortin during adult neurogenesis. *J. Comp. Neurol.* 467, 1–10.
- Campbell, S., MacQueen, G., 2004. The role of the hippocampus in the pathophysiology of major depression. *J. Psychiatry. Neurosci.* 29, 417–426.
- Catalano, R.D., Lannagan, T.R.M., Gorowiec, M., Denison, F.C., Norman, J.E., Jabbour, H.N., 2010. Prokineticins: novel mediators of inflammatory and contractile pathways at parturition? *Mol. Hum. Reprod.* 16, 311–319.
- Colucci, M., Cammarata, S., Assini, A., Croce, R., Clerici, F., Novello, C., Mazzella, L., Dagnino, N., Mariani, C., Tanganelli, P., 2006. The number of pregnancies is a risk factor for Alzheimer's disease. *Eur. J. Neurol.* 13, 1374–1377.
- Corbo, R.M., Gambina, G., Ulizzi, L., Monini, P., Broglio, E., Rosano, A., Scacchi, R., 2007. Combined effect of apolipoprotein e genotype and past fertility on age at onset of Alzheimer's disease in women. *Dement. Geriatr. Cogn. Disord.* 24, 82–85.
- Cui, J., Jothishankar, B., He, P., Staufienbiel, M., Shen, Y., Li, R., 2014. Amyloid precursor protein mutation disrupts reproductive experience-enhanced normal cognitive development in a mouse model of Alzheimer's disease. *Mol. Neurobiol.* 49, 103–112.
- Cuttler, C., Graf, P., Pawluski, J.L., Galea, L.A., 2011. Everyday life memory deficits in pregnant women. *Can. J. Exp. Psychol.* 65, 27–37.
- Darcy, J.M., Grzywacz, J.G., Stephens, R.L., Leng, I., Clinch, C.R., Arcury, T.A., 2011. Maternal depressive symptomatology: 16-month follow-up of infant and maternal health-related quality of life. *J. Am. Board. Fam. Med.* 24, 249–257.
- Darnaudéry, M., Perez-Martin, M., Del Favero, F., Gomez-Roldan, C., Garcia-Segura, L.M., Maccari, S., 2007. Early motherhood in rats is associated with a modification of hippocampal function. *Psychoneuroendocrinology* 32, 803–812.
- De Groot, R.H., Vuurman, E.F., Hornstra, G., Jolles, J., 2006. Differences in cognitive performance during pregnancy and early motherhood. *Psychol. Med.* 36, 1023–1032.
- De Weerth, C., Buitelaar, J.K., 2005. Physiological stress reactivity in human pregnancy - a review. *Neurosci. Biobehav. Rev.* 29, 295–312.
- Driscoll, I., Howard, S.R., Stone, J.C., Monfils, M.H., Tomanek, B., Brooks, W.M., Sutherland, R.J., 2006. The aging hippocampus: a multi-level analysis in the rat. *Neuroscience* 139, 1173–1185.
- Dulac, C., O'Connell, L.A., Wu, Z., 2014. Neural control of maternal and paternal behaviors. *Science* 345, 765–770.
- Ekdahl, C.T., Claassen, J.-H., Bonde, S., Kokaia, Z., Lindvall, O., 2003. Inflammation is detrimental for neurogenesis in adult brain. *Proc. Natl. Acad. Sci.* 100, 13632–13637.
- Eriksson, P.S., Perfilieva, E., Björk-Eriksson, T., Alborn, A.M., Nordborg, C., Peterson, D.A., Gage, F.H., 1998. Neurogenesis in the adult human hippocampus. *Nat. Med.* 4, 1313–1317.
- Furuta, M., Bridges, R.S., 2005. Gestation-induced cell proliferation in the rat brain. *Brain. Res. Dev. Brain. Res.* 156, 61–66.
- Galea, L.A.M., Leuner, B., Slattery, D.A., 2014. Hippocampal plasticity during the peripartum period: influence of sex steroids, stress and ageing. *J. Neuroendocrinol.* 26, 641–648.
- Galea, L.A.M., Ormerod, B.K., Sampath, S., Kostaras, X., Wilkie, D.M., Phelps, M.T., 2000. Spatial working memory and hippocampal size across pregnancy in rats. *Horm. Behav.* 37, 86–95.

- Galea, L.A.M., Roes, M.M., Dimech, C.J., Chow, C., Mahmoud, R., Lieblisch, S.E., Duarte-Guterman, P., 2018. Premarin has opposing effects on spatial learning, neural activation, and serum cytokine levels in middle-aged female rats depending on reproductive history. *Neurobiol. Aging*, 70, 291–307.
- Gatewood, J.D., Morgan, M.D., Eaton, M., McNamara, I.M., Stevens, L.F., MacBeth, A.H., Meyer, E.A.A., Lomas, L.M., Kozub, F.J., Lambert, K.G., Kinsley, C.H., 2005. Motherhood mitigates aging-related decrements in learning and memory and positively affects brain aging in the rat. *Brain. Res. Bull.* 66, 91–98.
- Groer, M.E., Jevitt, C., Ji, M., 2015. Immune changes and dysphoric moods across the postpartum. *Am. J. Reprod. Immunol.* 73, 193–198.
- Guerreiro, R.J., Santana, I., Brás, J.M., Santiago, B., Paiva, A., Oliveira, C., 2007. Peripheral inflammatory cytokines as biomarkers in Alzheimer's disease and mild cognitive impairment. *Neurodegener. Dis.* 4, 406–412.
- Haim, A., Julian, D., Albin-Brooks, C., Brothers, H.M., Lenz, K.M., Leuner, B., 2017. A survey of neuroimmune changes in pregnant and postpartum female rats. *Brain. Behav. Immun.* 59, 67–78.
- Hall, M.E., George, E.M., Granger, J.P., 2011. The heart during pregnancy. *Rev. Esp. Cardiol.* 64, 1045–1050.
- Hart, P.H., Vitti, G.F., Burgess, D.R., Whitty, G.A., Piccoli, D.S., Hamilton, J.A., 1989. Potential antiinflammatory effects of interleukin 4: suppression of human monocyte tumor necrosis factor alpha, interleukin 1, and prostaglandin E2. *Proc. Natl. Acad. Sci.* 86, 3803–3807.
- Helle, S., Lummaa, V., Jokela, J., 2004. Accelerated immunosenescence in preindustrial twin mothers. *Proc. Natl. Acad. Sci. U S A* 101, 12391–12396.
- Hendrick, V., Altschuler, L.L., Suri, R., 1998. Hormonal changes in the postpartum and implications for postpartum depression. *Psychosomatics* 39, 93–101.
- Hodes, G.E., Pfau, M.L., Leboeuf, M., Golden, S.A., Christoffel, D.J., Bregman, D., Rebusi, N., Heshmati, M., Aleyasin, H., Warren, B.L., Labonté, B., Horn, S., Lapidus, K.A., Stelzhammer, V., Wong, E.H.F., Bahn, S., Krishnan, V., Bolaños-Guzmán, C.A., Murrough, J.W., Merad, M., Russo, S.J., 2014. Individual differences in the peripheral immune system promote resilience versus susceptibility to social stress. *Proc. Natl. Acad. Sci.* 111, 16136–16141.
- Hoekzema, E., Barba-Müller, E., Pozzobon, C., Picado, M., Lucco, F., García-García, D., Soliva, J.C., Tobeña, A., Desco, M., Crone, E.A., Ballesteros, A., Carmona, S., Villarroya, O., 2016. Pregnancy leads to long-lasting changes in human brain structure. *Nat. Neurosci.* 20, 287–296.
- Holtan, S.G., Chen, Y., Kaimal, R., Creedon, D.J., Enninga, E.A.L., Nevala, W.K., Markovic, S.N., 2015. Growth modeling of the maternal cytokine milieu throughout normal pregnancy: macrophage-derived chemokine decreases as inflammation/counterregulation increases. *J. Immunol. Res.* 2015, 952571.
- Hoogland, I.C.M., Houbolt, C., van Westerloo, D.J., van Gool, W.A., van de Beek, D., 2015. Systemic inflammation and microglial activation: systematic review of animal experiments. *J. Neuroinflammation.* 12, 114.
- Karperien, A., Ahammer, H., Jelinek, H.F., 2013. Quantitating the subtleties of microglial morphology with fractal analysis. *Front. Cell. Neurosci.* 7, 3.
- Kinsley, C.H., Madonia, L., Gifford, G.W., Tureski, K., Griffin, G.R., Lowry, C., Williams, J., Collins, J., McLearn, H., Lambert, K.G., 1999. Motherhood improves learning and memory. *Nature* 402, 137–138.
- Korzhevskii, D.E., Kirik, O.V., 2016. Brain microglia and microglial markers. *Neurosci. Behav. Physiol.* 46, 284–290.
- Kreisel, T., Frank, M.G., Licht, T., Reshef, R., Ben-Menachem-Zidon, O., Baratta, M.V., Maier, S.F., Yirmiya, R., 2014. Dynamic microglial alterations underlie stress-induced depressive-like behavior and suppressed neurogenesis. *Mol. Psychiatry.* 19, 699–709.
- Kuhn, H.G., Dickinson-Anson, H., Gage, F.H., 1996. Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation. *J. Neurosci.* 16, 2027–2033.
- Lain, K.Y., Catalano, P.M., 2007. Metabolic changes in pregnancy. *Clin. Obstet. Gynecol.* 50, 938–948.
- Lee, J., Lee, Y., Yuk, D., Choi, D., Ban, S., Oh, K., Hong, J., 2008. Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment through enhancement of beta-amyloid generation. *J. Neuroinflammation* 5, 37.
- Leuner, B., Christian, M., Liron, N., Elizabeth, G., 2007. Maternal experience inhibits the production of immature neurons in the Hippocampus during the postpartum period through. *Hippocampus* 17, 434–442.
- Leuner, B., Gould, E., 2010. Dendritic growth in medial prefrontal cortex and cognitive flexibility are enhanced during the postpartum period. *J. Neurosci.* 30, 13499–13503.
- Lightman, S.L., Windle, R.J., Wood, S.A., Kershaw, Y.M., Shanks, N., Ingram, C.D., 2001. Peripartum plasticity within the hypothalamo-pituitary-adrenal axis. *Prog. Brain Res.* 133, 111–129.
- Loftis, J.M., Huckans, M., Morasco, B.J., 2010. Neuroimmune mechanisms of cytokine-induced depression: current theories and novel treatment strategies. *Neurobiol. Dis.* 37, 519–533.
- Louveau, A., Harris, T.H., Kipnis, J., 2015. Revisiting the mechanisms of CNS immune privilege. *Trends. Immunol.* 36, 569–577.
- Luo, X.-G., Chen, S.-D., 2012. The changing phenotype of microglia from homeostasis to disease. *Transl. Neurodegener.* 1, 9.
- Mahmoud, R., Wainwright, S.R., Chaiton, J.A., Lieblisch, S.E., Galea, L.A.M., 2016a. Ovarian hormones, but not fluoxetine, impart resilience within a chronic unpredictable stress model in middle-aged female rats. *Neuropharmacology* 107, 278–293.
- Mahmoud, R., Wainwright, S.R., Galea, L.A.M., 2016b. Sex hormones and adult hippocampal neurogenesis: regulation, implications, and potential mechanisms. *Front. Neuroendocrinol.* 41, 129–152.
- Miller, A.H., Haroon, E., Raison, C.L., Felger, J.C., 2014. Cytokine targets in the brain: impact on neurotransmitters and neurocircuits. *Depress. Anxiety.* 30, 297–306.
- Miller, A.H., Raison, C.L., 2015. The role of inflammation in depression: from evolutionary imperative to modern treatment target. *Nat. Rev. Immunol.* 16, 22–34.
- Milner, J.D., Orekov, T., Ward, J.M., Cheng, L., Torres-Velez, F., Junttila, I., Sun, G., Buller, M., Morris, S.C., Finkelman, F.D., Paul, W.E., 2010. Sustained IL-4 exposure leads to a novel pathway for hemophagocytosis, inflammation, and tissue macrophage accumulation. *Blood* 116, 2476–2483.
- Monje, M.L., 2003. Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302, 1760–1765.
- Mor, G., Cardenas, I., 2010. The immune system in pregnancy: a unique complexity. *Am. J. Reprod. Immunol.* 63, 425–433.
- Nacher, J., Alonso-Llosa, G., Rosell, D.R., McEwen, B.S., 2003. NMDA receptor antagonist treatment increases the production of new neurons in the aged rat hippocampus. *Neurobiol. Aging.* 24, 273–284.
- Nimmerjahn, A., Kirchhoff, F., Helmchen, F., 2005. Neuroscience: resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 308, 1314–1318.
- O'Hara, M.W., 2009. Postpartum depression: what we know. *J. Clin. Psychol.* 65, 1258–1269.
- Oatridge, A., Holdcroft, A., Saeed, N., Hajnal, J.V., Puri, B.K., Fusi, L., Bydder, G.M., 2002. Change in brain size during and after pregnancy: study in healthy women and women with preeclampsia. *Am. J. Neuroradiol.* 23, 19–26.
- Overstreet-Wadiche, L.S., Bromberg, D. a, Bensen, A.L., Westbrook, G.L., 2006. Seizures accelerate functional integration of adult-generated granule cells. *J. Neurosci.* 26, 4095–5103.
- Oxenkrug, G.F., 2011. Interferon-gamma-inducible kynurenines/pteridines inflammation cascade: implications for aging and aging-associated psychiatric and medical disorders. *J. Neural. Transm.* 118, 75–85.
- Parkhurst, C.N., Yang, G., Ninan, I., Savas, J.N., Yates, J.R., Lafaille, J.J., Hempstead, B.L., Littman, D.R., Gan, W.B., 2013. Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor. *Cell* 155, 1596–1609.
- Pawluski, J.L., Barakauskas, V.E., Galea, L.A.M., 2010. Pregnancy decreases oestrogen receptor alpha expression and pyknosis, but not cell proliferation or survival, in the hippocampus. *J. Neuroendocrinol.* 22, 248–257.
- Pawluski, J.L., Galea, L.A.M., 2007. Reproductive experience alters hippocampal neurogenesis during the postpartum period in the dam. *Neuroscience* 149, 53–67.
- Pawluski, J.L., Galea, L.A.M., 2006. Hippocampal morphology is differentially affected by reproductive experience in the mother. *J. Neurobiol.* 66, 71–81.
- Pawluski, J.L., Walker, S.K., Galea, L.A.M., 2006. Reproductive experience differentially affects spatial reference and working memory performance in the mother. *Horm. Behav.* 49, 143–149.
- Plümpe, T., Ehninger, D., Steiner, B., Klempin, F., Jessberger, S., Brandt, M., Römer, B., Rodriguez, G.R., Kronenberg, G., Kempermann, G., 2006. Variability of doublecortin-associated dendrite maturation in adult hippocampal neurogenesis is independent of the regulation of precursor cell proliferation. *BMC. Neurosci.* 7, 77.
- Posillico, C.K., Schwarz, J.M., 2016. An investigation into the effects of antenatal stressors on the postpartum neuroimmune profile and depressive-like behaviors. *Behav. Brain Res.* 298, 218–228.
- Prabhudas, M., Bonney, E., Caron, K., Dey, S., Erlebacher, A., Fazleabas, A., Fisher, S., Golos, T., Matzuk, M., McCune, J.M., Mor, G., Schulz, L., Soares, M., Spencer, T., Strominger, J., Way, S.S., Yoshinaga, K., Loreau, M., Naeem, S., Inchausti, P., 2015. Immune mechanisms at the maternal-fetal interface: perspectives and challenges. *Nat. Immunol.* 16, 237–243.
- Quan, N., Banks, W.A., 2007. Brain-immune communication pathways. *Brain. Behav. Immun.* 21, 727–735.
- Rannevik, G., Carlström, K., Jeppsson, S., Bjerre, B., Svanberg, L., 1986. A prospective long-term study in women from pre-menopause to post-menopause: changing profiles of gonadotrophins, oestrogens and androgens. *Maturitas* 8, 297–307.
- Rodríguez, M.I., Escames, G., López, L.C., López, A., García, J.A., Ortiz, F., Acuña-Castroviejo, D., 2007. Chronic melatonin treatment reduces the age-dependent inflammatory process in senescence-accelerated mice. *J. Pineal. Res.* 42, 272–279.
- Rolls, A., Schori, H., London, A., Schwartz, M., 2008. Decrease in hippocampal neurogenesis during pregnancy: a link to immunity. *Mol. Psychiatry.* 13, 468–469.
- Rosenblatt, J.S., Mayer, A.D., Giordano, A.L., 1988. Hormonal basis during pregnancy for the onset of maternal behavior in the rat. *Psychoneuroendocrinology* 13, 29–46.
- Rossant, J., Cross, J.C., 2001. Placental development: lessons from mouse mutants. *Nat. Rev. Genet.* 2, 538–548.
- Rutishauser, U., 2008. Polysialic acid in the plasticity of the developing and adult vertebrate nervous system. *Nat. Rev. Neurosci.* 9, 26–35.
- Sahay, A., Hen, R., 2007. Adult hippocampal neurogenesis in depression. *Nat. Neurosci.* 10, 1110–1115.
- Schloesser, R.J., Manji, H.K., Martinowich, K., 2009. Suppression of adult neurogenesis leads to an increased hypothalamo-pituitary-adrenal axis response. *Neuroreport* 20, 553–557.

- Scholzen, T., Gerdes, J., 2000. The Ki-67 protein: from the known and the unknown. *J. Cell. Physiol.* 182, 311–322.
- Schumacher, A., Costa, S.D., Zenclussen, A.C., 2014. Endocrine factors modulating immune responses in pregnancy. *Front. Immunol.* 5, 196.
- Schwarz, J.M., Sholar, P.W., Bilbo, S.D., 2012. Sex differences in microglial colonization of the developing rat brain. *J. Neurochem.* 120, 948–963.
- Setiawan, E., Wilson, A.A., Mizrahi, R., Rusjan, P.M., Miler, L., Rajkowska, G., Suridjan, I., Kennedy, J.L., Rekkas, P.V., Houle, S., Meyer, J.H., 2015. Role of translocator protein density, a marker of neuroinflammation, in the brain during major depressive episodes. *JAMA. Psychiatry.* 72, 268–275.
- Shimaoka, Y., Hidaka, Y., Tada, H., Amino, N., Nakamura, T., Murata, Y., Mitsuda, N., Morimoto, Y., 2000. Changes in cytokine production during and after normal pregnancy. *Am. J. Reprod. Immunol.* 44, 143–147.
- Shingo, T., Gregg, C., Enwere, E., Fujikawa, H., Hassam, R., Geary, C., Cross, J.C., Weiss, S., 2003. Pregnancy-stimulated neurogenesis in the adult female forebrain mediated by prolactin. *Science* 299, 117–120.
- Sierra, A., Beccari, S., Diaz-Aparicio, I., Encinas, J.M., Comeau, S., Tremblay, M.È., 2014. Surveillance, phagocytosis, and inflammation: how never-resting microglia influence adult hippocampal neurogenesis. *Neural. Plast.*
- Sierra, A., Encinas, J.M., Deudero, J.J.P., Chancey, J.H., Enikolopov, G., Overstreet-Wadiche, L.S., Tsirka, S.E., Maletic-Savatic, M., 2010. Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell* 7, 483–495.
- Slattery, D.A., Neumann, I.D., 2008. No stress please! Mechanisms of stress hypo-responsiveness of the maternal brain. *J. Physiol.* 586, 377–385.
- Snyder, J.S., Soumier, A., Brewer, M., Pickel, J., Cameron, H.A., 2011. Adult hippocampal neurogenesis buffers stress responses and depressive behaviour. *Nature* 476, 458–461.
- Sweatt, J.D., 2004. Hippocampal function in cognition. *Psychopharmacology (Berl)* 174, 99–110.
- vom Berg, J., Prokop, S., Miller, K.R., Obst, J., Kälin, R.E., Lopategui-Cabezas, I., Wegner, A., Mair, F., Schipke, C.G., Peters, O., Winter, Y., Becher, B., Heppner, F.L., 2012. Inhibition of IL-12/IL-23 signaling reduces Alzheimer's disease-like pathology and cognitive decline. *Nat. Med.* 18, 1812–1819.
- Workman, J.L., Barha, C.K., Galea, L.A.M., 2012. Endocrine substrates of cognitive and affective changes during pregnancy and postpartum. *Behav. Neurosci.* 126, 54–72.
- Workman, J.L., Raineki, C., Weinberg, J., Galea, L.A.M., 2015. Alcohol and pregnancy: effects on maternal care, HPA axis function, and hippocampal neurogenesis in adult females. *Psychoneuroendocrinology* 57, 37–50.
- Yau, S., Li, A., So, K., 2015. Involvement of adult hippocampal neurogenesis in learning and forgetting. *Neural Plast* 2015, 717958.