



## Correction to: Expression of miR-145 and Its Target Proteins Are Regulated by miR-29b in Differentiated Neurons

Abhishek Jauhari<sup>1,2</sup> · Tanisha Singh<sup>1,3</sup> · Sanjay Yadav<sup>1</sup>

Published online: 3 July 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

**Correction to: Mol Neurobiol (2018) 55:8978–8990**

<https://doi.org/10.1007/s12035-018-1009-9>

The original version of this article unfortunately contained an error. In Fig. 8, the image under section a) NTC, and b) NTC + PFT $\alpha$  are copied by mistake.

The corrected Fig. 8 is hereby given below. As there is no change in values of the figure, it will not change the result of paper.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s12035-018-1009-9>

---

✉ Sanjay Yadav  
sanjay@iitr.res.in


Abhishek Jauhari  
abhijauhari@gmail.com

Tanisha Singh  
tanishasingh186@gmail.com

<sup>1</sup> Developmental Toxicology Laboratory, Systems Toxicology and Health Risk Assessment Group, CSIR- Indian Institute of Toxicology Research (CSIR-IITR), Vishvigyan Bhawan, 31 Mahatma Gandhi Marg, Lucknow, Uttar Pradesh 226001, India

<sup>2</sup> Academy of Scientific and Innovative Research (AcSIR), CSIRIITR Campus, New Delhi, India

<sup>3</sup> Department of Biochemistry, School of Dental Sciences, Babu Banarasi Das University, Lucknow, Uttar Pradesh, India



**Fig. 8** Effect of overexpression of miR-145 and miR-29b on G1 phase arrest in SH-SY5Y cells. Flow cytometry-based PI uptake assay for cell cycle analysis of **a** SH-SY5Y cells transfected with NTC, **b** NTC exposed with pifithrin- $\alpha$  (PFT $\alpha$ ), **c** mimics of miR-29b (Syn-miR-29b), or **d** transfected with mimics of miR-145 or **e** transfected with mimics of

miR-29b and exposed with pifithrin- $\alpha$  (Syn-miR-29b + PFT $\alpha$ ). **f** Bar diagram of G1, S, and G2 phase percentages of SH-SY5Y cells. All values are the mean of three individual experiments. Significant changes are calculated by Student's t test. Pifithrin- $\alpha$  10  $\mu$ M for 24 h