



Original research article

# Fibroblast growth factor signaling mediates progenitor cell aggregation and nephron regeneration in the adult zebrafish kidney

Thomas F. Gallegos<sup>a</sup>, Caramai N. Kamei<sup>a</sup>, Michael Rohly<sup>b</sup>, Iain A. Drummond<sup>a,c,\*</sup>

<sup>a</sup> Massachusetts General Hospital, Nephrology Division, Boston, MA, 02129, USA

<sup>b</sup> Columbus State University, Columbus, 31907, Georgia

<sup>c</sup> Harvard Medical School Department of Genetics, Boston, MA, 02115, USA

## ARTICLE INFO

### Keywords:

Kidney  
Regeneration  
Adult kidney stem cell  
Nephron  
Fibroblast growth factor

## ABSTRACT

The zebrafish kidney regenerates after injury by development of new nephrons from resident adult kidney stem cells. Although adult kidney progenitor cells have been characterized by transplantation and single cell RNA seq, signals that stimulate new nephron formation are not known. Here we demonstrate that fibroblast growth factors and FGF signaling is rapidly induced after kidney injury and that FGF signaling is required for recruitment of progenitor cells to sites of new nephron formation. Chemical or dominant negative blockade of Fgfr1 prevented formation of nephron progenitor cell aggregates after injury and during kidney development. Implantation of FGF soaked beads induced local aggregation of *lhx1a:EGFP* + kidney progenitor cells. Our results reveal a previously unexplored role for FGF signaling in recruitment of renal progenitors to sites of new nephron formation and suggest a role for FGF signaling in maintaining cell adhesion and cell polarity in newly forming kidney epithelia.

## 1. Introduction

During growth or following acute injury, fish generate new kidney nephrons from a latent stem cell pool to expand or regenerate kidney function (Diep et al., 2011; Kamei et al., 2019; Reimschuessel, 2001; Zhou et al., 2010). During regeneration, new nephrons are integrated into the kidney structure by connecting to existing tubules in a process which replicates mesonephric development (Diep et al., 2011; McCampbell et al., 2015). The *lhx1a:EGFP* transgene marks kidney stem cells that have been shown to develop into nephrons after serial transplantation (Diep et al., 2011). In addition, a population of putative quiescent adult kidney stem cells has been identified in single cell RNA seq experiments based on a shared gene expression profile with mammalian metanephric nephron progenitor cells (Tang et al., 2017). After injury, these cells are stimulated to expand in number, migrate, and aggregate as small domed clusters on existing tubules to form nephron precursor structures (Diep et al., 2011; Kamei et al., 2019). New nephron aggregates form specifically on existing distal tubule nephron segments, implying growth and development signals are spatially restricted during kidney regeneration (Diep et al., 2011; Kamei et al., 2019). Recently we reported that Wnt signaling plays a major role in patterning new nephron aggregates and driving cell proliferation in extending nephrons (Kamei et al., 2019) however the identity of early acting signals that recruit

progenitor cells to sites of new nephron formation are unknown.

In this report, we show that multiple fibroblast growth factors (FGFs) are induced following acute kidney injury and that they are expressed specifically in tubular epithelial cells. We show that injury induced FGF signaling is required for nephrogenic aggregate formation and maintenance of aggregate cell-cell adhesion. FGFs are also sufficient to act as stem cell chemotactic factors in uninjured kidneys. Our results suggest that injury-induced FGF signaling provides a spatial, chemotactic cue to organize kidney stem cell-derived new nephrons at sites of integration and function with the kidney tubular architecture.

## 2. Results

### 2.1. FGF signaling is induced in response to acute kidney injury

Acute kidney injury by gentamicin injection in adult zebrafish induced the synchronous formation of many new nephrons marked by *lhx1a* expression at 7 days post-injury (dpi) (Fig. 1F and G). To assess the potential involvement of FGF/receptor tyrosine kinase (RTK) signaling in kidney regeneration we assayed expression of the FGFR/RTK response gene *dusp6*, a MAP kinase phosphatase, under kidney injury conditions. *dusp6* was not expressed in the naïve adult fish kidney (Fig. 1H), but was expressed in the nascent nephrogenic aggregates at 7 dpi, similar to *lhx1a*

\* Corresponding author. Nephrology Division / MGH 149-4.203, 149 13th Street, Charlestown, MA, 02129, USA.

E-mail address: [idrumsd@mg.harvard.edu](mailto:idrumsd@mg.harvard.edu) (I.A. Drummond).

<https://doi.org/10.1016/j.ydbio.2019.06.011>

Received 16 May 2019; Received in revised form 14 June 2019; Accepted 16 June 2019

Available online 18 June 2019

0012-1606/© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

(Fig. 1I), suggesting a role for FGF signaling during kidney regeneration. In histological sections, un-injured fish kidney showed no specific *dusp6* expression by in situ hybridization (Fig. 1J). By 3 days post injury, *dusp6* was broadly expressed throughout the kidney and was enriched in a subset of the gentamicin-injured renal tubules (Fig. 1K). At 5 dpi *dusp6* expression was restricted to single cells and cell clusters in the kidney mesenchyme adjacent to injured tubules (Fig. 1L). By 7 dpi the expression of *dusp6* was further restricted to cell clusters (Fig. 1M), identified as new nephron aggregates by co-immunolabelling with anti-Dusp6 and *lhx1a*:EGFP (Supplemental Figs. 1A–C). The specific FGFR1 inhibitor PD166866 reduced *dusp6* expression to sham-injured levels, confirming *dusp6* in situ labelling reflects localized FGF signaling (Supplemental Fig. 1D). *dusp6* expression after kidney injury suggests an early widespread activation of FGF signaling is followed by a stronger, more focal


signaling in aggregating nephron progenitor cells and new nephron aggregates.

## 2.2. Multiple FGF ligands and receptors are expressed following kidney injury

To identify FGFs and FGF receptors involved in kidney regeneration responses we conducted an RT-PCR screen following acute kidney injury (Fig. 2A). The FGF ligands *fgf3*, *fgf4*, *fgf5*, and *fgf10a* were upregulated as early as 3 days post-injury as were the FGF receptors *fgfr1* and *fgfr4* (Fig. 2A). Interestingly, the *fgfr1* gene, regulator of kidney tubulogenesis and cell adhesion (Trueb et al., 2013; Yang et al., 2016), was transiently down-regulated following injury (Fig. 2A). Other *fgf* and *fgfr* genes were either not detectable in kidney RNA samples or showed only minor changes in expression after injury (Supplemental Figs. 2A–C). Nephrogenic aggregates in regenerating zebrafish kidneys initially form adjacent to distal and collecting segments of renal tubules add *nana* (Diep et al., 2011; Kamei et al., 2015). To assess how Fgf ligand expression relates spatially to newly forming nephrons we assayed expression of the two most abundantly induced Fgf ligands, *fgf4* and *fgf10a*, by in situ hybridization. *fgf4* and *fgf10a* expression was concentrated in collecting ducts at 3 dpi (Fig. 2B and C). *fgf4* and *fgf10a* mRNA was localized to the apical cytoplasmic domain (Fig. 2B and C) as has been observed for other actively translated epithelial mRNAs (Kamei et al., 2019; Moor et al., 2017). To show that the in situ pattern for *fgf4* and *fgf10a* was not due to an "edge effect" and confirm the source of *fgf4* and *fgf10a* was pre-existing kidney epithelial tubules, we separated kidney tissue into intact tubule and interstitial single cell fractions and then assayed *fgf* ligand expression by qRT-PCR. Cell fractions from sham-injured kidneys showed no expression of *fgf* ligands (Fig. 2D and E), consistent with prior analyses of whole kidneys (Fig. 2A). Similarly, interstitial single cell fractions showed no expression of *fgf* ligand mRNA either before or three days after injury (Fig. 2D and E). However, tubule fractions from 3 dpi kidneys showed a 12.8-fold enrichment of *fgf4* and a 8.6-fold induction of *fgf10a* expression (Fig. 2D and E) compared to sham uninjured samples, confirming in situ hybridization results and identifying pre-existing tubules as the source of Fgf ligands induced by injury. The results suggest that Fgf ligand expression may provide a spatial cue for the formation new nephrons at sites of future attachment to the kidney collecting system.


## 2.3. FGF signaling is required for the formation of nephrogenic cell aggregates

To determine whether FGF signaling was required for kidney





(caption on next column)

**Figure 1.** *lhx1a* and *dusp6* expression marks new nephron formation in the regenerating adult zebrafish kidney. (A) An adult zebrafish with the shape of the kidney highlighted. (B) Diagram of the organization of the adult kidney. Nephrogenic stem cells (Nsc) are colored red and appear in the interstitium and along the renal tubules. Segments of the kidney tubular structure are: CD, collecting duct; Dt, distal tubule; Pt, proximal tubule; Gl, glomerulus. (C–E) Diagram of stem cell based nephrogenesis. Stem cells accumulate on the Dt segment and form aggregates of cells (C). The aggregate elongates into a tubular epithelium and connects to the Dt (D). The new nephron fully differentiates to form a functional nephron (E). (F) Whole mount in situ hybridization of control uninjured adult kidneys show no expression of *lhx1a*. (G) Seven days after gentamicin injury *lhx1a* was induced in aggregates of nephrogenic cells, red arrowheads. (H) *dusp6* was not expressed in controls but was induced in nephrogenic cells and cell aggregates at 7 dpi, red arrowheads (I). Scale Bars F–I are 500  $\mu$ m. (J) In histological sections, *dusp6* was not expressed in uninjured control kidneys. J–M empty arrowheads indicate *dusp6* negative tubules (K) *dusp6* was induced by kidney injury in the interstitium and some renal tubules at 3 dpi, in aggregating single cells adjacent to kidney tubules at 5 dpi (L), and highly expressed in nephrogenic aggregates at 7 dpi (M). In K–M, representative *dusp6* expressing structures are indicated with solid arrowheads. Scale Bars J–M are 50  $\mu$ m.



**Fig. 2. RT-PCR analysis identifies injury modulated FGF signaling molecules.** (A) RT-PCR assays revealed that *fgf3*, *fgf4*, *fgf5*, *fgf10a* and the receptors *fgfr1*, *fgfr4* were induced post-injury while the kinase domain deficient *fgfr11* gene exhibited reduced expression after injury. Histological sections of whole mount in situ hybridization 7 dpi injured kidneys show *fgf4* (B) and *fgf10a* (C) were expressed in the collecting duct epithelium. (B–C) Scale bars are 20 μm. In kidney cell fractionation experiments, *fgf4* (D) and *fgf10* (E) were induced in isolated tubules and not in single interstitial cells at 3dpi compared to sham injured kidneys. qPCR represents 3 biological replicates. Comparisons were made by Kruskal-Wallis test. ns indicates not-significant. Data are expressed as mean ± SD.




**Fig. 3. Inhibition of Fgf signaling prevents new nephron development.** (A) Injured zebrafish generate *lhx1a* + nephrogenic aggregates at 7 dpi by whole mount in situ hybridization. (B) Treatment with the FGFR1 inhibitor PD166866 blocks the appearance of *lhx1a* + nephrogenic aggregates in whole mount kidneys. (C) Quantification of *lhx1a* + nephrogenic aggregates in multiple whole mount kidneys demonstrates PD166866 completely blocks kidney regeneration. N = 7 fish per condition. Error bars indicate ±SD. p = 0.01 by Mann-Whitney T-test. (D) qRT-PCR quantitation of *lhx1a* expression shows PD166866 treatment blocks *lhx1a* expression post-injury. N = 6 fish per condition. Error bars indicate ±SD. \* indicates p = 0.02 by Kruskal-Wallis test. (E) Control non-heat shocked *Tg(hsp70:dnfgfr1)* transgenic fish show *lhx1a* + new nephron aggregates 7 days following gentamicin injury. (F) Heat shock induction of the dominant-negative Fgfr1 receptor blocked the development of *lhx1a* + new nephron aggregates. (G) Quantification of *lhx1a* + nephrogenic aggregates in multiple whole mount kidneys demonstrates that expression of the dominant-negative Fgfr1 receptor completely blocks kidney regeneration. Differences in injury responses (number of nephrogenic aggregates in A vs E) reflect strain dependent differences in sensitivity to gentamicin injury. N = 4 control and N = 5 heat shocked fish. Error bars indicate ±SD. p = 0.03 by Mann-Whitney T-test. In panels A,B,E,F scale bars are 100 μm.

regeneration, we blocked the FGF signaling pathway both pharmacologically and genetically by inactivating the *Fgfr1* receptor. In control regenerating kidneys, *lhx1a* + nephrogenic aggregates were present throughout the kidney at 7 dpi (Fig. 3A). Pharmacological inhibition of *Fgfr1* signaling using the compound PD166866 resulted in a nearly complete failure to form *lhx1a* + nephrogenic aggregates (Fig. 3B) with the number of *lhx1a* + aggregates reduced from an average of 77.9 per kidney to less than 2.4 ( $p = 0.01$ ) per kidney at 7 dpi (Fig. 3C). By qPCR, acute injury induced *lhx1a* expression 9.0-fold ( $p = 0.02$ ) relative to sham-injured kidneys (Fig. 3D). Treatment with PD166866 reduced *lhx1a* RNA expression to the level of sham-injured kidneys (Fig. 3D). Genetic perturbation of FGF signaling was performed using the *Tg(hsp70l:dnfgfr1-EGFP)* zebrafish line, which expresses a dominant negative *Fgfr1* receptor when heat shocked (Lee et al., 2005). Consistent with chemical inhibition of FGF signaling, dominant negative repression of *Fgfr* signaling blocked the appearance of *lhx1a* + nephrogenic aggregates normally induced by AKI (Fig. 3E, F). The severity of the loss of FGF signaling was equivalent to pharmacological inhibition, with the average number of nephrogenic aggregates reduced from 9.5 in injured kidneys to 0.2 ( $p = 0.03$ ) in injured and heat shocked *Tg(hsp70l:dnfgfr1-EGFP)* 7dpi kidneys (Fig. 3G). To determine whether inhibition of FGF signaling was actually preventing aggregate formation as opposed to blocking expression of *lhx1a*, we sectioned injured kidneys for histological analysis. Nephrogenic aggregates typically appear as dome-shaped basophilic cell clusters abutting kidney tubules (Supplemental Fig. 3A). In multiple *Fgfr* signaling inhibited kidneys we found no morphological evidence of nephrogenic aggregates (Supplemental Fig. 3B). The results suggest that *Fgf* signaling is not simply required to induce *lhx1a* gene expression in new nephrons but rather is required for the formation of nephrogenic aggregates as a prelude to nephron regeneration.

#### 2.4. FGF signaling is required for progenitor cell condensation at sites of nephron formation

The absence of nephrogenic aggregates in *Fgfr* inhibited kidneys suggested that progenitor cells may be lost or may fail to aggregate in the absence of FGF signals. To visualize the response of nephrogenic progenitor cells to *Fgfr* inhibition, we imaged *lhx1a:EGFP* + cells after blocking *Fgf* signaling during different time windows of regeneration. Inhibition of *Fgf* signaling from 3 to 5 dpi followed by imaging at 5 dpi delayed the formation of condensed *lhx1a:EGFP* + cell aggregates and caused the persistence of lamellipodia on clusters of *lhx1a:EGFP* + cells (Fig. 4 A, B). To better resolve cell scale and nuclei in the GFP + cell population, individual image channels are presented for Fig. 4B in Supplemental Fig. 4. Inhibition of *Fgf* signaling from 3 to 7 dpi resulted in complete failure of *lhx1a:EGFP* + cells to aggregate with loss of compact nephrogenic aggregates and the appearance of broadly distributed, single stellate GFP + cells and disorganized cell clumps (Fig. 4C and D). These data indicate that in the absence of *Fgf* signaling, *lhx1a:EGFP* + progenitor cells persist, but fail to migrate or aggregate at sites of regenerative new nephron formation. To test whether *Fgf* signaling was also required for maintenance of progenitor cell aggregates we inhibited *Fgf* signaling from 5 to 6 dpi after regenerative aggregates had formed as well as during mesonephric development, where similar *lhx1a:EGFP* + cell aggregates precede nephron differentiation (Diep et al., 2011, 2015). Nephrogenic cells form a rosette with a central apical ring, predicting the site of apical tight junctions in the new tubule lumen which extends in a proximal direction as the new nephron differentiates (Fig. 4E, inset). Short term *Fgfr* inhibition with PD166866 from 5 to 6 dpi induced the appearance of filopodia and lamellopodia which formed first on the most proximal cells which also lose apical *zo-1* localization (Fig. 4F). Ultimately polarity and cell aggregation is lost as cells appear to disperse randomly. In uninjured juvenile zebrafish, characteristic dome-shaped rosettes of *lhx1a:EGFP* + nephrogenic cells were numerous and prominent (Fig. 4G). Treatment with PD166866 for seven days disrupted pre-existing cell aggregates and promoted the appearance of



**Fig. 4. Disruption of nephrogenic cell aggregate formation in *Fgf* signaling inhibited kidneys.** (A) Adult *Tg(lhx1a:EGFP)* fish at 5 dpi exhibit condensed, dome-shaped cell aggregates on distal tubules. (B) Treatment of regenerating fish from 3 to 5 days post injury with the *Fgfr1* inhibitor PD166866 followed by imaging at 5 dpi showed delayed formation of tight adherent nephrogenic cell domes and the appearance of cell lamellopodia (arrowheads in B). (C–D) Tight, condensed *lhx1a:EGFP* + nephrogenic cell aggregates seen in control injured kidneys (C) did not form in fish treated with PD166866 from 3 to 7 dpi (D); instead *lhx1a:EGFP* + cells remained stellate-shaped and disorganized. (E) 6 dpi *lhx1a:EGFP* + nephrogenic cell aggregates display a rosette shape with a *zo-1* positive apical lumen (inset) extending in the direction of nephron growth (red arrowhead; proximal). (F) 24 h PD166866 treatment from day 5–6 induces filopodia and lamellopodia formation (white arrowheads) and reduction in apical *zo-1* localization (white arrow) with a ring of remnant *zo-1* only in distal cells (red arrowhead). (G) Juvenile *Tg(lhx1a:EGFP)* fish at 42 dpf produce nephrogenic aggregates in their developing kidneys identical to injury-induced, dome-shaped aggregates (arrowhead in G). (H) Treatment of juveniles from 35 to 42 dpf with PD166866 results in the loss of domed aggregates and appearance of single cells and stellate cell clusters (arrowheads in H), indicating a loss of adhesion. Scale bars in Z-projections (A–F) are 10  $\mu$ m and 100  $\mu$ m in (G, H).

dispersed single cells and stellate shaped clusters (Fig. 4H). Our results suggest that Fgf signaling is required for both cell recruitment and maintenance of cell adhesion and polarity in nephron progenitor cell rosettes.


To assess whether Fgf signaling was sufficient to induce kidney progenitor cell recruitment we implanted FGF soaked beads into an uninjured adult *lhx1a:EGFP* kidney in explant culture. After 24 h of *in vitro* culture, *lhx1a:EGFP* + progenitor cells were found to preferentially associate with FGF beads compared to BSA bead controls (Fig. 5A–C; D). FGF10, and to a lesser extent FGF4, induced recruitment of cells to the beads with FGF10 causing most cells to spread directly on the bead surface (Fig. 5C). Taken together, the data indicate that injury-induced Fgf expression in kidney tubules is sufficient and required for progenitor cell recruitment and rosette formation at sites of new nephron formation in regenerating zebrafish kidneys.

### 3. Discussion

The remarkable capacity of the zebrafish kidney to form new nephrons after injury is mediated by mobilization and differentiation of a resident population of *lhx1a:GFP*<sup>+</sup>, adult kidney stem/progenitor cells (Diep et al., 2011; Kamei et al., 2019). While serial transplantation studies provide strong support for *lhx1a:GFP* + cells acting as kidney progenitor cells (Diep et al., 2011), the signals that mobilize these cells and drive nephron differentiation were not known. We recently reported that Wnt signaling patterns new nephron aggregates into zones of cell proliferation and tubule invasion (Kamei et al., 2019). Our principle finding here is that acute kidney injury in adult zebrafish initiates expression of multiple FGF ligands and receptors which have an early and essential function in recruiting kidney progenitors to sites of new

nephron formation. In addition we find that continued FGFR signaling is required to maintain cell adhesion and polarity of nephron progenitor aggregates, revealing a role for FGF signaling in the earliest stages of mesenchyme to epithelial transformation that drives nephron formation. A similar function for FGF signaling in maintaining polarized epithelial rosettes is observed in the formation of hair cells in the zebrafish lateral line organ (Ma and Raible, 2009). Also, FGF signaling has been shown to mediate apico-basal polarity of the Drosophila mesoderm during gastrulation (Sun and Stathopoulos, 2018). Our findings establish FGF signaling as a key factor in cell polarization and define roles for FGF 4 and 10a as chemotactic factors in the context of kidney regeneration.

During mammalian kidney organogenesis, fibroblast growth factors 1, 2, 7, 8, 9, 10, 11, 12, 20 and receptors FGFR1, FGFR2 and FGFR11 are expressed in temporally and spatially restricted patterns consistent with discrete cell signaling roles (Barak et al., 2012; Bates, 2011; Brown et al., 2011; Gerber et al., 2009; Grieshammer et al., 2005; Ohuchi et al., 2000; Okazawa et al., 2015; Perantoni et al., 2005; Poladia et al., 2006; Qiao et al., 1999; Trueb et al., 2013). Branching morphogenesis of the ureteric epithelium requires FGF7 and FGF10 for full development of the collecting system (Ohuchi et al., 2000; Qiao et al., 1999). "Cap mesenchyme", a population mammalian nephron progenitor cells, is maintained by the activity of FGF9 and FGF20 which drive progenitor cell self-renewal at the periphery of the developing kidney (Barak et al., 2012; Brown et al., 2011). Both branching morphogenesis and progenitor self-renewal ultimately define nephron number and kidney organ size, with defects in FGF signaling leading to kidney hypodysplasia (Walker et al., 2016). FGF8 is expressed in the newly forming nephron epithelium and is required for survival and development of tubular epithelia (Grieshammer et al., 2005; Perantoni et al., 2005). FGF receptors acting in the mesenchyme (FGFR1/2 (Poladia et al., 2006)) and ureteric



**Fig. 5. Exogenous FGF beads are sufficient to recruit progenitor cells.** A) Stem cells marked by the transgene *lhx1a:eGFP* are not attracted to BSA soaked agarose beads after 24 h of culture. (B) Beads soaked with Fgf4 are more closely surrounded by renal stem cells, indicated by a decrease in cell to bead distance from an average of 28.8  $\mu\text{m}$ –17.75  $\mu\text{m}$ . (C) Beads soaked with FGF10 strongly attract renal stem cells which accumulate adjacent to the bead at a mean distance of 3.42  $\mu\text{m}$ . (D) Quantification of cell to bead distances after 24 h of culture demonstrate a chemoattractive role of FGF ligands on the recruitment renal stem cells. 3 biological replicates per condition. Error bars indicate  $\pm$ SD. Comparisons to BSA beads were made by Kruskal-Wallis test. ns indicates non-significant, starred comparison indicates  $p = 0.008$ . In Z-projection panels (A–C) scale bar is 25  $\mu\text{m}$ .

epithelium (FGFR2 (Zhao et al., 2004)) control multiple steps of kidney development with double mutants exhibiting a severe kidney aplasia phenotype (Poladia et al., 2006). Our results suggest a new role for epithelium-expressed FGFs as chemotactic factors recruiting and/or maintaining kidney mesenchymal cells at sites of new nephron formation. We found that zebrafish *fgf4* and *fgf10a* were expressed in kidney epithelia after injury and were sufficient, when applied ectopically in ligand soaked beads, to recruit *lhx1a*:GFP + kidney progenitor cells in uninjured kidneys. This finding is consistent with prior studies (Tang et al., 2017) where we found putative *six2a*, *eya2*, *osr1* + adult kidney progenitor cells show a roughly 16-fold enriched expression of *fgfr1a* and *fgfr2* over total kidney, suggesting adult kidney progenitor cells in uninjured kidneys are sensitive to Fgf. Although progenitor cell rosettes were not observed in our Fgf bead experiments this may be due to the relatively short, *in vitro* incubation period we used. The absence of new nephron cell aggregates in Fgfr inhibited zebrafish kidneys further argues that FGF signaling is required for progenitor cell recruitment. While the evidence for Fgf involvement in cell recruitment to tubules is strong, there must be additional factors controlling the size and location of nephron aggregates since progenitor cells are not spread evenly across Fgf-expressing epithelial tubules and instead form spaced cell clusters of uniform size.

Mammalian kidney cap mesenchyme, a cell population orthologous to the zebrafish *lhx1a*:GFP + nephron progenitor cells (Diep et al., 2011; Tang et al., 2017), is highly motile and dynamic during nephrogenesis (Combes et al., 2016; Lindstrom et al., 2018). In time lapse studies individual cells are observed to make transient filopodial contacts with the ureteric epithelium and then detach and migrate among other cap mesenchyme and stromal cells (Combes et al., 2016; O'Brien et al., 2018). This behavior suggests that cells destined to generate nephrons maintain a close but dynamic association with the inducing ureteric epithelium by a combination of attractive and repulsive cues (Combes et al., 2016; O'Brien et al., 2018). In mice, deficiency in *Robo2*, a putative repulsive cue in the ureteric tip, results in an ectopic association of cap mesenchyme with ureteric stalks (Wainwright et al., 2015), implying persistent cell association with the growing collecting system. Mutation in the ureteric bud tip morphogen *Wnt11* results in loss of stable attachment of mesenchymal cells to the UB tips and dispersal of cells from the cap mesenchyme niche (O'Brien et al., 2018). Our studies suggest FGFs may also act as chemoattractants during kidney development and regeneration. FGFs have broadly conserved roles in cell migration and adhesion (Bae et al., 2012; Boulet and Capecchi, 2012; Chi et al., 2006; Dalle Nogare and Chitnis, 2017; Hacohen et al., 1998; Li and Muneoka, 1999; Park et al., 1998; Roussigne et al., 2018). Blocking FGF signaling in our experiments allowed *lhx1a*:GFP + progenitor cells to disperse away from the distal nephron epithelium while FGF4 or FGF10 soaked beads exerted a strong chemotactic influence on kidney progenitor cells. Taken together, the data suggests that attractive cues like FGFs may maintain progenitor cells in close apposition to an inducing epithelium to promote efficient and localized production of new nephrons. Genetic analysis of this phenomenon in zebrafish will require new conditional *fgf* mutants as knockdown of *fgf4* results in strong left-right asymmetry defects (Yamauchi et al., 2009), *fgf10a* is required for fin development (Norton et al., 2005) and *fgf8* is essential for hindbrain patterning (Reifers et al., 1998).

Candidate mammalian FGF ligands known to stimulate migration and also expressed in the ureteric epithelium include FGF2, and FGF9 (Chi et al., 2006; Joannes et al., 2016; Lin et al., 2009; Song et al., 2016; Wang et al., 2017). Although both ligands are more commonly associated with mesenchymal cell survival and proliferation (Barak et al., 2012; Drummond et al., 1998), FGF ligands can have multiple functions in different cellular contexts. FGF receptor binding activates multiple signal transduction pathways including ERK and PI3K/Akt (Ornitz and Itoh, 2015). ERK activation is associated with cell survival and proliferation while PI3K signals cell survival and cell motility (Francavilla et al., 2013; Ornitz and Itoh, 2015). In HeLa cells FGF10 signals cell motility by inducing the

rapid phosphorylation of tyrosine 734 on FGFR2b, recruiting PI3K and SH3BP4 and leading to cell migration (Francavilla et al., 2013). Different FGFR splice forms may also lead to differential activation of signaling pathways (Sharma et al., 2015). Finally, expression of different negative feedback inhibitors (sprouty/ERK inhibition vs. Sprad/PI3K inhibition) can influence the cellular response to FGF receptor binding (Nutt et al., 2001; Sivak et al., 2005). Future studies should reveal how the cellular context of FGF signaling during kidney development and regeneration promotes cell aggregation at sites of new nephron formation.

## 4. Experimental procedures

### 4.1. Animals

Adult wild type TuAB and Tg(*hsp70l:dnfgfr1*-EGFP) zebrafish approximately one year of age and adult and juvenile zebrafish carrying the *lhx1a:eGFP* transgene were used for this study (Lee et al., 2005; Swanhart et al., 2010). Heat shocked animals were injured under standard protocol. After confirmation, they were moved to a rack within the fish facility in which water temperature is raised to 37 °C for 3 h per day. Control fish were maintained under standard maintenance conditions after injury. Experimentally induced acute kidney injury procedures were performed in accordance with an MGH approved IACUC protocol.

### 4.2. Chemicals

Acute kidney injury was induced by intraperitoneal injection of gentamicin (Sigma, Saint Louis, MO) at 80 mg/kg (2 mg/ml stock in PBS at 10 µl per 0.25g body mass) as previously described (Kamei et al., 2015). Sham injury was performed by injecting an equivalent volume of PBS intraperitoneally in place of gentamicin solution. After confirmation of kidney injury by visual identification of cast excretion at 1 day post injury (Kamei et al., 2015), *Fgfr1* signaling was inhibited by adding PD166866 (10 µM, Millipore Sigma, Burlington, MA) in DMSO (0.04% final concentration) to fish system water (Panek et al., 1998). Fish water was changed daily, with fresh DMSO or PD166866 added at the same concentration. The FGF inhibition protocol was maintained until animals were sacrificed at 7 days post-injury, except where otherwise noted.

### 4.3. In situ hybridization

In situ hybridizations were performed on adult kidney at the time post-injury described. Briefly, Digoxigenin labeled RNA probes for *lhx1a* and *dup6* were prepared from cDNA clones, while probes for *fgf4* and *fgf10a* were prepared by transcription from a PCR fragment amplified from genomic DNA as described previously (Thisse et al., 2004). Quantitation of *lhx1a* positive aggregates was performed on 5X images of whole mount tissue on the widest section of the kidney. Positive aggregates were marked and quantified in ImageJ. For histological sections, tissue from in situ hybridizations was embedded in JB4 blocks (Polysciences, Inc., Warrington, PA) and sectioned to a thickness of 10 µm. Sections were mounted in Permount (Sigma, Saint Louis, MO) and imaged on a Nikon Eclipse E800 microscope.

### 4.4. Kidney tubule isolation

Adult TuAB trunk kidneys (two per sample) were incubated at room temperature with 0.5 mg/ml Collagenase Type A (Sigma, Saint Louis, MO) in HBSS (Walkersville, MD) for 30 min on a rocking stage at room temperature. The tissue was vigorously vortexed for 3 min to fragment the tissue. The tube was left to stand for 5 min to allow the tubules to separate from single cells by gravity. The supernatant was passed through a 40 µm filter (Corning, Corning, NY), pelleted and lysed into Qiazol (Qiagen, Venlo, Netherlands) for RNA preparation. The tubular fraction was washed with HBSS and gravity collected again for 5 min, then lysed into Qiazol (Qiagen, Venlo, Netherlands) for RNA preparation.

Enrichment for hematopoietic cell types in single cell fraction RNA and kidney tubular epithelial cells in tubule fraction RNA was confirmed by qRT-PCR for *gata1* (single cells) and *cdh17* (tubule cells) (Supplemental Fig. 5).

#### 4.5. Immunofluorescence and microscopy

Tissue was fixed in 4% formaldehyde for 3 h then stained for indirect immunofluorescence. Primary antibodies used were chick anti-GFP (Abcam, Cambridge, UK) and mouse anti-Dusp6 (Sigma, Saint Louis, MO), and secondary antibodies were anti-chick IgG and anti-mouse IgG linked to Alexafluor 488 and 546 (Life Technologies, Carlsbad, CA). Nuclei were stained with Hoechst 33342 (Invitrogen, Carlsbad, CA). Confocal imaging was performed on a Zeiss LSM 510 microscope.

#### 4.6. Reverse-transcriptase PCR and qPCR assays

For screening expression of FGF related genes, RNA was harvested from three pooled kidney samples from adult TuAB kidneys, either un-injured or at the indicated time post-gentamicin induced AKI. cDNA was synthesized using Quantitect Reverse Transcription kit reagents (Qiagen, Venlo, Netherlands). Primers for initial RTPCR screening are presented in Supplemental Table 1. For qPCR, RNA harvested from individual kidneys was performed at the indicated time after gentamicin injury. RNA was harvested with Qiagen RNeasy columns and cDNA libraries were prepared using SuperScript IV (Invitrogen, Carlsbad, CA). qRT-PCR was performed on a StepOnePlus thermal cycler using Sybr Green reagents (Applied Biosystems, Foster City, CA). qRT-PCR results are expressed as fold change by  $\Delta\Delta C_T$ , with analysis described previously (Willems et al., 2008). qRT-PCR primers used were as follows: *gapdh*-forward 5'-cggagcaccaggtgtgtcca-3', *gapdh*-reverse 5'-agcaataccagcaccagcgtc-3', *lhx1a*-forward 5'-gctgcgagtgcaaatgtaac-3', *lhx1a*-reverse 5'-catcatgtgaagcagctcag-3', *dusp6*-forward 5'-tctatctcga ggggtgcttca-3', *dusp6*-reverse 5'-actcgtatcgcgaggagta-3', *fgf4*-forward 5'-ccggcgtacacaacgaaac-3', *fgf4*-reverse 5'-gaactgctcagatccgtaaagc-3', *fgf10a*-forward 5'-cgatcgtacagctacactcga-3', *fgf10a*-reverse 5'-agcttcagctcaatgccgaa-3'. Dendrograms of Fgf ligands and receptors were prepared using Phylip 3.695 drawtree program (Felsenstein, 1989).

#### 4.7. Progenitor cell motility assay

Kidneys were isolated from un-injured adult Tg(*lhx1a:eGFP*) fish for explant culture. The tissue was placed atop a polycarbonate filter, on a media equilibrated agarose raft. To deliver growth factors Affi-Gel Blue Gel (Bio-Rad, Hercules, CA) beads were incubated in mouse FGF4 or FGF10 (R&D Systems, Minneapolis, MN) or BSA (Sigma, Saint Louis, MO) at 500 ng/ml, then washed in PBS before implantation. Several beads were placed upon the widest part of the kidney by capillary action from a 1  $\mu$ l aspiration of washed beads. Cultures were maintained at 28 °C for 24 h before formaldehyde fixation and immunostaining. Culture media was prepared following previously described zebrafish explant culture (Cao and Poss, 2016), with amendments for culture without carbon dioxide supplementation. The culture medium was L-15 based (Thermo-fisher, Waltham, MA), with an addition of HEPES pH7.0–10 mM.

#### 4.8. Statistical analysis

All quantifications are represented with error bars indicating SD. All data represent three or more biological replicates. Individual comparisons were Mann-Whitney T-tests. Multiple comparisons were made using Kruskal-Wallis tests. All statistical analyses were performed and graphs prepared using Prism7 (GraphPad).

#### Author contributions

The project was conceived and designed by I.A.D and T.F.G; Experiments were performed by T.F.G, C.N.K, and M.R. The paper was written

by T.F.G and I.A.D.

#### Acknowledgements

The authors declare no competing interests. The authors thank Dr. Kenneth Poss for providing the Tg(*hsp70:dn-fgfr1*) transgenic line. We thank Dr. Neil Hukriede for the Tg(*lhx1a:eGFP*) transgenic line. This work was supported by NIH grant 5UH3DK107372-04 and grants from the Harvard Stem Cell Institute to I.A.D.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ydbio.2019.06.011>.

#### References

- Bae, Y.K., Trisnadi, N., Kadam, S., Stathopoulos, A., 2012. The role of FGF signaling in guiding coordinate movement of cell groups: guidance cue and cell adhesion regulator? *Cell Adhes. Migrat.* 6, 397–403.
- Barak, H., Huh, S.H., Chen, S., Jeanpierre, C., Martinovic, J., Parisot, M., Bole-Feysot, C., Nitschke, P., Salomon, R., Antignac, C., Ornitz, D.M., Kopan, R., 2012. FGF9 and FGF20 maintain the stemness of nephron progenitors in mice and man. *Dev. Cell* 22, 1191–1207.
- Bates, C.M., 2011. Role of fibroblast growth factor receptor signaling in kidney development. *Pediatr. Nephrol.* 26, 1373–1379.
- Boulet, A.M., Capecci, M.R., 2012. Signaling by FGF4 and FGF8 is required for axial elongation of the mouse embryo. *Dev. Biol.* 371, 235–245.
- Brown, A.C., Adams, D., de Caestecker, M., Yang, X., Friesel, R., Oxburgh, L., 2011. FGF/EGF signaling regulates the renewal of early nephron progenitors during embryonic development. *Development* 138, 5099–5112.
- Cao, J., Poss, K.D., 2016. Explant culture of adult zebrafish hearts for epicardial regeneration studies. *Nat. Protoc.* 11, 872–881.
- Chi, L., Itaranta, P., Zhang, S., Vainio, S., 2006. Sprouty2 is involved in male sex organogenesis by controlling fibroblast growth factor 9-induced mesonephric cell migration to the developing testis. *Endocrinology* 147, 3777–3788.
- Combes, A.N., Lefevre, J.G., Wilson, S., Hamilton, N.A., Little, M.H., 2016. Cap mesenchyme cell swarming during kidney development is influenced by attraction, repulsion, and adhesion to the ureteric tip. *Dev. Biol.* 418, 297–306.
- Dalle Nogare, D., Chitnis, A.B., 2017. A framework for understanding morphogenesis and migration of the zebrafish posterior Lateral Line primordium. *Mech. Dev.* 148, 69–78.
- Diep, C.Q., Ma, D., Deo, R.C., Holm, T.M., Naylor, R.W., Arora, N., Wingert, R.A., Bollig, F., Djordjevic, G., Lichman, B., Zhu, H., Ikenaga, T., Ono, F., Englert, C., Cowan, C.A., Hukriede, N.A., Handin, R.I., Davidson, A.J., 2011. Identification of adult nephron progenitors capable of kidney regeneration in zebrafish. *Nature* 470, 95–100.
- Diep, C.Q., Peng, Z., Ukah, T.K., Kelly, P.M., Daigle, R.V., Davidson, A.J., 2015. Development of the zebrafish mesonephros. *Genesis* 53, 257–269.
- Drummond, I.A., Mukhopadhyay, D., Sukhatme, V.P., 1998. Expression of fetal kidney growth factors in a kidney tumor line: role of FGF2 in kidney development. *Exp. Nephrol.* 6, 522–533.
- Felsenstein, J., 1989. PHYLIP - phylogeny inference package (version 3.2). *Cladistics* 5, 164–166.
- Francavilla, C., Rigbolt, K.T., Emdal, K.B., Carraro, G., Vernet, E., Bekker-Jensen, D.B., Streicher, W., Wikstrom, M., Sundstrom, M., Bellusci, S., Cavallaro, U., Blagoev, B., Olsen, J.V., 2013. Functional proteomics defines the molecular switch underlying FGF receptor trafficking and cellular outputs. *Mol. Cell* 51, 707–722.
- Gerber, S.D., Steinberg, F., Beyeler, M., Villiger, P.M., Trueb, B., 2009. The murine Fgfr1 receptor is essential for the development of the metanephric kidney. *Dev. Biol.* 335, 106–119.
- Grieshammer, U., Cebrian, C., Ilagan, R., Meyers, E., Herzlinger, D., Martin, G.R., 2005. FGF8 is required for cell survival at distinct stages of nephrogenesis and for regulation of gene expression in nascent nephrons. *Development* 132, 3847–3857.
- Hacohen, N., Kramer, S., Sutherland, D., Hiroimi, Y., Krasnow, M.A., 1998. Sprouty encodes a novel antagonist of FGF signaling that patterns apical branching of the *Drosophila* airways. *Cell* 92, 253–263.
- Joannes, A., Brayer, S., Besnard, V., Marchal-Somme, J., Jaillet, M., Mordant, P., Mal, H., Borie, R., Crestani, B., Mailloux, A.A., 2016. FGF9 and FGF18 in idiopathic pulmonary fibrosis promote survival and migration and inhibit myofibroblast differentiation of human lung fibroblasts in vitro. *Am. J. Physiol. Lung Cell Mol. Physiol.* 310, L615–L629.
- Kamei, C.N., Gallegos, T.F., Liu, Y., Hukriede, N., Drummond, I.A., 2019. Wnt Signaling Mediates New Nephron Formation during Zebrafish Kidney Regeneration, vol. 146. *Development*.
- Kamei, C.N., Liu, Y., Drummond, I.A., 2015. Kidney regeneration in adult zebrafish by gentamicin induced injury. *J. Vis. Exp.*, e51912.
- Lee, Y., Grill, S., Sanchez, A., Murphy-Ryan, M., Poss, K.D., 2005. Fgf signaling instructs position-dependent growth rate during zebrafish fin regeneration. *Development* 132, 5173–5183.
- Li, S., Muneoka, K., 1999. Cell migration and chick limb development: chemotactic action of FGF-4 and the AER. *Dev. Biol.* 211, 335–347.

- Lin, Y., Chen, L., Lin, C., Luo, Y., Tsai, R.Y., Wang, F., 2009. Neuron-derived FGF9 is essential for scaffold formation of Bergmann radial fibers and migration of granule neurons in the cerebellum. *Dev. Biol.* 329, 44–54.
- Lindstrom, N.O., De Sena Brandine, G., Tran, T., Ransick, A., Suh, G., Guo, J., Kim, A.D., Parvez, R.K., Ruffins, S.W., Rutledge, E.A., Thornton, M.E., Grubbs, B., McMahon, J.A., Smith, A.D., McMahon, A.P., 2018. Progressive recruitment of mesenchymal progenitors reveals a time-dependent process of cell fate acquisition in mouse and human nephrogenesis. *Dev. Cell* 45, 651–660 e654.
- Ma, E.Y., Raible, D.W., 2009. Signaling pathways regulating zebrafish lateral line development. *Curr. Biol.* 19, R381–R386.
- McCampbell, K.K., Springer, K.N., Wingert, R.A., 2015. Atlas of cellular dynamics during zebrafish adult kidney regeneration. *Stem Cell. Int.* 2015, 547636.
- Moor, A.E., Golan, M., Massasa, E.E., Lemze, D., Weizman, T., Shenhav, R., Baydatch, S., Mizrahi, O., Winkler, R., Golani, O., Stern-Ginossar, N., Itzkovitz, S., 2017. Global mRNA polarization regulates translation efficiency in the intestinal epithelium. *Science* 357, 1299–1303.
- Norton, W.H., Ledin, J., Grandel, H., Neumann, C.J., 2005. HSPG synthesis by zebrafish Ext2 and Extl3 is required for Fgf10 signalling during limb development. *Development* 132, 4963–4973.
- Nutt, S.L., Dingwell, K.S., Holt, C.E., Amaya, E., 2001. Xenopus Sprout2 inhibits FGF-mediated gastrulation movements but does not affect mesoderm induction and patterning. *Genes Dev.* 15, 1152–1166.
- O'Brien, L.L., Combes, A.N., Short, K.M., Lindstrom, N.O., Whitney, P.H., Cullen-McEwen, L.A., Ju, A., Abdelhalim, A., Michos, O., Bertram, J.F., Smyth, I.M., Little, M.H., McMahon, A.P., 2018. Wnt11 directs nephron progenitor polarity and motile behavior ultimately determining nephron endowment. *Elife* 7.
- Ohuchi, H., Hori, Y., Yamasaki, M., Harada, H., Sekine, K., Kato, S., Itoh, N., 2000. FGF10 acts as a major ligand for FGF receptor 2 IIIb in mouse multi-organ development. *BBRC (Biochem. Biophys. Res. Commun.)* 277, 643–649.
- Okazawa, M., Murashima, A., Harada, M., Nakagata, N., Noguchi, M., Morimoto, M., Kimura, T., Ornitz, D.M., Yamada, G., 2015. Region-specific regulation of cell proliferation by FGF receptor signaling during the Wolffian duct development. *Dev. Biol.* 400, 139–147.
- Ornitz, D.M., Itoh, N., 2015. The Fibroblast Growth Factor Signaling Pathway, vol. 4. Wiley Interdiscip Rev Dev Biol, pp. 215–266.
- Panek, R.L., Lu, G.H., Dahring, T.K., Batley, B.L., Connolly, C., Hamby, J.M., Brown, K.J., 1998. In vitro biological characterization and antiangiogenic effects of PD 166866, a selective inhibitor of the FGF-1 receptor tyrosine kinase. *J. Pharmacol. Exp. Ther.* 286, 569–577.
- Park, W.Y., Miranda, B., Lebeche, D., Hashimoto, G., Cardoso, W.V., 1998. FGF-10 is a chemotactic factor for distal epithelial buds during lung development. *Dev. Biol.* 201, 125–134.
- Perantoni, A.O., Timofeeva, O., Naillat, F., Richman, C., Pajni-Underwood, S., Wilson, C., Vainio, S., Dove, L.F., Lewandoski, M., 2005. Inactivation of FGF8 in early mesoderm reveals an essential role in kidney development. *Development* 132, 3859–3871.
- Poladia, D.P., Kish, K., Kutay, B., Hains, D., Kegg, H., Zhao, H., Bates, C.M., 2006. Role of fibroblast growth factor receptors 1 and 2 in the metanephric mesenchyme. *Dev. Biol.* 291, 325–339.
- Qiao, J., Uzzo, R., Obara-Ishihara, T., Degenstein, L., Fuchs, E., Herzlinger, D., 1999. FGF-7 modulates ureteric bud growth and nephron number in the developing kidney. *Development* 126, 547–554.
- Reifers, F., Bohli, H., Walsh, E.C., Crossley, P.H., Stainier, D.Y., Brand, M., 1998. Fgf8 is mutated in zebrafish acerebellar (ace) mutants and is required for maintenance of midbrain-hindbrain boundary development and somitogenesis. *Development* 125, 2381–2395.
- Reimschuessel, R., 2001. A Fish Model of Renal Regeneration and Development, vol. 42. ILAR journal/National Research Council, Institute of Laboratory Animal Resources, pp. 285–291.
- Roussigne, M., Wei, L., Tsingos, E., Kuchling, F., Alkobtawi, M., Tsalavouta, M., Wittbrodt, J., Carl, M., Blader, P., Wilson, S.W., 2018. Left/right asymmetric collective migration of parapineal cells is mediated by focal FGF signaling activity in leading cells. *Proc. Natl. Acad. Sci. U. S. A.* 115, E9812–E9821.
- Sharma, R., Beer, K., Iwanov, K., Schmohl, F., Beckmann, P.I., Schroder, R., 2015. The single fgf receptor gene in the beetle *Tribolium castaneum* codes for two isoforms that integrate FGF8- and Branchless-dependent signals. *Dev. Biol.* 402, 264–275.
- Sivak, J.M., Petersen, L.F., Amaya, E., 2005. FGF signal interpretation is directed by Sprouty and Spred proteins during mesoderm formation. *Dev. Cell* 8, 689–701.
- Song, Y.H., Zhu, Y.T., Ding, J., Zhou, F.Y., Xue, J.X., Jung, J.H., Li, Z.J., Gao, W.Y., 2016. Distribution of fibroblast growth factors and their roles in skin fibroblast cell migration. *Mol. Med. Rep.* 14, 3336–3342.
- Sun, J., Stathopoulos, A., 2018. FGF Controls Epithelial-Mesenchymal Transitions during Gastrulation by Regulating Cell Division and Apical Basal Polarity, vol. 145. Development.
- Swanhart, L.M., Takahashi, N., Jackson, R.L., Gibson, G.A., Watkins, S.C., Dawid, I.B., Hukriede, N.A., 2010. Characterization of an *lhx1a* transgenic reporter in zebrafish. *Int. J. Dev. Biol.* 54, 731–736.
- Tang, Q., Iyer, S., Lobbardi, R., Moore, J.C., Chen, H., Lareau, C., Hebert, C., Shaw, M.L., Neftel, C., Suva, M.L., Ceol, C.J., Bernards, A., Aryee, M., Pinello, L., Drummond, I.A., Langenau, D.M., 2017. Dissecting hematopoietic and renal cell heterogeneity in adult zebrafish at single-cell resolution using RNA sequencing. *J. Exp. Med.* 214, 2875–2887.
- Thisse, B., Heyer, V., Lux, A., Alunni, V., Degraeve, A., Seiliez, I., Kirchner, J., Parkhill, J.P., Thisse, C., 2004. Spatial and temporal expression of the zebrafish genome by large-scale in situ hybridization screening. *Methods Cell Biol.* 77, 505–519.
- Trueb, B., Amann, R., Gerber, S.D., 2013. Role of FGFR1 and other FGF signaling proteins in early kidney development. *Cell. Mol. Life Sci.* 70, 2505–2518.
- Wainwright, E.N., Wilhelm, D., Combes, A.N., Little, M.H., Koopman, P., 2015. ROBO2 restricts the nephrogenic field and regulates Wolffian duct-nephrogenic cord separation. *Dev. Biol.* 404, 88–102.
- Walker, K.A., Sims-Lucas, S., Bates, C.M., 2016. Fibroblast growth factor receptor signaling in kidney and lower urinary tract development. *Pediatr. Nephrol.* 31, 885–895.
- Wang, S., Lin, H., Zhao, T., Huang, S., Fernig, D.G., Xu, N., Wu, F., Zhou, M., Jiang, C., Tian, H., 2017. Expression and purification of an FGF9 fusion protein in *E. coli* and the effects of the FGF9 subfamily on human hepatocellular carcinoma cell proliferation and migration. *Appl. Microbiol. Biotechnol.* 101, 7823–7835.
- Willems, E., Leyns, L., Vandesompele, J., 2008. Standardization of real-time PCR gene expression data from independent biological replicates. *Anal. Biochem.* 379, 127–129.
- Yamauchi, H., Miyakawa, N., Miyake, A., Itoh, N., 2009. Fgf4 is required for left-right patterning of visceral organs in zebrafish. *Dev. Biol.* 332, 177–185.
- Yang, X., Steinberg, F., Zhuang, L., Bessey, R., Trueb, B., 2016. Receptor FGFR1 does not promote cell proliferation but induces cell adhesion. *Int. J. Mol. Med.* 38, 30–38.
- Zhao, H., Kegg, H., Grady, S., Truong, H.T., Robinson, M.L., Baum, M., Bates, C.M., 2004. Role of fibroblast growth factor receptors 1 and 2 in the ureteric bud. *Dev. Biol.* 276, 403–415.
- Zhou, W., Boucher, R.C., Bollig, F., Englert, C., Hildebrandt, F., 2010. Characterization of mesonephric development and regeneration using transgenic zebrafish. *Am. J. Physiol.* 299, F1040–F1047. Renal physiology.