



## Sexual differentiation of microglia

Alessandro Villa, Sara Della Torre, Adriana Maggi\*

Center of Excellence on Neurodegenerative Diseases and Dept of Pharmacological and Biomolecular Sciences, University of Milan, via Balzaretti, 9, Milan, Italy



### ARTICLE INFO

#### Keywords:

Neuroinflammation  
Sexual differentiation  
Neurodegeneration  
Development  
Adult brain

### ABSTRACT

Sex plays a role in the incidence and outcome of neurological illnesses, also influencing the response to treatments. Despite sexual differentiation of the brain has been extensively investigated, the study of sex differences in microglia, the brain's resident immune cells, has been largely neglected until recently. To fulfill this gap, our laboratory developed several tools, including cellular and animal models, which bolstered in-depth studies on sexual differentiation of microglia and its impact on brain physiology, as well as on the onset and progression of neurological disorders. Here, we summarize the current status of knowledge on the sex-dependent function of microglia, and report recent evidence linking these cells to the sexual bias in the susceptibility to neurological brain diseases.

### 1. Introduction

Microglia are a small minority of all the cells present in the Central Nervous System (CNS), about 10–15% of all brain cells (Del Rio-Hortega, 1965), and their function was believed to be restricted to the local immune defense of the CNS. More recently, a growing number of studies (Fig. 1) are demonstrating the large number of functions covered by these cells that are now considered to play a major role for brain maturation to maintain brain health.

Differently from the other cell types in the CNS, microglia do not originate from the ectoderm, but from myeloid precursors that, in early fetal development stages (embryonal day 8.5), migrate from the embryonic yolk sac to the brain (Kettenmann et al., 2011). Fate mapping analyses in mouse (Beers et al., 2006) and zebrafish (Herbomel et al., 1999) revealed that microglia originate from PU.1+ cells (Beers et al., 2006; Herbomel et al., 1999) and colonize the brain as CSF1R+ erythro-myeloid progenitor cells (Ginhoux et al., 2010; Villa et al., 2016). The number of cells migrating to the CNS is relatively small (Herbomel et al., 1999), but their ability to proliferate is sufficient to colonize the whole brain. This proliferative potential is maintained all through the life span induced by colony stimulating factor 1 (CSF1). The CSF1 receptor is expressed in microglia also in the adult mice and deletion of the *Csf1* gene or inhibition of the receptor activity results in a significant microglia loss (Elmore et al., 2014; Ginhoux et al., 2010). In the course of the embryo development, what is guiding microglia to the brain is not been clarified; in adults microglia may migrate to specific brain regions attracted by inflammatory stimuli. (Casano and Peri, 2015; Lenz et al., 2012; Schwarz et al., 2012). A characteristic of

microglia is their ability to acquire different morphologies, each of them mirroring a specific function (Benedusi et al., 2017). In the adult brain, microglia are found in different shapes: when highly ramified, microglia are patrolling the brain parenchyma in search of microorganisms, cells debris or deposits of misfolded proteins to be eliminated by engulfment and digestion; all these elements act as inflammatory stimuli to which microglia react by retracting all branches to take the amoeboid morphology characteristic of the inflammatory microglia. In this conformation microglia may engulf the noxious/toxic material and synthesize a vast array of inflammatory molecules (e.g. proinflammatory cytokines and chemokines, including tumor necrosis factor (TNF) and interleukin-1) aimed at opposing and destroying neurotropic microorganisms. Generally, this activated state is not maintained for long time, and microglia are able to revert to a deactivated state characterized by a novel change of morphology (towards the branched phenotype) and the synthesis and release of growth factors aimed at repairing the damage done in the course of the inflammatory reaction. We are still studying to identify the molecular mechanisms driving microglia through their differential functional roles and precise biomarkers for each stage of microglia activity.

Most relevant was the finding that microglia in their reactive form may trim neuronal dendritic spines; this suggested an involvement of these cells in the control of neuronal ability to communicate with each other in the adult brain (Kettenmann et al., 2013). In addition, it has been hypothesized that the double capacity of microglia to regulate neuronal communication and clear cell debris may be very relevant in the latest phases of brain development when the definitive neuronal circuitries are formed (Paolicelli et al., 2011). Indeed, different from

\* Corresponding author.

E-mail address: [adriana.maggi@unimi.it](mailto:adriana.maggi@unimi.it) (A. Maggi).

<https://doi.org/10.1016/j.yfrne.2018.11.003>

Received 19 September 2018; Received in revised form 7 November 2018; Accepted 24 November 2018

Available online 24 November 2018

0091-3022/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Number of publications containing the word microglia in the title or in the abstract.

Source: [dimensions.ai](https://dimensions.ai)

what found in adult, mature brain, microglia morphology in the immature CNS is mostly amoeboid and its gene expression includes both classical pro-inflammatory as well as alternative anti-inflammatory markers (Crain et al., 2013; Cunningham et al., 2013; Lenz et al., 2013). This phenotype, clearly different from that of the adult microglia, is likely associated with the specific functions exerted in the course of brain development and several authors demonstrated that in rodents, during the first week after birth, microglia eliminate redundant or apoptotic neurons and mold synaptic structures, participating in the shaping of the mature CNS (Mallat et al., 2005; Matcovitch-Natan et al., 2016).

The finding that, in addition to the specific immune mission, microglia play an active role in shaping brain circuitries and controlling neuronal activity raised questions concerning their relevance in the onset and progression of brain disorders.

## 2. Microglia in health and disease

A longstanding hypothesis has been that microglia had a detrimental role in selected neurological disorders (such as neurodegenerative diseases) because sustained neuroinflammatory processes can impair neuronal function and survival. Neurodegenerative diseases are generally characterized by an excessive production and accumulation of protein aggregates that constitute a trigger for microglia-dependent inflammatory processes; these deposits of aberrant proteins grow with the progression of the pathology and, in a long run, cause the recruitment of most microglia that - too busy in the removal of undesired material - cannot revert to the anti-inflammatory, reparatory, phenotype. The continuous and sustained release of neurotoxic molecules by microglia would damage the parenchyma as well as the other brain cells determining neuronal death (Rogers et al., 2002; Solito and Sastre, 2012). According to this hypothesis, McGeer et al. proposed the use of anti-inflammatory agents to limit the progression of these diseases (McGeer and McGeer, 1996). Afterwards, the growing awareness of the vast array of microglia functions in the brain helped in explaining the limited efficacy of anti-inflammatory therapies and compelled the research of novel therapeutic interventions aimed at maintaining or increasing microglia capabilities to repair damaged tissues and cells and to regulate aberrant neurotransmission (Akiyama et al., 2000; Hagino et al., 2004; Schenk and Yednock, 2002). However, the biology underlying the vast variety of microglia actions requires additional investigation because, as already mentioned, we start just now to understand that besides their anti-inflammatory actions, microglia serve a number of other functions not directly associated with inflammation, and possibly critical for the maintenance of neural functions (Salter and Beggs, 2014); the most relevant of such functions is the control of synaptic activity (Parkhurst et al., 2013; Tremblay et al., 2010). In the healthy CNS, microglia are not “resting” but are highly active in a

“surveillance and rapid response” state (Tremblay et al., 2011). Microglia processes scan the parenchyma continuously to get in touch with synapses; once in the proximity of a synapse, microglia sense the molecules released by neurons, or located at synaptic compartments, through specific receptors, such as the fractalkine receptor and the complement receptor 3 (Kettenmann et al., 2011; Koizumi et al., 2013), and react with the release of a wide repertoire of soluble factors such as BDNF, glycine, L-serine, TNF $\alpha$ , which affect neuronal processes, including basal neurotransmission and synaptic plasticity (Schafer et al., 2013; Yamasaki et al., 2014). For instance in case of excitatory synapses (Pocock and Kettenmann, 2007), microglia, once become aware of the extent of neuronal firing and synaptic function respond by extending their processes to be in direct contact with the active synapses (Chen et al., 2014; Tremblay et al., 2010; Wake et al., 2009); *in vivo* time-lapse imaging showed a decrease of firing after microglia enwrapping of the soma of highly active neurons (Li et al., 2012). Thus, microglia may tame the neuronal damage consequent to excessive synaptic activity by either releasing repair factors that limit the damage caused by a prolonged release of excitatory neurotransmitter or by engulfing and eliminating overactive dendritic spines. Another mechanism by which microglia participate in neuronal activities is by regulating adult neurogenesis as described in the hippocampus (Gemma and Bachstetter, 2013; Sierra et al., 2010), and this event might result in improving learning/memory processes (Parkhurst et al., 2013; Rogers et al., 2011).

These microglia physiological functions may be very important for the healthy functioning of the adult CNS. This new awareness led to the hypothesis that apart or in addition to their inflammatory functions, aberrations in non-inflammatory functions of microglia may contribute to the onset of neurodegenerative diseases (Tremblay et al., 2011). In this context, a systems biology approach applied on large-scale human brain-tissue sampling (Zhang et al., 2013), identified a remodeling in the complement-dependent signaling of microglia as the functional category most strongly associated with the pathophysiology of neurodegeneration. The anomalous upregulation of complement signaling could hyperactivate the synapse elimination pathways, thus triggering synaptic loss, an early hallmark of neurodegenerative diseases (Tremblay et al., 2011).

The importance of microglia in a neurodegenerative disorder such as AD became indisputable when genome-wide association studies (GWAS) reported a consistent number of genetic susceptibility loci for late-onset AD (LOAD), which were associated with the inflammatory pathway (Mhatre et al., 2015). One of the most striking findings was that a mutation (R47H) in TREM2 – a gene involved in the microglial phagocytic process - leads to a 3- to 4-fold increased risk of sporadic AD (Guerreiro et al., 2013). Accordingly, a recent study where the human mutated TREM2 was integrated into the genome of the 5XFAD murine model of AD showed significant impairment of microglia phagocytic activity (Song et al., 2018), thus suggesting a link between this microglial loss of function and AD. Since these pioneering studies, a profusion of novel investigations shifted the focus on the relevance of loss of microglia homeostasis in neuronal disorders. Keren-Shaul and colleagues (Keren-Shaul et al., 2017) using single-cell RNA-seq on the 5XFAD model of AD, discovered a novel microglia phenotype associated with neurodegenerative diseases (named DAM) and identified specific markers, spatial localization (primarily in the brain regions affected by the disease), and pathways associated with these cells (Keren-Shaul et al., 2017). This specific phenotype seems to trigger a protective mechanism aiming at containing and removing the neuronal damage (Deczkowska et al., 2018). Most relevant is that many of the genes associated with DAM are the same identified by GWAS studies on AD, including TREM2 (Guerreiro et al., 2013), that participates in DAM differentiation (Keren-Shaul et al., 2017). Further studies identified the presence of DAM in other ND models, such as tauopathy (Friedman et al., 2018), ALS (Friedman et al., 2018; Krasemann et al., 2017), MS (Keren-Shaul et al., 2017; Krasemann et al., 2017), as well as in aging

(Mrdjen et al., 2018). Moreover, the DAM expression pattern was detected also in postmortem brains from AD patients (Friedman et al., 2018; Keren-Shaul et al., 2017). These data suggest that microglia may acquire a specific phenotype in the presence of neurodegenerative cues necessary to slow down the progression of the disease.

Therefore, despite most of the evidence regarding the involvement of microglia in neurodegeneration was collected for AD, their engagement in the etiopathogenesis of other neurodegenerative diseases is most likely. Thus, the finding of these non-inflammatory functions of microglia will provide novel sights on microglia involvement in brain pathologies that diverge from the strict association with a gain of inflammatory activity; rather, CNS disorders could result from loss of selected non-inflammatory microglia functions.

### 3. Are microglia involved in brain sexual differentiation?

We have been aware of the fact that the brain is sexually differentiated for a long time and several morphological studies proved the existence of differences in the brain of females and males; the understanding of the mechanism responsible for such differentiation originates from pioneering experiments done in the second half of twentieth century when it was observed that the exposure to androgens early in the development accounted for major brain sex differences in terms of structure and function. Roger Gorski and colleagues (Gorski et al., 1978) initially identified sex differences in the preoptic area (POA) of the rat hypothalamus, a brain region generally implicated in the control of male reproductive behaviour. A cluster of cells in the POA, defined as the sexually dimorphic nucleus (SDN), was shown to be much larger in males than in females and involved in neural mechanisms necessary for sexual behaviour (Swaab et al., 2001). Subsequent studies extended the number of sexually dimorphic regions (including the anteroventral periventricular nucleus of the hypothalamus (AVPV, (Bloch et al., 1987)), and the spinal nucleus of the bulbocavernosus (SBN, (Breedlove and Arnold, 1983) in rat spinal cord) and demonstrated that the testosterone synthesized by the testes during embryogenesis in humans and neonatally in rodents (Clarkson and Herbison, 2016) is responsible for the masculinization of selected brain circuits that control sexually differentiated behaviours and physiological processes (Gorski et al., 1978; Lenz and McCarthy, 2010). In the absence of such organizing action induced by steroid hormones, the brain remains feminine as default program (Gorski et al., 1978; Lenz and McCarthy, 2010).

The sex differences in the CNS include sex-specific neuroanatomical features; neurons show regional differences in volume, cell number, connectivity, morphology, neurite complexity, dendritic length and spine number, but also transcriptional and epigenetic changes (McCarthy et al., 2009). Thus in the adult animals the neuronal response to stimuli is sexually differentiated: for instance in the rat hippocampus, the morphology of pyramidal neurons and stellate cells in response to visual stimuli vary with sex, with female rats raised in an enriched environment showing increased dendritic branching relative to males housed in the same environment (Juraska et al., 1985). In the same brain area, following repeated restraint stress, the number of apical branch points and dendritic length of the CA3c pyramidal neurons decreases in male, but not in female rats (Galea et al., 1997). Moreover, a strong sex difference exists in the long-term potentiation (LTP), due to differential synaptic NMDA receptor activation at perforant path in male and female rats, resulting in a more enduring LTP in males (Maren et al., 1994). GABA-mediated stimulation of neurons from the *substantia nigra* of juvenile rats produces sex-biased physiological effects leading to depolarization in males, but hyperpolarizing the same cells in females (Galanopoulou, 2005). Neurons are not the only brain cells undergoing sexual differentiation: astrocytes show a clear sexual dimorphism in terms of distribution (Collado et al., 1995; Garcia-Segura et al., 1988; Suarez et al., 1991), differentiation, primary process length and number (Amateau and McCarthy, 2002, 2004; Johnson et al., 2008) and function (Garcia-Segura et al., 1995; Kuo


et al., 2010; Mong and McCarthy, 1999; Suarez et al., 1992),

With regard to microglia most of the experiments done so far aimed at investigating sex differences in the course of CNS development. In the early postnatal development (P4), males have a higher number of microglia cells in brain regions involved in learning, memory and cognition processes (Schwarz et al., 2012); slight sex differences in cell size and phagocytic capacity were identified in the hippocampus across all developmental stages (Weinhard et al., 2018).

Studies in rodent showed that microglia maturation occurs slowly: just before birth microglia are close to be fully differentiated (Butovsky et al., 2014), but complete maturation is reached during the second postnatal week (Bennett et al., 2016; Matcovitch-Natan et al., 2016). Each stage in microglia development is associated with specific gene expression programs and regulatory networks (Maticovitch-Natan et al., 2016): in immature, embryonic microglia there is a major expression of proteins for the cell cycle and for chromatin remodelling, while in the adult microglia canonical transcription factors, such as EGR1 and SALL1, appear in the early post-natal stage and the expression levels rise with time. Other transcription factors, including JUN, FOS, MEF2A, and MAFB, are expressed in adult microglia only (Maticovitch-Natan et al., 2016). These transcriptional changes associated with developmental stages may be a reflection of the progressive changes occurring in the brain, as microglia cells are in constant communication with the other CNS cells through specific neurotransmitters, neurohormones, and neuromodulators, and are able to mold their function in response to the surrounding microenvironment (Crain et al., 2013). In the latest stages of CNS development, microglia morphology shifts from amoeboid to the ramified, quiescent structure typical of a healthy adult brain (Villa et al., 2016): this is one of the many changes in phenotype that characterize the transition between immature and adult microglia.

Hanamsagar and colleagues (Hanamsagar et al., 2017) profiled microglia transcriptome in the course of brain development and showed that from the transcriptional point of view, the maturation process has features resembling to the programs of pro-inflammatory activation typical of adult cells. Most relevant is that the temporal maturation steps follow distinct trajectories in males and in females. Starting from embryonic day 18 sex has a significant impact on microglia maturation that is delayed in males relative to females. In the presence of acute immune activation, such as following stimulation with LPS, it was observed an acceleration in male microglial development, while female microglia did not change its maturation stage (Hanamsagar et al., 2017). These data suggest that male microglia could be more sensitive to inflammatory events, which could be responsible for a faster aging of microglia and this could affect the risk of disorders (Hanamsagar et al., 2017).

We do not know how the genetic sex influences microglia maturation and the extent to which sex hormones are involved. Microglia expression of sex hormone receptors is relative to the stage of brain maturation (Villa et al., 2016): mRNA levels of the Estrogen Receptor  $\alpha$  (ER $\alpha$ ) are detectable in microglia obtained from P3 mice (Crain et al., 2013) and the content increases in adult mice (Crain et al., 2013). No sexual differences were observed in ER $\alpha$  microglia mRNA at any age (Crain et al., 2013; Sierra et al., 2008). ER $\beta$  expression was detected in primary microglia cultures from P0 newborns only (Saijo et al., 2011), and its expression becomes undetectable starting from P3 until adulthood (Crain et al., 2013; Sierra et al., 2008). Data on the expression of progesterone receptor (PR) and androgen receptor (AR) in the course of development show that microglia may express both (Quadros et al., 2007), although AR and PR do not appear to be expressed in microglia in adult mice (Sierra et al., 2008). Therefore, microglia may respond directly to the surge of testosterone that occurs perinatally and that could address microglia towards a male-specific pattern of maturation. The question to be raised here is whether microglia have a role in the shaping of neural interconnections occurring in brain development and whether the sex-specific differences in the maturation of microglia are relevant for the creation of the neuronal networks that characterize



**Fig. 2.** Adult microglia isolated from CX3CR1-GFP transgenic mice. Female microglia *in vitro* (upper left panel) show, by and large, a phenotype reminiscent of unactivated microglia in brain tissue (lower left panel), while male microglia morphology *in vitro* (upper right panel) is more reminiscent of the activated microglia phenotype (lower right panel). Scale bar: 20  $\mu$ m.

male and female brains. It is indeed increasingly recognized that throughout the development, the exposure to factors that permanently modify the function of microglia and immune system may severely impact on sexual behavior in adult life (Lenz et al., 2013). For instance, innate immune activation during development can lead to neurological outcomes in a sex-dependent manner (Schwarz et al., 2012), thus indicating the important role played by the endocrine-microglia communication in the structural organization of the brain during its maturation. A central role of microglia in the sexual differentiation of the brain was proposed by Lenz and colleagues (Lenz et al., 2013), which observed that perinatal treatments with minocycline – an inhibitor of microglial activity – prevented the masculinization of the brain that is normally induced by estradiol (Lenz et al., 2013). The pharmacological inhibition of microglia hampered the production of a pro-inflammatory molecule, the prostaglandin E2 (PGE2), which is synthesized in the POA following the neonatal testosterone surge, and is responsible for the establishment of male-specific neuroanatomical features in the brain (Amateau and McCarthy, 2004). A complete view of the mechanisms regulating brain sexual differentiation certainly requires more specific investigations; however, what we can conclude from the data reported so far is that during development microglia undergo a sex-dependent maturation process, and therefore studies carried out in cultures of neonatal microglia may not provide information translatable to adult microglia.

#### 4. Sex differences in adult microglia

Indeed, studies on the physiology of adult microglia have been limited so far; this is likely due to the difficulties of maintaining adult, fully mature, microglia cells in culture. The limited number of data available so far, however, showed a sex-specificity in the expression of classical or alternative activation markers (Crain et al., 2013; Weinhard et al., 2018), the expression of purinergic receptors (Crain and Watters, 2015), cell numbers (Mouton et al., 2002), distribution into the CNS (Lawson et al., 1990), response to exercise (Kohman et al., 2013) and to stress (Bollinger et al., 2016; Bollinger et al., 2017; Fonken et al., 2018). Quite peculiar is the sexual dimorphic involvement of microglia in neuropathic pain signalling: while in male mice pain sensitivity is mediated by specific microglia-neuronal signalling pathways triggered by the activation of P2X4R on spinal microglia (Beggs et al., 2012), further experiments have shown that these cells do not participate in pain processing in female animals (Mapplebeck et al., 2017; Sorge et al., 2015; Taves et al., 2016). Similarly, social interactions in adolescents are influenced by microglia in males only, through a mechanism mediated by the complement C3, resulting in the engulfment and lysosomal elimination of spines expressing neuronal dopamine D1 receptors in the nucleus accumbens (Kopeck et al., 2018).

These reports strongly support the idea that sex differences in microglia in the brain may be more extensive than just a difference in number, morphology, or specific functions, but may also be phenotypically distinct.

Recently, by taking advantage of the novel technique to isolate pure

population of microglia from the brain of adult mice developed in our laboratory (Pepe et al., 2014; Villa et al., 2018), we carried out a whole genome RNA-seq analysis to compare the transcriptomes of microglia in male and female mice (Villa et al., 2018). When we compared the abundance of the transcripts of microglia from the two sexes (by applying a threshold of 0.01 to p-values) we found more than 500 differentially expressed genes (DEGs) (about 200 more expressed in males, and 350 more expressed in females). The functional analyses of the genes more expressed in males revealed a marked homogeneity: most of the DEGs showed a strict association with inflammatory processes, including regulation of cell migration and cytokine production; molecular signature analysis of transcription factors (TFs) identified NF- $\kappa$ B as the TF most involved in the regulation of the genes more expressed in males (Villa et al., 2018). Analogously, a subsequent paper reported for male microglia a transcriptomic profile skewed towards the pro-inflammatory activation (Guneykaya et al., 2018). The higher NF- $\kappa$ B activity was confirmed using the NF- $\kappa$ B-reporter mouse recently generated in our laboratory and carrying the luciferase gene under the control of a NF- $\kappa$ B-responsive synthetic promoter (Rizzi et al., 2017). In this mouse model, luciferase activity in microglia purified from the brain of both sexes was 2.4-fold higher in males (Villa et al., 2018). This male-specific grade of activation of the transcription factor seems to be a feature characteristic of microglia, since whole-body, *in vivo* imaging in unstimulated conditions showed a similar basal level of NF- $\kappa$ B transcriptional activation in males and females (Villa et al., 2018).

Less straight-forward phenotype resulted from the analysis of the microglia genes more expressed in females: functional analyses identified ontogenies associated with morphogenesis, development or cytoskeleton organization. More informative was the molecular signature analysis of TFs, that identified proteins related to the inhibition of inflammatory response and promotion of repair mechanisms (Villa et al., 2018). Taken together, these data support the idea of a differential reactivity in microglia originating from the two sexes, strengthened by morphological (Fig. 2.) and biochemical studies carried out in primary cultures of microglia isolated from adult male and female mice, grown *in vitro* with neuron/astrocyte mixed cells (Villa et al., 2018). Thus, microglia cells appear to be sexually differentiated and to maintain sex-specific features independently by the microenvironment: this vision was further supported by the finding that microglia retain their sex-related characteristics even after transplantation in the male brain (Villa et al., 2018).

The molecular mechanisms leading to this sex dichotomy are still uncertain; our current hypothesis, corroborated by preliminary results, is that epigenetic modifications take place shortly after birth, when the surge of testosterone occurs (Gorski et al., 1978; McCarthy et al., 2009). We believe that this is the case because we experimentally mimicked the masculinization process by treating female mice with repeated neonatal injection of E<sub>2</sub> (Wu et al., 2009). The microglia isolated from the adult masculinized females showed a sex-bias even if the results obtained were not conclusive as the microglia of adult, masculinized females did not show a signature of gene expression completely superimposable with genetic males (Villa et al., 2018). Of course several factors may influence microglia function in adult animals, therefore more studies are required to fully understand the mechanisms leading adult microglia to acquire the male/female phenotype; among these we believe that gonadal steroids as well as neurosteroids may influence microglia as demonstrated by our as well as other groups in male and female animals treated with inflammatory stimuli as LPS in the presence/absence of estradiol (Loram et al., 2012; Vegeto et al., 2006; Vegeto et al., 2001).

Most sex-dependent differences in mammals are associated with reproductive functions and maintained in evolution; it is premature to speculate whether this is the case also for microglia. Data mining of datasets reporting the human transcriptomes of male and female brains at different stages of development (including adulthood), such as The Human Brain Transcriptome (HBT) (Kang et al., 2011), or the more

recent BRAINSPAN dataset (<http://www.brainspan.org/static/download.html>), led to the conclusion that in the postnatal male brain there is a high expression of genes associated with microglia phagocytic and immune function (Prilutsky et al., 2017; Werling et al., 2016); moreover, genes involved in synaptic pruning were upregulated in males at prenatal stages and are downregulated in males postnatally (Prilutsky et al., 2017). Likewise, the Bilbo group compared the development-associated gene expression patterns observed in murine microglia to the BRAINSPAN dataset: similar patterns were observed also in humans despite the heterogeneity of brain tissue, since the expression of microglia-specific group of genes increased with age (Hanamsagar et al., 2017). They also observed that male microglial transcriptome was more developmentally mature than female microglia. Interestingly, environmental factors seem to play a role in triggering the developmental programs of microglial cells in a sex-dependent manner, being the maternal microbiome likely to influence the transcriptional maturation of microglia in the fetal brain, especially in males (Thion et al., 2018). Conversely, female microglia are more prone to responding to microbiome changes in adulthood, showing profound changes in microglial transcriptomic signatures after acute and chronic microbiome depletion (Thion et al., 2018). The main sex-biased features of microglia identified so far are summarized in Fig. 3.

These findings may have relevance for the understanding of the sex-related differences in the susceptibility or progression to brain disorders as the lifespan exposure to infectious agents may have a greater effect on males than females; moreover, activation of microglia during prenatal development could lead to an increased pace in reaching maturity of male microglia (Hanamsagar et al., 2017). It is important to underline that more studies should be done to support these findings as transcriptomics analysis of purified human cortical microglia from *post-mortem* samples did not find significant sex-related differences (Galaturo et al., 2017).

## 5. Microglia sex difference and brain disorders

A sex difference in the incidence, severity, and/or progression has been reported for several neurological diseases (Villa et al., 2016). For example, AD has a higher prevalence in women above 65 years old (1.6–3:1 ratio compared to men), and also progresses with a greater cognitive deterioration (Plassman et al., 2011; Seshadri et al., 1997). Men have a higher incidence of Parkinson's disease (PD) (3.5:1 compared to women) and the disease has a slower progression in women (Balderechi et al., 2000; Elbaz et al., 2002). Females have a lower incidence of stroke (which depends on age as well), however they display poorer outcomes and suffer a more precipitous decline in function following stroke compared to males (Roy-O'Reilly and McCullough, 2014). Depression is prevalent in females (Altemus et al., 2014), while schizophrenia (McGrath et al., 2004) and autism (Werling and Geschwind, 2013) are prevalent in males. As a general rule, with some exception (e.g. Parkinson's disease), epidemiological studies indicate that disorders that emerge early in life are more common in males, whereas disorders that emerge later in life (at adolescence or beyond) are more common in females.

The reasons underlying these sex-related differences are still unknown as their origin could be genetic, associated to brain sexual differentiation and/or circulating sex steroids, all factors able to influence the activities of neurons, astrocytes, and microglia (Arnold, 2009; Gillies et al., 2004; Gorski et al., 1978; Joel et al., 2015; Joel and McCarthy, 2017; Li et al., 2014; Schwarz and McCarthy, 2008).

Even if it is plausible that microglia could be involved, it is premature to hypothesize the extent to which these cells could play a role. Initial studies are claiming a role for microglia in the sex prevalence of selected neurological diseases. A number of studies reported a pro-inflammatory activation of microglia following acute and chronic stress [extensively reviewed in (Calcina et al., 2016)], and the recent work of Bollinger and colleagues (Bollinger et al., 2016; Bollinger et al., 2017)




Fig. 3. Main sex-biased features of microglia. The schematic cartoon reports some of the most emblematic features of female (left side) or male (right side) microglia.

highlighted a marked sex difference in microglial density, morphology, and immune factor expression across corticolimbic circuitry in stressed rats, which is possibly linked to the women's high vulnerability to stress-linked psychological disorders (Riecher-Rossler, 2017). Werling and colleagues (Werling et al., 2016) showed that autism spectrum disorder (ASD) is associated with a marked upregulation of sex-specific genes (genes generally more expressed in healthy males); among them there was a significant enrichment of microglia activation markers (Werling et al., 2016). The implication of these findings is that the sex-specific incidence of autism might be associated with the higher microglia reactivity, typical of males, responsible for alterations in neuronal connectivity leading to the manifestation of the disease prevalent in males. Parallel analyses on the same datasets showed that in the course of development microglia mature faster in males, leading to think that autistic individuals have an exaggerated development of microglia than healthy controls (Hanamsagar et al., 2017). A similar aberrant microglia phenotype was observed in AD tissue compared to controls (Hanamsagar et al., 2017) even if AD is prevalent in females. However, it is known that with ageing females experience a higher level of microglial activation compared to age-matched males, with possible implications on the development of neurodegenerative diseases (Mangold et al., 2017). Indeed, it is quite difficult to dissect the cellular and molecular mechanisms underpinning different brain pathologies when their development occurs in decades or, worse, it is rooted in brain development. The study of acute diseases (like stroke or brain injury) may provide a better understanding on the role played by the different neural cells, microglia included. Studies in traumatic brain injury (Acaz-Fonseca et al., 2015) or middle cerebral artery occlusion (Bodhankar et al., 2015), indicated that microglia from female mice show a lower degree of inflammatory activation following the insult.

In these studies, however, it is important to take into account the presence of circulating sex steroids known to influence microglia inflammatory response and progression through the different stages ensuing the inflammatory reaction (Vegeto et al., 2001; Villa et al., 2015). To circumvent this problem and to have a clearer vision of the role of sex of microglia in the neuroinflammatory process, we recently set up a methodology by which we can transplant adult microglia and evaluate its behaviour in a setting (physiological or diseased) different from the original. Using this methodology, we investigated the relevance of microglia sex in acute stroke. After permanent middle cerebral artery

occlusion (pMCAO) (Villa et al., 2018) naïve male mice develop a larger injury than females, modelling stroke outcomes seen in men and women (Murphy et al., 2004; Villa et al., 2018). However, we observed that when microglia isolated from female mice was transplanted in males prior to pMCAO, there was a significant reduction of the lesioned area and a higher expression of the anti-inflammatory marker Ym1 around the damaged area. This clearly demonstrated the major role of microglia in containing the hypoxic damage and provided a strong evidence that this feature is more efficacious in female than in male microglia, independently from the levels of circulating estrogens (Villa et al., 2018).

## 6. Conclusions

This review intended to summarize the progress of the recent years in the understanding of microglia biology and its major role in healthy as well as diseased brain. The results obtained so far highlight microglia as major players in the brain physio-pathology and the complexity of activities that these cells may display.

We believe that one of the major progresses is associated with the possibility to study adult microglia with methodologies enabling 1. to identify and label these cells with specific genetic markers, 2. to isolate from the adult brains highly (> 95%) pure populations of these cells (Pepe et al., 2014; Villa et al., 2018). This latter is particularly relevant considering the complexity of functions covered by these cells and their phenotypic evolution in the course of life. These studies are demonstrating the major limitations of the classical *in vitro* studies carried out with microglia obtained from immature brains and not stratified for sex, as the stage of maturation of these cells is very relevant for their final functions. Major efforts should therefore be done to improve the methodologies to grow mature microglia *in vitro*: some progress has been recently done from our group by using co-cultures of microglia and neurons where neurons appear to enable microglia to maintain their specific phenotype (at least with regard to the sex of origin) (Villa et al., 2018), but of course more studies should be done and more biomarkers should be generated to enable us to correctly stage microglia both *in vitro* and *in vivo*. Our prediction is that transplantation studies combined with animal genetics will have a major impact on defining the role of microglia in sustaining/containing the progression of neurological and neuropsychiatric diseases: we will be able to

observe the fate and activity of cells extracted from diseased brains in healthy brains and *vice-versa* in young, adult or aged animals. The advances obtained so far in our understanding of microglia functions might only be the tip of a big iceberg and this must be an incentive for further studies on microglia particularly for their potential major role as novel target for the numerous brain disorders waiting for a cure.

## Acknowledgments

Funding: This work was supported by the European Union's grants ERC-2012-ADG322977-Ways and Seventh Framework Programme (FP7/2007–2013) under grant agreement no. 278850 (INMiND).

## References

- Acaz-Fonseca, E., Duran, J.C., Carrero, P., Garcia-Segura, L.M., Arevalo, M.A., 2015. Sex differences in glia reactivity after cortical brain injury. *Glia* 63, 1966–1981. <https://doi.org/10.1002/glia.22867>.
- Akiyama, H., Barger, S., Barnum, S., Bradt, B., Bauer, J., Cole, G.M., Cooper, N.R., Eikelenboom, P., Emmerling, M., Fiebich, B.L., Finch, C.E., Frautschi, S., Griffin, W.S., Hampel, H., Hull, M., Landreth, G., Lue, L., Mrak, R., Mackenzie, I.R., McGeer, P.L., O'Banion, M.K., Pachter, J., Pasinetti, G., Plata-Salamán, C., Rogers, J., Rydel, R., Shen, Y., Streit, W., Strohmeyer, R., Tooyoma, I., Van Muiswinkel, F.L., Veerhuis, R., Walker, D., Webster, S., Wegrzyniak, B., Wenk, G., Wyss-Coray, T., 2000. Inflammation and Alzheimer's disease. *Neurobiol. Aging* 21, 383–421.
- Altemus, M., Sarvaiya, N., Neill Epperson, C., 2014. Sex differences in anxiety and depression clinical perspectives. *Front. Neuroendocrinol.* 35, 320–330. <https://doi.org/10.1016/j.yfrne.2014.05.004>.
- Amateau, S.K., McCarthy, M.M., 2002. Sexual differentiation of astrocyte morphology in the developing rat preoptic area. *J. Neuroendocrinol.* 14, 904–910.
- Amateau, S.K., McCarthy, M.M., 2004. Induction of PGE2 by estradiol mediates developmental masculinization of sex behavior. *Nat. Neurosci.* 7, 643–650. <https://doi.org/10.1038/nn1254>.
- Arnold, A.P., 2009. The organizational-activational hypothesis as the foundation for a unified theory of sexual differentiation of all mammalian tissues. *Horm. Behav.* 55, 570–578. <https://doi.org/10.1016/j.yhbeh.2009.03.011>.
- Baldereschi, M., Di Carlo, A., Rocca, W.A., Vanni, P., Maggi, S., Perisintono, E., Grigoletto, F., Amaducci, L., Inzitari, D., 2000. Parkinson's disease and parkinsonism in a longitudinal study: two-fold higher incidence in men. ILSA Working Group. *Italian Longitudinal Study on Aging. Neurology* 55, 1358–1363.
- Beers, D.R., Henkel, J.S., Xiao, Q., Zhao, W., Wang, J., Yen, A.A., Siklos, L., McKercher, S.R., Appel, S.H., 2006. Wild-type microglia extend survival in PU.1 knockout mice with familial amyotrophic lateral sclerosis. *Proc. Natl. Acad. Sci. USA* 103, 16021–16026. <https://doi.org/10.1073/pnas.0607423103>.
- Beggs, S., Trang, T., Salter, M.W., 2012. P2X4R+ microglia drive neuropathic pain. *Nat. Neurosci.* 15, 1068–1073. <https://doi.org/10.1038/nn.3155>.
- Benedusi, V., Della Torre, S., Mitro, N., Caruso, D., Oberto, A., Tronel, C., Meda, C., Maggi, A., 2017. Liver ERalpha regulates AgRP neuronal activity in the arcuate nucleus of female mice. *Sci. Rep.* 7, 1194. <https://doi.org/10.1038/s41598-017-01393-0>.
- Bennett, M.L., Bennett, F.C., Liddelow, S.A., Ajami, B., Zamanian, J.L., Fernhoff, N.B., Mullinyawe, S.B., Bohlen, C.J., Adil, A., Tucker, A., Weissman, I.L., Chang, E.F., Li, G., Grant, G.A., Hayden Gephart, M.G., Barres, B.A., 2016. New tools for studying microglia in the mouse and human CNS. *Proc. Natl. Acad. Sci. USA* 113, E1738–E1746. <https://doi.org/10.1073/pnas.1525528113>.
- Bloch, G.J., Babcock, A.M., Gorski, R.A., Micevych, P.E., 1987. Cholecystokinin stimulates and inhibits lordosis behavior in female rats. *Physiol. Behav.* 39, 217–224.
- Bodhankar, S., Lapato, A., Chen, Y., Vandenbark, A.A., Saugstad, J.A., Offner, H., 2015. Role for microglia in sex differences after ischemic stroke: importance of M2. *Metab. Brain Dis.* 30, 1515–1529. <https://doi.org/10.1007/s11011-015-9714-9>.
- Bollinger, J.L., Bergeon Burns, C.M., Wellman, C.L., 2016. Differential effects of stress on microglial cell activation in male and female medial prefrontal cortex. *Brain Behav. Immun.* 52, 88–97. <https://doi.org/10.1016/j.bbi.2015.10.003>.
- Bollinger, J.L., Collins, K.E., Patel, R., Wellman, C.L., 2017. Behavioral stress alters corticolic microglia in a sex- and brain region-specific manner. *PLoS ONE* 12, e0187631. <https://doi.org/10.1371/journal.pone.0187631>.
- Breedlove, S.M., Arnold, A.P., 1983. Hormonal control of a developing neuromuscular system. II. Sensitive periods for the androgen-induced masculinization of the rat spinal nucleus of the bulbocavernosus. *J. Neurosci.* 3, 424–432.
- Butovsky, O., Jedrychowski, M.P., Moore, C.S., Cialic, R., Lanser, A.J., Gabrieli, G., Koelsperger, T., Dake, B., Wu, P.M., Doykan, C.E., Fanek, Z., Liu, L., Chen, Z., Rothstein, J.D., Ransohoff, R.M., Gygi, S.P., Antel, J.P., Weiner, H.L., 2014. Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nat. Neurosci.* 17, 131–143. <https://doi.org/10.1038/nn.3599>.
- Calcina, M.A., Bonsall, D.R., Bloomfield, P.S., Selvaraj, S., Baricello, T., Howes, O.D., 2016. Stress and neuroinflammation: a systematic review of the effects of stress on microglia and the implications for mental illness. *Psychopharmacology* 233, 1637–1650. <https://doi.org/10.1007/s00213-016-4218-9>.
- Casano, A.M., Peri, F., 2015. Microglia: multitasking specialists of the brain. *Dev. Cell* 32, 469–477. <https://doi.org/10.1016/j.devcel.2015.01.018>.
- Chen, Z., Jalabi, W., Hu, W., Park, H.J., Gale, J.T., Kidd, G.J., Bernatowicz, R., Gossman, Z.C., Chen, J.T., Dutta, R., Trapp, B.D., 2014. Microglial displacement of inhibitory synapses provides neuroprotection in the adult brain. *Nat. Commun.* 5, 4486. <https://doi.org/10.1038/ncomms5486>.
- Clarkson, J., Herbison, A.E., 2016. Hypothalamic control of the male neonatal testosterone surge. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 371, 20150115. <https://doi.org/10.1098/rstb.2015.0115>.
- Collado, P., Beyer, C., Hutchison, J.B., Holman, S.D., 1995. Hypothalamic distribution of astrocytes is gender-related in Mongolian gerbils. *Neurosci. Lett.* 184, 86–89.
- Crain, J.M., Nikodemova, M., Watters, J.J., 2013. Microglia express distinct M1 and M2 phenotypic markers in the postnatal and adult central nervous system in male and female mice. *J. Neurosci. Res.* 91, 1143–1151. <https://doi.org/10.1002/jnr.23242>.
- Crain, J.M., Watters, J.J., 2015. Microglial P2 Purinergic Receptor and Immunomodulatory Gene Transcripts Vary By Region, Sex, and Age in the Healthy Mouse CNS. *Transcr. Open Access* 3. <https://doi.org/10.4172/2329-8936.1000124>.
- Cunningham, C.L., Martínez-Cerdano, V., Noctor, S.C., 2013. Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *J. Neurosci.* 33, 4216–4233. <https://doi.org/10.1523/JNEUROSCI.3441-12.2013>.
- Deczkowska, A., Keren-Shaul, H., Weiner, A., Colonna, M., Schwartz, M., Amit, I., 2018. Disease-associated microglia: a universal immune sensor of neurodegeneration. *Cell* 173, 1073–1081. <https://doi.org/10.1016/j.cell.2018.05.003>.
- Del Rio-Hortega, R., 1965. *Microglia. Cytology and cellular Pathology of the Nervous System*, pp. 483–535.
- Elbaz, A., Bower, J.H., Maraganore, D.M., McDonnell, S.K., Peterson, B.J., Ahlskog, J.E., Schaid, D.J., Rocca, W.A., 2002. Risk tables for parkinsonism and Parkinson's disease. *J. Clin. Epidemiol.* 55, 25–31.
- Elmore, M.R., Najafi, A.R., Koike, M.A., Dagher, N.N., Spangenberg, E.E., Rice, R.A., Kitazawa, M., Matusow, B., Nguyen, H., West, B.L., Green, K.N., 2014. Colony-stimulating factor 1 receptor signaling is necessary for microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron* 82, 380–397. <https://doi.org/10.1016/j.neuron.2014.02.040>.
- Fonken, L.K., Frank, M.G., Gaudet, A.D., D'Angelo, H.M., Daut, R.A., Hampson, E.C., Ayala, M.T., Watkins, L.R., Maier, S.F., 2018. Neuroinflammatory priming to stress is differentially regulated in male and female rats. *Brain Behav. Immun.* 70, 257–267. <https://doi.org/10.1016/j.bbi.2018.03.005>.
- Friedman, B.A., Srinivasan, K., Ayalon, G., Meilandt, W.J., Lin, H., Huntley, M.A., Cao, Y., Lee, S.H., Haddick, P.C.G., Ngu, H., Modrusan, Z., Larson, J.L., Kaminker, J.S., van der Brug, M.P., Hansen, D.V., 2018. Diverse brain myeloid expression profiles reveal distinct microglial activation states and aspects of Alzheimer's disease not evident in mouse models. *Cell Rep.* 22, 832–847. <https://doi.org/10.1016/j.celrep.2017.12.066>.
- Galanopoulou, A.S., 2005. GABA receptors as broadcasters of sexually differentiating signals in the brain. *Epilepsia* 46 (Suppl 5), 107–112. <https://doi.org/10.1111/j.1528-1167.2005.01007.x>.
- Galatro, T.F., Holtman, I.R., Lerario, A.M., Vainchtein, I.D., Brouwer, N., Sola, P.R., Veras, M.M., Pereira, T.F., Leite, R.E.P., Moller, T., Wes, P.D., Sogayar, M.C., Laman, J.D., den Dunnen, W., Pasqualucci, C.A., Oba-Shinjo, S.M., Boddeke, E., Marie, S.K.N., Eggen, B.J.L., 2017. Transcriptomic analysis of purified human cortical microglia reveals age-associated changes. *Nat. Neurosci.* 20, 1162–1171. <https://doi.org/10.1038/nn.4597>.
- Galea, L.A., McEwen, B.S., Tanapat, P., Deak, T., Spencer, R.L., Dhabhar, F.S., 1997. Sex differences in dendritic atrophy of CA3 pyramidal neurons in response to chronic restraint stress. *Neuroscience* 81, 689–697.
- García-Segura, L.M., Duenas, M., Busiguina, S., Naftolin, F., Chowen, J.A., 1995. Gonadal hormone regulation of neuronal-glia interactions in the developing neuroendocrine hypothalamus. *J. Steroid Biochem. Mol. Biol.* 53, 293–298.
- García-Segura, L.M., Suarez, I., Segovia, S., Tranque, P.A., Cales, J.M., Aguilera, P., Olmos, G., Guillemon, A., 1988. The distribution of glial fibrillary acidic protein in the adult rat brain is influenced by the neonatal levels of sex steroids. *Brain Res.* 456, 357–363.
- Gemma, C., Bachstetter, A.D., 2013. The role of microglia in adult hippocampal neurogenesis. *Front. Cell. Neurosci.* 7, 229. <https://doi.org/10.3389/fncel.2013.00229>.
- Gillies, G.E., Murray, H.E., Dexter, D., McArthur, S., 2004. Sex dimorphisms in the neuroprotective effects of estrogen in an animal model of Parkinson's disease. *Pharmacol. Biochem. Behav.* 78, 513–522. <https://doi.org/10.1016/j.pbb.2004.04.022>.
- Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M.F., Conway, S.J., Ng, L.G., Stanley, E.R., Samokhvalov, I.M., Merad, M., 2010. Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science* 330, 841–845. <https://doi.org/10.1126/science.1194637>.
- Gorski, R.A., Gordon, J.H., Shryne, J.E., Southam, A.M., 1978. Evidence for a morphological sex difference within the medial preoptic area of the rat brain. *Brain Res.* 148, 333–346.
- Guerreiro, R., Wojtas, A., Bras, J., Carrasquillo, M., Rogava, E., Majounie, E., Cruchaga, C., Sassi, C., Kauwe, J.S., Younkin, S., Hazrati, L., Collinge, J., Pocock, J., Lashley, T., Williams, J., Lambert, J.C., Amouyel, P., Goate, A., Rademakers, R., Morgan, K., Powell, J., St George-Hyslop, P., Singleton, A., Hardy, J., Alzheimer Genetic Analysis, G., 2013. TREM2 variants in Alzheimer's disease. *N. Engl. J. Med.* 368, 117–127. <https://doi.org/10.1056/NEJMoa1211851>.
- Gunevskaya, D., Ivanov, A., Hernandez, D.P., Haage, V., Wojtas, B., Meyer, N., Maricos, M., Jordan, P., Buonfiglioli, A., Gielniewski, B., Ochocka, N., Comert, C., Friedrich, C., Artiles, L.S., Kaminska, B., Mertins, P., Beule, D., Kettenmann, H., Wolf, S.A., 2018. Transcriptional and translational differences of microglia from male and female brains. 2773–2783 e2776. *Cell Rep.* 24. <https://doi.org/10.1016/j.celrep.2018.08.001>.
- Hagino, Y., Kariura, Y., Manago, Y., Amano, T., Wang, B., Sekiguchi, M., Nishikawa, K., Aoki, S., Wada, K., Noda, M., 2004. Heterogeneity and potentiation of AMPA type of glutamate receptors in rat cultured microglia. *Glia* 47, 68–77. <https://doi.org/10.1002/glia.20100>.

- 1002/glia.20034.
- Hanamsagar, R., Alter, M.D., Block, C.S., Sullivan, H., Bolton, J.L., Bilbo, S.D., 2017. Generation of a microglial developmental index in mice and in humans reveals a sex difference in maturation and immune reactivity. *Glia* 65, 1504–1520. <https://doi.org/10.1002/glia.23176>.
- Herbomel, P., Thisse, B., Thisse, C., 1999. Ontogeny and behaviour of early macrophages in the zebrafish embryo. *Development* 126, 3735–3745.
- Joel, D., Berman, Z., Tavor, I., Wexler, N., Gaber, O., Stein, Y., Shefi, N., Pool, J., Urchs, S., Margulies, D.S., Liem, F., Hanggi, J., Jancke, L., Assaf, Y., 2015. Sex beyond the genitalia: the human brain mosaic. *Proc. Natl. Acad. Sci. USA* 112, 15468–15473. <https://doi.org/10.1073/pnas.1509654112>.
- Joel, D., McCarthy, M.M., 2017. Incorporating sex as a biological variable in neuropsychiatric research: where are we now and where should we be? *Neuropsychopharmacology* 42, 379–385. <https://doi.org/10.1038/npp.2016.79>.
- Johnson, R.T., Breedlove, S.M., Jordan, C.L., 2008. Sex differences and laterality in astrocyte number and complexity in the adult rat medial amygdala. *J. Comp. Neurol.* 511, 599–609. <https://doi.org/10.1002/cne.21859>.
- Juraska, J.M., Fitch, J.M., Henderson, C., Rivers, N., 1985. Sex differences in the dendritic branching of dentate granule cells following differential experience. *Brain Res.* 333, 73–80.
- Kang, H.J., Kawasawa, Y.I., Cheng, F., Zhu, Y., Xu, X., Li, M., Sousa, A.M., Pletikos, M., Meyer, K.A., Sedmak, G., Guennel, T., Shin, Y., Johnson, M.B., Krsnik, Z., Mayer, S., Fertuzinhos, S., Umlauf, S., Lisgo, S.N., Vortmeyer, A., Weinberger, D.R., Mane, S., Hyde, T.M., Huttner, A., Reimers, M., Kleinman, J.E., Sestan, N., 2011. Spatio-temporal transcriptome of the human brain. *Nature* 478, 483–489. <https://doi.org/10.1038/nature10523>.
- Keren-Shaul, H., Spinrad, A., Weiner, A., Matcovitch-Natan, O., Dvir-Szternfeld, R., Ulland, T.K., David, E., Baruch, K., Lara-Astaiso, D., Toth, B., Itzkovitz, S., Colonna, M., Schwartz, M., Amit, I., 2017. A unique microglia type associated with restricting development of Alzheimer's disease. 1276-1290 e1217. *Cell* 169. <https://doi.org/10.1016/j.cell.2017.05.018>.
- Kettenmann, H., Hanisch, U.K., Noda, M., Verkhratsky, A., 2011. Physiology of microglia. *Physiol. Rev.* 91, 461–553. <https://doi.org/10.1152/physrev.00011.2010>.
- Kettenmann, H., Kirchhoff, F., Verkhratsky, A., 2013. Microglia: new roles for the synaptic stripper. *Neuron* 77, 10–18. <https://doi.org/10.1016/j.neuron.2012.12.023>.
- Kohman, R.A., Bhattacharya, T.K., Wojcik, E., Rhodes, J.S., 2013. Exercise reduces activation of microglia isolated from hippocampus and brain of aged mice. *J. Neuroinflammation* 10, 114. <https://doi.org/10.1186/1742-2094-10-114>.
- Koizumi, S., Ohsawa, K., Inoue, K., Kohsaka, S., 2013. Purinergic receptors in microglia: functional modal shifts of microglia mediated by P2 and P1 receptors. *Glia* 61, 47–54. <https://doi.org/10.1002/glia.22358>.
- Kopec, A.M., Smith, C.J., Ayre, N.R., Sweat, S.C., Bilbo, S.D., 2018. Microglial dopamine receptor elimination defines sex-specific nucleus accumbens development and social behavior in adolescent rats. *Nat. Commun.* 9, 3769. <https://doi.org/10.1038/s41467-018-06118-z>.
- Krasemann, S., Madore, C., Cialic, R., Baufeld, C., Calcagno, N., El Fatimy, R., Beckers, L., O'Loughlin, E., Xu, Y., Fanek, Z., Greco, D.J., Smith, S.T., Tweet, G., Humulock, Z., Zrzavy, T., Conde-Sanroman, P., Gacias, M., Weng, Z., Chen, H., Tjon, E., Mazaheri, F., Hartmann, K., Madi, A., Ulrich, J.D., Glatzel, M., Worthmann, A., Heeren, J., Budnik, B., Lemere, C., Ikezu, T., Heppner, F.L., Litvak, V., Holtzman, D.M., Lassmann, H., Weiner, H.L., Ochando, J., Haass, C., Butovsky, O., 2017. The TREM2-APOE pathway drives the transcriptional phenotype of dysfunctional microglia in neurodegenerative diseases. 566–581 e569. *Immunity* 47. <https://doi.org/10.1016/j.immuni.2017.08.008>.
- Kuo, J., Hamid, N., Bondar, G., Dewing, P., Clarkson, J., Micevych, P., 2010. Sex differences in hypothalamic astrocyte response to estradiol stimulation. *Biol. Sex Differ.* 1, 7. <https://doi.org/10.1186/2042-6410-1-7>.
- Lawson, L.J., Perry, V.H., Dri, P., Gordon, S., 1990. Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience* 39, 151–170.
- Lenz, K.M., McCarthy, M.M., 2010. Organized for sex - steroid hormones and the developing hypothalamus. *Eur. J. Neurosci.* 32, 2096–2104. <https://doi.org/10.1111/j.1460-9568.2010.07511.x>.
- Lenz, K.M., Nugent, B.M., Haliyur, R., McCarthy, M.M., 2013. Microglia are essential to masculinization of brain and behavior. *J. Neurosci.* 33, 2761–2772. <https://doi.org/10.1523/JNEUROSCI.1268-12.2013>.
- Lenz, K.M., Nugent, B.M., McCarthy, M.M., 2012. Sexual differentiation of the rodent brain: dogma and beyond. *Front. Neurosci.* 6, 26. <https://doi.org/10.3389/fnins.2012.00026>.
- Li, R., Cui, J., Shen, Y., 2014. Brain sex matters: estrogen in cognition and Alzheimer's disease. *Mol. Cell. Endocrinol.* 389, 13–21. <https://doi.org/10.1016/j.mce.2013.12.018>.
- Li, Y., Du, X.F., Liu, C.S., Wen, Z.L., Du, J.L., 2012. Reciprocal regulation between resting microglial dynamics and neuronal activity in vivo. *Dev. Cell* 23, 1189–1202. <https://doi.org/10.1016/j.devcel.2012.10.027>.
- Loram, L.C., Sholar, P.W., Taylor, F.R., Wiesler, J.L., Babb, J.A., Strand, K.A., Berkelhammer, D., Day, H.E., Maier, S.F., Watkins, L.R., 2012. Sex and estradiol influence glial pro-inflammatory responses to lipopolysaccharide in rats. *Psychoneuroendocrinology* 37, 1688–1699. <https://doi.org/10.1016/j.psyneuen.2012.02.018>.
- Mallat, M., Marín-Teva, J.L., Chéret, C., 2005. Phagocytosis in the developing CNS: more than clearing the corpses. *Curr. Opin. Neurobiol.* 15, 101–107.
- Mangold, C.A., Wronowski, B., Du, M., Masser, D.R., Hadad, N., Bixler, G.V., Brucklacher, R.M., Ford, M.M., Sonntag, W.E., Freeman, W.M., 2017. Sexually divergent induction of microglial-associated neuroinflammation with hippocampal aging. *J. Neuroinflammation* 14, 141. <https://doi.org/10.1186/s12974-017-0920-8>.
- Mapplebeck, J.C., Beggs, S., Salter, M.W., 2017. Molecules in pain and sex: a developing story. *Mol. Brain* 10, 9. <https://doi.org/10.1186/s13041-017-0289-8>.
- Maren, S., De Oca, B., Fanselow, M.S., 1994. Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP and contextual learning. *Brain Res.* 661, 25–34.
- Matcovitch-Natan, O., Winter, D.R., Giladi, A., Vargas Aguilar, S., Spinrad, A., Sarrazin, S., Ben-Yehuda, H., David, E., Zelada Gonzalez, F., Perrin, P., Keren-Shaul, H., Gury, M., Lara-Astaiso, D., Thaiss, C.A., Cohen, M., Bahar Halpern, K., Baruch, K., Deczkowska, A., Lorenzo-Vivas, E., Itzkovitz, S., Elinav, E., Sieweke, M.H., Schwartz, M., Amit, I., 2016. Microglia development follows a stepwise program to regulate brain homeostasis. *Science* 353, aad8670. <https://doi.org/10.1126/science.aad8670>.
- McCarthy, M.M., Auger, A.P., Bale, T.L., De Vries, G.J., Dunn, G.A., Forger, N.G., Murray, E.K., Nugent, B.M., Schwarz, J.M., Wilson, M.E., 2009. The epigenetics of sex differences in the brain. *J. Neurosci.* 29, 12815–12823. <https://doi.org/10.1523/JNEUROSCI.3331-09.2009>.
- McGeer, P.L., McGeer, E.G., 1996. Anti-inflammatory drugs in the fight against Alzheimer's disease. *Ann. N. Y. Acad. Sci.* 777, 213–220.
- McGrath, J., Saha, S., Welham, J., El Saadi, O., MacCauley, C., Chant, D., 2004. A systematic review of the incidence of schizophrenia: the distribution of rates and the influence of sex, urbanicity, migrant status and methodology. *BMC Med* 2, 13. <https://doi.org/10.1186/1741-7015-2-13>.
- Mhatre, S.D., Tsai, C.A., Rubin, A.J., James, M.L., Andreasson, K.I., 2015. Microglial malfunction: the third rail in the development of Alzheimer's disease. *Trends Neurosci.* 38, 621–636. <https://doi.org/10.1016/j.tins.2015.08.006>.
- Mong, J.A., McCarthy, M.M., 1999. Steroid-induced developmental plasticity in hypothalamic astrocytes: implications for synaptic patterning. *J. Neurobiol.* 40, 602–619.
- Mouton, P.R., Long, J.M., Lei, D.L., Howard, V., Jucker, M., Calhoun, M.E., Ingram, D.K., 2002. Age and gender effects on microglia and astrocyte numbers in brains of mice. *Brain Res.* 956, 30–35.
- Mrdjen, D., Pavlovic, A., Hartmann, F.J., Schreiner, B., Utz, S.G., Leung, B.P., Lelios, I., Heppner, F.L., Kipnis, J., Merkler, D., Greter, M., Becher, B., 2018. High-dimensional single-cell mapping of central nervous system immune cells reveals distinct myeloid subsets in health, aging, and disease. 380–395 e386. *Immunity* 48. <https://doi.org/10.1016/j.immuni.2018.01.011>.
- Murphy, S.J., McCullough, L.D., Smith, J.M., 2004. Stroke in the female: role of biological sex and estrogen. *ILAR J.* 45, 147–159.
- Paolicelli, R.C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira, T.A., Guiducci, E., Dumas, L., Ragozzino, D., Gross, C.T., 2011. Synaptic pruning by microglia is necessary for normal brain development. *Science* 333, 1456–1458. <https://doi.org/10.1126/science.1202529>.
- Parkhurst, C.N., Yang, G., Nanan, I., Savas, J.N., Yates 3rd, J.R., Lafaille, J.J., Hempstead, B.L., Littman, D.R., Gan, W.B., 2013. Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor. *Cell* 155, 1596–1609. <https://doi.org/10.1016/j.cell.2013.11.030>.
- Pepe, G., Calderazzi, G., De Maglie, M., Villa, A.M., Vegeto, E., 2014. Heterogeneous induction of microglia M2a phenotype by central administration of interleukin-4. *J. Neuroinflammation* 11, 211. <https://doi.org/10.1186/s12974-014-0211-6>.
- Plassman, B.L., Langa, K.M., McCammon, R.J., Fisher, G.G., Potter, G.G., Burke, J.R., Steffens, D.C., Foster, N.L., Giordani, B., Unverzagt, F.W., Welsh-Bohmer, K.A., Heeringa, S.G., Weir, D.R., Wallace, R.B., 2011. Incidence of dementia and cognitive impairment, not dementia in the United States. *Ann. Neurol.* 70, 418–426. <https://doi.org/10.1002/ana.22362>.
- Pocock, J.M., Kettenmann, H., 2007. Neurotransmitter receptors on microglia. *Trends Neurosci.* 30, 527–535. <https://doi.org/10.1016/j.tins.2007.07.007>.
- Prilutsky, D., Kho, A.T., Feiglin, A., Hammond, T., Stevens, B., Kohane, I.S., 2017. Sexual dimorphism of complement-dependent microglial synaptic pruning and other immune pathways in the developing brain. *bioRxiv*. <https://doi.org/10.1101/204412>.
- Quadros, P.S., Pfau, J.L., Wagner, C.K., 2007. Distribution of progesterone receptor immunoreactivity in the fetal and neonatal rat forebrain. *J. Comp. Neurol.* 504, 42–56. <https://doi.org/10.1002/cne.21427>.
- Riecher-Rossler, A., 2017. Sex and gender differences in mental disorders. *Lancet Psychiat.* 4, 8–9. [https://doi.org/10.1016/S2215-0366\(16\)30348-0](https://doi.org/10.1016/S2215-0366(16)30348-0).
- Rizzi, N., Rebecchi, M., Levandis, G., Ciana, P., Maggi, A., 2017. Identification of novel loci for the generation of reporter mice. e37. *Nucleic Acids Res.* 45. <https://doi.org/10.1093/nar/gkw1142>.
- Rogers, J., Strohmeyer, R., Kovelowski, C.J., Li, R., 2002. Microglia and inflammatory mechanisms in the clearance of amyloid beta peptide. *Glia* 40, 260–269. <https://doi.org/10.1002/glia.10153>.
- Rogers, J.T., Morganti, J.M., Bachstetter, A.D., Hudson, C.E., Peters, M.M., Grimmig, B.A., Weeber, E.J., Bickford, P.C., Gemma, C., 2011. CX3CR1 deficiency leads to impairment of hippocampal cognitive function and synaptic plasticity. *J. Neurosci.* 31, 16241–16250. <https://doi.org/10.1523/JNEUROSCI.3667-11.2011>.
- Roy-O'Reilly, M., McCullough, L.D., 2014. Sex differences in stroke: the contribution of coagulation. *Exp. Neurol.* 259, 16–27. <https://doi.org/10.1016/j.expneurol.2014.02.011>.
- Saijo, K., Collier, J.G., Li, A.C., Katzenellenbogen, J.A., Glass, C.K., 2011. An ADIOL-ERbeta-CtBP transrepression pathway negatively regulates microglia-mediated inflammation. *Cell* 145, 584–595. <https://doi.org/10.1016/j.cell.2011.03.050>.
- Salter, M.W., Beggs, S., 2014. Sublime microglia: expanding roles for the guardians of the CNS. *Cell* 158, 15–24. <https://doi.org/10.1016/j.cell.2014.06.008>.
- Schafer, D.P., Lehrman, E.K., Stevens, B., 2013. The “quad-partite” synapse: microglia-synapse interactions in the developing and mature CNS. *Glia* 61, 24–36. <https://doi.org/10.1002/glia.22389>.
- Schenk, D.B., Yednock, T., 2002. The role of microglia in Alzheimer's disease: friend or foe? *Neurobiol. Aging* 23 677–679; discussion 683–674.

- Schwarz, J.M., McCarthy, M.M., 2008. Steroid-induced sexual differentiation of the developing brain: multiple pathways, one goal. *J. Neurochem.* 105, 1561–1572. <https://doi.org/10.1111/j.1471-4159.2008.05384.x>.
- Schwarz, J.M., Sholar, P.W., Bilbo, S.D., 2012. Sex differences in microglial colonization of the developing rat brain. *J. Neurochem.* 120, 948–963. <https://doi.org/10.1111/j.1471-4159.2011.07630.x>.
- Seshadri, S., Wolf, P.A., Beiser, A., Au, R., McNulty, K., White, R., D'Agostino, R.B., 1997. Lifetime risk of dementia and Alzheimer's disease. The impact of mortality on risk estimates in the Framingham Study. *Neurology* 49, 1498–1504.
- Sierra, A., Encinas, J.M., Deudero, J.J., Chancey, J.H., Enikolopov, G., Overstreet-Wadiche, L.S., Tsirka, S.E., Maletic-Savatic, M., 2010. Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell* 7, 483–495. <https://doi.org/10.1016/j.stem.2010.08.014>.
- Sierra, A., Gottfried-Blackmore, A., Milner, T.A., McEwen, B.S., Bulloch, K., 2008. Steroid hormone receptor expression and function in microglia. *Glia* 56, 659–674. <https://doi.org/10.1002/glia.20644>.
- Solito, E., Sastre, M., 2012. Microglia function in Alzheimer's disease. *Front. Pharmacol.* 3, 14. <https://doi.org/10.3389/fphar.2012.00014>.
- Song, W.M., Joshita, S., Zhou, Y., Ulland, T.K., Gilfillan, S., Colonna, M., 2018. Humanized TREM2 mice reveal microglia-intrinsic and -extrinsic effects of R47H polymorphism. *J. Exp. Med.* 215, 745–760. <https://doi.org/10.1084/jem.20171529>.
- Sorge, R.E., Mapplebeck, J.C., Rosen, S., Beggs, S., Taves, S., Alexander, J.K., Martin, L.J., Austin, J.S., Sotocinal, S.G., Chen, D., Yang, M., Shi, X.Q., Huang, H., Pilon, N.J., Bilan, P.J., Tu, Y., Klip, A., Ji, R.R., Zhang, J., Salter, M.W., Mogil, J.S., 2015. Different immune cells mediate mechanical pain hypersensitivity in male and female mice. *Nat. Neurosci.* 18, 1081–1083. <https://doi.org/10.1038/nn.4053>.
- Suarez, I., Bodega, G., Rubio, M., Fernandez, B., 1991. Sexual dimorphism in the distribution of glial fibrillary acidic protein in the supraoptic nucleus of the hamster. *J. Anat.* 178, 79–82.
- Suarez, I., Bodega, G., Rubio, M., Fernandez, B., 1992. Sexual dimorphism in the hamster cerebellum demonstrated by glial fibrillary acidic protein (GFAP) and vimentin immunoreactivity. *Glia* 5, 10–16. <https://doi.org/10.1002/glia.440050103>.
- Swaab, D.F., Chung, W.C., Kruijver, F.P., Hofman, M.A., Ishunina, T.A., 2001. Structural and functional sex differences in the human hypothalamus. *Horm. Behav.* 40, 93–98. <https://doi.org/10.1006/hbeh.2001.1682>.
- Taves, S., Berta, T., Liu, D.L., Gan, S., Chen, G., Kim, Y.H., Van de Ven, T., Laufer, S., Ji, R.R., 2016. Spinal inhibition of p38 MAP kinase reduces inflammatory and neuropathic pain in male but not female mice: Sex-dependent microglial signaling in the spinal cord. *Brain Behav. Immun.* 55, 70–81. <https://doi.org/10.1016/j.bbi.2015.10.006>.
- Thion, M.S., Low, D., Silvin, A., Chen, J., Grisel, P., Schulte-Schrepping, J., Blecher, R., Ulas, T., Squarzone, P., Hoeffel, G., Couplier, F., Siopi, E., David, F.S., Scholz, C., Shihui, F., Lum, J., Amoyo, A.A., Larbi, A., Poidinger, M., Buttgerit, A., Lledo, P.M., Greter, M., Chan, J.K.Y., Amit, I., Beyer, M., Schultze, J.L., Schlitzer, A., Pettersson, S., Ginhoux, F., Garel, S., 2018. Microbiome influences prenatal and adult microglia in a sex-specific manner. 500–516 e516. *Cell* 172. <https://doi.org/10.1016/j.cell.2017.11.042>.
- Tremblay, M.E., Lowery, R.L., Majewska, A.K., 2010. Microglial interactions with synapses are modulated by visual experience. *PLoS Biol.* 8, e1000527. <https://doi.org/10.1371/journal.pbio.1000527>.
- Tremblay, M.E., Stevens, B., Sierra, A., Wake, H., Bessis, A., Nimmerjahn, A., 2011. The role of microglia in the healthy brain. *J. Neurosci.* 31, 16064–16069. <https://doi.org/10.1523/JNEUROSCI.4158-11.2011>.
- Vegeto, E., Belcredito, S., Ghisletti, S., Meda, C., Eteri, S., Maggi, A., 2006. The endogenous estrogen status regulates microglia reactivity in animal models of neuroinflammation. *Endocrinology* 147, 2263–2272. <https://doi.org/10.1210/en.2005-1330>.
- Vegeto, E., Bonincontro, C., Pollio, G., Sala, A., Viappiani, S., Nardi, F., Brusadelli, A., Viviani, B., Ciana, P., Maggi, A., 2001. Estrogen prevents the lipopolysaccharide-induced inflammatory response in microglia. *J. Neurosci.* 21, 1809–1818.
- Villa, A., Gelosa, P., Castiglioni, L., Cimino, M., Rizzi, N., Pepe, G., Lolli, F., Marcelllo, E., Sironi, L., Vegeto, E., Maggi, A., 2018. Sex-specific features of microglia from adult mice. *Cell Rep.* 23, 3501–3511. <https://doi.org/10.1016/j.celrep.2018.05.048>.
- Villa, A., Rizzi, N., Vegeto, E., Ciana, P., Maggi, A., 2015. Estrogen accelerates the resolution of inflammation in macrophagic cells. *Sci. Rep.* 5, 15224. <https://doi.org/10.1038/srep15224>.
- Villa, A., Vegeto, E., Poletti, A., Maggi, A., 2016. Estrogens, neuroinflammation, and neurodegeneration. *Endocr. Rev.* 37, 372–402. <https://doi.org/10.1210/er.2016-1007>.
- Wake, H., Moorhouse, A.J., Jinno, S., Kohsaka, S., Nabekura, J., 2009. Resting microglia directly monitor the functional state of synapses in vivo and determine the fate of ischemic terminals. *J. Neurosci.* 29, 3974–3980. <https://doi.org/10.1523/JNEUROSCI.4363-08.2009>.
- Weinhard, L., Neniskyte, U., Vadisiute, A., di Bartolomei, G., Ayygun, N., Riviere, L., Zonfrillo, F., Dymecki, S., Gross, C., 2018. Sexual dimorphism of microglia and synapses during mouse postnatal development. *Dev. Neurobiol.* 78, 618–626. <https://doi.org/10.1002/dneu.22568>.
- Werling, D.M., Geschwind, D.H., 2013. Sex differences in autism spectrum disorders. *Curr. Opin. Neurol.* 26, 146–153. <https://doi.org/10.1097/WCO.0b013e32835ee548>.
- Werling, D.M., Parikshak, N.N., Geschwind, D.H., 2016. Gene expression in human brain implicates sexually dimorphic pathways in autism spectrum disorders. *Nat. Commun.* 7, 10717. <https://doi.org/10.1038/ncomms10717>.
- Wu, M.V., Manoli, D.S., Fraser, E.J., Coats, J.K., Tollkuhn, J., Honda, S., Harada, N., Shah, N.M., 2009. Estrogen masculinizes neural pathways and sex-specific behaviors. *Cell* 139, 61–72. <https://doi.org/10.1016/j.cell.2009.07.036>.
- Yamasaki, R., Lu, H., Butovsky, O., Ohno, N., Rietsch, A.M., Cialic, R., Wu, P.M., Doykan, C.E., Lin, J., Cotleur, A.C., Kidd, G., Zorlu, M.M., Sun, N., Hu, W., Liu, L., Lee, J.C., Taylor, S.E., Uehlein, L., Dixon, D., Gu, J., Floruta, C.M., Zhu, M., Charo, I.F., Weiner, H.L., Ransohoff, R.M., 2014. Differential roles of microglia and monocytes in the inflamed central nervous system. *J. Exp. Med.* 211, 1533–1549. <https://doi.org/10.1084/jem.20132477>.
- Zhang, B., Gaiteri, C., Bodea, L.G., Wang, Z., McElwee, J., Podtezhnikov, A.A., Zhang, C., Xie, T., Tran, L., Dobrin, R., Fluder, E., Clurman, B., Melquist, S., Narayanan, M., Suver, C., Shah, H., Mahajan, M., Gillis, T., Mysore, J., MacDonald, M.E., Lamb, J.R., Bennett, D.A., Molony, C., Stone, D.J., Gudnason, V., Myers, A.J., Schadt, E.E., Neumann, H., Zhu, J., Emilsson, V., 2013. Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* 153, 707–720. <https://doi.org/10.1016/j.cell.2013.03.030>.