



British Mycological  
Society promoting fungal science

journal homepage: [www.elsevier.com/locate/fbr](http://www.elsevier.com/locate/fbr)



## Review

# Killer toxin-like chitinases in filamentous fungi: Structure, regulation and potential roles in fungal biology



Georgios TZELEPIS<sup>a,\*</sup>, Magnus KARLSSON<sup>b</sup>

<sup>a</sup>Department of Plant Biology, Uppsala BioCenter and Linnean Center for Plant Biology, Swedish University of Agricultural Sciences, Box 7080, S-75007 Uppsala, Sweden

<sup>b</sup>Department of Forest Mycology and Plant Pathology, Uppsala BioCenter, Swedish University of Agricultural Sciences, Box 7026, S-75007 Uppsala, Sweden

### ARTICLE INFO

#### Article history:

Received 6 September 2018

Received in revised form

13 November 2018

Accepted 21 November 2018

#### Keywords:

C-group

Chitinases

Killer toxin

LysM domain

Mycoparasitism

### ABSTRACT

Fungal chitinases are hydrolytic enzymes responsible for degradation of chitin. Chitinases are involved in several aspects of fungal biology, including cell wall remodelling during hyphal growth, conidial germination, autolysis, mycoparasitism and nutrient acquisition. They are divided into three distinct phylogenetic groups; A, B and C. Chitinases from the C group show structural similarities with the killer toxin zymocin produced by the yeast *Kluyveromyces lactis* and it is speculated that they have a similar function in filamentous ascomycetes, by facilitating penetration of toxins into cells of competing individuals. Genome analyses show that certain fungal species with a mycoparasitic lifestyle contain high numbers of killer toxin-like chitinases, compared with specialized saprotrophs and plant pathogens. Recent developments within this research field have revealed considerable variation in the modular structure and regulation of killer toxin-like chitinases, suggesting more diverse roles than merely fungal-fungal interactions. In this review, we summarize the current knowledge about this intriguing class of chitinases, including their modular structure, evolution, gene regulation, and functional analyses in mycoparasitic as well as in saprotrophic species. We also propose important questions for future research.

© 2018 British Mycological Society. Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

Fungal cells are surrounded by a dynamic matrix, the cell wall, which provides the appropriate strength for protection against environmental stress conditions, such as osmotic pressure, toxins, mechanical injuries and parasite attack, while at the same time allowing flexibility, and rigidity

(Bowman and Free, 2006, Keegstra, 2010, Latge, 2007). The fungal cell wall mainly consists of chitin, mannans,  $\beta$ -1,3-glucan and proteins (Latge, 2007). Chitin is a biopolymer that consists of  $\beta$ -1,4-linked N-acetyl-D-glucosamine (GlcNAc) units. Chitin is not restricted to fungi, but is also found in the cuticle of insects and certain nematodes (Merzendorfer and Zimoch, 2003, Hill et al., 1991), in crustacean and mollusc

\* Corresponding author.

E-mail address: [Georgios.Tzelepis@slu.se](mailto:Georgios.Tzelepis@slu.se) (G. Tzelepis).

<https://doi.org/10.1016/j.fbr.2018.11.001>

1749-4613/© 2018 British Mycological Society. Published by Elsevier Ltd. All rights reserved.

shells (Kurita, 2006, Peters, 1972), and in protozoa and algae (Mulisch, 1993, Kapaun and Reisser 1995), thereby making it the second most distributed polymer in nature after cellulose (Gortari and Hours, 2013). However, it is absent from vertebrates and plants, which makes chitin a suitable target for drugs and pesticides (Spindler et al., 1990, Chaudhary et al., 2013). Plants can also recognize chitin from invading fungi by specific receptors and thereby activate basal defence responses leading to pattern-triggered immunity (PTI) (Boller and Felix, 2009, Newman et al., 2013). Chitin only constitutes 10–20 % of the total dry biomass in filamentous fungi, while it is even lower in yeasts (1–2 %) (Bartnicki-Garcia, 1968, de Nobel et al., 2000, Klis et al., 2002). Despite the relatively low proportion of chitin in fungal cell walls, it plays a crucial role for cell wall functioning by balancing plasticity and rigidity (Specht et al., 1996).

Chitin is hydrolysed by chitinases (EC.3.2.1.14), which cleave the  $\beta$ -1,4 glycan bond creating chitin oligomers or dimers (Gooday, 1990). Chitinases are present in a wide range of organisms, eukaryotic and prokaryotic, playing a crucial and vital role in chitin recycling in both marine and terrestrial ecosystems (Keyhani and Roseman, 1999). Fungal genomes contain a plethora of genes putatively encoding chitinolytic enzymes (Karlsson and Stenlid, 2008, Gruber and Seidl-Seiboth, 2012) that play multiple roles in fungal biology, including cell wall remodelling during growth, autolysis, hydrolysis of exogenous chitin for nutrient acquisition and attack of the cell wall of competing fungi for environmental niches (Kuranda and Robbins, 1991, Shin et al., 2009, Baker et al., 2009, Dunkler et al., 2005, Boldo et al., 2009). The most extreme form of competition can be found in mycoparasitic fungi that utilize chitinases to attack their fungal prey (Mamarabadi et al., 2008, Gruber and Seidl-Seiboth, 2012). According to their amino acid sequences, chitinases are classified in two distinct glycoside hydrolase families; 18 (GH18) and 19 (GH19) (Henrissat, 1991 Lombard et al., 2014). The latter is present in plant and bacteria, while fungal chitinases exclusively belong to GH18 (Karlsson and Stenlid, 2009). Furthermore, chitinases can be classified according to their preferred cleavage patterns into exochitinases, which cleave the chitin polymer from the exposed ends and endochitinases, which cleave the polymer at random positions (van Aalten et al., 2001, Horn et al., 2006). Phylogenetically, GH18 fungal chitinases are divided into three distinct groups (A, B and C) that can be further subdivided into subgroups (A-II, A-IV, A-V, B-I to B-VI, C-I, and C-II) (Seidl et al., 2005, Karlsson and Stenlid, 2008, Karlsson et al., 2016). Fungal GH18 enzymes from the B-V subgroup encode endo- $\beta$ -N-acetylglucosaminidases (Stals et al., 2010, Tzelepis et al., 2014a, 2017).


Chitinases in group C are predicted to be large proteins with a molecular weight higher than 200 kDa and show sequence and domain similarity to the  $\alpha/\beta$  subunits of the secreted zymocin killer toxin, which is produced by the dairy yeast *Kluyveromyces lactis* (Stark and Boyd, 1986, Magliani et al., 1997). The  $\alpha$ -subunit in zymocin displays exochitinase activity, thus degrading the cell wall chitin in competing yeasts, and together with the  $\beta$ -subunit facilitates the penetration of a third subunit ( $\gamma$ ) into the cytoplasm (Butler et al., 1991). The  $\gamma$ -subunit is a tRNase toxin responsible for inhibiting the

proliferation of *Saccharomyces cerevisiae* by arresting cell division in the G3 phase (Stark et al., 1990). The  $\alpha$ - and  $\beta$ -subunits are encoded and translated as a single polypeptide that is subsequently cleaved by a protease to produce two separate proteins, while the  $\gamma$ -subunit is encoded by a separate gene (Magliani et al., 1997). Because of this similarity with zymocin, -group C chitinases are also referred to as killer toxin-like chitinases, and are hypothesized to function in fungal antagonism by permeabilization of fungal cell walls to allow penetration of antifungal molecules (Seidl et al., 2005). The aim of this review article is to summarize the current knowledge regarding the group C killer toxin-like chitinases in filamentous fungi and to identify important knowledge gaps that should be addressed in future research.

## 2. Phylogeny and modular structure of killer toxin-like chitinases

Group C chitinases have been identified only in filamentous ascomycetes (Karlsson and Stenlid, 2009). Based on phylogenetic analyses of the GH18 module, group C chitinases are divided into two distinct subgroups, i.e. C-I and C-II (Seidl et al. 2005, Karlsson and Stenlid 2008). This sequence diversification of the GH18 module is also reflected in differences in the modular structure of the members of the two subgroups. Members from both subgroups typically contain a GH18 module with an intact catalytic DXXDXDXE motif indicating that they are active chitinolytic enzymes (van Aalten et al., 2001, Gruber et al., 2011a,b, Tzelepis et al., 2012, 2014b). They also typically have a N-terminal endoplasmic reticulum-targeting signal peptide, indicating that they are secreted out of the cell (Gruber et al., 2011a,b, Tzelepis et al., 2012, 2014b). A few group C chitinases, such as Chi18-1 from *Trichoderma reesei* and TAC6/TVC6 from the mycoparasites *T. atroviride* and *T. virens*, respectively, carry amino acid substitutions in the catalytic motif that may render them inactive (Karlsson and Stenlid, 2009, Gruber et al., 2011a). Subgroup C-I chitinases are further characterized by the presence of chitin-binding family 18 carbohydrate-binding modules (CBM18) together with the GH18 catalytic module located at the N-terminal of the protein (Fig. 1A). In contrast, subgroup C-II chitinases contain both CBM18 and peptidoglycan/chitin-binding CBM50 LysM modules, located at the N-terminal side of the centrally located GH18 module (Fig. 1A). Analysis of predicted 3D structures of zymocin and the C-II subgroup gh18-8 from *Neurospora crassa* showed high structural similarities between zymocin and C-II subgroup chitinases (Fig. 1B).

CBM18 modules are shown to bind to chitin (Boraston et al., 2004) and may function to increase the adherence of -group C chitinases to insoluble chitin substrates (Seidl-Seiboth et al., 2014). CBM50 modules were first studied in prokaryotic organisms and display binding affinity towards both chitin and peptidoglycans (Buist et al., 2008). They were subsequently described in many eukaryotic organisms including plants and fungi (Zhang et al., 2009, Seidl et al., 2005). In phytopathogenic fungi, LysM effector proteins may facilitate pathogen infections by binding and thus masking small chitin fragments from the host immune system, as is the case with the Ecp6 effector from the tomato pathogen *Cladosporium fulvum* (de



**Fig. 1 – Modular architecture of fungal group C killer toxin-like chitinases. A) Domain structures of zymocin and fungal C group chitinases as predicted by SMART software. B) 3D protein model of zymocin and *Neurospora crassa* gh18-8 (C-II) chitinase as predicted by raptorX software. Their modular structures contain the chitinase catalytic domain (GH18), a single secretion peptide (SP) and one or two chitin-binding family 18 carbohydrate-binding modules (CBM18). Chitinases at the C-II subgroup contain two additional LysM peptidoglycan-binding family 50 carbohydrate-binding modules (CBM50). In some cases, additional modules are identified including transmembrane domains (TM) and the Hce2 homologue of the Ecp2 effector. Domain structure is drawn schematically and not to scale.**

Jonge *et al.*, 2010). In addition, the TAL6 CBM50 module-containing protein from the mycoparasitic species *T. atroviride* is shown to inhibit germination of its own spores, thereby expanding the role of LysM proteins to regulation of fungal developmental processes (Seidl-Seiboth *et al.*, 2013). The exact role of the CBM50 LysM modules present in group C chitinases has not been studied, but may function to enhance binding to the cell walls of fungal antagonists. In addition to CBM18 and CBM50 modules, certain subgroup C-II chitinases in *Aspergillus nidulans* and *N. crassa* are predicted to contain one or multiple transmembrane motifs in the C-terminal part (Tzelepis *et al.*, 2012, 2014b), indicating that these proteins are likely localized

in the plasma membrane such as gh18-8 in *N. crassa* and chi2-3 in *A. nidulans* (Fig. 1A).

Interestingly, Stergiopoulos *et al.*, (2012) reported that a small protein termed Hce2, which is homologous to the *C. fulvum* Ecp2 effector, is fused to the C-terminal of certain C-II group chitinases (Fig. 1A). These GH18-Hce2 proteins are found more frequently in saprotrophic species and in pathogens of humans, insects and mycoparasites, rather than in plant pathogens (Stergiopoulos *et al.*, 2012). The Ecp2 effector protein was previously identified in phytopathogenic species such as *Mycosphaerella graminicola* and *M. fijiensis* (Stergiopoulos *et al.*, 2010), and was shown to be a virulence

factor during plant pathogenesis (Lauge *et al.*, 1997). Although the exact function of the Ecp2 effector is not known, it is speculated that it triggers necrosis in host plants, similar to Avr2 in *C. fulvum* (Stergiopoulos *et al.*, 2010). In addition, the Chi18-10 group C chitinase from *T. reesei* also contains an epidermal growth factor-1-like module (EGF-1), shown to be involved in protein-protein interactions (Wouters *et al.*, 2005), at the C-terminal of the protein (Seidl *et al.*, 2005). The functions of the Hce2 and EGF-1 modules - are unknown, but they may represent toxic peptides playing a role in antagonist interactions, in analogy with the zymocin  $\gamma$ -subunit.

### 3. Evolution and genomic organization of killer toxin-like chitinases

The highest numbers of group C chitinases are found in *T. virens* (15 genes), *A. nidulans* (11 genes), the truffle mycoparasite *Tolypocladium ophioglossoides* (10 genes) and *T. atroviride* (9 genes) (Karlsson and Stenlid 2008, Karlsson *et al.*, 2018). Evolutionary analyses among *Trichoderma* species with predominantly saprophytic (*T. reesei*) or mycoparasitic (*T. atroviride* and *T. virens*) lifestyles, indicated selection for gene contractions and expansions, respectively. Specifically, *T. reesei* contains only four C-group chitinase genes, while *T. atroviride* and *T. virens* display an increased number of this type of chitinases, containing nine and fifteen genes, respectively as mentioned above (Ihrmark *et al.*, 2010). High numbers of C-group chitinases in *Trichoderma* thus appears to be related with aggressiveness and broad host range mycoparasitism (Kubicek *et al.*, 2011). However, recent genome sequencing of more mycoparasitic species suggests that this correlation may not hold true, as the number of C-group killer toxin-like chitinases range from two to fifteen genes in different species (Karlsson *et al.*, 2018), indicating reliance on different mechanisms during the mycoparasitic attack (Karlsson *et al.*, 2018). Selection for increased numbers of C-group chitinases is also implicated in other soil-borne ascomycetes including *A. nidulans* and *Uncinocarpus reesii* (Karlsson and Stenlid 2008). Furthermore, low levels of sequence identity between C-group chitinases (Ihrmark *et al.*, 2010), low number of orthologs (only five orthologs are identified between *T. atroviride* and *T. virens*) (Gruber *et al.*, 2011b) and the localization in small non-syntenic clusters at scaffold ends (Gruber *et al.*, 2011b), suggests that killer toxin-like chitinases evolve through a birth-and-death process followed by sequence diversification (Seidl-Seiboth *et al.*, 2014).

Interestingly, C-group chitinase genes are frequently clustered with genes containing only one or multiple CBM50 LysM motifs but not any catalytic domains (Kubicek *et al.*, 2011, Gruber *et al.*, 2011a). The function of these CBM50 LysM proteins remain unclear, but expression data show that they are perfectly co-regulated with their respective C-group chitinase genes in *T. atroviride*, indicating a role in fungal-fungal interactions, similar to LysM effectors in plant pathogens (de Jonge and Thomma, 2009). The LysM protein TAL6, whose gene is associated with the group C *tac6* chitinase gene in *T. atroviride*, showed strong inhibition activity of *T. atroviride* conidial germination but not against conidia from other

species, suggesting a role in regulation of fungal growth (Seidl-Seiboth *et al.*, 2013).

### 4. Regulation of killer toxin-like chitinase genes

The transcriptional patterns of subgroup C-I and C-II chitinases have been studied thoroughly in saprotrophic model fungi such as *N. crassa* and *A. nidulans* but also in several mycoparasitic species. A summary of the available gene regulation data is presented in Table 1. In *N. crassa*, the C-II *gh18-6* and *gh18-8* genes were induced during interactions with the ascomycete *Fusarium sporotrichioides* as compared to growth on carbon rich media, while the *gh18-8* gene was induced during carbon starvation conditions (Tzelepis *et al.*, 2012). However, different expression patterns were observed upon interactions with other fungal species. For instance, the *gh18-6* gene was induced during interactions with *Botrytis cinerea*, but not during confrontation with the basidiomycete *R. solani*, whereas *gh18-8* was induced during *N. crassa* self-interaction (Tzelepis *et al.*, 2012). The third group C gene in *N. crassa*, *gh18-9* from subgroup C-I, was induced during carbon starvation conditions but not during fungal-fungal interactions (Tzelepis *et al.*, 2012).

In *A. nidulans*, the four subgroup C-II genes *chiC2-1*, *chiC2-2*, *chiC2-3* and *chiC2-4* were all highly induced during interactions with *B. cinerea* and *R. solani*, while no induction was observed upon interaction with the oomycete *Phytophthora niederhauerii* that lacks chitin in the cell wall (Tzelepis *et al.*, 2014b). *ChiC2-2* and *chiC2-3* were also up-regulated during interactions with *F. sporotrichioides*. In contrast with the induced expression during interactions with living *R. solani*, all genes were down-regulated during growth on media where *R. solani* cell wall material was used as the sole carbon source. All genes were expressed, but not induced, during growth on chitin (Tzelepis *et al.*, 2014b).

Expression patterns of group C chitinase genes in the aggressive mycoparasite *T. atroviride* have also been studied. Seidl *et al.*, (2005) showed that the *gh18-10* gene (later renamed to *tac3*) was induced during confrontation with *R. solani* mycelium and during growth on media containing *R. solani* cell wall material. However, in a later study, the eight group C chitinase genes in *T. atroviride* *tac1-tac8* (including *tac3*) were all shown to be induced during confrontation with *B. cinerea*, but not with *R. solani* or during *T. atroviride* self-interactions (Gruber *et al.*, 2011a). No expression was also observed on media containing *R. solani* cell wall material, except for the *tac6* gene (Gruber *et al.*, 2011a). Four genes, *tac2*, *tac3*, *tac6* and *tac7*, were shown to be expressed on chitin-containing media (Gruber *et al.*, 2011a). A possible involvement of the *tac2* and *tac6* gene products in hyphal growth and network formation comes from the induced expression in the central part of the mycelial colony but not in the peripheral zone (Gruber *et al.*, 2011b). As mentioned previously, several *tal* CBM50 LysM genes that are clustered together with group C chitinases in the *T. atroviride* genome were also co-regulated with *tac* chitinase gene expression during interactions with *B. cinerea* (Gruber *et al.*, 2011b).

**Table 1 – List of induced genes encoding for putative C-group chitinases in filamentous fungal species.**

Gene name	Species	Subgroup	Conditions	Reference
gh18-6	<i>N. crassa</i>	C-II	Interactions with <i>F.sporotrichoides</i> Interactions with <i>B. cinerea</i>	Tzelepis et al., (2012)
gh18-8	<i>N. crassa</i>	C-II	Interactions with <i>F.sporotrichoides</i> <i>N. crassa</i> self-Interactions Carbon starvation	Tzelepis et al., (2012)
gh18-9	<i>N. crassa</i>	C-I	Carbon starvation	Tzelepis et al., (2012)
chiC2-1	<i>A. nidulans</i>	C-II	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> <i>R. solani</i> cell wall material	Tzelepis et al., (2014b)
chiC2-2	<i>A. nidulans</i>	C-II	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> Interactions with <i>F.sporotrichoides</i>	Tzelepis et al., (2014b)
chiC2-3	<i>A. nidulans</i>	C-II	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> Interactions with <i>F.sporotrichoides</i>	Tzelepis et al., (2014b)
chiC2-4	<i>A. nidulans</i>	C-II	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i>	Tzelepis et al., (2014b)
tac1	<i>T. atroviride</i>	C-I	Interactions with <i>B. cinerea</i>	Gruber et al., (2011a)
tac2	<i>T. atroviride</i>	C-I	Interactions with <i>B. cinerea</i> Chitin-containing media	Gruber et al., (2011a)
tac3	<i>T. atroviride</i>	C-II	Interactions with <i>B. cinerea</i> Chitin-containing media	Gruber et al., (2011a)
tac4	<i>T. atroviride</i>	C-I	Interactions with <i>B. cinerea</i>	Gruber et al., (2011a)
tac5	<i>T. atroviride</i>	C-I	Interactions with <i>B. cinerea</i>	Gruber et al., (2011a)
tac6	<i>T. atroviride</i>	C-I	Interactions with <i>B. cinerea</i> <i>R. solani</i> cell wall material Chitin-containing media	Gruber et al., (2011a)
tac7	<i>T. atroviride</i>	C-II	Interactions with <i>B. cinerea</i> Chitin-containing media	Gruber et al., (2011a)
tac8	<i>T. atroviride</i>	C-I	Interactions with <i>B. cinerea</i>	Gruber et al., (2011a)
tvc2	<i>T. virens</i>	C-I	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> <i>B. cinerea</i> cell wall material <i>T. virens</i> cell wall material Colloidal chitin Hyphal network formation	Gruber et al., (2011b)
tvc3	<i>T. virens</i>	C-II	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> <i>B. cinerea</i> cell wall material <i>T. virens</i> cell wall material Colloidal chitin Carbon starvation Hyphal network formation	Gruber et al., (2011b)
tvc4	<i>T. virens</i>	C-I	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> <i>B. cinerea</i> cell wall material <i>R. solani</i> cell wall material <i>T. virens</i> cell wall material Colloidal chitin Crude chitin Carbon starvation Conidia formation Hyphal network formation	Gruber et al., (2011b)
tvc5	<i>T. virens</i>	C-I	<i>B. cinerea</i> cell wall material Colloidal chitin Carbon starvation Hyphal network formation	Gruber et al., (2011b)
tvc6	<i>T. virens</i>	C-I	<i>B. cinerea</i> cell wall material Colloidal chitin Carbon starvation Hyphal network formation	Gruber et al., (2011b)

(continued on next page)

Table 1 (continued)

Gene name	Species	Subgroup	Conditions	Reference
tvc7	<i>T. virens</i>	C-II	<i>B. cinerea</i> cell wall material <i>R. solani</i> cell wall material <i>T. virens</i> cell wall material Colloidal chitin Crude chitin Carbon starvation Hyphal network formation	Gruber et al., (2011b)
tvc9	<i>T. virens</i>	C-II	<i>B. cinerea</i> cell wall material <i>R. solani</i> cell wall material <i>T. virens</i> cell wall material Hyphal network formation Carbon starvation	Gruber et al., (2011b)
tvc10	<i>T. virens</i>	C-II	Interactions with <i>B. cinerea</i> Interactions with <i>R. solani</i> <i>B. cinerea</i> cell wall material Colloidal chitin Crude chitin Carbon starvation Hyphal network formation	Gruber et al., (2011b)
tvc11	<i>T. virens</i>	?	<i>B. cinerea</i> cell wall material Hyphal network formation	Gruber et al., (2011b)
tvc12	<i>T. virens</i>	C-I	<i>B. cinerea</i> cell wall material Carbon starvation Hyphal network formation	Gruber et al., (2011b)
tvc14	<i>T. virens</i>	C-II	<i>B. cinerea</i> cell wall material Hyphal network formation	Gruber et al., (2011b)
chiC1	<i>C. rosea</i>	C-I	Interactions with <i>B. cinerea</i> Interactions with <i>F. graminearum</i> colloidal chitin	Nygren et al., (2018) Tzelepis et al., (2015)
TOPH09828	<i>T. ophioglossoides</i>	?	Truffle mycoparasitism	Quandt et al., (2016)
94536	<i>M. thermophila</i>	C-II	Plant straw material	Kolbusz et al. (2014)

Investigating the expression profiles of group C chitinases in the mycoparasitic species *T. virens* showed that the regulation of these genes is complex and divergent from *T. atroviride* (Gruber et al., 2011b). Although *T. virens* possess fifteen group C chitinases, only four genes (*tvc2*, *tvc3*, *tvc4* and *tvc10*) were induced during confrontation with *B. cinerea* and *R. solani* compared with when *T. virens* was grown alone on solid medium, but notably, not compared with a *T. virens* self-interaction (Gruber et al., 2011b). In contrast, eleven out of the fifteen genes were up-regulated during growth on medium containing *B. cinerea* cell wall material, while four of them were induced on medium containing *R. solani* cell wall material and five during growth on *T. virens* cell walls (Gruber et al., 2011b). Even differences in chitin quality seem to influence group C chitinase gene expression in *T. virens*; seven genes were induced during growth on colloidal chitin versus three on crude chitin. Regulation by developmental stage appear to be important for group C chitinases in *T. virens* as all genes, except for the four putative pseudogenes *tvc1*, *tvc8*, *tvc13* and *tvc15*, were induced during hyphal network formation. *Tvc4* was also induced in conidia (Gruber et al., 2011b) (Table 1).


Gene expression of group C chitinases has also been studied in other mycoparasitic species. The subgroup C-I gene *chiC1* in *Clonostachys rosea* was induced during confrontation with *F. graminearum*, but not during interactions with *B. cinerea* (Nygren et al., 2018). However, neither *chiC1* nor the subgroup C-II gene *chiC2* were induced when co-cultivated with *F.*

*graminearum* in liquid cultures (Tzelepis et al., 2015), again emphasizing that the developmental stage and physiological status may influence the regulation of group C chitinase genes. The *chiC1* gene was induced during growth on medium containing chitin as the sole carbon source (Tzelepis et al., 2015). A group C chitinase gene was up-regulated in the truffle mycoparasite *T. ophioglossoides* during growth on truffle tissue (Quandt et al., 2016) (Table 1). A putative group C chitinase gene was also induced in the thermophilic fungus *Myceliophthora thermophila* during growth on straw material from a range of plants (Kolbusz et al., 2014).

To summarize, a majority of group C chitinase genes are induced during interspecific interactions with other fungi (Table 1), supporting their possible roles as killer toxin-like facilitators of toxic compounds into the cells of competing individuals. However, induction of certain killer toxin-like chitinase genes upon growth on chitin or during self-interactions indicates that their chitinolytic activity has found additional uses in cell wall remodelling, autolysis and nutrient acquisition (Fig. 2).

## 5. Functional analyses of killer toxin-like chitinases

Structural modelling of the GH18 module of TAC2 from *T. atroviride* indicated that group C chitinases are processive exochitinases, due to the presence of a [SA]XGGW motif indicative of



**Fig. 2 – Proposed function of group C killer toxin-like chitinases during fungal-fungal interactions. A)** Killer toxin-like chitinases are secreted from the producer, bind to cell wall chitin of the antagonist through their CBM18 or CBM50 modules and facilitate the penetration of secondary metabolites or the Hce2 protein into the cytoplasm of the antagonist. The Hce2 protein is presumably released from certain group C chitinases by hitherto unknown proteases. Secreted group C chitinases can also be involved in degradation of exogenous chitin for nutrition purposes. **B)** Killer toxin-like chitinases with transmembrane domains are presumably located in the cell wall of the producer and would require physical contact with the antagonist cell wall for their activity. Additional function in cell wall remodelling, hyphal branching or exogenous chitin degradation can be speculated. Dashed arrows show the dynamic flow of the molecules. Figure was created with BioRender.

processivity and due to a deep and narrow catalytic cleft (Gruber *et al.*, 2011a). Data derived from phenotypic analyses of group C chitinase gene deletion strains are limited. In *N. crassa*, deletion of the *gh18-6* and *gh18-8* genes did not have any impact on colony growth or morphology, conidiation, biomass production or responses to abiotic stress conditions (Tzelepis *et al.*, 2012). Similarly, no differences in colony morphology or growth were observed in *A. nidulans* *chiC2-1*, *chiC2-2*, *chiC2-3* or *chiC2-4* deletion strains (Tzelepis *et al.*, 2014b). Lack of phenotypes is possibly attributed to functional redundancy among the multiple chitinases present in these species. However, deletion of these genes led to increased *A. nidulans* biomass production, while deletion of *chiC2-1*, *chiC2-2* and *chiC2-4* led to more resistant phenotypes in abiotic stress conditions (Tzelepis *et al.*, 2014b). Interestingly, deletion of the *chiC2-2* gene, which carries the Hce2 peptide at the C-terminal, resulted in a mutant with reduced *in vitro* antagonistic ability

(Tzelepis *et al.*, 2014b), suggesting a specific role of Hce2 in fungal-fungal interactions.

Functional analyses of killer toxin-like chitinases have also been conducted in mycoparasitic species such as *C. rosea* and *T. atroviride*. Deletion of the *chiC2* gene in *C. rosea* has an impact on its *in vitro* antagonistic ability (Tzelepis *et al.*, 2015). Specifically, the *chiC2* deletion strain had a reduced inhibitory activity against *B. cinerea* and *R. solani*, but not against *F. graminearum* (Tzelepis *et al.*, 2015). However, the biocontrol ability of *C. rosea* against *B. cinerea* remains unaffected in the  $\Delta$ *chiC2* strain (Tzelepis *et al.*, 2015). Interestingly, this deletion strain also shows a reduced conidiation rate as compared to wild type (Tzelepis *et al.*, 2015). Similarly, deletion of the *tac6* gene in *T. atroviride* is reported to result in altered conidiation patterns (Seidl-Seiboth *et al.*, 2014). However, as mentioned previously, the *tac6* gene carries mutations in its predicted catalytic motif that may result in loss of its chitinolytic ability (Gruber *et al.*, 2011a).

## 6. Concluding remarks

Based on their sequence similarity with the yeast killer toxin zymocin, C-group chitinases have been hypothesized to function in fungal-fungal interactions by permeabilizing antagonist cell walls to facilitate diffusion of toxic compounds (Fig. 2A), either as secreted enzymes or as enzymes bound to the plasma membrane (Fig. 2B). Induction of many killer toxin-like chitinase genes during interspecific fungal interactions and functional studies in *A. nidulans* and *C. rosea* support this idea. However, more detailed studies of the molecular function of these proteins and their interaction with the fungal cell wall is needed before the killer toxin-like chitinase facilitated toxin entry hypothesis can be confirmed. On the other hand, regulation by developmental stage and nutritional stimuli and phenotypic effects on growth, conidiation and stress tolerance in *A. nidulans*, *C. rosea* and *T. atroviride* imply more diverse roles of killer toxin-like chitinases. It is important to investigate whether these examples constitute evidence of specific activities performed by killer toxin-like chitinases, or merely reflects the difficulties of the producers to handle and protect themselves against the action of these, presumably dangerous, enzymes. It is likely that co-localized, co-regulated CBM50 LysM proteins may bind to cell wall chitin of the producer in order to protect against the hydrolytic action of killer toxin-like chitinases. However, this hypothesis needs functional validation. Related to this question is whether killer toxin-like chitinases are proteolytically cleaved after secretion out of the fungal cell, in analogy with the  $\alpha/\beta$  subunit of zymocin. If so, this could release presumably toxic peptides such as Hce2 and EGF-1 (Fig. 2), again in analogy with the zymocin  $\gamma$  subunit. In addition, this proteolytic cleavage may also release additional CBM50 LysM modules from subgroup C-II chitinases that may protect the producer from its own killer toxin-like chitinases. This post-translational release of protective proteins from the chitinase itself would represent the optimal solution for co-regulating CBM50 LysM proteins and killer toxin-chitinases and be an elegant way of minimizing self-inflicted damage. Finally, the rapid evolution of killer toxin-like chitinases, both in gene copy number changes and sequence divergence, may indicate co-evolution with their cell wall targets or with interfering proteins (proteases or inhibitors) produced by the antagonist in an effector arms race manner. Identification of these killer toxin-like chitinase interfering factors, if they exist, would open up new possibilities in agricultural and medical biotechnology. For example, compounds that specifically interfere with the defence against killer toxin-like chitinases in phytopathogenic fungi may be used to augment the efficacy of fungal biological control agents such as *Trichoderma* spp. and *C. rosea*.

## Conflict of interest

Both authors agree with the submission and they declare no conflict of interest.

## Acknowledgements

We acknowledge financial support from the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS, grant nos. 942-2015-368 and 942-2015-1128) and from the Centre for Biological Control (CBC) at the Swedish University of Agricultural Sciences.

## REFERENCES

- Baker, L.G., Specht, C.A., Lodge, J.K., 2009. Chitinases are essential for sexual development but not vegetative growth in *Cryptococcus neoformans*. *Eukaryot. Cell* 8, 1692–1705.
- Bartnicki-Garcia, S., 1968. Cell wall chemistry, morphogenesis, and taxonomy of fungi. *Annu. Rev. Microbiol.* 22, 87–108.
- Boldo, J.T., Junges, A., Do Amaral, K.B., Staats, C.C., Vainstein, M.H., Schrank, A., 2009. Endochitinase CHI2 of the biocontrol fungus *Metarhizium anisopliae* affects its virulence toward the cotton stainer bug *Dysdercus peruvianus*. *Curr. Genet.* 55, 551–560.
- Boller, T., Felix, G., 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu. Rev. Plant Biol.* 60, 379–406.
- Boraston, A.B., Bolam, D.N., Gilbert, H.J., Davies, G.J., 2004. Carbohydrate-binding modules: fine-tuning polysaccharide recognition. *Biochem. J.* 382, 769–781.
- Bowman, S.M., Free, S.J., 2006. The structure and synthesis of the fungal cell wall. *Bioessays* 28, 799–808.
- Buist, G., Steen, A., Kok, J., Kuipees, O.R., 2008. LysM, a widely distributed protein motif for binding to peptidoglycans. *Mol. Microbiol.* 68, 838–847.
- Butler, A.R., Odonnell, R.W., Martin, V.J., Gooday, G.W., Stark, M.J.R., 1991. *Kluyveromyces lactis* toxin has an essential chitinase activity. *Eur. J. Biochem.* 199, 483–488.
- Chaudhary, P.M., Tupe, S.G., Deshpande, M.V., 2013. Chitin synthase inhibitors as antifungal agents. *Mini Rev. Med. Chem.* 13, 222–236.
- de Jonge, R., Thomma, B.P.H.J., 2009. Fungal LysM effectors: extinguishers of host immunity? *Trends Microbiol.* 17, 151–157.
- de Jonge, R., van Esse, H.P., Kombrink, A., et al., 2010. Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science* 329, 953–955.
- de Nobel, H., van den Ende, H., Klis, F.M., 2000. Cell wall maintenance in fungi. *Trends Microbiol.* 8, 344–345.
- Dunkler, A., Walther, A., Specht, C.A., Wendland, J., 2005. *Candida albicans* CHT3 encodes the functional homolog of the Cts1 chitinase of *Saccharomyces cerevisiae*. *Fungal Genet. Biol.* 42, 935–947.
- Gooday, G.W., 1990. Physiology of microbial degradation of chitin and chitosan. *Biodegradation* 1, 177–190.
- Gortari, M.C., Hours, R.A., 2013. Biotechnological processes for chitin recovery out of crustacean waste: A mini review. *Electron. J. Biotechnol.* 16.
- Gruber, S., Vaaje-Kolstad, G., Matarese, F., Lopez-Mondejar, R., Kubicek, C.P., Seidl-Seiboth, V., 2011a. Analysis of subgroup C of fungal chitinases containing chitin-binding and LysM modules in the mycoparasite *Trichoderma atroviride*. *Glycobiology* 21, 122–133.
- Gruber, S., Kubicek, C.P., Seidl-Seiboth, V., 2011b. Differential regulation of orthologous chitinase genes in mycoparasitic *Trichoderma* species. *Appl. Environ. Microbiol.* 77, 7217–7226.
- Gruber, S., Seidl-Seiboth, V., 2012. Self versus non-self: fungal cell wall degradation in *Trichoderma*. *Microbiology-Sgm* 158, 26–34.

- Henrissat, B., 1991. A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem. J.* 280, 309–316.
- Hill, D.E., Fetterer, R.H., Urban, J.F., 1991. *Ascaris*-Suum: stage-specific differences in lectin binding to the larval cuticle. *Exp. Parasitol.* 73, 376–383.
- Horn, S.J., Sorbotten, A., Synstad, B., Sikorski, P., Sorlie, M., Varum, K.M., Eijsink, V.G.H., 2006. Endo/exo mechanism and processivity of family 18 chitinases produced by *Serratia marcescens*. *FEBS J.* 273, 491–503.
- Ihrmark, K., Asmail, N., Ubhayasekera, W., Melin, P., Stenlid, J., Karlsson, M., 2010. Comparative molecular evolution of *Trichoderma* chitinases in response to mycoparasitic interactions. *Evol. Bioinform.* 6, 1–25.
- Kapaun, E., Reisser, W., 1995. A chitin-like glycan in the cell wall of a *Chlorella* sp (Chlorococcales, Chlorophyceae). *Planta* 197, 577–582.
- Karlsson, M., Atanasova, L., Funck Jensen, D., Zeilinger, S., 2018. Necrotrophic mycoparasites and their genomes. In: Heitman, J., Howlett, B., Crous, P., Stukenbrock, E., James, T., Gow, N. (Eds.), *The Fungal Kingdom*. ASM Press, Washington DC, pp. 1005–1026.
- Karlsson, M., Stenlid, J., Lindahl, B., 2016. Functional differentiation of chitinases in the white-rot fungus *Phanerochaete chrysosporium*. *Fungal Ecol.* 22, 52–60.
- Karlsson, M., Durling, M.B., Choi, et al., 2015. Insights on the evolution of mycoparasitism from the genome of *Clonostachys rosea*. *Genome Biol. Evol.* 7, 465–480.
- Karlsson, M., Stenlid, J., 2009. Evolution of family 18 glycoside hydrolases: Diversity, domain-structures and phylogenetic relationships. *J. Mol. Microbiol. Biotechnol.* 16, 208–223.
- Karlsson, M., Stenlid, J., 2008. Comparative evolutionary histories of the fungal chitinase gene family reveal non-random size expansions and contractions due to adaptive natural selection. *Evol. Bioinform.* 4, 47–60.
- Keegstra, K., 2010. Plant cell walls. *Plant Physiol.* 154, 483–486.
- Keyhani, N.O., Roseman, S., 1999. Physiological aspects of chitin catabolism in marine bacteria. *BBA Gen. Subj.* 1473, 108–122.
- Klis, F.M., Mol, P., Hellingwerf, K., Brul, S., 2002. Dynamics of cell wall structure in *Saccharomyces cerevisiae*. *FEMS Microbiol. Rev.* 26, 239–256.
- Kolbusz, M.A., Di Falco, M., Ishmael, N., Marqueteau, S., Moisan, M.C., Baptista, C.D.S., Powlowski, J., Tsang, A., 2014. Transcriptome and exoproteome analysis of utilization of plant-derived biomass by *Myceliophthora thermophila*. *Fungal Genet. Biol.* 72, 10–20.
- Kubicek, C.P., Herrera-Estrella, A., Seidl-Seiboth, V., et al., 2011. Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of *Trichoderma*. *Genome Biol.* 12, R40.
- Kuranda, M.J., Robbins, P.W., 1991. Chitinase is required for cell separation during growth of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 266, 19758–19767.
- Kurita, K., 2006. Chitin and chitosan: Functional biopolymers from marine crustaceans. *Mar. Biotechnol.* 8, 203–226.
- Latge, J.P., 2007. The cell wall: a carbohydrate armor for the fungal cell. *Mol. Microbiol.* 66, 279–290.
- Lauge, R., Joosten, M.H.A.J., Vandenackerveken, G.F.J.M., Vandenbroek, H.W.J., de Wit, P.J.G.M., 1997. The *in planta*-produced extracellular proteins ECP1 and ECP2 of *Cladosporium fulvum* are virulence factors. *Mol. Plant Microbe Interact.* 10, 725–734.
- Lombard, V., Ramulu, H.G., Drula, E., Coutinho, P.M., Henrissat, B., 2014. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* 42, D490–D495.
- Magliani, W., Conti, S., Gerloni, M., Bertolotti, D., Polonelli, L., 1997. Yeast killer systems. *Clin. Microbiol. Rev.* 10, 369–400.
- Mamarabadi, M., Jensen, B., Lubeck, M., 2008. Three endochitinase-encoding genes identified in the biocontrol fungus *Clonostachys rosea* are differentially expressed. *Curr. Genet.* 54, 57–70.
- Merzendorfer, H., Zimoch, L., 2003. Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J. Exp. Biol.* 206, 4393–4412.
- Mulisch, M., 1993. Chitin in protistan organisms; distribution, synthesis and deposition. *Eur. J. Protistol.* 29, 1–18.
- Newman, M.A., Sundelin, T., Nielsen, J.T., Erbs, G., 2013. MAMP (microbe-associated molecular pattern) triggered immunity in plants. *Front. Plant Sci.* 4, 139.
- Nygren, K., Dubey, M., Zapparata, A., Iqbal, M., Tzelepis, G.D., Durling, M.B., Jensen Funck, D., Karlsson, M., 2018. The mycoparasitic fungus *Clonostachys rosea* responds with both common and specific gene expression during interspecific interactions with fungal prey. *Evol. Appl.* 11, 931–949.
- Peters, W., 1972. Occurrence of chitin in mollusca. *Comp. Biochem. Physiol.* 41, 541–544.
- Quandt, C.A., Di, Y.M., Elser, J., Jaiswal, P., Spatafora, J.W., 2016. Differential expression of genes involved in host recognition, attachment, and degradation in the mycoparasite *Tolyposcladium ophioglossoides*. *G3* 6, 731–741.
- Seidl, V., Huemer, B., Seiboth, B., Kubicek, C.P., 2005. A complete survey of *Trichoderma* chitinases reveals three distinct subgroups of family 18 chitinases. *FEBS J.* 272, 5923–5939.
- Seidl-Seiboth, V., Ihrmark, K., Druzhinina, I., Karlsson, M., 2014. Molecular evolution of *Trichoderma* chitinases. In: Gupta, V., Schmoll, M., Herrera-Estrella, A., Upadhyay, R.S., Druzhinina, I., Tuohy, M. (Eds.), *Biotechnology and Biology of Trichoderma*, vol. 5. Elsevier., pp. 67–78.
- Seidl-Seiboth, V., Zach, S., Frischmann, A., Spadiut, O., Dietzsch, C., Herwig, C., Ruth, C., Rodler, A., Jungbauer, A., Kubicek, C.P., 2013. Spore germination of *Trichoderma atroviride* is inhibited by its LysM protein TAL6. *FEBS J.* 280, 1226–1236.
- Shin, K.S., Kwon, N.J., Kim, Y.H., Park, H.S., Kwon, G.S., Yu, J.H., 2009. Differential roles of the *ChiB* chitinase in autolysis and cell death of *Aspergillus nidulans*. *Eukaryot. Cell* 8, 738–746.
- Specht, C.A., Liu, Y.L., Robbins, P.W., Bulawa, C.E., Iartchouk, N., et al., 1996. The *chsD* and *chsE* genes of *Aspergillus nidulans* and their roles in chitin synthesis. *Fungal Genet. Biol.* 20, 153–167.
- Spindler, K.D., Spindlerbarth, M., Londershausen, M., 1990. Chitin metabolism; a target for drugs against parasites. *Parasitol. Rev.* 76, 283–288.
- Stals, I., Samyn, B., Sergeant, K., White, T., Hoorelbeke, K., Coorevits, A., Devreese, B., Claeysens, M., Piens, K., 2010. Identification of a gene coding for a deglycosylating enzyme in *Hypocrea jecorina*. *FEMS Microbiol. Lett.* 303, 9–17.
- Stark, M.J.R., Boyd, A., 1986. The killer toxin of *Kluyveromyces lactis*; characterization of the toxin subunits and identification of the genes which encode them. *EMBO J.* 5, 1995–2002.
- Stark, M.J.R., Boyd, A., Mileham, A.J., Romanos, M.A., 1990. The plasmid-encoded killer system of *Kluyveromyces lactis*: a review. *Yeast* 6, 1–29.
- Stergiopoulos, I., Kourmpetis, Y.A., Slot, J.C., Bakker, F.T., de Wit, P.J.G.M., Rokas, A., 2012. In silico characterization and molecular evolutionary analysis of a novel superfamily of fungal effector proteins. *Mol. Biol. Evol.* 29, 3371–3384.
- Stergiopoulos, I., Van den Burg, H.A., Okmen, B., Beenen, H.G., Van Liere, S., Kema, G.H.J., de Wit, P.J.G.M., 2010. Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. *Proc. Natl. Acad. Sci. U.S.A.* 107, 7610–7615.
- Tzelepis, G., Karlsson, M., Suzuki, T., 2017. Deglycosylating enzymes acting on N-glycans in fungi: Insights from a genome survey. *BBA Gen. Subj.* 1861, 2551–2558.
- Tzelepis, G., Dubey, M., Jensen Funck, D., Karlsson, M., 2015. Identifying glycoside hydrolase family 18 genes in the mycoparasitic fungal species *Clonostachys rosea*. *Microbiology-Sgm* 161, 1407–1419.

- Tzelepis, G., Hosomi, A., Tanim, J.H., Hirayama, H., Dubey, M., Jensen Funck, D., Suzuki, T., Karlsson, M., 2014a. Endo- $\beta$ -N-acetylglucosamidases (ENGases) in the fungus *Trichoderma atroviride*: Possible involvement of the filamentous fungi-specific cytosolic ENGase in the ERAD process. *Biochem. Biophys. Res. Commun.* 449, 256–261.
- Tzelepis, G., Melin, P., Stenlid, J., Jensen Funck, D., Karlsson, M., 2014b. Functional analysis of the C-II subgroup killer toxin-like chitinases in the filamentous ascomycete *Aspergillus nidulans*. *Fungal Genet. Biol.* 64, 58–66.
- Tzelepis, G., Melin, P., Jensen Funck, D., Stenlid, J., Karlsson, M., 2012. Functional analysis of glycoside hydrolase family 18 and 20 genes in *Neurospora crassa*. *Fungal Genet. Biol.* 49, 717–730.
- van Aalten, D.M.F., Komander, D., Synstad, B., Gaseidnes, S., Peter, M.G., Eijsink, V.G.H., 2001. Structural insights into the catalytic mechanism of a family 18 exo-chitinase. *Proc. Natl. Acad. Sci. U.S.A.* 98, 8979–8984.
- Wouters, M.A., Rigoutsos, I., Chu, C.K., Feng, L.L., Sparrow, D.B., Dunwoodie, S.L., 2005. Evolution of distinct EGF domains with specific functions. *Protein Sci.* 14, 1091–1103.
- Zhang, X.C., Cannon, S.B., Stacey, G., 2009. Evolutionary genomics of LysM genes in land plants. *BMC Evol. Biol.* 9, 183.