



## Letter to the editor

**Responses to “Comments on Botta et al. (2018). Potentially active spoilage bacteria community during the storage of vacuum packaged beefsteaks treated with aqueous ozone and electrolysed water. International Journal of Food Microbiology, 266, 337–345”**

Dear Editor,

Herein our response to criticisms posed by the Authors to our scientific paper [Botta et al. \(2018\)](#): Potentially active spoilage bacteria community during the storage of vacuum packaged beefsteaks treated with aqueous ozone and electrolysed water. *International Journal of Food Microbiology*, 266, 337–345.

Despite the first author is included within the inventors of the European Patent EP 2207415 B1, we want (the company owning the patent and us) to clarify that the current electrolyzed water (EW) generator named EVA system® has been improved from the first prototype patented, especially it has being modified in several efficiency aspects. For instance, it is known that important parameters for efficacy of the produced EW, such as oxidation reduction potential (ORP), free chlorine (AC) and pH can be significantly changed by the electrical current value, flow rate, electrode material, electrolysis time and amperage of EW generator ([Rahman et al., 2016](#)). Therefore, we confirm that EVA System® provided by De Nora S.P.A. produces a mother solution of 4000 ppm of AC at pH 9.00 and a KCl residual of 1%, which was obtained electrolyzing a brine of 15 g/L. More information on the EVA System® and its technical sheet can be easily and freely found at the company website for a further confirmation of this set up and machine features.

Besides this, we strongly disagree about the Author's criticisms over the terminology. To best of our knowledge the term “electrolyzed water” includes basic electrolyzed water (BEW), acid electrolyzed water (AEW), neutral electrolyzed water (NEW) and slightly acid electrolyzed water (SAEW). Most of the EW machines can be divided into 2 types: machines (i) with 2-cell chambers separated by diaphragms that produce separately from NaCl brine the AEW (HOCl, -OCl, HCl, O<sub>2</sub>, Cl<sub>2</sub>) and BEW (NaOH, H<sub>2</sub>), respectively from anode and cathode compartments; and machines (ii) without diaphragm that produce in a single chamber NEW and SAEW, respectively from NaCl or HCl (alone or in combination with NaCl) ([Rahman et al., 2016](#)). In this frame the “Electrochemically activated solutions” (ECAS), also called “oxidizing waters” or “anolytes” by the Authors ([Ferro et al., 2018](#)) should refer to the AEW, which (by the way) has shown limited efficacy on utensils, food products, and surfaces owing to various factors, the most important of which include the type of surface, presence of organic matter, and type of tape water used ([Rahman et al., 2016](#)). In any case, it is not our intention to debate here the pros and cons shown in scientific literature by the different type of EW.

As far as the final free chlorine of the “net solution” is concerned, the AC has been measured by iodometric titration ([APHA, 1992](#)) before and after the dilution with distilled water ([Laureano et al., 2016](#); [Yaseen et al., 2016](#)). In relation to these measurements the “net solution” were freshly prepared for each experiment at 100 ppm of free chlorine (pH ~ 7.0; thus a NEW). We are aware of the pH effect on free chlorine concentration, the decrease of chlorine due the interaction with organic matter and along the time (investigation on the chlorine decomposition in different storage condition is currently ongoing for this NEW), as well as the differential effectiveness existing between hypochlorous acid and hypochlorite ([Athayde et al., 2017](#); [Rahman et al., 2016](#)).

Anyway, we do not want go into the merits of pH variations and dilutions hypothesized by the authors, because are based on their personal experience and not supported by data. Moreover, we believe we have provided all information necessary to repeat the preparation of electrolyzed water in the Material and Methods section, which has already been peer reviewed, as all the article.

Regarding the choice of the salt, we simply preferred to avoid any unwanted biostatic/biocide activity by any residual NaCl during the subsequent storage ([Sharan et al., 2010](#)). Moreover the potential residual NaCl has been proven more corrosive than KCl on some type of steels (grid containing the beefsteaks during treatments) ([Enestam et al., 2013](#)).

As rightly stated by the Authors, in three of the studies cited ([Ding et al., 2010](#); [Jadeja and Hung, 2014](#); [Tango et al., 2014](#)) different types of EW or their combination have been effective in reducing bacterial count of meat surfaces, contrariwise to our outcomes, even if only [Jadeja and Hung \(2014\)](#) have tested a neutral EW. On the other hand we cannot compare results from [Bosilevac et al. \(2005\)](#) to ours, since they sprayed beef hides with BEW and AEW at 52 and 60 °C respectively, and high temperature can significantly improve the sanitizing efficacy of EW ([Cao et al., 2009](#); [Fabrizio and Cutter, 2003](#)). Anyway, in literature we can also find studies that highlight a limited or absent reduction of microbial loads on beef carcass sprayed with EW (from 60 to 250 ppm of free chlorine) ([Kalchayanand et al., 2008](#); [Veasey and Muriana, 2016](#)).

As far as the ozone is concerned, it is surprising the Authors' attempt to demonstrate the efficacy of aqueous ozone (AO) in the sanitization/decontamination of meat by citing [Rice et al. \(2002\)](#) (a magazine article dealing on the use of ozone in food industry as surface sanitizer and its legislation in USA) and another magazine article dealing the use of AO

<https://doi.org/10.1016/j.ijfoodmicro.2018.10.017>

and ozone in winery environments (Hampson et al., 2000). It is also noteworthy that in the report proposed (Hallam and Hiscock, 2006), which collects references on this topic and does not show original data, some of the cited studies confirm the efficacy of AO in decontamination of meat/beef carcass (Bosilevac et al., 2005; Gorman et al., 1995) and others a limited or absent activity (Castillo et al., 2003; Reagan et al., 1996).

Regardless from the number of references demonstrating or not the efficacy of AO and EW, in the Table 1 of our article (Botta et al., 2018) at day 0 the value refer to the beefsteak before the treatments, contrariwise of what argue by the Authors and in agreement to what is reported in our Material and Methods section: “The samplings were performed before the treatments, for each treatment and each batch, on the first day and after 5, 9 and 15 days of storage at 4 °C”. Unlike most of the studied cited above, in our study we wanted to check the efficacy of the two treatments along the shelf life of the product on the market, which thus starts from the first day after the packaging of the meat (day 1). The authors focused their attention on some “not reasonable” differences between EW/AO treated and untreated meat, but neither significantly lower nor significantly higher values can be observed between them looking at the microbial and pH dynamics shown in Table 1.

Concerning the experimental set up, both treatments timing and concentrations have been chosen to avoid the risk of discoloration, which is reported also by the same Authors as an effective side effect, especially for AO. The reasons of our choices also reported in the Discussion part: “However, the limiting factors for applying AO and EW to raw beef remain undoubtedly their concentrations and exposure times, which were here chosen on the basis of the acceptability of the colour of the meat after treatments and considering the effectiveness of treatments performed by spraying AO and EW on different meats and seafood products.”

Finally, we firmly disagree with the biases expressed by the Authors: “It is obviously difficult to imagine what the reason for the failure was but, certainly, there were some inaccuracies as to how the experiments were executed.”

As all experimental methods applied in small-scale processing line, this approach can be improved and implemented, but however the experiments performed in Botta et al. (2018) were executed with the maximum scientific rigor. It is strange, from our point of view, that Authors consider a limited/non-significant effects of the two experimental sanitizing treatments as an unacceptable “failure”, which was “certainly” caused by inaccuracies that however are “difficult to imagine”. The failure is the state or condition of not meeting a desirable or intended objective, in opposite of success. Would a significant effect of the treatments has been the success? In our opinion, this is not an independent scientific approach. The objective of our work, considering all its possible methodological limitations, was not to confirm at any cost the efficacy of these two specific treatments with EW and AO, but to define their action on meat microbiota dynamics by means of culture-dependent and -independent approaches along the shelf-life. The overall evaluation of EW or AO as meat sanitizer need undoubtedly further investigation, and the final judgment on the suitability of these treatments is absolutely beyond the purpose of this work. In light of this we cannot consider our outcomes detrimental for the body of knowledge of this subject.

Despite the Author consider our results not relevant in relation to the initial “failure” of the treatments, we believed that the RNA-based analysis integrated with the volatilome has helped to unravel the complexity of the potentially active microbiota, in this way expanding

the current knowledge on the spoilage dynamics of vacuum packaged beefsteaks (as reported in the conclusions). This because the spoilage dynamics of raw meat have mainly been studied with DNA-based HTS approaches so far, which cannot be associated to the volatilome only produced by active microbiota.

## References

- APHA, 1992. APHA Method 4500-Cl: Standard Methods for the Examination of Water and Wastewater.
- Athayde, D.R., Flores, D.R.M., da Silva, J.S., Genro, A.L.G., Silva, M.S., Klein, B., Mello, R., Campagnol, P.C.B., Wagner, R., de Menezes, C.R., Barin, J.S., Cichoski, A.J., 2017. Application of electrolyzed water for improving pork meat quality. *Food Res. Int.* 100, 757–763. <https://doi.org/10.1016/j.foodres.2017.08.009>.
- Bosilevac, J.M., Shackelford, S.D., Brichta, D.M., Koohmaraie, M., 2005. Efficacy of ozonated and electrolyzed oxidative waters to decontaminate hides of cattle before slaughter. *J. Food Prot.* 68, 1393–1398. <https://doi.org/10.4315/0362-028X-68.7.1393>.
- Botta, C., Ferruccio, I., Cavallero, M.C., Riva, S., Giordano, M., Cocolin, L., 2018. Potentially active spoilage bacteria community during the storage of vacuum packaged beefsteaks treated with aqueous ozone and electrolyzed water. *Int. J. Food Microbiol.* 0–1. <https://doi.org/10.1016/j.ijfoodmicro.2017.10.012>.
- Cao, W., Zhu, Z.W., Shi, Z.X., Wang, C.Y., Li, B.M., 2009. Efficiency of slightly acidic electrolyzed water for inactivation of *Salmonella enteritidis* and its contaminated shell eggs. *Int. J. Food Microbiol.* 130, 88–93. <https://doi.org/10.1016/j.ijfoodmicro.2008.12.021>.
- Castillo, A., McKenzie, K.S., Lucia, L.M., Acuffi, G.R., 2003. Ozone treatment for reduction of *Escherichia coli* O157:H7 and *Salmonella serotype typhimurium* on beef carcass surfaces. *J. Food Prot.* 66, 775–779. <https://doi.org/10.4315/0362-028X-66.5.775>.
- Ding, T., Rahman, S.M.E., Purev, U., Oh, D.H., 2010. Modelling of *Escherichia coli* O157:H7 growth at various storage temperatures on beef treated with electrolyzed oxidizing water. *J. Food Eng.* 97, 497–503. <https://doi.org/10.1016/j.jfoodeng.2009.11.007>.
- Enestam, S., Bankiewicz, D., Tuiremo, J., Mäkelä, K., Hupa, M., 2013. Are NaCl and KCl equally corrosive on superheater materials of steam boilers? *Fuel* 104, 294–306. <https://doi.org/10.1016/j.fuel.2012.07.020>.
- Fabrizio, K.A., Cutter, C.N., 2003. Stability of electrolyzed oxidizing water and its efficacy against cell suspensions of *Salmonella typhimurium* and *Listeria monocytogenes*. *J. Food Prot.* 66, 1379–1384. <https://doi.org/10.4315/0362-028X-66.8.1379>.
- Ferro, S., Amorico, T., Deo, P., 2018. Role of food sanitising treatments in inducing the “viable but nonculturable” state of microorganisms. *Food Control.* <https://doi.org/10.1016/j.foodcont.2018.04.016>.
- Gorman, B.M., Sofos, J.N., Morgan, J.B., Schmidt, G.R., Smith, G.C., 1995. Evaluation of hand-trimming, various sanitizing agents, and hot water spray-washing as decontamination interventions for beef brisket adipose tissue. *J. Food Prot.* 58, 899–907. <https://doi.org/10.4315/0362-028X-58.8.899>.
- Hallam, V., Hiscock, D., 2006. *Food Science Australia.* <http://www.meatupdate.csiro.au/new/Ozonated%20Wate>.
- Hampson, B.B., Science, F., Obispo, S.L., 2000. Use of Ozone for Winery and Environmental Sanitation.
- Jadeja, R., Hung, Y.C., 2014. Efficacy of near neutral and alkaline pH electrolyzed oxidizing waters to control *Escherichia coli* O157:H7 and *Salmonella typhimurium* DT 104 from beef hides. *Food Control* 41, 17–20. <https://doi.org/10.1016/j.foodcont.2013.12.030>.
- Kalchayanand, N., Arthur, T.M., Bosilevac, J.M., Brichta-Harhay, D.M., Guerini, M.N., Wheeler, T.L., Koohmaraie, M., 2008. Evaluation of various antimicrobial interventions for the reduction of *Escherichia coli* O157:H7 on bovine heads during processing. *J. Food Prot.* 71, 621–624.
- Laureano, J., Giacosa, S., Rio Segade, S., Torchio, F., Craverio, F., Gerbi, V., Englezos, V., Carboni, C., Cocolin, L., Rantsiou, K., Faroni, L.R.D., Rolle, L., 2016. Effects of continuous exposure to ozone gas and electrolyzed water on the skin hardness of table and wine grape varieties. *J. Texture Stud.* 47, 40–48. <https://doi.org/10.1111/jtxs.12158>.
- Rahman, S.M.E., Khan, I., Oh, D., 2016. Electrolyzed Water as a Novel Sanitizer in the Food Industry: Current Trends and Future Perspectives. 15. pp. 471–490. <https://doi.org/10.1111/1541-4337.12200>.
- Reagan, J.O., Acuff, G.R., Buege, D.R., Buyck, M.J., Dickson, J.S., Katster, C.L., Marsden, J.L., Morgan, J.B., Nickelson, R., Smith, G.C., Sofos, J.N., 1996. Trimming and washing of beef carcasses as a method of improving the microbiological quality of meat. *J. Food Prot.* 59, 751–756. <https://doi.org/10.4315/0362-028X-59.7.751>.
- Rice, R.G., Graham, D., Lowe, M.T., 2002. Recent ozone applications in food processing and sanitation. *Food Saf. Mag.* October/November. <https://www.foodsafetymagazine.com/magazine-archives/october-november-2002/recent-ozone-applications-in-food-processing-and-sanitation/>.
- Sharan, R., Chhibber, S., Attri, S., Reed, R.H., 2010. Inactivation and sub-lethal injury of

- Escherichia coli* in a copper water storage vessel: effect of inorganic and organic constituents. *Antonie van Leeuwenhoek. Int. J. Gen. Mol. Microbiol.* 98, 103–115. <https://doi.org/10.1007/s10482-010-9435-3>.
- Tango, C.N., Mansur, A.R., Kim, G.H., Oh, D.H., 2014. Synergetic effect of combined fumaric acid and slightly acidic electrolysed water on the inactivation of food-borne pathogens and extending the shelf life of fresh beef. *J. Appl. Microbiol.* 117, 1709–1720. <https://doi.org/10.1111/jam.12658>.
- Veasey, S., Muriana, P., 2016. Evaluation of electrolytically-generated hypochlorous acid (“electrolyzed water”) for sanitation of meat and meat-contact surfaces. *Foods* 5, 42. (1–15). <https://doi.org/10.3390/foods5020042>.
- Yaseen, T., Damara, E., Albanese, P., Ricelli, A., Carboni, C., Ferri, V., D’Onghia, A., 2016. Effect of electrolyzed water and continuous ozone exposure on some aspects of clementine quality during conservation. In: Ippolito, A., Sanzani, S., Wisniewski, M., Droby, S. (Eds.), III International Symposium on Postharvest Pathology: Using Science to Increase Food, <https://doi.org/10.17660/ActaHortic.2016.1141.59>.

Cristian Botta, Ilario Ferrocino, Maria Chiara Cavallero,  
Simonetta Riva, Manuela Giordano, Luca Cocolin\*

Department of Agriculture, Forestry and Food Sciences, University of Torino,  
Italy

E-mail address: [lucasimone.cocolin@unito.it](mailto:lucasimone.cocolin@unito.it) (L. Cocolin).

---

\* Corresponding author.