



Access to a high resource environment protects against accelerated maturation following early life stress: A translational animal model of high, medium and low security settings

Arielle R. Strzelewicz^a, Evelyn Ordoñez Sanchez^b, Alejandro N. Rondón-Ortiz^a, Anthony Raneri^c, Sydney T. Famularo^b, Debra A. Bangasser^b, Amanda C. Kentner^{c,*}

^a School of Pharmacy, Massachusetts College of Pharmacy and Health Sciences, Boston, MA 02115, United States

^b Department of Psychology, Temple University, Philadelphia, PA 19122, United States

^c School of Arts & Sciences, Health Psychology Program, Massachusetts College of Pharmacy and Health Sciences, Boston, MA 02115, United States

ARTICLE INFO

Keywords:

Early life stress
Limited bedding
Western diet
Animal models
Social economic status
Environmental enrichment
Precocious puberty
Kisspeptin
Maternal care

ABSTRACT

Early life exposure to a low security setting, characterized by a scarcity of resources and limited food access, increases the risk for psychiatric illness and metabolic dysfunction. We utilized a translational rat model to mimic a low security environment and determined how this manipulation affected offspring behavior, metabolism, and puberty. Because food insecurity in humans is associated with reduced access to healthy food options the “low security” rat manipulation combined a Western diet with exposure to a limited bedding and nesting manipulation (WD-LB). In this setting, dams were provided with limited nesting materials during the pups' early life (P2-P10). This manipulation was contrasted with standard rodent caging (SD) and environmental enrichment (EE), to model “medium security” and “high security” environments, respectively. To determine if transitioning from a low to high security environment improved outcomes, some juvenile WD-LB offspring were exposed to EE. Maternal care was impacted by these environments such that EE dams engaged in high quality care when on the nest, but spent less time on the nest than SD dams. Although WD-LB dams excessively chased their tails, they were very attentive to their pups, perhaps to compensate for limited resources. Offspring exposed to WD-LB only displayed subtle changes in behavior. However, WD-LB exposure resulted in significant metabolic dysfunction characterized by increased body weight, precocious puberty and alterations in the hypothalamic kisspeptin system. These negative effects of WD-LB on puberty and weight regulation were mitigated by EE exposure. Collectively, these studies suggest that both compensatory maternal care and juvenile enrichment can reduce the impact of a low security environment. Moreover, they highlight how utilizing diverse models of resource (in)stability can reveal mechanisms that confer vulnerability and resilience to early life stress.

1. Introduction

Both basic and clinical research suggest that early-life stressors (e.g. abuse, neglect) can modify brain development and make an individual prone to mental illness and metabolic dysfunction in later life (Spencer et al., 2017; Walker et al., 2017; Yam et al., 2016; Pervanidou and Chrousos, 2012; Avishai-Eliner et al., 2001). For example, exposure to early-life stress increases the risk for mental illnesses, such as mood and depressive disorders, anxiety disorders, and disruptive behavior disorders (Afari et al., 2014; Green et al., 2010; Heim et al., 2008; Chapman et al., 2004; Anda et al., 2002). Moreover, early-life adversity has a positive association with metabolic dysfunctions including metabolic syndrome and the occurrence of precocious puberty (Cowan and

Richardson, 2018; Pervanidou and Chrousos, 2012; Björntorp, 2008). To fully understand the relationship between early-life stress and these later life outcomes, it is necessary to develop ecologically relevant animal models in order to derive their mechanistic underpinnings. This step will be imperative for the design and testing of translational treatments and preventative methods.

The hypothalamic pituitary adrenal (HPA) axis is a fundamental feature of the stress response (Jacobson and Sapolsky, 1991). Normally each component of this axis develops simultaneously, connecting into a functional network. However severely stressful events during the perinatal period can cause them to develop out of sync, resulting in life-long modifications in how the organism responds to stressful experiences (Heim et al., 2008; Levine, 1994). Even a short period of stressor

* Corresponding author.

E-mail address: amanda.kentner@mcphs.edu (A.C. Kentner).

<https://doi.org/10.1016/j.yhbeh.2019.01.003>

Received 8 September 2018; Received in revised form 18 December 2018; Accepted 10 January 2019

Available online 02 March 2019

0018-506X/© 2019 Elsevier Inc. All rights reserved.

exposure during the first week of life in rodents can cause behavioral deviations and a vulnerability to physiological disruptions that parallel the clinical symptoms associated with early-life adversity (Walker et al., 2017; Champagne and Curley, 2007). Moreover, the hypothalamic-pituitary-gonadal (HPG) axis is also susceptible to early life programming, connecting the downstream effects of early adverse experiences on reproductive functioning and stress responsivity to a larger interactive network (Kentner and Pittman, 2010).

Puberty is the process of hormones orchestrating physiological changes that turns an organism from its sexually immature to its sexually mature state. During early development, kisspeptin in particular plays a role in the secretion of sex steroids (e.g. luteinizing hormone and follicle stimulating hormone) and the onset of puberty (Kauffman et al., 2007; Skorupskaite et al., 2014). Precocious puberty is when an organism experiences puberty significantly earlier than its species predicted time. In humans, the occurrence of precocious puberty is indicated by significant pubertal markers such as the development of secondary sex characteristics (e.g. pubic hair and breast development) before age 8 or menarche before age 9 (Kaplowitz and Hoffman, 2018; Barker and Kappy, 2011; Cesario & Hughes, 2007). Causes of precocious puberty include infection or trauma directed to the parts of the brain that control reproduction (Cowan and Richardson, 2018; Kaplowitz and Hoffman, 2018), in addition to early life stress (Li et al., 2014; Kelly et al., 2017; Virdis et al., 1998). Precocious puberty is associated with short adult stature, emotional distress (i.e. depression), and other central nervous system abnormalities (Kaplowitz and Hoffman, 2018; Cesario & Hughes, 2007; Chalumeau et al., 2002; Angold and Worthman, 1993). Both males and females can experience precocious puberty but it affects girls 10 times more frequently (Cesario & Hughes, 2007). Thus, precocious puberty could be considered a sex-dependent characteristic of humans who undergo early-life stress. Precocious puberty due to early life stress has also been reported in animal models of maternal separation and litter isolation (Cowan and Richardson, 2018; Kentner et al., 2018; Grassi-Oliveira et al., 2016). Notably, sex-dependent precocious puberty is preventable by probiotic treatment (Cowan and Richardson, 2018) and sensory enrichment can also delay reproductive accelerated maturation precipitated by early-life adversity (Kentner et al., 2018).

Parental care has a critical influence on offspring development and negative experiences mediated through the parent-offspring relationship can significantly impair developmental outcomes. In the rodent laboratory, the dynamics of these relationships can be manipulated experimentally by changing the circumstances of the nest environment (Perry et al., 2018; Walker et al., 2017; McLaughlin et al., 2016; Connors et al., 2015; Kenny et al., 2014; Champagne and Curley, 2007; Levine, 2001; Gilles et al., 1996). One commonly used model of early-life adversity is the limited bedding model (LB) which involves reducing the availability of bedding materials that a dam uses to construct a nest for herself and her pups (Perry et al., 2018; Walker et al., 2017; Rice et al., 2008; Gilles et al., 1996). The lack of materials is a maternal stressor as it decreases the dam's ability to construct a satisfactory nest (Perry et al., 2018; Walker et al., 2017) which influences the way that she interacts with her offspring (Rice et al., 2008; Champagne and Curley, 2007). Dams housed with limited bedding have lower nest construction quality scores, have poorer nursing habits, and have been reported to step-on and rough handle their pups (Perry et al., 2018; Heun-Johnson and Levitt, 2016; Sullivan and Holman, 2010; Ivy et al., 2008). These maternal behavioral patterns show that the LB paradigm can cause care fragmentation and maltreatment (Perry et al., 2018; Walker et al., 2017; Gilles et al., 1996). There is evidence to support that this adversity causes long-term negative consequences on cognitive development, motor skills, and socialization of offspring (Gilles et al., 1996).

As it stands, the LB protocol can be translated to the experience of a low resource or 'insecure' environment for humans. As established earlier, poor access to resources impacts how a dam interacts with her

pups and this is also true of human parents. Socioeconomic status can affect parental care quality. For example, high economic pressure has been positively correlated to the use of punitive or authoritarian parenting (Leinonen et al., 2003). These parenting strategies are associated with conduct disorders and disruptive behavior disorders in children (Stormshak et al., 2010). Additionally, lower socioeconomic status is related to low birth weight, asthma, deficient language development, and fewer pre-academic skills (Burchinal et al., 2000; Aber et al., 1997). The rodent LB protocol is most useful for modeling an impoverished environment because, in addition to the stress from lack of physical resources, it can cause behavioral modifications in the mother known to affect a child's development and adult outcomes.

On the other hand, environmental enrichment (EE) in the animal laboratory can translate to a high resource/high security situation and has been shown to rescue the effects of insecure "stressful" environments (Francis et al., 2002; Bredy et al., 2003; Bredy et al., 2004). Indeed, previous studies have shown that environmental complexity in early life can mitigate the effects of a number of early-life stressors including inflammation, poor maternal care, and neonatal brain injury (Kentner et al., 2016; Schneider et al., 2006; Bredy et al., 2004; Bredy et al., 2003; Pedrini Schuch et al., 2016; Durán-Carabali et al., 2018), highlighting its potential to mitigate the effects of LB, which to our knowledge has not been directly explored previously. Quantifying/mapping the therapeutic benefits of this housing condition has translational value in that enrichment protocols have been successfully utilized in clinical settings (Janssen et al., 2014; White et al., 2015; Woo and Leon, 2013; Woo et al., 2015), underscoring its acceptability and feasibility for use with patients.

Importantly, EE has also been shown to effect the quality of maternal care (Connors et al., 2015; Welberg et al., 2006; Sale et al., 2004; Durán-Carabali et al., 2018), but these effects have been underexplored. While there is some research focusing on defined periods of enrichment exposure (e.g. either pre- or postnatally; Cancedda et al., 2004; Rosenfeld and Weller, 2012), the number of studies evaluating dams in lifelong (or a combined pre- and postnatal) EE exposure are limited (Connors et al., 2015; Welberg et al., 2006; MacRae et al., 2015; Durán-Carabali et al., 2018). Moreover, some using this protocol have utilized co-parenting in the homecage, making the individual contribution of rodent dams difficult to assess. For this reason, we were interested in evaluating maternal care quality across the continuum of resource rich and poor animal laboratory conditions. Given the strength of maternal care quality in shaping offspring outcomes, it is important to understand differences in parent-offspring interactions as a function of environmental complexity. Environmental enrichment is an enhanced laboratory condition promoting the expression of species typical behaviors, when designing animal models for translational research it is imperative to consider environmental complexity and its impact on neurobiological indices when trying to understand both the underlying etiology of disease and neurotypical development.

One imperative for improving animal models of early-life stressors is to simulate multiple adverse childhood experiences (ACEs). The 2016 National Survey of Children's Health reported that one in ten children experience three or more ACEs and that poverty is one of the most commonly experienced stressors (Sacks and Murphey, 2018). Experiencing multiple ACEs is associated with increased risk for both metabolic and psychiatric diseases in adulthood (Sacks and Murphey, 2018; Afari et al., 2014; Pervanidou and Chrousos, 2012; Heim et al., 2008; Chapman et al., 2004; Anda et al., 2002). While there are many early-life stressors that could be incorporated into the LB paradigm, diet is of special interest considering the association of both food insecurity and obesity with a variety of metabolic and psychological outcomes (Spencer et al., 2017; Yam et al., 2016). Food insecurity is the condition of ones' diet being restricted in terms of variety, nutrition, and amount of food (Eicher-Miller and Zhao, 2018). Children who are food insecure are more likely to have poor outcomes including, but not limited to, generally poor health and greater number of hospitalizations,




Fig. 1. A) Timeline of study and housing conditions. B) Multilevel environmental enrichment cage. C) Representative picture of the WD-LB cage and bedding floor (inset).

psychosocial and behavioral problems, worse developmental outcomes, and are more likely to suffer from childhood obesity (Gundersen and Kreider, 2009). When examining the relationship between food security and low-income women it was found that food insecurity and diet quality were inversely related; those who were identified as food insecure had lower levels of perceived neighborhood safety, had a higher body mass index, and had lower access to healthy food options (Sanjeevi et al., 2018). A Western diet (WD) is characterized by high levels of refined sugars and saturated fats, and low fiber content which are all associated with obesity (Francis and Stevenson, 2012). In animal models, high-fat and sugar diets can cause impairments of the hippocampus and dysregulation of the HPA axis (Boitard et al., 2015; Maniam et al., 2015). In humans there is a higher occurrence of psychological disturbances (e.g. mood, anxiety, somatoform and eating disorders) among adolescents who are obese in addition to reduced hippocampal volume, which is associated with depression (Kalyan-Masih et al., 2016; Björntorp, 2008; Campbell et al., 2004; Britz et al., 2000). On its own early-life stress causes increased vulnerability to metabolic syndromes which is only exacerbated by food insecurity (Yam et al., 2016; Tilburg et al., 2010). For example, early-life stress induced by the LB model paired with a WD in male rats was associated with higher body fat and a positive correlation between white adipose tissue and object recognition scores indicative of cognitive impairments (Yam et al., 2016).

In the present study, we adapted rodent caging systems to simulate low, medium, and high security environments and evaluated their effects on maternal care and offspring development. The ‘insecure’ housing was modelled by a combination of WD-LB exposure, while standard rodent caging served as a ‘medium security’ environment. In

contrast, a high resource environment was modelled by housing a subset of animals in EE. Between these settings we compared differences in maternal care and tested the hypothesis that early life stress caused by a low resource scenario would result in poor maternal care and associated metabolic dysfunction, including precocious puberty in offspring. Finally, we tested whether weaning into EE could rescue some of the adverse outcomes associated with the early-life stress protocol.

2. Methods

2.1. Animals and housing

The experiment was approved by the MCPHS Institutional Animal Care and Use Committee and was carried out in compliance with the recommendations from the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Thirty-six female and twelve male virgin Sprague-Dawley rats were acquired from Charles River (Wilmington, MA) and habituated in same-sex pairs; vivarium kept at 20 °C on a twelve-hour light/dark cycle (0700–1900 light) in larger sized cages (51 × 41 × 22 cm). These cages were a one level cage with access to corn bedding, one plastic tube, one chew bone, Nestlets® (Ancare) and ad libitum access to food and water. After a two-week habitation period, a subset of female ($n = 14$) and male ($n = 4$) animals were allocated to a Western diet (WD; LabDiet® 5TJN). These females were assigned to later undergo LB housing (WD-LB; described below) in order to model a low resource or ‘insecure’ housing condition. A separate set of female ($n = 22$) and male ($n = 6$) rats were maintained on the standard diet (LabDiet® 5001). All animals were

maintained on their respective diet for four weeks, prior to breeding, and remained on their assigned diet until weaning on postnatal day (P) 22. One week prior to breeding ten females fed the standard diet were moved into EE (91.5 × 64 × 159 cm; Critter Nation, Muncie IN), maintaining their same-sex dyads. This EE cohort made up our ‘high resource security’ group. The EE housing units were multilevel cages with access to bedding, one tube, one chew bone, ample Nestlets® and toys. The location and type of toys used were changed two times weekly in order to stimulate novelty. The remaining standard chow fed females ($n = 12$) represented the medium or ‘middle class’ resource control group (standard housed; SD). Animals were weighed once weekly at 12 pm. A timeline of the procedures can be found in Fig. 1A. In line with initiatives to improve the reporting of experimental methods, we have completed the adapted reporting table from Kentner et al. (2018) and provided it as Supplementary Table 1.

2.2. Breeding and delivery

Breeding consisted of pairing one male with two females until pregnancy was verified by increased weight gain and the observation of visible teats. Pregnant females were kept in pairs until approximately gestational day (G)18, at which point they were housed individually in order to prevent the mixing of pups between litters. With respect to the WD-LB and SD groups, pregnant dams were placed into individual standard sized one level cages (27 × 48 × 20 cm). For the pregnant EE animals, a divider was built into the home cage so that litters could be separated – one litter housed in the top portion and one in the bottom until weaning. Each cage section had two levels (see Fig. 1B). Toys were taken away from EE animals on G18 and returned on P14 in order to prevent the risk of pup injury during the early neonatal period. Day of birth was designated as P1; on P2 litters from all housing conditions were culled to $n = 12$ (equal number of males and females where possible). Animals remained undisturbed until the morning of P10 at which point all cages were cleaned. In line with initiatives to reduce the number of animals in research (National Research Council, 2011), on P14, a small subset of male and female animals ($n = 2$) from the WD-LB litters were quickly administered a single dose of lipopolysaccharide (LPS; *Escherichia coli*; serotype O26:B6, L-3755; Sigma, St Louis, MO; 100 mg/kg) so they could be included as part of a separate study (unpublished). It has previously been shown that this treatment does not disrupt maternal care (Spencer et al., 2006). One male and female animal from each litter was evaluated in the measurements described below. A list of offspring group designations can be found in Table 1.

2.3. Limited bedding model

On the late afternoon of P2 a stainless-steel grate was placed into the cage of WD fed animals (adapted from McLaughlin et al., 2016) creating the WD-LB housing group. In our case, the stainless steel grates sat on top of the bedding, restricting access of the dams to their bedding (see Fig. 1C). Additionally, a single piece of paper towel was added in lieu of Nestlets® (Gilles et al., 1996). Cages were left undisturbed until the morning of P10 at which point the stainless-steel grates were removed and litters placed into fresh clean cages with the same level of access to nesting materials as the SD group.

Table 1
Offspring group designations.

Group Name	Description
SD	Control male and female offspring fed a standard diet and housed in standard laboratory cages.
WD-LB	Male and female offspring fed a western diet and housed in standard laboratory cages. Animals were exposed to reduced bedding beginning on the late afternoon of P2 through until the morning of P10.
WD-LB- EE	Male and female offspring fed a western diet and housed in standard laboratory cages. Animals were exposed to reduced bedding beginning on the late afternoon of P2 through until the morning of P10. These animals were placed into environmental enrichment at weaning on P22.
EE	Male and female offspring fed standard diet housed in enriched multilevel cages; offspring were given novel enrichment toys twice weekly starting on P14.

2.4. Maternal care observations

The first week of life is a critical time for rodents (Champagne and Curley, 2007). Between P3-P6 passive home cage maternal behavior observations were conducted twice daily; once in the light phase (07:45–10:00 h) and once in the dark (19:45–22:00 h) to account for the nocturnality of rodents (Champagne and Curley, 2007). Each observational period consisted of five one-minute observations, with five minutes of no observation occurring between each one-minute bin. Recorded maternal behaviors included the frequency of pup retrievals, dam licking pup, passive/low crouch nursing, active/high crouch nursing, nest building/digging in bedding, dam self-grooms, dam sleeping, dam eating/drinking, dam chasing her tail, and total time on nest (seconds).

2.5. Assessment of puberty onset

Beginning on P25, female rats were evaluated daily for vaginal openings (VO) until all animals were scored as open. Animals were scored by their appearance as being either closed, or having a complete vaginal opening (Cowan and Richardson, 2018; Kentner et al., 2018; Grassi-Oliveira et al., 2016).

2.6. Open field test

To evaluate the presence of anxiety-like behavior (Crawley, 2007), an open field test was conducted in dams on P22 and offspring on P50. Thirty minutes prior to testing animals were allowed to habituate to the testing room. Dams were placed into a black, square box (72 cm × 72 cm × 36 cm) and video recorded for 10 min. Juvenile male and female offspring were similarly evaluated in a smaller arena (40 cm × 40 cm × 28 cm) and video recorded for 5 min. Between each recording session all of the equipment was thoroughly cleaned with Quatriside TB. Videos were then scored through an automated behavioral monitoring software program (Cleversys TopScan, Reston, VA) and evaluated for the percent of time (seconds) spent in the center of the arena $[(\text{total time in center})/(\text{total time in center} + \text{total time in perimeter}) * 100]$ and total distance travelled (cm) in the arena.

2.7. Social preference test

Immediately following the open field test, which was used as a habituation period to the test arena, the social preference for a novel conspecific versus an inanimate object (Crawley, 2007) was evaluated. During this assessment, juvenile offspring were placed into the center of the arena and video recorded for 5 min. During this period animals had the choice to visit either a same sex/size/age novel conspecific or an object, each confined within a small wire cup on opposite ends of the arena. Placement of the novel conspecific and the object were counterbalanced between tests and experimental animals groups. Between each recording all of the equipment was thoroughly cleaned with Quatriside TB. Using the software program Stop Watch+ (Atlanta, Georgia), videos were then scored by two blinded observers with an established interrater reliability of > 90%. The variables scored were the duration of time (seconds) spent with and frequency of visits to

either the same sex/size/age novel conspecific, or object. From these variables, a social preference score was calculated and expressed as a preference ratio arena $[(\text{total time with novel rat})/(\text{total time with novel rat} + \text{total time with object}) \times 100]$ where scores > 0.5 indicated a preference for the novel rat compared to the object.

2.8. Tissue collection

Two hours after behavioral testing dams and their offspring underwent a 20-minute restraint stress procedure. During this period, animals were individually restrained in a soft open-ended plastic pastry bag (Daymark) and immediately euthanized with a mixture of Ketamine/Xylazine (40–80 mg/kg, i.p. 5–10 mg/kg, i.p.). Cardiac blood was collected in EDTA tubes and plasma collected. Brains were immediately collected and placed on a clean petri dish set on wet ice. Whole hypothalamus was quickly dissected from the freshly collected tissue and frozen on dry ice. All tissues were stored in -70°C for future processing.

2.9. ELISA

Plasma corticosterone and leptin levels were analyzed in duplicate by ELISA using the manufacturer's instructions. With respect to the plasma corticosterone ELISA we followed the small sample assay protocol provided with the testing kit (ADI-900-097, Enzo Life Sciences, Farmingdale, NY, USA). The minimum detectable concentration was 26.99 pg/mL, and the intra and inter-assay coefficients of variation were 6.6% and 7.8%, respectively. The minimum detectable leptin concentration (abcam ab100773, Abcam) was 30 pg/mL, intra and inter-assay coefficients of variation were $< 10\%$ and $< 12\%$, respectively.

2.10. PCR

Total RNA was extracted from frozen hypothalamus with TRIzol reagent (Thermo Fisher). Isolated RNA was quantified utilizing a NanoDrop 2000 spectrophotometer (Thermo Fisher). cDNA synthesis and qPCR were performed using TaqMan assay probes to evaluate *Kiss1*, *Kiss1R*, *Gnrh1*, *Gnrhr* and *Gapdh* (Rn00710914_m1, Rn00576940_m1, Rn00562754_m1, Rn00578981_m1 and Rn01775763_g1, respectively) in a StepOne Plus station (Applied Biosystems) as described previously (Kentner et al., 2018). Each sample was analyzed in duplicate and relative gene expression levels were evaluated using the $\Delta\Delta\text{Ct}$ method with *Gapdh* as the housekeeping gene. Data are presented as mean expression relative to CON-nSaline animals, which have been normalized to 1 for each gene of interest.

2.11. Westerns

Microdissections of the medial prefrontal cortex (mPFC), centered on the infralimbic and prelimbic regions with the ventral anterior cingulate cortex included (A/P = 3.24 mm and M/L = 0 mm from Bregma, D/V = -3.2 mm from the surface of the brain) were taken using a trephine from flash froze tissue. Tissue samples were homogenized in RIPA buffer (R0278, Sigma) and Halt™ protease and phosphatase inhibitor cocktail (ThermoFisher, 78440), spun at $40,000 \times g$ for 30 min at 4°C , and the supernatant assayed with the BCA protein Assay (ThermoFisher, 23227). Samples (30 μg) were loaded on 4–15% Mini-PROTEAN® TGX™ Precast Protein Gels (BioRad, 456-1084). Gels were run for 35 min, at 200 V in $1 \times$ Tris/Glycine/SDS buffer (BioRad, 1610771). Then gels were transferred to a PVDF membrane (Immobilon-FL, IPFL00010) in $1 \times$ Tris/Glycine buffer (BioRad, 1610734) for 1.5 h at 55 V. After transfer, membranes were stained with the REVERT Total Protein Stain Kit (BioRad, 926-11016) for total protein normalization according to manufacturer's instructions and immediately imaged with Odyssey FC Imaging System using Image

Studio Life Version 4.0. Then membranes were blocked with Odyssey Blocking Buffer (927-50000) for 1 h. Afterwards, membranes were probed with an anti-synaptophysin antibody, clone SY38 (1:1000, Millipore, MAB5258) diluted in 1:1 Odyssey Blocking Buffer:TBS overnight at room temperature. Membranes were washed three times for 5 min with TBST and then probed with anti-mouse IRDye 800CW (1:20,000, LiCor) 4 h at room temperature. After three additional 5 min TBST washes, membranes were placed in TBS before imaging with Odyssey FC Imaging System. After scanning, Odyssey Infrared Imaging software quantified the integrated intensity of each band and determined molecular weights based on Biorad Precision Plus Protein Standards (1610374). The ratio of synaptophysin to total protein for the lane was calculated (Kirshner and Gibbs, 2018). For the figures, each channel of the image was adjusted for brightness and contrast individually using the Odyssey Infrared Imaging software and adjustments were applied equally to the entire image.

2.12. Statistical analyses

Statistics were performed using the software package Statistical Software for the Social Sciences (SPSS) version 21.0 (IBM, Armonk, NY) or GraphPad Prism (version 7.0), in the case of the survival curves described below. Alpha levels were set to $p < 0.05$ for all omnibus tests. A one- or two-way repeated measures ANOVA was used to evaluate maternal (baseline, weeks 1–4, P22) and offspring body weights (P22, P29, P36, P43, P50) respectively. In each case, time was the repeated factor while either housing or housing and sex were the independent variables, as appropriate. The Greenhouse–Geisser correction procedure was applied due to violations to the assumption of sphericity for repeated measure designs. For maternal behavior, Chronbach's alpha was calculated and interrater reliability between two raters was found to be $> 80\%$. Due to the severity of skewness (Shapiro-Wilks) for maternal and offspring behaviors and plasma corticosterone levels, non-parametric Kruskal-Wallis tests (non-parametric equivalent to the one-way ANOVA) were employed. Pairwise comparisons were made using conservative adjusted alpha correction values. Day of VO and female offspring hypothalamic gene expression were evaluated using one way-ANOVAs. With respect to parametric data, LSD post hocs were applied except where there were fewer than three levels, in which case pairwise *t*-tests and Levene's (applied in the occurrence of unequal variances on the post hoc assessments) were utilized. The time to VO was also assessed, using the Log-rank (Mantel Cox) test, and plotted as a survival curve. For applicable offspring analyses, if there were no sex differences, then male and female data were collapsed together for analyses and presentation. Data were graphically expressed as mean \pm SEM.

3. Results

3.1. Maternal body weights

A repeated measures ANOVA revealed a significant interaction between time and housing for maternal body weights ($F(3.956, 59.337) = 4.743$, $p = 0.002$; Fig. 2). LSD post hocs did not show a difference across housing groups for maternal body weight at either baseline or P22, however significant group differences were observed across the course of the study with WD rats showing significantly greater weight gain compared to SD controls (Week 2: $p = 0.001$; Weeks 3 and 4: $p = 0.0001$) and EE animals (Week 3: $p = 0.05$; Week 4: $p = 0.023$), validating the western diet treatment. EE females also had significantly more weight gain than SD rats across several points of the study (Week 1: $p = 0.008$; Week 2: $p = 0.023$; Week 4: $p = 0.006$).

3.2. Maternal care

With respect to P3–6 maternal behavior, although there were no




Fig. 2. Validation of Western diet on body weight gain (grams) across time. Baseline, weeks 1–4 (prior to breeding), and P22 (weaning). Data expressed as mean \pm SEM; $n = 8$ –10 per group. Various symbols are employed (e.g., \$, &, *) to delineate differences between groups where * $p < 0.05$, ** $p < 0.001$, *** $p < 0.0001$, difference between SD and WD; #indicates differences between SD and EE; &indicates difference between WD and EE.

treatment differences with respect to time on nest (seconds) during the light phase observation period ($X^2(2) = 4.535$, $p = 0.104$; Fig. 3A), Kruskal-Wallis tests revealed a significant main effect of housing during the night ($X^2(2) = 12.545$, $p = 0.002$; Fig. 3B) and for the combined total of the daylight and nighttime observations ($X^2(2) = 11.319$, $p = 0.003$; Fig. 3C) on this measure. Follow-up tests demonstrated that SD (dark: $p = 0.021$; light & dark: $p = 0.013$) and WD-LB (dark:

$p = 0.002$; light & dark: $p = 0.007$) dams both spent more time on nest than EE mothers.

During the light phase, there was a significant main effect of housing for licking/grooming ($X^2(2) = 12.763$, $p = 0.002$), active nursing ($X^2(2) = 7.539$, $p = 0.023$), passive nursing ($X^2(2) = 7.954$, $p = 0.019$), maternal self-grooming ($X^2(2) = 14.779$, $p = 0.001$), maternal eating and drinking ($X^2(2) = 6.396$, $p = 0.041$), and tail chasing ($X^2(2) = 7.495$, $p = 0.024$) behaviors (Fig. 3D). Post hoc analysis revealed that WD-LB dams licked their pups more frequently than both SD ($p = 0.004$) and EE ($p = 0.012$) mothers. WD-LB dams also groomed themselves more than those reared in SD cages ($p = 0.0001$). EE animals engaged in active nursing significantly more often than SD dams ($p = 0.019$) and SD dams engaged in passive nursing significantly more frequently than EE dams ($p = 0.016$). Finally, WD-LB animals demonstrated more tail chase behaviors than EE dams ($p = 0.024$).

For observations during the dark phase, there were differences in the frequency of licking/grooming ($X^2(2) = 10.478$, $p = 0.005$), active nursing ($X^2(2) = 6.491$, $p = 0.039$), passive nursing ($X^2(2) = 10.897$, $p = 0.004$), and total nursing ($X^2(2) = 13.772$, $p = 0.001$) behaviors (Fig. 3E). Pairwise comparisons revealed that WD-LB dams licked their pups significantly more than both EE ($p = 0.014$) and SD ($p = 0.015$) mothers. EE animals engaged in fewer bouts of passive nursing compared to SD ($p = 0.014$) and WD-LB ($p = 0.007$) animals. The same pattern was also true for the frequency of total nursing behaviors (WD-LB vs SD: $p = 0.026$; WD-LB vs EE: $p = 0.001$).

For the combined day and nighttime observation periods there were differences in licking/grooming pups ($X^2(2) = 16.716$, $p = 0.0001$), active nursing ($X^2(2) = 10.235$, $p = 0.006$), passive nursing ($X^2(2) = 10.067$, $p = 0.007$), total nursing ($X^2(2) = 9.844$, $p = 0.007$),




Fig. 3. Maternal behavior observations from standard (SD), Western diet-limited bedding (WD-LB), and environmentally enriched (EE) rat dams as a function of postnatal days in the light phase (left panels), dark phase (middle panels) and light and dark phases combined (right panels). Panels A–C display total time on nest (seconds). Panels D–F display the frequency of licking, active and passive nursing, total nursing, maternal self-grooming, maternal eating and drinking and tail chase behaviors. All maternal behavior data are expressed as mean \pm SEM; $n = 8$ –10 per group. Various symbols are employed (e.g., \$, &, *) to delineate differences between groups where * $p < 0.05$, ** $p < 0.01$, *** $p < 0.0001$ indicates differences between SD and WD-LB, #indicates differences between SD and EE. &indicates differences between WD-LB and EE.

maternal self-grooming ($X^2(2) = 6.627, p = 0.036$), and tail chase ($X^2(2) = 6.337, p = 0.042$) behaviors (Fig. 3F). Post hoc demonstrated that WD-LB dams licked their pups significantly more often than both SD ($p = 0.001$) and EE ($p = 0.002$) mothers. For active nursing, SD dams engaged in significantly fewer bouts than both WD-LB ($p = 0.036$) and EE ($p = 0.01$) animals. SD dams engaged in significantly more passive nursing behaviors than the EE ($p = 0.006$) dams. For total nursing behaviors, EE animals did not nurse as frequently as either the SD ($p = 0.019$) or WD-LB ($p = 0.013$) animals. With regards to maternal self-grooming, there were no significant differences across housing groups. Finally, WD-LB animals chased their tails significantly more than EE ($p = 0.042$) dams.

3.3. Maternal open field test and stress-induced plasma corticosterone

No significant differences were observed across housing groups on measures of P22 maternal anxiety-like behavior and locomotor activity (open field test) or on maternal stress-induced (20 min restraint stressor) plasma corticosterone ($p > 0.05$, data not shown).

3.4. Offspring body weights

A two way repeated measures design (housing by sex) revealed a sex by time interaction ($F(2.199, 142.921) = 79.720, p = 0.0001; n = 8-10$; Fig. 4AB). Post hoc showed that male offspring were heavier than females ($p < 0.0001$) at P29, P36, P43, and P50. There was also a housing by time interaction ($F(6.596, 142.921) = 2.451, p = 0.023$; Fig. 4AB). WD-LB animals were heavier than SD and EE offspring at weaning on P22 (SD: $p = 0.0001$; EE: $p = 0.0001$) and on P29 (SD: $p = 0.0001$; EE: $p = 0.0009$). WD-LB-EE animals had significantly higher body weights than SD ($p = 0.0001$) and EE ($p = 0.0001$), but were not different than WD-LB offspring on P22. On P29, EE offspring weighed more than SD rat ($p = 0.017$). By P29, WD-LB-EE rats weighed significantly less than WD-LB offspring, suggesting that enrichment was able to quickly rescue the metabolic disruptions induced by a high fat diet. There were no differences in weight across the housing groups by P36, supporting the existence of a resilient weight regulation mechanism that may have turned on after the cessation of the western diet on P22.

3.5. Offspring behaviors

Non-parametric Kruskal-Wallis tests did not uncover sex or housing differences for either percent time spent in the center of the open field or for locomotor activity level ($n = 8-10; p > 0.05$, data not shown). Non-parametric analysis revealed significant main effects of sex on the frequency of visits to the novel rat ($X^2(1) = 16.667, p = 0.0001$; Fig. 4C) and object ($X^2(1) = 14.553, p = 0.0001$; Fig. 4D) in the social preference test, in that females made a higher number of visits overall ($p < 0.05$). There was also a main effect of housing for the total frequency of visits that female offspring made to the novel object ($X^2(3) = 17.970, p = 0.043$; Fig. 4D) in that female SD rats visited the novel object more than WD-LB-EE rats. Kruskal-Wallis tests uncovered a main effect of housing on the social preference test ($X^2(3) = 8.282, p = 0.041$; Fig. 4E), but follow-up tests did not confirm differences as a function of treatment ($p > 0.05$).

3.6. Precocious puberty, hypothalamic gene expression, and plasma leptin concentrations

One way ANOVA showed a significant effect of housing on pubertal maturation ($F(3, 34) = 3.235, p = 0.034; n = 8-10$, Fig. 5A). LSD follow up tests revealed that full vaginal opening occurred significantly earlier in WD-LB female offspring compared to SD ($p = 0.025$), EE ($p = 0.040$), and WD-LB-EE ($p = 0.007$) rats. A more in depth evaluation of time to vaginal opening using the Log-rank (Mantel-Cox) test

revealed a significant effect of housing ($X^2(3) = 11.83, p = 0.008$; Fig. 5B). Comparisons of the survival curves revealed significant differences in day of vaginal opening (DVO) between SD (DVO: P38) vs. WD-LB (DVO: P36; $X^2(1) = 4.685, p = 0.0304$), WD-LB vs. WD-LB-EE (DVO: P38.5; $X^2(1) = 8.225, p = 0.0041$), as well as WD-LB vs. EE (DVO: P38; $X^2(1) = 4.628, p = 0.0315$). Animals from all treatment groups displayed full vaginal openings by P43 (Fig. 5B).

The kisspeptin/gonadotropin releasing hormone (GnRh) system is an important pathway integral to puberty initiation (Kauffman et al., 2007; Skorupskaitė et al., 2014) so we evaluated the P50 expression of several related hypothalamic gene targets ($n = 7-10$). One way ANOVAs did not indicate differences as a function of housing for *Gnrh1* or for its receptor *Gnrhr* ($p > 0.05$, Fig. 5C). However, there were significant main effects of housing for *Kiss1* ($F(3,30) = 3.089, p = 0.043$; Fig. 5C) and *Kiss1r* expression ($F(3, 30) = 7.708, p = 0.022$; Fig. 5C). LSD follow-up tests showed that WD-LB female rats had significantly lower levels of *Kiss1* compared to SD rats ($p = 0.006$). Environmental enrichment did not completely rescue this effect (WD-LB vs WD-LB-EE: $p = 0.052$). However, WD-LB-EE animals were not significantly different from SD rats on this measure ($p > 0.05$), suggesting some modest attenuation following enrichment. Hypothalamic *Kiss1r* was significantly elevated in WD-LB-EE animals compared to both SD ($p = 0.005$) and WD-LB ($p = 0.016$) females. This suggests that the elevated levels of *Kiss1r* following EE may have compensated in part for the depleted *Kiss1* levels associated with WD-LB housing.

Given evidence that plasma leptin concentrations are correlated with precocious puberty we elected to measure this hormone in our female animals (Sominsky et al., 2016). A one-way ANOVA revealed a significant housing effect on plasma leptin concentrations ($F(3, 25) = 3.172; p = 0.042$; Fig. 5D). LSD post hoc tests demonstrated that WD-LB female rats had higher plasma leptin levels compared to WD-EE ($p = 0.007$) and EE ($p = 0.033$) animals, respectively. There were no significant differences between SD and WD-LB females ($p > 0.05$).

3.7. Offspring plasma corticosterone

No significant differences were observed across experimental groups on measures of stress-induced (20 min restraint stressor) plasma corticosterone for P50 offspring ($n = 7-10, p > 0.05$, data not shown).

3.8. Levels of synaptophysin in the mPFC

The mPFC continues to develop through puberty, and this development is characterized by alterations in synapses (Drzewiecki et al., 2016). To determine whether sex or housing influenced mPFC maturation we evaluated synaptophysin, a marker of synapses used to assess cortical synaptic development (Glantz et al., 2007). Values that exceeded 2 standard deviations above or below the group mean were considered outliers and dropped from the analysis. No significant differences were observed in synaptophysin levels in mPFC ($n = 8-10, p > 0.05$, Fig. 6).

4. Discussion

The present experiments were designed to assess the impact of high, medium, and low security environments on maternal care quality and, in tandem, to evaluate the maturation and neurobehavioral outcomes of male and female offspring. The results show that resource insecurity and environmental complexity differentially affect maternal behavior, but not along a continuum ranging across defined categories from 'low' to 'high' quality, as one might expect. Instead, circadian timing greatly influenced the temporal pattern of behavioral expression, unveiling potential strengths and weaknesses in maternal care quality across the housing conditions. Offspring born to low resource WD-LB dams were resilient to substantive disruptions in behavior and mPFC plasticity.




Fig. 4. Offspring body weights (grams) for male (A) and female (B) animals. C) Social preference ratios for male and female (collapsed) offspring. Values > 0.05 indicate that the experimental rat spent more time visiting with the novel rat compared to the novel object. The total frequency of visits made to the D) novel rat and E) novel object in the social preference test. All data are expressed as mean ± SEM, n = 8–10. Various symbols are employed (e.g., a, b, c) to delineate differences between groups where ^ap < 0.05, ^{aa}p < 0.01, ^{aaa}p < 0.001 indicates differences between males and females; ^bindicates differences between SD and WD-LB, ^cindicates differences between WD-LB and EE, ^dindicates differences between WD-LB and WD-LB-EE, ^eindicates differences between WD-LB-EE and EE, ^findicates differences between SD and WD-LB-EE, and ^gindicates differences between SD and EE.

Only subtle perturbations were observed in the measures evaluated here, and were limited to differences in exploration of the novel object by female WD-LB rats in the social preference test. This did not translate into impairments of species typical preferences for social novelty. Instead, consequences of early life stress were most apparent by metabolic changes such as increased body weight and accelerated puberty, in addition to alterations in the levels of the kisspeptin gene and expression of its receptor. Importantly, placement into EE during the early juvenile period fast-tracked the regulation of body weight after cessation of the Western diet in WD-LB offspring. Moreover, the juvenile enrichment intervention prevented the precocious puberty associated with a history of early life stress. Given that enrichment is a translationally valid intervention for both animals and humans (Woo and Leon, 2013; Woo et al., 2015; Kentner et al., 2018), this study offers further insight into additional unexplored uses for this tool. This is in addition to highlighting the utility of developing and incorporating

clinically relevant animal models into our basic research.

While animal models can be used as a translational approach to evaluate the neurobehavioral effects of early-life stress, or even positive early life experiences, it is important to acknowledge their strengths and shortcomings in terms of test validity. For example, construct, face, and predictive validity satisfy the three-criteria required for an animal disease model: congruent risk factors, symptoms, and efficacy of treatment, respectively (Estes and McAllister, 2016; Belzung and Lemoine, 2011). Although early-life exposure to stressors do not always lead to a disease state, these adverse experiences do increase the risk for mental illnesses (e.g. depression, anxiety) in later life (Kumsta et al., 2017; Burkholder et al., 2016; Bentall et al., 2012; Green et al., 2010; Heim et al., 2008; Chapman et al., 2004). Employing animal models of early-life stress such as maternal separation or LB meet construct validity in that they mimic risk factors such as neglect or abuse. Moreover, they demonstrate face validity in that specific symptoms of mental




Fig. 5. A) Day of full vaginal opening for female offspring. B) Survival curves of full vaginal opening for each group expressed across time. C) Relative hypothalamic gene expression of *Gnrh1*, *Gnrhr1*, *Kiss1*, *Kiss1R* normalized to SD controls. D) Plasma leptin levels (pg/mL) on postnatal day 50. All data are expressed as mean \pm SEM. Various symbols are employed (e.g., a, b, c) to delineate differences between groups where ^a $p < 0.05$, ^{aa} $p < 0.01$, ^{aaa} $p < 0.001$ indicates differences between WD-LB and SD; ^bindicates differences between WD-LB and EE, ^cindicates differences between WD-LB and WD-LB-EE; ^dindicates differences between SD and WD-LB-EE.

illnesses (e.g. impaired emotional reactivity, disrupted HPA regulation, cognitive disturbances, anxiety-like behavior) can be observed in animals that have experienced these stressors during the perinatal period (Grassi-Oliveira et al., 2016; Leussis et al., 2012; Raineki et al., 2012; Chatterjee et al., 2007; Gonzalez et al., 2001; Fernandez-Teruel et al., 1991).

Animal models of early-life stress can also be used to predict treatment efficacy. For example, in humans, skin-to-skin contact in the form of Kangaroo Care has been successful in treating the adverse effects of premature birth and improving parent-infant attachment when compared to traditional treatment (Charpak et al., 2016; Baley, 2015;

Tessier et al., 2003). Similarly, when rat pups are isolated or artificially reared in the laboratory, maternal tactile contact can be simulated by stroking the animal with a paintbrush to emulate licking and grooming behaviors (Chatterjee et al., 2007). Experimentally this has led to improvements in a number of neural proteins integral to development: higher densities of synaptophysin, neural cell adhesion molecule, growth associated protein 43, brain-derived neurotrophic factor and glucocorticoid receptor expression, when compared to control animals (Chatterjee et al., 2007; Kentner et al., 2018). Artificial rearing can have an intergenerational, detrimental behavioral effect. Artificially reared mothers will engage in the same maternal behaviors as control mothers but they will do so for less time, and additionally these dams will show an increase of non-maternal behaviors such as digging, biting at the cage floor, and chasing their own tail (Lomanowska and Melo, 2016; Barrett and Fleming, 2011; Burton et al., 2007; Gonzalez et al., 2001). Tactile stimulation during artificial rearing can improve the quality of maternal behaviors that artificially reared mothers provide to their own offspring, compared to artificially reared mothers that did not receive tactile stimulation (Gonzalez et al., 2001). For human infants, similar tactile contact in the form of Kangaroo Care has short-term improvements in birthweight and survival, and long-term improvements in academic success, and higher IQ (Charpak et al., 2016; Baley, 2015; Tessier et al., 2003). For human mothers there are improvements in breast milk production, reduced feelings of hopelessness, and better infant attachment (which also extends to fathers; Baley, 2015).

In a similar vein, children who were institutionalized and then moved to foster care showed improvements in cognitive, motor, and behavioral domains (as measured by the Bayley Scale of Infant Development) as well improvements in intellectual functioning in verbal and performance domains (as measured by the Weschler Preschool Primary Scale of Intelligence) when compared to their peers who remained institutionalized (Nelson III et al., 2007). When observing the differences between institutionalized vs never institutionalized children's brains MRI revealed that those with a history of institutionalization had significantly reduced cortical gray matter volume (Sheridan et al., 2012). Foster care is superior to institutionalization because of better caregiver contact, better access to resources, and a higher-quality environment all of which are beneficial to cognitive development (Sheridan et al., 2012). Physical contact from their caregivers to their children is essential to the child's development. Even in healthy infants, skin-to-skin contact with parents has been shown to stabilize body temperature and blood glucose concentrations, decreases crying, and supports cardiorespiratory stability (Feldman-Winter and Goldsmith, 2016). In animal models, EE has been demonstrated to reverse the negative effects of processive and systemic stressors (Champagne and Curley, 2016; Kentner et al., 2016; Vivineto et al., 2013; Schneider et al., 2006; Bredy et al., 2004). Thus, improvements to one's environment can lead to better outcomes in both humans and rats. While it is important to model the effects of stress in animals, it is also important to model the effects of EE so that efficacy of potential interventions can be established. Indeed, in the present study, we moved juvenile LB animals into EE to measure whether the metabolic consequences of early life stress could be rescued. Overall, each animal model of early-life stress will have its own particular strengths and weaknesses, but the most translational ones will meet several of these tests of validity.

With respect to the current study, the maternal care quality of EE dams was similar to what our laboratory has previously reported (Connors et al., 2015), except here we evaluated individual litters in isolation, as opposed to as part of a co-parenting design. This allowed us to better control for the contributions of individual dams on their litters, which is important given reports that dams may 'take-turns' caring for pups in a colony nest setting (Sale et al., 2004). It should be noted that in these co-parenting conditions one dam could not sufficiently meet the needs of both litters on her own, and it is unlikely that pups would receive more maternal attention overall. However, one great

Our WD-LB exposed offspring were resilient to behavioral perturbations, as has been similarly reported by others who exposed animals to a maternal high fat diet (Zieba et al., 2018). We only observed minor sex-specific disruptions in the frequency of novel object exploration in the social preference test. In contrast to the negligible behavioral effects reported here, animals exposed to limited resource conditions typically display robust deficits in attachment learning and impaired social behaviors, depressive and anxiety-like behaviors and disrupted maternal care (Moriceau et al., 2009; Rainekei et al., 2010; Rainekei et al., 2012; Dalle Molle et al., 2012; Brunson et al., 2005; Rincón-Cortés and Sullivan, 2016; Roth et al., 2009), although it appears that whether or not aberrant behavior is observed may be task specific (Ivy et al., 2010) or even dependent on age of testing or animal strain (Maniam et al., 2016; Brunson et al., 2005; Dalle Molle et al., 2012).

An important point to consider is that our LB animals were exposed to a WD preconceptionally and maintained on this diet across early development, until weaning, at which point it was replaced with regular chow. A post-weaning high fat diet has been demonstrated to reverse the anxiety-like effects associated with LB (Maniam et al., 2016), suggesting it may have value as an intervention tool (Maniam et al., 2016; Machado et al., 2013). Here, our data indicate that a Western style diet may also have *preventative* effects against the consequences of LB exposure. Indeed, maternal high-fat diet mitigated a variety of negative outcomes (e.g. anxiety, cognitive and social impairments, nociception and HPA dysregulation) associated with maternal separation including downregulation of *Bdnf* and *Crh* mRNA, spine loss and dendritic atrophy (Rincel et al., 2016; Rincel et al., 2018). This could account for the small impact that WD-LB housing had on offspring brain and behavior in the present work. Importantly, maternal high fat diet exposure is associated with increased maternal care (Purcell et al., 2011; Rincel et al., 2016) implicating an indirect pathway by which diet may influence offspring development. While our own WD-LB dams spent relatively less time in active nursing postures, compared to EE mothers, they engaged in higher rates of licking and grooming which conceivably may have counteracted the behavioral effects of early life stress through programming of the HPA-axis (Liu et al., 1997). This is important to note because although EE dams had lower total contact time with their nest, they made-up for this by participating in more active nursing postures. Together, this reminds us that despite the available resources no parent is perfect and for the most part the kids are (behaviorally) alright.

Although offspring of WD-LB animals appeared to be behaviorally resilient to the early life stress, metabolic consequences of this condition were more apparent. Specifically, these animals had higher body weights at weaning and female offspring displayed precocial puberty and downregulated *Kiss1* mRNA expression in the hypothalamus. Mutations and deletions of *Kiss1* or its receptor are accompanied by impairments in reproductive functioning, including an interruption of pubertal development, associated with reductions in GnRh (Kauffman et al., 2007). Moreover, prepubertal administration of kisspeptin initiates precocious puberty, as indicated by LH release and advanced vaginal openings (Navarro et al., 2004). Therefore, we were surprised to measure diminished expression of *Kiss1* mRNA in our WD-LB animals. Indeed, we had anticipated that elevated levels of this gene would stimulate GnRh secretion, resulting in the accelerated pubertal onset associated with WD-LB. Since we evaluated whole hypothalamus, it should be considered that we overlooked the tissue specific nature regulating the expression of these genes. For example, sex steroids inhibit or stimulate *Kiss1* levels in a region specific manner (see Kauffman et al., 2007). We may have observed the predicted gene expression patterns (e.g. higher *Kiss1*) if we had targeted localized regions of the hypothalamus. It should also be noted that we collected tissue following puberty initiation and we likely missed a critical window. Future work will need to include time-dependent sampling methods in order to capture a more accurate developmental picture of how early life stress

affects the GnRH-Kisspeptin system and its influence on puberty. In addition to that, potential compensatory mechanisms such as differences in receptor sensitivity and regulatory-feedback systems will need to be evaluated. Finally, one must remember that the proportion of mRNA does not necessarily correlate to protein levels as this relationship is coordinated by several nuanced biological processes (Payne, 2015; Vogel and Marcotte, 2012).

In medicine, the pathophysiology of precocial puberty typically only considers physical trauma, genetic predisposition, illness or some physical cause for the accelerated maturation (Kaplowitz and Hoffman, 2018; Chalumeau et al., 2002; Strauss and Barbieri, 2009). However, early life stress is beginning to garner more attention as being a precipitator of early pubertal onset (Li et al., 2014; Kelly et al., 2017; Cowan and Richardson, 2018; Viridis et al., 1998). Recently, we observed precocious puberty in a rodent model of the neonatal intensive care unit (mimicking reduced parental contact, nosocomial infection, and medical manipulations). However, in this early life stress model we measured reduced *GnRh* receptor, as opposed to *Kiss1* (Kentner et al., 2018), though this finding is vulnerable to the methodological limitations listed above. Alternatively, these data suggest that different early life stressors may result in similar physiological outcomes (in this case accelerated puberty) that are mediated via a diverse set of mechanistic pathways. This finding is similar to observations that males and females can have similar behavioral outputs mediated by different mechanisms (e.g. Sorge et al., 2015) and the future mapping of this will be interesting to follow. Even though the genes between our early life stress models are differentially affected, they are both suggestive of suppressed gonadotropin levels and possibly a peripheral GnRh-independent mechanism (Barker and Kappy, 2011). The centrally measured changes in GnRh-Kisspeptin genes could consequently be due to disruptions in HPG feedback mechanisms but the underlying programming effects of early life challenges are mostly unclear (Kentner and Pittman, 2010).

We show that environmental enrichment, in the form of a more complex laboratory caging system, prevented several central and metabolic disruptions following early life stress. Indeed, enhanced environmental stimulation accelerated the restoration of body weight to levels of SD regular chow fed rats when access to the WD food ceased. This is a natural metabolic defence mechanism that occurs when regular chow is restored in rodents (Siersbaek et al., 2017; Fisher et al., 2018; Raun et al., 2007), and EE accelerated this process. Moreover, the prepubertal experience of enhanced laboratory stimulation was associated with elevated levels of hypothalamic *Kiss1r* in WD-LB-EE females, in addition to the normalization of species typical pubertal onset. Interventions in the form of sensory stimulation replacement and microbiome manipulation have also counteracted the occurrence of precocial puberty as a consequence of early life adversity (Kentner et al., 2018; Cowan and Richardson, 2018). In the low resource stress model used here, the pattern of changes in the kisspeptin system are suggestive of a compensatory mechanism to counteract the reduction in *Kiss1* observed in WD-LB exposed rats. While *Kiss1r* was also lower in SD compared to WD-LB-EE females, SD rats had elevated expression of *Kiss1*, indicating that hormone levels in these animals may have been sufficient to maintain homeostasis and typical pubertal development. In general, we hypothesize that the enrichment conferred protection in WD-LB-EE females, maintaining an adequate balance between the ligands and receptors of the kisspeptin system, preventing the pubertal timing disturbances associated with early life stress. What needs to be determined is whether this animal model of low resource availability interrupts factors such as sexual motivation and behavior, or even fertility, and if EE might prevent or even restore these systems. As mentioned, future work will also need to address potential changes in the trajectory of kisspeptin and GnRh associated genes expression across development and address the issue of tissue specificity.

In sum, these studies highlight the utility to develop better translational models of low, medium, and high insecurity environments. Our

results suggest that the negative consequences of a high insecurity environment can be mitigated by parental care and environmental enrichment in the juvenile period. As these non-pharmacological interventions and their neurobiological effects are underexplored in the basic literature, employing these models can lead to novel insights into mechanisms that promote stress resilience.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.yhbeh.2019.01.003>.

Acknowledgements

The authors would like to extend their thanks to both Amanda Speno and Isabella Connors for technical assistance. This project was funded in part by NIMH under Award Number R15MH114035 (to ACK) and by the National Science Foundation (NSF CAREER IOS-1552416 to D.A.B). The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the financial supporters.

Author contributions

A.C.K. and D.A.B designed and supervised the study, A.R.S., E.O., A.R.-O., S.T.F., and A.R. ran the experiments, A.C.K., A.R.S., D.A.B., and E.O., S.T.F., analyzed the data and A.R.S., and A.C.K wrote the manuscript.

Disclosures and potential conflict of interests

None.

References

- Aber, J.L., Bennett, N.G., Conley, D.C., Li, J., 1997. The effects of poverty on child health and development. *Annu. Rev. Public Health* 18, 463–483. <https://doi.org/10.1146/annurev.publhealth.18.1.463>.
- Afari, N., Ahumada, S.M., Wright, L.J., Mostoufi, S., Golnari, G., Reis, V., Cuneo, J.G., 2014. Psychological trauma and functional somatic syndromes: a systematic review and meta-analysis. *Psychosom. Med.* 76, 1–20. <https://doi.org/10.1097/PSY.000000000000010>.
- Anda, R.F., Whitfield, C.I., Felitti, V.J., Chapman, D., Edwards, V.J., Dube, S.R., Williamson, D.F., 2002. Adverse childhood experiences, alcoholic parents, and later risk of alcoholism and depression. *Psychiatr. Serv.* 1001–1009. <https://doi.org/10.1176/appi.ps.53.8.1001>.
- Angold, A., Worthman, C.W., 1993. Puberty onset of gender differences in rates of depression: a developmental, epidemiologic and neuroendocrine perspective. *J. Affect. Disord.* 29 (2–3), 145–158. [https://doi.org/10.1016/0165-0327\(93\)90029-j](https://doi.org/10.1016/0165-0327(93)90029-j).
- Avishai-Eliner, S., Gilles, E., Eghbal-Ahmad, M., Bar-El, Y., Baram, T.Z., 2001. Altered regulation of gene and protein expression of hypothalamic-pituitary-adrenal axis components in an immature rat model of chronic stress. *J. Neuroendocrinol.* 13, 799–807.
- Baenninger, L.P., 1967. Comparison of behavioural development in socially isolated and grouped rats. *Anim. Behav.* 15, 312–323.
- Baley, J., 2015. Skin-to-skin care for term and preterm infants in the neonatal ICU. *Pediatrics* 136, 596–599. <https://doi.org/10.1542/peds.2015-2335>.
- Barker, J.M., Kappy, M.S., 2011. Precocious puberty in girls. *Bernmans Pediat. Decis. Mak.* 168–172. <https://doi.org/10.1016/b978-0-323-05405-8.00051-6>.
- Barrett, J., Fleming, A.S., 2011. Annual research review: all mothers are not created equal: neural and psychobiological perspectives on mothering and the importance of individual differences. *J. Child Psychol. Psychiatry* 52, 368–397. <https://doi.org/10.1111/j.1469-7610.2010.02306.x>.
- Belzung, C., Lemoine, M., 2011. Criteria of validity for animal models of psychiatric disorders: focus on anxiety disorders and depression. *Biol. Mood Anxiety Disord.* 1–9. <https://doi.org/10.1186/2045-5380-1-9>.
- Bentall, R.P., Wickham, S., Shevlin, M., Varese, F., 2012. Do specific early-life adversities lead to specific symptoms of psychosis? A study from the 2007 the adult psychiatric morbidity survey. *Schizophr. Bull.* 38, 734–740. <https://doi.org/10.1093/schbul/sbs049>.
- Björntorp, P., 2008. Do stress reactions cause abdominal obesity and comorbidities? *Obes. Rev.* 2, 73–86. <https://doi.org/10.1046/j.1467-789x.2001.00027.x>.
- Boitard, C., Maroun, M., Tantot, F., Cavaroc, A., Sauviant, J., Marchand, A., ... Ferreira, G., 2015. Juvenile obesity enhances emotional memory and amygdala plasticity through glucocorticoids. *J. Neurosci.* 35, 4092–4103. <https://doi.org/10.1523/JNEUROSCI.3122-14.2015>.
- Bredy, T.W., Humpartzoomian, R.A., Meaney, M.J., 2003. Partial reversal of the effect of maternal care on cognitive function through environmental enrichment. *Neuroscience* 118, 171–576. [https://doi.org/10.1016/S0306-4522\(02\)00918-1](https://doi.org/10.1016/S0306-4522(02)00918-1).
- Bredy, T.W., Zhang, T.Y., Grant, R.J., Diorio, J., Meaney, M.J., 2004. Peripubertal environmental enrichment reverses the effects of maternal care on hippocampal development and glutamate receptor subunit expression. *Eur. J. Neurosci.* 20, 1355–1362. <https://doi.org/10.1111/j.1460-9568.2004.03599.x>.
- Britz, B., Siegfried, W., Ziegler, A., Lamertz, C., Herpertz-Dahlmann, B.M., Remschmidt, H., ... Hebebrand, J., 2000. Rates of psychiatric disorders in a clinical study group of adolescents with extreme obesity and in obese adolescents ascertained via a population based study. *Int. J. Obes.* (12), 1707–1714.
- Brunson, K.L., Kramár, E., Lin, B., Chen, Y., Colgin, L.L., Yanagihara, T.K., et al., 2005. Mechanisms of late-onset cognitive decline after early-life stress. *J. Neurosci.* 25, 9328–9338.
- Burchinal, M.R., Peisner-Feinberg, E., Bryant, D.M., Clifford, R., 2000. Children's social and cognitive development and child-care quality: testing for differential associations related to poverty, gender, or ethnicity. *Appl. Dev. Sci.* 4, 149–165. https://doi.org/10.1207/S1532480XADS0403_4.
- Burkholder, A.R., Koss, K.J., Hostinar, C.E., Johnson, A.E., Gunnar, M.R., 2016. Early life stress: effects on the regulation of anxiety expression in children and adolescents. *Soc. Dev.* (4), 777–793. <https://doi.org/10.1111/sode.12170>.
- Burton, C.L., Chatterjee, D., Chatterjee-Chakraborty, M., Lovic, V., Grella, S.L., Steiner, M., Fleming, A.S., 2007. Prenatal restraint stress and motherless rearing disrupts expression of plasticity markers and stress-induced corticosterone release in adult female Sprague-Dawley rats. *Brain Res.* 1158, 28–38. <https://doi.org/10.1016/j.brainres.2007.05.003>.
- Campbell, S., Marriott, M., Nahmias, C., MacQueen, G.M., 2004. Lower hippocampal volume in patients suffering from depression: ameta-analysis. *Am. J. Psychiatr.* 161, 598–607. <https://doi.org/10.1176/appi.ajp.161.4.598>.
- Cancedda, L., Putignano, E., Sale, A., Berardi, N., Maffei, L., 2004. Acceleration of visual system development by environmental enrichment. *J. Neurosci.* 24, 4840–4848.
- Cesario, S.K., Hughes, L.A., 2007. Precocious puberty: a comprehensive review of literature. *J. Obstet. Gynecol. Neonatal Nurs.* 36, 263–274.
- Chalumeau, M., Chemaitilly, W., Trivin, C., Adan, L., Breart, G., Brauner, R., 2002. Central precocious puberty in girls: an evidence-based diagnosis tree to predict central nervous system abnormalities. *Pediatrics* 109 (1), 61–67. <https://doi.org/10.1542/peds.109.1.61>.
- Champagne, F.A., Curley, J.P., 2007. Studying the epigenetic influence of maternal care in Rodents. In: Crawley, J.N. (Ed.), *What's Wrong with My Mouse? Strategies for Rodent Behavior Phenotyping*. Society for Neuroscience, San Diego, pp. 71–80.
- Champagne, F.A., Curley, J.P., 2016. Plasticity of the maternal brain across the lifespan. *New Dir. Child Adolesc. Dev.* (153), 9–21. <https://doi.org/10.1002/cad.20164>.
- Chapman, D.P., Whitfield, C.L., Felitti, V.J., Dube, S.R., Edwards, V.J., Anda, R.F., 2004. Adverse childhood experiences and the risk of depressive disorders in adulthood. *J. Affect. Disord.* 82, 217–225. <https://doi.org/10.1016/j.jad.2003.12.013>.
- Charpak, N., Tessier, R., Ruiz, J.G., Hernandez, J.T., Uriza, F., Marin, J., ... Maldonado, D., 2016. Twenty-year follow-up of kangaroo mother care versus traditional care. *Pediatrics* 139. <https://doi.org/10.1542/peds.2016-2063>.
- Chatterjee, D., Chatterjee-Chakraborty, M., Rees, S., Cauchi, J., Medeiros, C.B., Fleming, A.S., 2007. Maternal isolation alters the expression of neural proteins during development: 'stroking' stimulation reverses these effects. *Brain Res.* 1158, 11–27. <https://doi.org/10.1016/j.brainres.2007.04.069>.
- Connors, E.J., Migliore, M.M., Pillsbury, S.L., Shaik, A.N., Kentner, A.C., 2015. Environmental enrichment models a naturalistic form of maternal separation and shapes the anxiety response patterns of offspring. *Psychoneuro* 52, 153–167. <https://doi.org/10.1016/j.psyneuen.2014.10.021>.
- Cowan, C.S., Richardson, R., 2018. Early-life stress leads to sex-dependent changes in pubertal timing in rats that are reversed by a probiotic formulation. *Dev. Psychobiol.* <https://doi.org/10.1002/dev.21765>.
- Crawley, J.N., 2007. Social behavior tests for mice. In: Crawley, J.N. (Ed.), *What's Wrong with My Mouse? Strategies for Rodent Behavior Phenotyping*. Society for Neuroscience, San Diego, pp. 63–70.
- Dalle Molle, R., Portella, A., Goldani, M., Kapczynski, F., Leistner-Segal, S., Salum, G., et al., 2012. Associations between parenting behavior and anxiety in a rodent model and a clinical sample: relationship to peripheral BDNF levels. *Transl. Psychiatry* 2 (11), e195.
- Drzewiecki, C.M., Willing, J., Juraska, J.M., 2016. Synaptic number changes in the medial prefrontal cortex across adolescence in male and female rats. A role for pubertal onset. *Synapse* 70, 361–368.
- Durán-Carabali, L.E., Arceo, D.M., Odorczyk, F.K., Reichert, L., Cordeiro, J.L., Sanches, E.F., Freitas, L.D., Dalmaz, C., Pagnussat, A., Netto, C.A., 2018. Prenatal and early postnatal environmental enrichment reduce acute cell death and prevent neurodevelopment and memory impairments in rats submitted to neonatal hypoxia ischemia. *Mol. Neurobiol.* 55, 3627–3641.
- Eicher-Miller, H.A., Zhao, Y., 2018. Evidence for the age-specific relationship of food insecurity and key dietary outcomes among US children and adolescents. *Cambridge Univ. Press* 31, 98–113. <https://doi.org/10.1017/S0954422417000245>.
- Estes, M.L., McAllister, K.A., 2016. Maternal immune activation: implications for neuropsychiatric disorders. *Science* 353, 772–777. <https://doi.org/10.1126/science.aag3194>.
- Feldman-Winter, L., Goldsmith, J.P., 2016. Safe sleep and skin-to-skin care in the neonatal period for healthy term newborns. *Am. Acad. Pediatr.* 138, 1–10. <https://doi.org/10.1542/peds.2016-1889>.
- Fernandez-Teruel, A., Escorihuela, R.M., Driscoll, P., Tobena, A., Battig, K., 1991. Infantile (handling) stimulation and behavior in young roman high- and low-avoidance rats. *Physiol. Behav.* 50, 563–565. [https://doi.org/10.1016/0031-9384\(91\)90546-Z](https://doi.org/10.1016/0031-9384(91)90546-Z).
- Fisher, I.P., Irmiler, M., Meyer, C.W., Sachs, S.J., Neff, F., Hrabě de Angelis, M., Beckers, J., Tschöp, M.H., Hofmann, S.M., Ussar, S., 2018. A history of obesity leaves and

- inflammatory fingerprint in liver and adipose tissue. *Int. J. Obes.* 42, 507–517.
- Francis, D.D., Diorio, J., Plotsky, P.M., Meaney, M.J., 2002. Environmental enrichment reverses the effects of maternal separation on stress reactivity. *J. Neurosci.* 22 (18), 7840–7843. <https://doi.org/10.1523/JNEUROSCI.22-18-07840.2002>.
- Francis, H., Stevenson, R., 2012. The longer-term impacts of Western diet on human cognition and the brain. *Appetite* 63, 112–128. <https://doi.org/10.1016/j.appet.2012.12.018>.
- Gilles, E.E., Schultz, L., Baram, T.Z., 1996. Abnormal corticosterone regulation in an immature rat model of continuous chronic stress. *Pediatr. Neurol.* 15, 114–119. [https://doi.org/10.1016/0887-8994\(96\)00153-1](https://doi.org/10.1016/0887-8994(96)00153-1).
- Glantz, L.A., Gilmore, J.H., Hamer, R.M., Lieberman, J.A., Jarskog, L.F., 2007. Synaptophysin and postsynaptic density protein 95 in the human prefrontal cortex from mid-gestation into early adulthood. *Neuroscience* 149, 582–591.
- Gonzalez, A., Lovic, V., Ward, G.R., Wainwright, P.E., Fleming, A.S., 2001. Intergenerational effects of complete maternal deprivation and replacement stimulation on maternal behavior and emotionality in female rats. *Dev. Psychobiol.* 38, 11–32. [https://doi.org/10.1002/1098-2302\(2001\)38:1<11::AID-DEV2>3.0.CO;2-B](https://doi.org/10.1002/1098-2302(2001)38:1<11::AID-DEV2>3.0.CO;2-B).
- Grassi-Oliveira, R., Honeycutt, J.A., Holland, F.H., Ganguly, P., Brenhouse, H.C., 2016. Cognitive impairment effects of early life stress in adolescents can be predicted with early biomarkers: impacts of sex, experience, and cytokines. *Psychoneuroendocrinology* 71, 19–30. <https://doi.org/10.1016/j.psyneuen.2016.04.016>.
- Green, J.F., McLaughlin, K.A., Berglund, P.A., Gruber, M.J., Sampson, N.A., Zaslavsky, A.M., Kessler, R.C., 2010. Childhood adversities and adult psychopathology in the National Comorbidity Survey Replication (NCS-R) I: associations with onset of DSM-IV disorders. *Arch. Gen. Psychiatry* 67, 1–20. <https://doi.org/10.1001/archgenpsychiatry.2009.186>.
- Gundersen, C., Kreider, B., 2009. Bounding the effects of food insecurity on children's health outcomes. *J. Health Econ.* 28, 971–983. <https://doi.org/10.1016/j.jhealeco.2009.06.012>.
- Hayes, L.D., 2000. To nest communally or not to nest communally: a review of rodent communal nesting and nursing. *Anim. Behav.* 59, 677–688.
- Heim, C., Newport, D.J., Mletzko, T., Miller, A.H., Nemeroff, C.B., 2008. The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology* 33, 693–710. <https://doi.org/10.1016/j.psyneuen.2008.03.008>.
- Heun-Johnson, H., Levitt, P., 2016. Early-life stress paradigm transiently alters maternal behaviors, dam-pup interactions, and offspring vocalizations in mice. *Front. Behav. Neurosci.* 10, 1–18. <https://doi.org/10.3389/fnbeh.2016.00142>.
- Hurst, J.L., Barnard, C.J., Nevison, C.M., West, C.D., 1997. Housing and welfare in laboratory rats: welfare implications of isolation and social contact among caged males. *Anim. Welf.* 6, 329–347.
- Ivy, A.S., Brunson, K.L., Sandman, C., Baram, T.Z., 2008. Dysfunctional nurturing behavior in rat dams with limited access to nesting material: a clinically relevant model for early-life stress. *Neuroscience* 154, 1132–1142. <https://doi.org/10.1016/j.neuroscience.2008.04.019>.
- Ivy, A.S., Rex, C.S., Chen, Y., Dubé, C., Maras, P.M., Grigoriadis, D.E., et al., 2010. Hippocampal dysfunction and cognitive impairments provoked by chronic early-life stress involve excessive activation of CRH receptors. *J. Neurosci.* 30 (39), 13005–13015.
- Jacobson, L., Sapolsky, R., 1991. The role of the Hippocampus in feedback regulation of the hypothalamic-pituitary-adrenocortical Axis. *Endocr. Rev.* 12, 118–134. <https://doi.org/10.1210/edrv-12-2-118>.
- Janssen, H., Ada, L., Bernhardt, J., McElduff, P., Pollack, M., Nilsson, M., Spratt, N.J., 2014. An enriched environment increases activity in stroke patients undergoing rehabilitation in a mixed-rehabilitation unit: a pilot non-randomized controlled trial. *Disabil. Rehabil.* 36, 255–262.
- Kalyan-Masih, P., Vega-Torres, J.D., Miles, C., Haddad, E., Rainsbury, S., Baghchechi, M., ... Figueroa, J., 2016. Western high-fat diet consumption during adolescence increases susceptibility to traumatic stress while selectively disrupting hippocampal and ventricular volumes. *Cogn. Behav. Ther.* 3, 1–24. <https://doi.org/10.1523/ENEURO.0125-16.2016>.
- Kaplowitz, P.B., Hoffman, R.P., 2018, July 05. Precocious puberty: background, pathophysiology, epidemiology. from. <https://emedicine.medscape.com/article/924002-overview#a5>, Accessed date: 10 August 2018.
- Kauffman, A.S., Clifton, D.K., Steiner, R.A., 2007. Emerging ideas about kisspeptin-GPR54 signaling in the neuroendocrine regulation of reproduction. *Trends Neurosci.* 30 (10), 504–511. <https://doi.org/10.1016/j.tins.2007.08.001>.
- Kelly, Y., Zilanawala, A., Sacker, A., Hiatt, R., Viner, R., 2017. Early puberty in 11-year-old girls: millennium cohort study findings. *Arch. Dis. Child.* 102, 232–237.
- Kenny, S.L., Wright, L.D., Green, A., Mashoodh, R., Perrot, T.S., 2014. Expression of maternal behavior and activation of the bed nucleus of the stria terminalis during predatory threat exposure: modulatory effects of transport stress. *Physiol. Behav.* 123, 148–155. <https://doi.org/10.1016/j.physbeh.2013.08.024>.
- Kentner, A.C., Bilbo, A.D., Brown, A.S., Hsiao, E.Y., McAllister, A.K., Meyer, U., Pearce, B.D., Pletnikov, M.V., Yolken, R.H., Bauman, M.D., 2018. Maternal immune activation: reporting guidelines to improve the rigor, reproducibility, and transparency of the model. *Neuropsychopharmacology*. <https://doi.org/10.1038/s41386-018-0185-7>.
- Kentner, A.C., Cryan, J.F., Brummelte, S., 2018. Resilience priming: translational models for understanding resiliency and adaptation to early-life adversity. *Dev. Psychobiol.* <https://doi.org/10.1002/dev.21775>. (in press).
- Kentner, A.C., Khoury, A., Queiroz, E.L., MacRae, M., 2016. Environmental enrichment rescues the effects of early life inflammation on markers of synaptic transmission and plasticity. *Brain Behav. Immun.* 57, 151–160. <https://doi.org/10.1016/j.bbi.2016.03.013>.
- Kentner, A.C., Pittman, Q.J., 2010. Minireview: early-life programming by inflammation of the neuroendocrine system. *Endocrinology* 151 (10), 4602–4606. <https://doi.org/10.1210/en.2010-0583>.
- Kentner, A.C., Scalia, S., Shin, J., Migliore, M.M., Rondon-Ortiz, A.N., 2018. Targeted sensory enrichment interventions protect against behavioral and neuroendocrine consequences of early life stress. *Psychoneuroendocrinology* 98, 74–85.
- Kirshner, Z.Z., Gibbs, R.B., 2018. Use of the REVERT® total protein stain as a loading control demonstrates significant benefits over the use of housekeeping proteins when analyzing brain homogenates by Western blot: an analysis of samples representing different gonadal hormone states. *Mol. Cell. Endocrinol.* 473, 156–165.
- Kumsta, R., Schlotz, W., Golm, D., Moser, D., Kennedy, M., Knights, N., ... Sonuga-Barke, E., 2017. HPA axis dysregulation in adult adoptees twenty years after severe institutional deprivation in childhood. *Psychoneuroendocrinology* 88, 196–202. <https://doi.org/10.1016/j.psyneuen.2017.09.021>.
- Leinonen, J.A., Solantaus, T.S., Punamaki, R.-L., 2003. Social support and the quality of parenting under economic pressure and workload in Finland: the role of family structure and parental gender. *J. Fam. Psychol.* 17, 409–418. <https://doi.org/10.1037/0893-3200.17.3.409>.
- Leussis, M.P., Freund, N., Brenhouse, H.C., Thompson, B.S., Andersen, S.L., 2012. Depressive-like behavior in adolescents after maternal separation: sex differences, controllability, and GABA. *Dev. Neurosci.* 34, 210–217. <https://doi.org/10.1159/000339162>.
- Levine, S., 1994. The ontogeny of the hypothalamic-pituitary-adrenal axis: the influence of maternal factors. *Ann. N. Y. Acad. Sci.* 30, 275–288.
- Levine, S., 2001. Primary social relationships influence the development of the hypothalamic-pituitary-adrenal axis in rats. *Physiol. Behav.* 73, 255–260. [https://doi.org/10.1016/S0031-9384\(01\)00496-6](https://doi.org/10.1016/S0031-9384(01)00496-6).
- Li, L., Denholm, R., Power, C., 2014. Child maltreatment and household dysfunction: associations with pubertal development in a British birth cohort. *Int. J. Epidemiol.* 43 (4), 1163–1173. <https://doi.org/10.1093/ije/dyu071>.
- Liu, D., Diorio, J., Tannenbaum, B., Caldji, C., Francis, D., Freedman, A., ... Meaney, M.J., 1997. Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. *Science* 227, 1659–1662. <https://doi.org/10.1126/science.277.5332.1659>.
- Lomanowska, A.M., Melo, A.I., 2016. Deconstructing the function of maternal stimulation in offspring development: insights from the artificial rearing model in rats. *Horm. Behav.* 77, 224–236. <https://doi.org/10.1016/j.yhbeh.2015.05.017>.
- Luebke, A.M., Mancini, K.J., Kiel, E.J., Spangler, B.R., Semlak, J.L., Fussner, L.M., 2016. Dimensionality of helicopter parenting and relations to emotional, decision-making, and academic functioning in emerging adults. *Assessment* 1–17. <https://doi.org/10.1177/1073191116665907>.
- Machado, T.D., Dalle Molle, R., Laureano, D.P., Portella, A.K., Werlang, I.C.R., da Silva Benetti, C., Noschang, C., Silveira, P.P., 2013. Early life stress is associated with anxiety, increased stress responsivity and preference for 'comfort foods' in adult female rats. *Stress* 16, 549–556.
- MacRae, M., Macrina, T., Khoury, A., Migliore, M.M., Kentner, A.C., 2015. Tracing the trajectory of behavioral impairments and oxidative stress in an animal model of neonatal inflammation. *Neuroscience* 298, 455–466.
- Maniatis, J., Antoniadis, C.P., Le, V., Morris, M.J., 2016. A diet high in fat and sugar reverses anxiety-like behavior induced by limited nesting in male rats: impacts on hippocampal markers. *Psychoneuroendocrinology* 68, 202–209.
- Maniatis, J., Antoniadis, C.P., Morris, M.J., 2015. Sugar consumption produces effects similar to early life stress exposure on hippocampal markers of neurogenesis and stress response. *Front. Mol. Neurosci.* 19, 1–10. <https://doi.org/10.3389/fnmol.2015.00086>.
- McLaughlin, R.J., Verlez, S., Gray, J.M., Hill, M.N., Walker, C.-D., 2016. Inhibition of anandamide hydrolysis dampens the neuroendocrine response to stress in neonatal rats subjected to suboptimal rearing conditions. *J. Biol. Stress* 19, 114–124. <https://doi.org/10.3109/10253890.2015.1117448>.
- Moriceau, S., Shionoya, K., Jakubs, K., Sullivan, R.M., 2009. Early-life stress disrupts attachment learning: the role of amygdala corticosterone, locus ceruleus corticotropin releasing hormone, and olfactory bulb norepinephrine. *J. Neurosci.* 29 (50), 15745–15755.
- National Research Council, 2011. Guide for the Care and Use of Laboratory Animals, 8th edn. National Academy of Sciences, Washington, DC.
- Navarro, V.M., Fernández-Fernández, R., Castellano, J.M., Roa, J., Mayen, A., Barreiro, M.L., Gaytan, F., Aguilar, E., Pinilla, L., Dieguez, C., Tena-Sempere, M., 2004. Advanced vaginal opening and precocious activation of the reproductive axis by KiSS-1 peptide: the endogenous ligand of GPR54. *J. Physiol.* 561, 379–386.
- Nelson III, C.A., Zeannah, C.H., Fox, N.A., Marshall, P.J., Smyke, A.T., Guthrie, D., 2007. Cognitive recovery in socially deprived young children: the Bucharest early intervention project. *Science* 318, 1937–1940. <https://doi.org/10.1126/science.1143921>.
- Payne, S.H., 2015. The utility of protein and mRNA correlation. *Trends Biochem. Sci.* 40, 1–3.
- Pedrin Schuch, C., Diaz, R., Deckmann, I., Roja, J., Deniz, B., Pereira, L., 2016. Early environmental enrichment affects neurobehavioral development and prevents brain damage in rats submitted to neonatal hypoxia-ischemia. *Neurosci. Lett.* 617, 101–107. <https://doi.org/10.1016/j.neulet.2016.02.015>.
- Perry, R.E., Finegood, E.D., Braren, S.H., Dejesou, M.L., Putrino, D.F., Wilson, D.A., et al., Blair, C., 2018. Developing a neurobehavioral animal model of adversity: drawing cross-species connections between environments of scarcity-poverty, parenting quality, and infant outcome. *Dev. Psychopathol.* 2, 1–20. <https://doi.org/10.1017/S095457941800007X>.
- Pervanidou, P., Chrousos, G.P., 2012. Metabolic consequences of stress during childhood and adolescence. *Metabolism* 61, 611–619. <https://doi.org/10.1016/j.metabol.2011.03.013>.

- 10.005.
- Purcell, R.H., Sun, B., Pass, L.L., Power, M.L., Moran, T.H., Tamashiro, K.L.K., 2011. Maternal stress and high-fat diet effect on maternal behavior, milk composition, and pup ingestive behavior. *Physiol. Behav.* 104, 474–479.
- Raineki, C., Cortés, M.R., Belhoue, L., Sullivan, R.M., 2012. Effects of early life abuse differ across development: infant social behavior deficits are followed by adolescent depressive-like behaviors mediated by the amygdala. *J. Neurosci.* 32, 7758–7765. <https://doi.org/10.1523/JNEUROSCI.5843-11.2012>.
- Raineki, C., Moriceau, S., Sullivan, R.M., 2010. Developing a neurobehavioral animal model of infant attachment to an abusive caregiver. *Biol. Psychiatry* 67, 1137–1145. <https://doi.org/10.1016/j.biopsych.2009.12.019>.
- Raun, K., von Voss, P., Gotfredsen, C.F., Golozubova, V., Rolin, B., Knudsen, L.B., 2007. Liraglutide, a long-acting glucagon-like peptide-1 analog reduces body weight and food intake in obese candy-fed rats, whereas a dipeptidyl peptidase-IV inhibitor, Vildagliptin, does not. *Diabetes* 56, 8–15.
- Rice, C.J., Sandman, C.A., Lenjavi, M.R., Baram, T.Z., 2008. A novel mouse model for acute and long-lasting consequence of early life stress. *Endocrinology* 149, 4892–4900. <https://doi.org/10.1210/en.2008-0633>.
- Rincel, M., Lépinay, A.L., Delage, P., Fioramonti, J., Théodorou, V.S., Layé, S., Darnaudéry, M., 2016. Maternal high-fat diet prevents developmental programming by early-life stress. *Transl. Psychiatry* 29, 1–9. <https://doi.org/10.1038/tp.2016.235>.
- Rincel, M., Lépinay, A.L., Janthakhin, Y., Soudain, G., Yvon, S., Da Silva, S., Joffre, C., Aubert, A., Séré, A., Layé, S., Theodorou, V., Ferreira, G., Darnaudéry, A., 2018. Maternal high fat diet and early life stress differentially modulate spine density and dendritic morphology in the medial prefrontal cortex of juvenile and adult rats. *Brain Struct. Funct.* 223, 883–895.
- Rincón-Cortés, M., Sullivan, R., 2016. Emergence of social behavior deficit, blunted corticolimbic activity and adult depression-like behavior in a rodent model of maternal maltreatment. *Transl. Psychiatry* 6 (10), e930.
- Rosenfeld, A., Weller, A., 2012. Behavioral effects of environmental enrichment during gestation in WKY and Wistar rats. *Behav. Brain Res.* 233, 245–255.
- Roth, T.L., Lubin, F.D., Funk, A.J., Sweatt, J.D., 2009. Lasting epigenetic influence of early-life adversity on the BDNF gene. *Biol. Psychiatry* 65 (9), 760–769.
- Sacks, V., Murphey, D., 2018. The prevalence of adverse childhood experiences, nationally, by state, and by race or ethnicity. In: *Child Trends*.
- Sale, A., Putignano, E., Cancedda, L., Landi, S., Cirulli, F., Berardi, N., Maffei, L., 2004. Enriched environment and acceleration of visual system development. *Neuropharmacology* 47, 649–660.
- Sanjeevi, N., Freeland-Graves, J., Hersh, M., 2018. Food insecurity, diet quality and body mass index of women participating in the supplemental nutrition assistance program: the role of intrapersonal, home environment, community and social factor. *Appetite* 125, 109–117. <https://doi.org/10.1016/j.appet.2018.01.036>.
- Schneider, T., Turczak, J., Przewlocki, R., 2006. Environmental enrichment reverses behavioral alterations in rats prenatally exposed to valproic acid: issues for a therapeutic approach in autism. *Neuropsychopharmacology* 31, 36–46. <https://doi.org/10.1038/sj.npp.1300767>.
- Schultz, L.A., Lore, R.K., 1993. Communal reproductive success in rats (*Rattus norvegicus*): effects of group composition and prior social experience. *J. Comp. Psychol.* 107 (2), 216–222.
- Sheridan, M.A., Fox, N.A., Zeanah, C.H., McLaughlin, K.A., Nelson III, C.A., 2012. Variation in neural development as a result of exposure to institutionalization early in childhood. *Proc. Natl. Acad. Sci. U. S. A.* 109, 12927–12932. <https://doi.org/10.1073/pnas.1200041109>.
- Siersbaek, M., Varticovski, L., Yang, S., Baek, S., Nielsen, R., Mandrup, S., Hager, G., Chung, J.H., Grøntved, L., 2017. High fat diet-induced changes of mouse hepatic transcription and enhancer activity can be reversed by subsequent weight loss. *Sci. Rep.* 7, 40220.
- Skorupkaite, K., George, J.T., Anderson, R.A., 2014. The kisspeptin-GnRH pathway in human reproductive health and disease. *Hum. Reprod. Update* 20 (4), 485–500. <https://doi.org/10.1093/humupd/dmu009>.
- Sominsky, L., Ziko, I., Soch, A., Smith, J.T., Spencer, S.J., 2016. Neonatal overfeeding induces early decline of the ovarian reserve: implications for the role of leptin. *Mol. Cell. Endocrinol.* 431, 25–35.
- Sorge, R.E., Mapplebeck, J.C.S., Rosen, S., Beggs, S., Taves, S., Alexander, J.K., Martin, L.J., Austin, J.-A., Sotocinal, S.G., Chen, D., Yang, M., ... Mogil, J.S., 2015. Different immune cells mediate mechanical pain hypersensitivity in male and female mice. *Nat. Neurosci.* 18, 1081–1083.
- Spencer, S.J., Korosi, A., Layé, S., Shukitt-Hale, B., Barrientos, R., 2017. Food for thought: how nutrition impacts cognition and emotion. *NPJ Sci. Food* 7, 1–8. <https://doi.org/10.1038/s41538-017-0008-y>.
- Spencer, S.J., Martin, S., Mouihate, A., Pittman, Q.J., 2006. Early-life immune challenge: defining a critical window for effects on adult responses to immune challenge. *Neuropsychopharmacology* 31, 1910–1918.
- Stormshak, E.A., Bierman, K.L., McMahon, R.J., Lengua, L.J., 2010. Parenting practices and child disruptive behavior problems in elementary school. *J. Clin. Child Psychol.* 29, 17–29. https://doi.org/10.1207/S15374424jccp2901_3.
- Strauss, J.F., Barbieri, R.L. (Eds.), 2009. *Yen and Jaffe's Reproductive Endocrinology: Physiology, Pathophysiology, and Clinical Management*, 6th edition. Elsevier, China.
- Sullivan, R.M., Holman, P.J., 2010. Transitions in sensitive period attachment learning in infancy: the role of corticosterone. *Neurosci. Biobehav. Rev.* 34, 835–844. <https://doi.org/10.1016/j.neubiorev.2009.11.010>.
- Tessier, R., Cristo, M.B., Velez, S., Giron, M., Nadeau, L., de Calume, F., ... Charpak, N., 2003. Kangaroo mother care: a method for protecting high-risk low-birth-weight and premature infants against developmental delay. *Infant Behav. Dev.* 26, 384–397. [https://doi.org/10.1016/S0163-6383\(03\)00037-7](https://doi.org/10.1016/S0163-6383(03)00037-7).
- Tilburg, M.A., Runyan, D.K., Zolotor, A.J., Graham, A.J., Dubowitz, J.C., Litrownik, A.J., 2010. Unexplained gastrointestinal symptoms after abuse in a prospective study of children at risk for abuse and neglect. *Ann. Fam. Med.* 8, 134–140. <https://doi.org/10.1370/afm.1053>.
- Vahl, T.P., Ulrich-Lai, Y.M., Ostrander, M.M., Dolgas, C.M., Elfers, E.E., Seeley, R.J., D'Alessio, D.A., Herman, J.P., 2005. Comparative analysis of ACTH and corticosterone sampling methods in rats. *Am. J. Physiol. Endocrinol. Metab.* 289, E823–E828.
- Virdis, R., Street, M.E., Zampolli, M., Radetti, G., Pezzini, B., Benelli, M., Ghizzoni, L., Volta, C., 1998. Precocious puberty in girls adopted from developing countries. *Arch. Dis. Child.* 78, 152–154. <https://doi.org/10.1136/adc.78.2.152>.
- Vivinetto, A.L., Suárez, M.M., Rivarola, M.A., 2013. Neurobiological effects of neonatal maternal separation and post-weaning environmental enrichment. *Behav. Brain Res.* 240, 110–118. <https://doi.org/10.1016/j.bbr.2012.11.014>.
- Vogel, C., Marcotte, E.M., 2012. Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nat. Rev. Genet.* 227–232.
- Walker, C.-D., Bath, K.G., Joels, M., Korosi, A., Larauche, M., Lucassen, P.J., ... Baram, T.Z., 2017. Chronic early life stress induced by limited bedding and nesting (LBN) material in rodents: critical considerations of methodology, outcomes and translational potential. *Int. J. Biol. Stress* 20, 421–448. <https://doi.org/10.1080/10253890.2017.1343296>.
- Welberg, L., Thiruvikraman, K.V., Plotsky, P.M., 2006. Combined pre- and postnatal environmental enrichment programs the HPA axis differentially in male and female rats. *Psychoneuroendocrinology* 31 (5), 553–564. <https://doi.org/10.1016/j.psyneuen.2005.11.011>.
- White, J.H., Bartley, E., Janssen, H., Jordan, L.A., Spratt, N., 2015. Exploring stroke survivor experience of participation in an enriched environment: a qualitative study. *Disabil. Rehabil.* 37, 593–600. <https://doi.org/10.3109/09638288.2014.935876>.
- Woo, C.C., Donnelly, J.H., Steinberg-Epstein, R., Leon, M., 2015. Environmental enrichment as a therapy for autism: a clinical trial replication and extension. *Behav. Neurosci.* 129 (4), 412–422. <https://doi.org/10.1037/bne0000068>.
- Woo, C.C., Leon, M., 2013. Environmental enrichment as an effective treatment for autism: a randomized controlled trial. *Behav. Neurosci.* 127 (4), 487–497. <https://doi.org/10.1037/a0033010>.
- Yam, K.Y., Nannick, E.F., Abbink, M.R., la Fleur, S.E., Schipper, L., van den Beukel, J.C., ... Korosi, A., 2016. Exposure to chronic early-life stress lastingly alters the adipose tissue, the leptin system, and changes the vulnerability to western-style diet later in life in mice. *Psychoneuroendocrinology* 77, 186–195. <https://doi.org/10.1016/j.psyneuen.2016.12.012>.
- Zieba, J., Uddin, G.M., Youngson, N.A., Karl, T., Morris, M.J., 2018. Long-term behavioural effects of maternal obesity in C57BL/6 mice. *Physiol. Behav.* <https://doi.org/10.1016/j.physbeh.2018.11.004>. in press.