



## Note

## A novel microbiological medium for the growth of periodontitis associated pathogens

Niall A. Hickey, Kathryn A. Whitehead, Liliana Shalamanova, Jonathan A. Butler, Rebecca L. Taylor\*

Department of Life Sciences, Faculty of Science and Engineering, Manchester Metropolitan University, Chester Street, Manchester M1 5GD, UK



## ARTICLE INFO

## Keywords:

Chronic periodontitis  
Growth medium  
Oral microbiology  
Cell culture

## ABSTRACT

A novel microbiological medium designed to be more representative of gingival crevicular fluid. Chosen representative periodontal microorganisms showed good growth with minimal effect on human cell viability. This will enable more comparisons between different periodontitis associated organisms and their potential role in host health and systemic disease.

Chronic Periodontitis (CP), an infection of the periodontium, leading to the formation of periodontal pockets and tooth loss if untreated (Hajishengallis, 2015). It is the most common infectious inflammatory disease worldwide affecting humans (Singh et al., 2015) with 45% of English adults having at least one periodontal pocket (Heidari et al., 2015). CP is of particular interest as it can be detrimental to the health of the patient and a risk factor for myriad diseases such as Alzheimer's, (Stein et al., 2012) cardiovascular disease (Dietrich et al., 2008) and diabetes mellitus (Engelbreton et al., 2013). Subgingival bacteria are the main aetiological agents behind the inflammatory processes seen in CP and induce dysregulation of the normal oral microbiota (Wahid et al., 2013) and an increased flow of gingival crevicular fluid (GCF), an inflammatory exudate that bathes the gingival crevice and favours the growth of anaerobic periodontal microorganisms (Winning et al., 2015). A key contributor to CP is *Porphyromonas gingivalis* (Wade, 2013), however, it appears that no singular group of organism is responsible for causing CP and it is likely that polymicrobial synergy and dysbiosis of multiple periodontitis associated pathogens play a role in modulation of the disease (Hajishengallis and Lamont, 2012). Many of these periodontitis associated pathogens are difficult to culture *in vitro*, which is partly compounded by the lack of available broad host growth media. Periodontal pathogens are usually fastidious anaerobes and rely on nutrients secreted by other bacteria during synergism along with specific microbial and non-microbial nutrient sources (Sharma, 2010). The aim of this study was to develop a growth medium that can facilitate the culture of a wide selection of periodontitis associated microorganisms and enable further study of potential systemic disease links *via* tissue culture based

investigations. With the growth requirements of many of these organisms being specifically evolved to the unique environment of the gingival crevice, emulating components of the GCF can allow a developed medium to be more representative of this environmental interface and allow more representative growth of periodontal microorganisms.

Bacterial strains were maintained in growth media as recommended by the manufacturer. *Actinomyces israelii* (NCTC 12972), *Campylobacter showae* (NCTC 12843), *Fusobacterium nucleatum* subsp. *Fusifforme* (11326) and *Prevotella intermedia* (NCTC 13070) were cultivated in anaerobic conditions (80% N<sub>2</sub>–10% H<sub>2</sub>–10% CO<sub>2</sub>) in a Baker-Ruskin Concept Plus anaerobic incubator (Ruskin, UK) at 37 °C for 48 h. *P. gingivalis* (NCTC 11834) was cultivated in an anaerobic incubator for 72 h. *Rothia dentocariosa* (NCTC 10917), *Streptococcus constellatus* (NCTC 11325) and *Streptococcus sanguinis* (NCTC 7863) were cultivated in carbon rich conditions (5% CO<sub>2</sub>) in a CO<sub>2</sub> incubator (LEEC, UK) at 37 °C for 48 h.

The base medium utilised was Basal Medium Mucin (BMM) as it has successfully been shown to emulate saliva for the growth of dental plaque biofilms in an 'artificial mouth' system (Wong and Sissions, 2001). The constituents of BMM contained: 2.5 g/L partially purified pig gastric mucin, 10 g/L proteose peptone, 5 g/L trypticase peptone, 5 g/L yeast extract, 2.5 g/L KCl, 1 mMol/L urea, 1 mMol/L arginine, 1 mg/L of haemin and 1 mg/L of menadione, all purchased from Sigma Aldrich, UK.

The new growth medium, Artificial Gingival Crevicular Fluid – Periodontal (AGCF-P) was composed of BMM with multiple modifications. From BMM, 1 mg/mL of haemin and menadione were omitted and 5% horse blood (TCS Biosciences) was added once cooled to 50 °C

\* Corresponding author.


E-mail address: [r.taylor@mmu.ac.uk](mailto:r.taylor@mmu.ac.uk) (R.L. Taylor).

<https://doi.org/10.1016/j.mimet.2019.105647>

Received 13 May 2019; Received in revised form 23 May 2019; Accepted 29 May 2019

Available online 30 May 2019

0167-7012/ © 2019 Elsevier B.V. All rights reserved.



**Fig. 1.** Growth curves of periodontitis associated pathogens cultured in AGCF-P broth. A) *Actinomyces israelii* B) *Fusobacterium nucleatum* C) *Porphyromonas gingivalis* D) *Parvimonas micra* E) *Streptococcus constellatus* F) *Streptococcus sanguinis* ( $n = 3$ ).

after sterilisation by autoclave. This was designed to emulate the inorganic and organic constituents of GCF which are notably similar to plasma, as infiltration and degradation of red blood cells occurring during CP (Ferguson, 2006). The pH of the medium was adjusted to reflect the actual pH of GCF during CP ( $7.96 \pm 0.1$ ) (Bickel et al., 1985). Buffering of pH was performed using the phosphate buffer system of Potassium Phosphate monobasic anhydrous and Sodium Phosphate dibasic heptahydrate. Cysteine was added as a reducing agent to further mimic the reducing capacity and support the growth of anaerobic organisms. The full composition of AGCF-P was: 10 g/L proteose peptone, 5 g/L trypticase peptone, 5 g/L yeast extract, 2.5 g/L KCl, 2.5 g/L partially purified pig gastric mucin, 0.5 g/L L-cysteine, 0.17 g/L L-arginine, 0.18 g/L L-urea, 1.68 g/L Potassium Phosphate monobasic anhydrous, 41.31 g/L Sodium Phosphate dibasic heptahydrate and 5% horse blood. If creating solid media 12 g/L of agar no 3 was added. All reagents were procured from Sigma Aldrich, UK with the exception of trypticase peptone (Scientific Laboratory Supplies, UK) and horse blood.

The growth of the bacterial strains on AGCF-P agar was determined by streaking a single isolated colony onto an AGCF-P agar plate. This plate was sub-cultured again to ensure continuity and photographed (not shown). Growth of bacterial strains in liquid culture was tested by inoculating a single colony into 5 mL of AGCF-P broth and grown for

the NCTC recommended duration; growth was demonstrated by inoculating an AGCF-P agar plate and further incubation. For growth curves, three overnight cultures of the chosen microorganism were adjusted to an absorbance of  $1.0 \pm 0.05$  at  $540\text{nm}$  obtained using a spectrophotometer (Jenway, UK). One millilitre per 100 mL of culture was inoculated into three sterile culture flasks with liquid AGCF-P growth media equilibrated overnight. Growth was assessed at 0, 2, 4, 6, 24, and 48 h or daily for 5 day for *P. gingivalis* and at each time point colony forming units per mL (CFU/mL) were quantified using the Miles-Misra droplet method and viability was determined (Miles et al., 1938).

Human Gingival Fibroblast (HGF-1) primary cells (LGC standards, UK) were cultured in Dulbecco Modified Eagle Medium (DMEM) with Glutamine and Glucose (Lonza, UK), supplemented with 10% foetal calf serum (Lonza, UK), 50  $\mu\text{g/mL}$  streptomycin (Lonza, UK) and 50  $\mu\text{g/mL}$  penicillin (Lonza, UK). Immortalised Human Kidney Proximal Tubule cells (HK-2) (ATCC, UK) were cultured in 50:50 glucose free DMEM (Invitrogen, UK) and Hams F-12 medium (Lonza, UK) supplemented with 10% foetal calf serum (Lonza, UK), 50  $\mu\text{g/mL}$  streptomycin (Lonza, UK), 50  $\mu\text{g/mL}$  penicillin (Lonza, UK) and 2.5 mMol glutamine (Lonza, UK). Cells were growth-arrested in serum free cell culture media, in the absence of foetal calf serum.

To assess the effect of AGCF-P bacterial growth medium on mammalian cell viability, the AGCF-P medium was incubated for 5 days in




Fig. 2. Cell viability of HGF-1 cells after 24 h (A) or 48 h (B) of incubation with 1:20 dilutions of AGCF-P supernatants collected over 5 days. Viability is presented as percentage change from the control (C: cells exposed to serum-free cell culture medium) and error bars represent SEM ( $n = 3$ ).




Fig. 3. Cell viability of HK-2 cells after 24 h (A) or 48 h (B) of incubation with 1:20 dilutions of AGCF-P supernatants collected over 5 days. Viability is presented as percentage change from the control (C: cells exposed to serum-free cell culture medium) and the error bars represent SEM ( $n = 3$ ).

anaerobic conditions without bacterial inoculation. At 0, 1, 2, 3, 4 and 5 days aliquots of the AGCF-P medium were plated out to check for bacterial contamination. Viability testing was conducted as detailed in (Slate et al., 2019), with a two-way ANOVA with multiple comparisons used to determine significant differences between the cell viability control and different time points of AGCF-P.

The growth of periodontal associated pathogens was assessed in liquid and solid AGCF-P media by inoculation or streaking with a single colony of the chosen organisms. *A. israelii*, *C. showae*, *F. nucleatum* subsp. *Fusifforme*, *P. intermedia*, *P. gingivalis*, *R. dentocariosa*, *S. constellatus* and *S. sanguinis* all showed good growth in liquid and solid cultures even with multiple subcultures (data not shown). Growth curves of representative examples of periodontal pathogens were conducted over 2 days in AGCF-P broth or 5 days for *P. gingivalis*, to demonstrate growth (Fig. 1) with inoculation achieved by adjusting absorbance to an optical density of 1.0 at 600<sup>nm</sup> using spectroscopy, phosphate buffered saline and Suba Seal rubber septas (Sigma-Aldrich, UK). All organisms reached a good level of growth after 24 h, averaging at 10<sup>7</sup> CFU/mL indicating that the media was sufficiently nutritious to support the representative selection of periodontal microorganisms.

The effect of the AGCF-P medium on the cell viability of HGF-1 (Fig. 2), and HK-2 (Fig. 3) was assessed by exposing the cells to 1:20 dilutions of AGCF-P supernatants. These supernatants were collected after incubation for a 5-day period in anaerobic conditions, in order to account for abiotic degradation of the medium under bacterial culture conditions. Although the cell viability of HGF-1 was slightly increased (Fig. 2), there was no significant effect of AGCF-P supernatants on the viability of these cells or HK-2 cells after 24 h (Fig. 2A and 3A) or 48 h (Fig. 2B and 3B) of incubation, as compared to control cells exposed to serum-free cell culture medium only. The mild positive effect of AGCF-P

on the viability of primary gingival fibroblasts can be attributed to an increased cell proliferation due to nutrients provided by AGCF-P.

Mimicking the chemical components of the physiological environment is a successful strategy that has been used to cultivate particularly fastidious or uncultivable organisms and also can influence the bacterial products present (Vartoukian et al., 2010). Emulating some of the elements of the GCF should allow for more representative growth of disease-related organisms as alternative nutrient sources available may stimulate regulatory changes in gene expression profiles leading to the production of environment specific bacterial products, such as metabolites and secreted proteins (Balleza et al., 2009). AGCF-P has been shown to cultivate a selection of periodontal organisms and could be used to test their secreted products in human cells without the growth media confounding the results, facilitating the development of *in vitro* and *ex vivo* models of CP. The production of more biologically relevant conditions furthers research into the pathogenesis of CP, enabling wide-ranging comparisons between organisms to identify key microbial products and their effect on the human host.

#### Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

#### Conflict of interests

The authors report no conflict of interest.

## References

- Balleza, E., López-Bojorquez, L.N., Martínez-Antonio, A., Resendis-Antonio, O., Lozada-Chávez, I., Balderas-Martínez, Y.I., et al., 2009. Regulation by transcription factors in bacteria: beyond description. *FEMS Microbiol. Rev.* 33 (1), 133–151.
- Bickel, M., Munoz, J.L., Giovannini, P., 1985. Acid-base properties of human gingival crevicular fluid. *J. Dent. Res.* 64 (10), 1218–1220.
- Dietrich, T., Jimenez, M., Krall Kaye, E.A., Vokonas, P.S., Garcia, R.L., 2008. Age-dependent associations between chronic periodontitis/Edentulism and risk of coronary heart disease. *Circulation.* 117 (13), 1668.
- Engelbreton, S.P., Hyman, L.G., Michalowicz, B.S., et al., 2013. The effect of nonsurgical periodontal therapy on hemoglobin a1c levels in persons with type 2 diabetes and chronic periodontitis: a randomized clinical trial. *JAMA.* 310 (23), 2523–2532.
- Ferguson, D.B., 2006. Oral Bioscience. Authors Online Limited.
- Hajishengallis, G., 2015. Periodontitis: from microbial immune subversion to systemic inflammation. *Nat. Rev. Immunol.* 15 (1), 30–44.
- Hajishengallis, G., Lamont, R.J., 2012. Beyond the red complex and into more complexity: the polymicrobial synergy and dysbiosis (PSD) model of periodontal disease etiology. *Mol Oral Microbiol* 27 (6), 409–419.
- Heidari, E., Banerjee, A., Newton, J.T., 2015. Oral health status of non-phobic and dentally phobic individuals; a secondary analysis of the 2009 Adult Dental Health Survey. *Br. Dent. J.* 219 (9), E9.
- Miles, A.A., Misra, S.S., Irwin, J.O., 1938. The estimation of the bactericidal power of the blood. *Epidemiol. Infect.* 38 (6), 732–749.
- Sharma, A., 2010. Virulence mechanisms of *Tannerella forsythia*. *Periodontol* 2000 54 (1), 106–116.
- Singh, S.K., Harding, A., Poole, S., Kesavalu, L., Crean, S., 2015. *Porphyromonas gingivalis* periodontal infection and its putative links with Alzheimer's disease. *Mediat. Inflamm.* 2015, 137357.
- Slate, A.J., Shalamanova, L., Akhidime, I.D., Whitehead, K.A., 2019. Rhenium and yttrium ions as antimicrobial agents against multidrug resistant *Klebsiella pneumoniae* and *Acinetobacter baumannii* biofilms. *Letts. Appl. Microbiol.*
- Stein, P.S., Steffen, M.J., Smith, C., Jicha, G., Ebersole, J.L., Abner, E., et al., 2012. Serum antibodies to periodontal pathogens are a risk factor for Alzheimer's disease. *Alzheimers Dement.* 8 (3), 196–203.
- Vartoukian, S.R., Palmer, R.M., Wade, W.G., 2010. Strategies for culture of 'unculturable' bacteria. *FEMS Microbiol. Lett.* 309 (1), 1–7.
- Wade, W.G., 2013. The oral microbiome in health and disease. *Pharmacol. Res.* 69 (1), 137–143.
- Wahid, A., Chaudhry, S., Ehsan, A., Butt, S., Ali Khan, A., 2013. Bidirectional relationship between Chronic Kidney Disease & Periodontal Disease. *Pak. J. Med. Sci.* 29 (1), 211–215.
- Winning, L., Patterson, C.C., Cullen, K.M., Stevenson, K.A., Lundy, F.T., Kee, F., et al., 2015. The association between subgingival periodontal pathogens and systemic inflammation. *J. Clin. Periodontol.* 42 (9), 799–806.
- Wong, L., Sissions, C.H., 2001. A comparison of human dental plaque microcosm biofilms grown in an undefined medium and a chemically defined artificial saliva. *Arch. Oral Biol.* 46 (6), 477–486.