



# The Role of RNA in HIV-1 Vif-Mediated Degradation of APOBEC3H

Jiayi Wang<sup>1,2,3,4</sup>, Jordan T. Becker<sup>1,2,3,4</sup>, Ke Shi<sup>1,2,3</sup>, Kate V. Lauer<sup>1,2,3,4</sup>, Daniel J. Salamango<sup>1,2,3,4</sup>, Hideki Aihara<sup>1,2,3</sup>, Nadine M. Shaban<sup>1,2,3,4</sup> and Reuben S. Harris<sup>1,2,3,4,5</sup>

**1 - Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, Minneapolis, MN 55455, USA**

**2 - Masonic Cancer Center, University of Minnesota, Minneapolis, MN 55455, USA**

**3 - Institute for Molecular Virology, University of Minnesota, Minneapolis, MN 55455, USA**

**4 - Center for Genome Engineering, University of Minnesota, Minneapolis, MN 55455, USA**

**5 - Howard Hughes Medical Institute, University of Minnesota, Minneapolis, MN 55455, USA**

**Correspondence to Nadine M. Shaban and Reuben S. Harris:** Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, Minneapolis, MN 55455, USA. Fax: +1 612 625 2163. [nmshaban@umn.edu](mailto:nmshaban@umn.edu), [rsh@umn.edu](mailto:rsh@umn.edu)

<https://doi.org/10.1016/j.jmb.2019.09.014>

**Edited by Thomas J. Smith**

## Abstract

As many as five members of the APOBEC3 family of DNA cytosine deaminases are capable of inhibiting HIV-1 replication by deaminating viral cDNA cytosines and interfering with reverse transcription. HIV-1 counteracts restriction with the virally encoded Vif protein, which forms a hybrid ubiquitin ligase complex that directly binds APOBEC3 enzymes and targets them for proteasomal degradation. APOBEC3H (A3H) is unique among family members by dimerization through cellular and viral duplex RNA species. RNA binding is required for localization of A3H to the cytoplasmic compartment, for efficient packaging into nascent HIV-1 particles and ultimately for effective virus restriction activity. Here we compared wild-type human A3H and RNA binding-defective mutants to ask whether RNA may be a factor in the functional interaction with HIV-1 Vif. We used structural modeling, immunoblotting, live cell imaging, and split green fluorescence protein (GFP) reconstitution approaches to assess the capability of HIV-1 Vif to promote the degradation of wild-type A3H in comparison to RNA binding-defective mutants. The results combined to show that RNA is not strictly required for Vif-mediated degradation of A3H, and that RNA and Vif are likely to bind this single-domain DNA cytosine deaminase on physically distinct surfaces. However, a subset of the results also indicated that the A3H degradation process may be affected by A3H protein structure, subcellular localization, and differences in the constellation of A3H interaction partners, suggesting additional factors may also influence the fate and functionality of this host-pathogen interaction.

© 2019 Elsevier Ltd. All rights reserved.

## Introduction

The human APOBEC3 family consists of seven members, namely APOBEC3A (A3A), APOBEC3B (A3B), APOBEC3C (A3C), APOBEC3D (A3D), APOBEC3F (A3F), APOBEC3G (A3G), and APOBEC3H (A3H) (reviewed by Refs. [1–3]). These enzymes have all exhibited single-stranded DNA (ssDNA) cytosine deamination activity in a variety of assays and are expressed at varying levels in the

cytoplasmic compartment of different cell types (an exception being the predominantly nuclear A3B). At least five of these deaminases, A3C/D/F/G/H, are expressed in CD4-positive T lymphocytes and capable of restricting the replication of HIV-1 through deamination-dependent and deamination-independent mechanisms. Virus restriction minimally requires nucleic acid-binding activity to enter viral particles (RNA), deaminate viral cDNA replication intermediates (DNA), and impede reverse

transcription (RNA and DNA). The virion infectivity factor, Vif, provides the primary APOBEC3 counter-defense mechanism for HIV-1 and related lentiviruses (reviewed by Refs. [1–3]). Vif heterodimerizes with the cellular transcription cofactor CBF- $\beta$  (core binding factor- $\beta$ ) and nucleates the assembly of an E3 ubiquitin ligase complex that directly binds and targets APOBEC3 proteins for polyubiquitination and proteasomal degradation [4–11]. Mutagenesis studies have shed light on some of the molecular determinants of the APOBEC3-Vif interaction. For instance, residue F39 of Vif is specifically required for interacting with A3H because changing it to valine results in a separation-of-function variant that still degrades A3G and A3F but no longer degrades A3H [12,13]. For A3H,  $\alpha$ 3 and  $\alpha$ 4 helices have been implicated in binding to Vif with D121K on  $\alpha$ 4 conferring resistance to HIV-1 Vif-mediated degradation [14–19].

RNA has multiple roles in the antiviral activities of APOBEC3 enzymes. First, RNA binding is a key determinant in cytoplasmic localization of several APOBEC3 family members [20–26]. The clearest example is human A3H where disruption of RNA-binding residues causes a simultaneous disruption of cytoplasmic localization [24,27]. Similar observations have been made for various A3G and A3F mutants, but the precise RNA-binding mechanism has yet to be determined for these related enzymes [25,28,29]. Second, RNA binding is required for the packaging of A3F, A3G, and A3H into viral particles and subsequent antiviral activity [27,30–40]. In particular, CLIP-seq (cross-linking immunoprecipitation coupled to next generation sequencing) experiments with A3F, A3G, and A3H have indicated preferential binding to viral RNA over cellular RNA, which is likely to facilitate packaging into virions [39]. Last, but not least, a proportion of the DNA deaminase-independent activities of A3F, A3G, and A3H can be attributed to the RNA-binding properties of these enzymes, exerting a steric block to DNA synthesis by the viral reverse transcriptase [27,41–48].

Multiple A3H crystal structures have combined to demonstrate an evolutionarily conserved dimerization mechanism mediated by duplex RNA [27,49,50]. Conserved structural elements in A3H create an extensive, positively charged patch that binds strongly to duplex RNA. For human A3H, the RNA-binding region is comprised of amino acids in loop 1 (L1 R18), loop 7 (L7 H114 and W115), and  $\alpha$ -helix 6 ( $\alpha$ 6 R171, A172, R175, R176, and R179) [27]. Amino acid substitutions of several of these residues (R18E, H114A, W115A, A172E, R175/6EE, and R179E) cause perturbations in cytoplasmic localization, encapsidation, and Vif-deficient HIV-1 restriction [24,27]. Moreover, a human A3H triple mutant,


R115A, R175E, and R176E, lacks all of these activities and is also monomeric and fully defective for binding to duplex RNA [27].

Here, we investigated the role of RNA in HIV-1 Vif-mediated degradation of human A3H. First, immunoblots of cellular extracts were used to compare the Vif susceptibility of wild-type (WT) A3H and a panel of mutants with substitutions of one or more amino acids implicated in binding to duplex RNA. Second, live-cell imaging approaches were used to quantify the kinetics of Vif-mediated degradation of WT A3H versus select RNA binding-defective mutants. Finally, a split green fluorescence protein (GFP) reconstitution system was used to show that HIV-1 Vif is not capable of disrupting RNA-mediated A3H dimerization in living cells. Overall, the results of these experiments support a model in which Vif targets a surface on A3H that is distinct from the duplex RNA-binding region.

## Results

### Structures of A3H in complex with RNA

The x-ray structures of human, chimpanzee, and pig-tailed macaque A3H revealed a novel dimerization mechanism in which two monomers are united by binding to opposing sides of the same duplex RNA [27,49,50] (human A3H ribbon schematic in Fig. 1a and a superposition of the three structures in Fig. 1b). Several residues in A3H loop 1, loop 7, and helix 6 combine to bind duplex RNA, including a loop 7 tryptophan residue from each protomer that stacks with RNA nucleobases (W115 in each species' enzyme; amino acid positions shaded red in Fig. 1a and shown by alignment in Fig. 1c). Mutating these residues in human A3H compromises RNA binding and dimerization and, surprisingly, results in elevated ssDNA cytosine deamination activity [27]. For instance, a human A3H triple mutant with W115A in loop 7 and R175E and R176E in helix 6 has a monomeric size exclusion profile and a large increase in ssDNA deaminase activity [27]. This triple mutant combines amino acid substitutions that individually cause defects in cytoplasmic localization, packaging into nascent HIV-1 particles, and restricting the infectivity of Vif-deficient HIV-1 virions, which are all metrics of duplex RNA-binding activity [27]. Interestingly, the duplex RNA-binding domain is located near the predicted HIV-1 Vif-binding surface of human A3H, which, prior studies have shown, requires amino acids in helices 3 and 4 including D121 [14,16,19] (positions shaded yellow in Fig. 1d; considered again below and in Discussion).




**Fig. 1. Structures of A3H in complex with RNA.** (a) X-ray structure of human A3H in complex with duplex RNA showing loop 1, loop 7, and  $\alpha$ -helix 6 interacting with duplex RNA (ribbon schematic of PDB ID 6B0B). Positions of conserved RNA-interacting residues are shaded red. (b) Superposition of human (blue; PDB ID 6B0B), chimpanzee (green; PDB ID 5Z98), and pig-tailed macaque (beige; PDB ID 5W3V) A3H structures. (c) Partial amino acid sequence alignment of the loop 7 and  $\alpha$ -helix 6 regions of human, chimpanzee, and pig-tailed macaque A3H proteins. Conserved RNA-interacting residues are highlighted in red. (d) A3H structure with yellow highlights for previously identified Vif-interacting residues including D121. The orientation is similar to those in the bottom of panel (a) and panel (b) to facilitate comparisons.

### A subset of A3H RNA-binding residues are required for functional interaction with Vif

Most naturally occurring A3H variants including haplotype II are sensitive to Vif-mediated degradation [12,13,15,16,18,51,52]. The WT construct used here is A3H haplotype II splice variant 1–183. To ask whether RNA binding–defective proteins are susceptible or resistant to Vif-mediated degradation, a panel of mutants was cotransfected into 293T cells with 0.5  $\mu$ g or 1.0  $\mu$ g of HIV-1 LAI Vif (WT) or a degradation-defective mutant (F39V) and, after a 48-h incubation, protein levels were analyzed using immunoblotting whole cell extracts (representative blots in Fig. 2a with quantification in b). As controls, expression of WT Vif decreases levels of WT A3H as well as those of an A3H E56A catalytic mutant by approximately 50%, and Vif F39V is defective in WT A3H degradation (similar to previous observations [15,27]). As an additional control, A3H D121K resists degradation as evidenced by similar band intensities with WT Vif and the F39V mutant (again similar to previous observations [16,19]).

In comparison, A3H mutants with single amino acid substitutions of residues implicated in RNA binding showed either Vif-sensitive or Vif-resistant phenotypes. These different phenotypes were even observed for mutants within the same surface region of the protein. For instance, A3H helix 6 mutants A172E, R175E, and R179E showed a WT-like Vif sensitivity, whereas R176E appeared fully resistant to Vif-mediated degradation (representative blots in Fig. 2a with quantification in 2b). A3H loop 7 mutants, H114A and W115A, showed completely resistant phenotypes, whereas A3H loop 1 R18E retained a WT-like Vif sensitivity. Based on the single amino acid substitution mutant phenotypes, not surprisingly, the monomeric A3H triple mutant was also resistant to Vif-mediated degradation, and this property was unchanged by adding an E56A catalytic glutamate substitution. Similar results were obtained for the Vif protein of a different HIV-1 strain (IIIB) with previously demonstrated hyperfunctional amino acid substitutions [13] showing strong degradation of WT A3H and little degradation activity with the monomeric A3H triple mutant (Fig. 2c). These additional results are




**Fig. 2. Vif susceptibility of human A3H RNA binding-defective mutants.** (a) Immunoblots of 293T cells expressing WT A3H or the indicated mutants, together with empty vector (-), HIV-1 LAI Vif, or a F39V derivative. A3H was detected using an anti-A3H antibody, and Vif was detected using an anti-HA antibody. Anti- $\alpha$ -Tubulin was used as a loading control. (b) A bar graph quantifying immunoblot data from panel (a) and from an independent experiment (not shown). The A3H level for each construct is normalized to that of the corresponding empty vector control. The mean and difference between two independent experiments are shown. The dashed line represents level of sensitivity of WT A3H to Vif. (c) Immunoblots of 293T cells expressing WT A3H or the indicated mutants, together with empty vector (-), HIV-1 IIIIB-derived hyper-Vif (F39, H48, EKGE60-63) or hypo-Vif (V39, N48, GDAK60-63) with the same antibodies as in panel (a). (d) A structural depiction of human A3H residues required (red) or dispensable (orange) for functional interaction with HIV-1 Vif.

notable because the hyperfunctional Vif protein caused degradation of almost all cellular A3H and even showed partial activity against A3H D121K (all results in this panel are in comparison to a Vif variant, hypo, that has F39V as aforementioned and no A3H degradation activity as reported previously [13,51]). These results are summarized by color-coding phenotypes of these mutants on the surface of the three-dimensional structure of A3H with RNA (Fig. 2d).

#### RNA-binding mutants of A3H are degraded less efficiently by HIV-1 Vif

We next used live-cell fluorescent video microscopy to measure the degradation kinetics of fluorescently tagged A3H at a single-cell level during

HIV-1 infection. HeLa cell lines stably expressing WT or mutant A3H fused to the C-terminus of a yellow fluorescent protein (YFP-A3H) were generated and then infected with a replication-defective HIV-1 construct that, after integration into host cell genomic DNA, expresses mCherry as a reporter protein (MOI = 0.5). WT A3H and the Vif-resistant derivative D121K variant, each localized to cytoplasmic and nucleolar compartments, as reported [17,24,27,48,53]. The RNA binding-defective A3H triple mutant showed compromised cytoplasmic localization and appeared predominantly nuclear, also as reported [24,27]. As expected, WT LAI Vif expressed from the virus at the same time as the mCherry marker caused the degradation of WT A3H (representative image in Fig. 3a and Supplemental



**Fig. 3. The degradation kinetics of human A3H and key mutants.** (a) Representative images of HeLa cells stably expressing WT or mutant A3H fused to the C-terminus of a YFP before infection or 25-h post infection with an HIV-1 LAI mCherry reporter virus (WT Vif; MOI = 0.5). Parallel cultures were also infected with a reporter virus expressing LAI Vif F39V as a negative control (images not shown). Cells circled with dashed lines became infected as evidenced by mCherry expression. The scale bar at the upper right corner represents 10  $\mu$ m. Also see Supplemental movies S1–2 and S5–6. (b) Representative images of HeLa cells stably expressing WT or mutant A3H fused to the C-terminus of a YFP before infection or 16-h post infection with an HIV-1 NL4-3 mCherry reporter virus (hyper-Vif; MOI = 0.5). Parallel cultures were also infected with a reporter virus expressing hypo-Vif as a negative control (images not shown). Cells circled with dashed lines became infected as evidenced by mCherry expression. The scale bar at the upper right corner represents 10  $\mu$ m. Also see Supplemental movies S3–4 and S7–8. (c) Quantification of the rate of A3H degradation (change in YFP fluorescence) of a minimum of 10 infected cells representing each of the indicated conditions (mean  $\pm$  SD). (d) Flow cytometry quantification of the relative MFI of YFP-A3H in HeLa cells infected under the same conditions as panels (a–b). Data were normalized by setting the postinfection MFI of each Vif F39V or hypo-Vif reaction to 100%.

movie S1). The WT YFP-A3H fluorescence intensity declined at a relative rate of approximately 10% per hour (Fig. 3c). In comparison, both the YFP-A3H-D121K control and the RNA binding–defective triple mutant were resilient to degradation by LAI Vif (representative image in Fig. 3a, quantification in c, and Supplemental movie S2). As an additional control, infection of the same cells with an HIV-1 construct encoding Vif-F39V did not cause a reduction in fluorescence of any forms of A3H (images not shown; quantification summarized in Fig. 3c). These single-cell results are consistent with the immunoblot data described earlier using LAI Vif and untagged WT A3H.

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jmb.2019.09.014>.


Additional live cell fluorescent imaging experiments were done with a set of HIV-1 NL4-3 mCherry reporter viruses encoding either hyper- or hypo-Vif (variants described earlier and, notably, NL4-3 Vif and IIIB Vif have identical amino acid sequences). We observed that HIV-1 NL4-3 mCherry hyper-Vif not only degraded WT YFP-A3H (Supplemental movie S3), as expected, but it also triggered the degradation of YFP-A3H-D121K and the YFP-A3H triple mutant (Supplemental movie S4), albeit with slower kinetics (representative image in Fig. 3b and quantification in 3c). Infection by HIV-1 NL4-3 mCherry hypo-Vif did not reduce levels of any form of A3H (Fig. 3c). These single-cell imaging results were confirmed using flow cytometric quantification of the median fluorescence intensity (MFI) of the YFP (A3H) signal in infected cells 48 h post infection (Fig. 3d). Taken together, these real-time imaging data showed that the A3H RNA binding–defective mutant is less susceptible to HIV-1 Vif-mediated degradation compared with WT A3H. However, it can still be degraded by HIV-1 NL4-3–derived hyper-Vif (but not hypo-Vif), suggesting that the RNA binding–defective mutant of A3H is still capable of interacting with hyper-Vif and, importantly, that RNA may not be essential for a functional interaction to occur between Vif and A3H.

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jmb.2019.09.014>.

### A split GFP system to study A3H dimerization in living cells

To be able to study A3H dimerization in living cells, a split fluorescence system was designed in which one A3H protomer was N-terminally tagged with GFP  $\beta$ -strands 1–10 (GFP1–10) and the second with GFP  $\beta$ -strand 11 (GFP11) plus an extended flexible linker. Based on prior reports with other proteins including various homo- and heterodimerization partners (reviewed in Refs. [54,55]) and structural data from our group and others [27,49,50], efficient GFP reconstitution would only

be expected after A3H dimerization through duplex RNA (schematic in Fig. 4a). Indeed, a robust GFP signal was observed in both the cytoplasmic compartment and in nucleoli, after cotransfection of



**Fig. 4. A split GFP system to study A3H dimerization in living cells.** (a) A schematic of the split GFP system for A3H dimerization. One A3H construct is fused to GFP  $\beta$ -strand 1–10 (GFP1–10) and the other to  $\beta$ -strand 11 (GFP11). GFP reconstitution occurs upon duplex RNA–mediated dimerization of A3H. (b) Representative fluorescent microscopy images of 293T cells expressing split GFP constructs of the indicated A3s. An mCherry plasmid was included as a cotransfection control. The scale bar represents 20  $\mu$ m. (c) Flow cytometry profiles of the same cells as in panel (b). The MFI is shown for the whole mCherry-positive population, and the percentages of GFP-negative and GFP-positive cells are included at the bottom left and right, respectively (gates based on untransfected 293T cells analyzed in parallel).

293T cells with these constructs (Fig. 4b and c). Using flow cytometry, more than 70% of transfected mCherry-positive cells (cotransfection marker) were GFP-positive with MFI more than 20-fold higher than negative controls (Fig. 4c). Moreover, transfection of either construct alone failed to emit fluorescence (not shown), and similarly tagged A3A constructs also failed to reconstitute strong GFP signal (Fig. 4b and c). The latter negative result was expected because A3A is monomeric in living cells and it also behaves as a monomer under most conditions *in vitro* [56–58] (although A3A dimerization has been reported under some conditions *in vitro* [59]). As further validation of the robustness of this system, A3H-D121K yielded strong GFP signal similar to WT A3H, and the RNA binding–defective triple mutant (W115A, R175E, R176E) was indistinguishable from A3A (Fig. 4b and c). These results indicated that the split GFP reconstitution system can be used as a quantitative measure of A3H dimerization in living cells.

#### Vif overexpression does not interfere with A3H-mediated dimerization

We next asked whether HIV-1 IIIB Vif N48H, previously reported to degrade WT A3H but not cause cell cycle arrest [60,61], is capable of interfering with A3H dimerization using the split GFP system. In one scenario, Vif may bind to A3H regardless of its dimerization status, and if so, the GFP fluorescence signal should be unaffected by the presence of Vif (top schematic in Fig. 5a). Alternatively, Vif may intercept A3H before binding RNA, prevent dimerization, and lead to diminished GFP fluorescence reconstitution (bottom schematic in Fig. 5a).

To distinguish between these possibilities, 293T cells were cotransfected with the same panel of split GFP constructs as aforementioned together with Vif N48H or an empty vector control. Each reaction also included an mCherry expression vector as a transfection control. After a 32-h incubation period, one set of reactions was treated for 16 h with MG132 and the parallel set was untreated, and then all reactions were analyzed in parallel using immunoblotting and flow cytometry (Fig. 5b and c). Similar to results presented aforementioned in Fig. 2, in the absence of MG132, Vif triggered a reduction in WT A3H levels but did not affect the A3H RNA binding–defective triple mutant (W115A R175E R176E) or Vif-resistant A3H-D121K. This result is supported by flow cytometry where an MFI reduction of fourfold was observed (representative flow cytometry histograms in Fig. 5c and duplicate experiments averaged in the bar graph in the top part of Fig. 5b).


MG132 was included in the experiment to prevent A3H proteasomal degradation and facilitate quantification of the dimerization phenotype. Interestingly, the dimerization potential of WT A3H was unaffected by Vif under these conditions, as evidenced by

statistically indistinguishable MFIs (representative flow cytometry histograms in Fig. 5c and duplicate experiments averaged in the bar graph in the top part of Fig. 5b). It is worth noting that the lack of GFP reconstitution by the A3H triple mutant is likely due to a failure to dimerize through RNA and not to lower expression levels because MG132 treatment caused an increase in steady-state levels of this mutant to near those of WT A3H [49] (compare lanes 3, 7, 11, and 15 in Fig. 5b) and its GFP MFI remained approximately tenfold lower (representative flow cytometry histograms in Fig. 5c and duplicate experiments averaged in the bar graph in the top part of Fig. 5b). These results combined to indicate that the dimerization status of A3H is unlikely to be affected by the functional interaction with Vif.

## Discussion

RNA plays essential roles in the dimerization, subcellular localization, and antiviral functions of A3H. Here, we address whether the RNA bound by A3H has a role in HIV-1 Vif-mediated degradation. Immunoblot results with a panel of RNA-binding mutants were somewhat inconclusive with some A3H mutants resisting Vif and others maintaining WT levels of sensitivity (Fig. 2). The Vif resistance of an RNA binding–defective triple mutant of A3H was confirmed by imaging A3H degradation in living cells, but observations with a hyperfunctional Vif variant indicated that even this mutant could be targeted for degradation (albeit at an approximately four-fold slower rate than WT A3H; Fig. 3). A split GFP system enabled asking whether Vif might compete with the same surface as duplex RNA and interfere with A3H dimerization (Fig. 4). These results combined to favor a model in which the Vif and RNA-binding surfaces of A3H are physically distinct. For instance, even under conditions of proteasome inhibition to prevent A3H degradation, Vif did not have the capability to alter the dimerization potential of WT A3H.

As confirmed here and demonstrated in prior studies,  $\alpha$  helix 4 including residue D121 is essential for A3H degradation by HIV-1 Vif [14–19]. However, the hyper-Vif variant could still trigger the degradation of the D121K mutant, suggesting that this Vif protein may have a greater conformational flexibility and/or recognize a slightly different overall surface than the Vif proteins from HIV-1 IIIB/NL4-3 and HIV-1 LAI. Subcellular localization and/or as-yet-unidentified binding partners might also play a role in the observed resistance of the subset of RNA binding–defective mutants to Vif-mediated degradation. For instance, cytoplasmic A3H was degraded at a faster rate than nucleolar A3H (e.g., residual nucleolar A3H is evident in Fig. 3a and b in the presence of WT Vif and hyper-Vif, respectively; also



see Supplemental movies S1 and S3). Moreover, several of the RNA binding–defective mutants, including the triple mutant, tended to accumulate at higher levels in the nuclear compartment including the nucleoli, and this mislocalization likely also contributed to the overall degree of Vif resistance (particularly in immunoblots of whole cell lysates). Taken together with our prior localization studies, the RNA binding–defective mutants that are more nuclear also tend to be more resistant to Vif (and those that are cytoplasmic are more sensitive) [24,27]. It is therefore reasonable to postulate that the binding of Vif to CBF- $\beta$  immediately post translation and the assembly of a large >200 kDa E3 ligase complex hinders Vif from accessing the nuclear compartment.

These results are also sensible from practical and evolutionary perspectives. Recent structural, biophysical, and functional studies have indicated that duplex RNA is required for A3H to package from cytoplasmic sites into assembling viral particles and also for deamination-independent restriction of virus replication [27,49,50,62]. The massive molar excess of structured RNA in cells would make it difficult for Vif to compete for the same binding surface of A3H. Therefore, it is likely that the HIV-1 Vif protein evolved to avoid competition with RNA and bind directly to a distinct A3H surface. Thus, Vif may have evolved the potential to degrade both monomeric as well as RNA-dimerized forms of cellular A3H and thereby maintain the upper hand in counteracting restriction.

## Materials and Methods

### APOBEC3 expression constructs

The constructs expressing intron-containing WT, E56A, H114A, W115A, R175E, R176E, W115A/R175E/R176E, R18E, A172E, and R179E huA3H haplotype II in pcDNA3.1(+) (Invitrogen) have been reported [27]. Other mutants of A3H (D121K and E56A/W115A/R175E/R176E) within the same backbone were made using site-directed mutagenesis and verified using Sanger sequencing (primer sequences available on request). Retroviral transduction vectors encoding YFP fused to A3H (haplotype II), and mutants were generated by subcloning SYFP2 from SYFP2-C1 (Addgene #22878) into

pcDNA3.1 plasmids containing A3H using NheI and KpnI restriction cut sites followed by transfer into MigR1-derived simple retroviral vector using MluI and MfeI cut sites. The split GFP system used in this study was based on a previous system [63] with minor modifications. A gblock (IDT)-encoding GFP  $\beta$ -strand 11 (GFP11) followed by a 27 amino acid flexible linker was ordered and cloned into the previously reported pcDNA3.1(+) construct of WT A3H protein to the N-termini [27]. A ClaI restriction site was added in between GFP11 and the linker. GFP  $\beta$ -strand 1–10 (GFP1–10) was PCR amplified and cloned in place of GFP11 to make the GFP1–10 construct.

### HIV-1 and Vif expression constructs

The lentiviral Vif protein from HIV-1 IIIB (EU541617) was codon optimized (GenScript Corp) and cloned into pVR1012 with a C-terminal hemagglutinin (HA) tag using Sall and BamHI. The Vif protein from HIV-1 LAI (1102247D) was cloned into pVR1012 with a C-terminal HA tag using Sall and NotI. F39V mutant of HIV-1 LAI Vif, hypovariant (F39V), N48H, and hypervariant (N48H, GDAK60-63EKGE) of HIV-1 IIIB Vif were made using site-directed mutagenesis and verified using Sanger sequencing (primer sequences available on request). HIV-1 NL4-3 mCherry reporter viruses were generated by subcloning hypo- and hyper-Vif–encoding segments from IIIB infectious molecular clones [64] into a NL4-3 E-R-mCherry full-length reporter virus plasmid [65] using AgeI and Sall cut sites. HIV-1 LAI mCherry reporter viruses were generated by transferring the mCherry reporter cassette from pRGH (NIH ARP #12427) into a previously described LAI infectious molecular clone using BlnI and XhoI cut sites. LAI F39V was generated by site-directed mutagenesis. All plasmids were confirmed using restriction digestion and Sanger sequencing.

### Cell lines

293T cells were maintained at 37 °C and 5% CO<sub>2</sub> in Dulbecco's modified Eagle medium (DMEM) containing 10% fetal bovine serum (FBS) and 1% penicillin–streptomycin (P/S). HeLa cells were maintained at 37 °C and 5% CO<sub>2</sub> in DMEM supplemented with 10% FBS, 1% P/S, and 1% GlutaMAX.

### Vif degradation experiments

293T cells were transfected with WT or mutant pcDNA3.1(+)-A3H, along with 500 or 1000 ng

**Fig. 5. Vif does not interfere with A3H dimerization in the split GFP system.** (a) Models for Vif interaction with A3H. The top schematic depicts Vif binding to a distinct surface on A3H (predicted to have no interference in GFP reconstitution), whereas the bottom schematic depicts Vif competing with RNA for A3H (interference in GFP reconstitution). (b) Immunoblots of 293T cells expressing A3A, A3H, or the indicated A3H mutants, together with empty vector (-) or HIV-1 IIIB Vif N48H  $\pm$  MG132. A3H was detected using an anti-A3H antibody and Vif was detected using an anti-HA antibody. Anti- $\alpha$ -Tubulin was used as a loading control. A bar graph is shown earlier the immunoblots for quantification of flow cytometry data from panel (c) and from an independent experiment not shown. The MFI level of cells of each indicated condition is normalized to A3A in the absence of Vif or MG132. The mean and difference between two independent experiments are shown. (c) Flow cytometry profiles of 293T cells expressing the indicated A3 and Vif construct ( $\pm$ MG132). Each profile is labeled as in Fig. 4c.

pVR1012 HIV-1 LAI Vif WT/F39V-HA (Fig. 2a), or 50 or 100 ng pVR1012 HIV-1 IIIB hyper-/hypo-Vif (Fig. 2c) or empty vector, using Transit-LT1 (Mirus). The following amounts of A3H were transfected: 25 ng of WT, D121K, E56A, H114A, R175E, R18E, and A172E, 100 ng of R179E, and 200 ng of W115A, R176E, W115A/R175E/R176E, and E56A/W115A/R175E/R176E. Empty vector is added to equalize the amounts of total DNA. After 48 h, cells were harvested for immunoblot analysis.

### Immunoblotting experiments

Cell lysates were prepared by resuspension of washed cell pellets directly in 2.5× Laemmli sample buffer. Proteins were separated using discontinuous sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to polyvinylidene difluoride (PVDF) membranes (Millipore). HA-tagged A3 proteins were detected using monoclonal mouse anti-HA (BioLegend #MMS-101P). A3H proteins were detected using polyclonal rabbit anti-A3H (Novus #NBP1-91682). Tubulin was detected using a monoclonal mouse anti- $\alpha$ -Tubulin antibody (Covance #MMS-489P). Immunoblots were quantified using ImageStudio.

### Live-cell fluorescent imaging

Reporter viruses were produced in 293T cells by cotransfection with VSV-G to generate pseudotyped single cycle infectious virus. Viral inoculum was harvested at 48 h post transfection and passed through a 0.45- $\mu$ m syringe filter. HeLa YFP-A3H stable cells were plated at 20,000 cells per well of an ibidi  $\mu$ -well slide (ibidi GmbH) in 200  $\mu$ L of Fluorobrite DMEM (Thermo Fisher Scientific) supplemented with 10% FBS, 1% GlutaMAX, 1% P/S, and 100 mM HEPES. Cells were infected at an MOI of 0.5 infectious units per cell with 2  $\mu$ g/mL polybrene. Live-cell fluorescent video microscopy was performed as previously described [65], beginning at 12 h after addition of virus, and a multicolor image was captured every 60 min (mCherry, YFP, and DAPI) for up to 40 h post infection. All images were processed and analyzed using FIJI/ImageJ2 [66]. Briefly, regions of interest (ROIs) were drawn around whole single cells based on YFP-A3H image and around nuclei of those cells based on DAPI in every frame of live-cell movies. Cytoplasmic fluorescence was quantified by subtracting nuclear from whole-cell integrated density (IntDen). The cytoplasmic YFP and whole-cell mCherry fluorescence were normalized to each single cell's value at the beginning of each movie. Rate of change in YFP (showing YFP-A3H degradation) was quantified for the period of 5 h following detection of mCherry.

### Split GFP experiments

293T cells were transfected with 100 ng GFP1–10 plasmids, 25 ng GFP11 plasmids, 500 ng of HIV-1 IIIB Vif N48H-HA or empty vector, and 100 ng mCherry plasmid. Cells were treated with MG132 at a final concentration of 2  $\mu$ M at 32 h and subjected to analyses using fluorescence imaging, flow cytometry, and immunoblotting at 48 h post transfections. Cells were treated with NucBlue live-cell

stain (Invitrogen) at 37 °C for 30 min for DAPI (4',6'-diamidino-2-phenylindole) staining before imaging. Images were collected using a Nikon inverted Ti-E deconvolution microscope. A fraction of the cells was resuspended in phosphate-buffered saline (PBS) with 1 mM ethylenediaminetetraacetic acid (EDTA) in preparation for flow cytometry. GFP and mCherry fluorescence were measured on a BD FACS Canto II flow cytometer. Data were analyzed using FlowJo flow cytometry analysis software (version 8.8.6). The fraction of GFP+ cells and median fluorescence intensity (MFI) of all transfected (mCherry-positive) cells was quantified for each sample. The remaining cells were prepared for immunoblotting.

### Sequence alignments

All amino acid sequences were aligned using Clustal Omega. The amino acid sequences of human A3H, chimpanzee A3H, and pig-tailed macaque A3H correspond to GenBank accession number [ACK77775.1](#), PDB ID 5Z98\_A, and PDB ID 5W3V\_A, respectively.

### Ethics approval and consent to participate

There are no ethical considerations related to this work.

### Consent for publication

Not applicable.

### Availability of Data and Materials

All data generated or analyzed during this study are included in this published article and its supplementary information files.

---

### Acknowledgments

This work was supported by NIAID R37 AI064046 (to R.S.H.), NIGMS R01 GM118000 (to R.S.H. and H.A.), and NIGMS R01 GM118047 (to H.A.). J.W. received partial salary support from the University of Minnesota Graduate School, Interdisciplinary Doctoral Fellowship. D.J.S. received partial salary support from the University of Minnesota Craniofacial Research Training (MinnCResT) program (NIH T90DE022732). R.S.H. is the Margaret Harvey Schering Land Grant Chair for Cancer Research, a Distinguished University McKnight Professor, and an Investigator of the Howard Hughes Medical Institute. The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: pRGH-WT (#12427) from Ivan

Sadowski and Viviana Simon. Microscopy imaging and analysis was performed at the University Imaging Centers, University of Minnesota.

## Author Contributions

J.W., N.M.S., and R.S.H. conceived and designed the studies. J.W. performed Vif degradation experiments and split GFP studies. N.M.S., K.S., and H.A. provided structural analysis. J.T.B. performed live-cell imaging experiments. J.W., J.T.B., K.V.L., N.M.S., D.J.S., and N.M.S. created DNA constructs. R.S.H., H.A., and N.M.S. supervised the studies. J.W., N.M.S., and R.S.H. drafted the manuscript with contributions from all authors in figure preparation and revisions.

## Conflicts of Interest

R.S.H. is a cofounder, shareholder, and consultant of ApoGen Biotechnologies Inc. H.A. is a consultant of ApoGen Biotechnologies Inc. The other authors have no conflicts to declare.

Received 10 June 2019;

Received in revised form 4 September 2019;

Accepted 16 September 2019

Available online 16 October 2019

### Keywords:

APOBEC3H;

DNA cytosine deaminase;

HIV-1 Vif;

Subcellular localization;

Vif-APOBEC interaction

### Abbreviations used:

A3, APOBEC3; A3H, APOBEC3H; CBF- $\beta$ , core binding factor- $\beta$ ; GFP, green fluorescence protein; HA, hemagglutinin; ssDNA, single-stranded DNA; Vif, virion infectivity factor; WT, wild-type.

## References

- [1] B.A. Desimie, K.A. Delviks-Frankenberry, R.C. Burdick, D.F. Qi, T. Izumi, V.K. Pathak, Multiple APOBEC3 restriction factors for HIV-1 and one Vif to rule them all, *J. Mol. Biol.* 426 (2014) 1220–1245.
- [2] R.S. Harris, J.P. Dudley, APOBECs and virus restriction, *Virology* 479 (2015) 131–145.
- [3] Y. Nakano, H. Aso, A. Soper, E. Yamada, M. Moriwaki, G. Juarez-Fernandez, Y. Koyanagi, K. Sato, A conflict of interest: the evolutionary arms race between mammalian APOBEC3 and lentiviral Vif, *Retrovirology* 14 (2017).
- [4] S.G. Conticello, R.S. Harris, M.S. Neuberger, The Vif protein of HIV triggers degradation of the human antiretroviral DNA deaminase APOBEC3G, *Curr. Biol.* 13 (2003) 2009–2013.
- [5] R. Mariani, D. Chen, B. Schrofelbauer, F. Navarro, R. Konig, B. Bollman, C. Munk, H. Nymark-McMahon, N.R. Landau, Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif, *Cell* 114 (2003) 21–31.
- [6] M. Marin, K.M. Rose, S.L. Kozak, D. Kabat, HIV-1 Vif protein binds the editing enzyme APOBEC3G and induces its degradation, *Nat. Med.* 9 (2003) 1398–1403.
- [7] A.M. Sheehy, N.C. Gaddis, M.H. Malim, The antiretroviral enzyme APOBEC3G is degraded by the proteasome in response to HIV-1 Vif, *Nat. Med.* 9 (2003) 1404–1407.
- [8] K. Stopak, C. de Noronha, W. Yonemoto, W.C. Greene, HIV-1 Vif blocks the antiviral activity of APOBEC3G by impairing both its translation and intracellular stability, *Mol. Cell* 12 (2003) 591–601.
- [9] X. Yu, Y. Yu, B. Liu, K. Luo, W. Kong, P. Mao, X.F. Yu, Induction of APOBEC3G ubiquitination and degradation by an HIV-1 Vif-Cul5-SCF complex, *Science* 302 (2003) 1056–1060.
- [10] S. Jäger, D.Y. Kim, J.F. Hultquist, K. Shindo, R.S. LaRue, E. Kwon, M. Li, B.D. Anderson, L. Yen, D. Stanley, C. Mahon, J. Kane, K. Franks-Skiba, P. Cimermancic, A. Burlingame, A. Sali, C.S. Craik, R.S. Harris, J.D. Gross, N.J. Krogan, Vif hijacks CBF-beta to degrade APOBEC3G and promote HIV-1 infection, *Nature* 481 (2011) 371–375.
- [11] W. Zhang, J. Du, S.L. Evans, Y. Yu, X.F. Yu, T-cell differentiation factor CBF-beta regulates HIV-1 Vif-mediated evasion of host restriction, *Nature* 481 (2011) 376–379.
- [12] M. Binka, M. Ooms, M. Steward, V. Simon, The activity spectrum of Vif from multiple HIV-1 subtypes against APOBEC3G, APOBEC3F, and APOBEC3H, *J. Virol.* 86 (2012) 49–59.
- [13] E.W. Refsland, J.F. Hultquist, E.M. Luengas, T. Ikeda, N.M. Shaban, E.K. Law, W.L. Brown, C. Reilly, M. Emerman, R.S. Harris, Natural polymorphisms in human APOBEC3H and HIV-1 Vif combine in primary T lymphocytes to affect viral G-to-A mutation levels and infectivity, *PLoS Genet.* 10 (2014) e1004761.
- [14] M. Nakashima, S. Tsuzuki, H. Awazu, A. Hamano, A. Okada, H. Ode, M. Maejima, A. Hachiya, Y. Yokomaku, N. Watanabe, H. Akari, Y. Iwatani, Mapping region of human restriction factor APOBEC3H critical for interaction with HIV-1 Vif, *J. Mol. Biol.* 429 (2017) 1262–1276.
- [15] A. Harari, M. Ooms, L.C. Mulder, V. Simon, Polymorphisms and splice variants influence the antiretroviral activity of human APOBEC3H, *J. Virol.* 83 (2009) 295–303.
- [16] A. Zhen, T. Wang, K. Zhao, Y. Xiong, X.F. Yu, A single amino acid difference in human APOBEC3H variants determines HIV-1 Vif sensitivity, *J. Virol.* 84 (2010) 1902–1911.
- [17] M.M. Li, M. Emerman, Polymorphism in human APOBEC3H affects a phenotype dominant for subcellular localization and antiviral activity, *J. Virol.* 85 (2011) 8197–8207.
- [18] X. Wang, A. Abudu, S. Son, Y. Dang, P.J. Venta, Y.H. Zheng, Analysis of human APOBEC3H haplotypes and anti-human immunodeficiency virus type 1 activity, *J. Virol.* 85 (2011) 3142–3152.
- [19] M. Ooms, M. Letko, V. Simon, The structural interface between HIV-1 Vif and human APOBEC3H, *J. Virol.* 91 (2017) e02289.

- [20] S.L. Kozak, M. Marin, K.M. Rose, C. Bystrom, D. Kabat, The anti-HIV-1 editing enzyme APOBEC3G binds HIV-1 RNA and messenger RNAs that shuttle between polysomes and stress granules, *J. Biol. Chem.* 281 (2006) 29105–29119.
- [21] S. Gallois-Montbrun, B. Kramer, C.M. Swanson, H. Byers, S. Lynham, M. Ward, M.H. Malim, Antiviral protein APOBEC3G localizes to ribonucleoprotein complexes found in P bodies and stress granules, *J. Virol.* 81 (2007) 2165–2178.
- [22] T. Izumi, R. Burdick, M. Shigemi, S. Plisov, W.S. Hu, V.K. Pathak, Mov10 and APOBEC3G localization to processing bodies is not required for virion incorporation and antiviral activity, *J. Virol.* 87 (2013) 11047–11062.
- [23] P.K. Phalora, N.M. Sherer, S.M. Wolinsky, C.M. Swanson, M.H. Malim, HIV-1 replication and APOBEC3 antiviral activity are not regulated by P bodies, *J. Virol.* 86 (2012) 11712–11724.
- [24] D.J. Salamango, J.T. Becker, J.L. McCann, A.Z. Cheng, Ö. Demir, R.E. Amaro, W.L. Brown, N.M. Shaban, R.S. Harris, APOBEC3H subcellular localization determinants define zipcode for targeting HIV-1 for restriction, *Mol. Cell. Biol.* 38 (2018) e00356.
- [25] M.D. Stenglein, H. Matsuo, R.S. Harris, Two regions within the amino-terminal half of APOBEC3G cooperate to determine cytoplasmic localization, *J. Virol.* 82 (2008) 9591–9599.
- [26] M.J. Wichroski, G.B. Robb, T.M. Rana, Human retroviral host restriction factors APOBEC3G and APOBEC3F localize to mRNA processing bodies, *PLoS Pathog.* 2 (2006) e41.
- [27] N.M. Shaban, K. Shi, K.V. Lauer, M.A. Carpenter, C.M. Richards, D. Salamango, J. Wang, M.W. Lopresti, S. Banerjee, R. Levin-Klein, W.L. Brown, H. Aihara, R.S. Harris, The antiviral and cancer genomic DNA deaminase APOBEC3H is regulated by an RNA-mediated dimerization mechanism, *Mol. Cell* 69 (2018) 75–86.
- [28] H. Fukuda, S. Li, L. Sardo, J.L. Smith, K. Yamashita, A.D. Sarca, K. Shirakawa, D.M. Standley, A. Takaori-Kondo, T. Izumi, Structural determinants of the APOBEC3G N-terminal domain for HIV-1 RNA association, *Front Cell Infect Microbiol.* 9 (2019) 129.
- [29] T. Kouno, E.M. Luengas, M. Shigematsu, S.M. Shandilya, J. Zhang, L. Chen, M. Hara, C.A. Schiffer, R.S. Harris, H. Matsuo, Structure of the Vif-binding domain of the antiviral enzyme APOBEC3G, *Nat. Struct. Mol. Biol.* 22 (2015) 485–491.
- [30] L. Apolonia, R. Schulz, T. Curk, P. Rocha, C.M. Swanson, T. Schaller, J. Ule, M.H. Malim, Promiscuous RNA binding ensures effective encapsidation of APOBEC3 proteins by HIV-1, *PLoS Pathog.* 11 (2015) e1004609.
- [31] H.P. Bogerd, B.R. Cullen, Single-stranded RNA facilitates nucleocapsid: APOBEC3G complex formation, *RNA* 14 (2008) 1228–1236.
- [32] S. Cen, F. Guo, M. Niu, J. Saadatmand, J. Deffassieux, L. Kleiman, The interaction between HIV-1 Gag and APOBEC3G, *J. Biol. Chem.* 279 (2004) 33177–33184.
- [33] H. Huthoff, M.H. Malim, Identification of amino acid residues in APOBEC3G required for regulation by human immunodeficiency virus type 1 Vif and virion encapsidation, *J. Virol.* 81 (2007) 3807–3815.
- [34] M.A. Khan, S. Kao, E. Miyagi, H. Takeuchi, R. Goila-Gaur, S. Opi, C.L. Gipson, T.G. Parslow, H. Ly, K. Strebler, Viral RNA is required for the association of APOBEC3G with human immunodeficiency virus type 1 nucleoprotein complexes, *J. Virol.* 79 (2005) 5870–5874.
- [35] A. Schafer, H.P. Bogerd, B.R. Cullen, Specific packaging of APOBEC3G into HIV-1 virions is mediated by the nucleocapsid domain of the gag polyprotein precursor, *Virology* 328 (2004) 163–168.
- [36] E.S. Svarovskaia, H. Xu, J.L. Mbisa, R. Barr, R.J. Gorelick, A. Ono, E.O. Freed, W.S. Hu, V.K. Pathak, Human apolipoprotein B mRNA-editing enzyme-catalytic polypeptide-like 3G (APOBEC3G) is incorporated into HIV-1 virions through interactions with viral and nonviral RNAs, *J. Biol. Chem.* 279 (2004) 35822–35828.
- [37] T. Wang, C. Tian, W. Zhang, K. Luo, P.T. Sarkis, L. Yu, B. Liu, Y. Yu, X.F. Yu, 7SL RNA mediates virion packaging of the antiviral cytidine deaminase APOBEC3G, *J. Virol.* 81 (2007) 13112–13124.
- [38] T. Wang, C. Tian, W. Zhang, P.T. Sarkis, X.F. Yu, Interaction with 7SL RNA but not with HIV-1 genomic RNA or P bodies is required for APOBEC3F virion packaging, *J. Mol. Biol.* 375 (2008) 1098–1112.
- [39] A. York, S.B. Kutluay, M. Errando, P.D. Bieniasz, The RNA binding specificity of human APOBEC3 proteins resembles that of HIV-1 nucleocapsid, *PLoS Pathog.* 12 (2016) e1005833.
- [40] V. Zennou, D. Perez-Caballero, H. Gottlinger, P.D. Bieniasz, APOBEC3G incorporation into human immunodeficiency virus type 1 particles, *J. Virol.* 78 (2004) 12058–12061.
- [41] K. Bélanger, M.A. Langlois, RNA-binding residues in the N-terminus of APOBEC3G influence its DNA sequence specificity and retrovirus restriction efficiency, *Virology* 483 (2015) 141–148.
- [42] K.N. Bishop, M. Verma, E.Y. Kim, S.M. Wolinsky, M.H. Malim, APOBEC3G inhibits elongation of HIV-1 reverse transcripts, *PLoS Pathog.* 4 (2008) e1000231.
- [43] K. Gillick, D. Pollpeter, P. Phalora, E.Y. Kim, S.M. Wolinsky, M.H. Malim, Suppression of HIV-1 infection by APOBEC3 proteins in primary human CD4(+) T cells is associated with inhibition of processive reverse transcription as well as excessive cytidine deamination, *J. Virol.* 87 (2013) 1508–1517.
- [44] Y. Iwatani, D.S. Chan, F. Wang, K.S. Maynard, W. Sugiura, A.M. Gronenborn, I. Rouzina, M.C. Williams, K. Musier-Forsyth, J.G. Levin, Deaminase-independent inhibition of HIV-1 reverse transcription by APOBEC3G, *Nucleic Acids Res.* 35 (2007) 7096–7108.
- [45] T. Kobayashi, Y. Koizumi, J.S. Takeuchi, N. Misawa, Y. Kimura, S. Morita, K. Aihara, Y. Koyanagi, S. Iwami, K. Sato, Quantification of deaminase activity-dependent and -independent restriction of HIV-1 replication mediated by APOBEC3F and APOBEC3G through experimental-mathematical investigation, *J. Virol.* 88 (2014) 5881–5887.
- [46] M. Morse, R. Huo, Y. Feng, I. Rouzina, L. Chelico, M.C. Williams, Dimerization regulates both deaminase-dependent and deaminase-independent HIV-1 restriction by APOBEC3G, *Nat. Commun.* 8 (2017) 597.
- [47] E.N. Newman, R.K. Holmes, H.M. Craig, K.C. Klein, J.R. Lingappa, M.H. Malim, A.M. Sheehy, Antiviral function of APOBEC3G can be dissociated from cytidine deaminase activity, *Curr. Biol.* 15 (2005) 166–170.
- [48] M. Ooms, S. Majdak, C.W. Seibert, A. Harari, V. Simon, The localization of APOBEC3H variants in HIV-1 virions determines their antiviral activity, *J. Virol.* 84 (2010) 7961–7969.
- [49] T. Matsuoka, T. Nagae, H. Ode, H. Awazu, T. Kurosawa, A. Hamano, K. Matsuoka, A. Hachiya, M. Imahashi, Y. Yokomaku, N. Watanabe, Y. Iwatani, Structural basis of chimpanzee APOBEC3H dimerization stabilized by double-stranded RNA, *Nucleic Acids Res.* 46 (2018) 10368–10379.
- [50] J.A. Bohn, K. Thummar, A. York, A. Raymond, W.C. Brown, P.D. Bieniasz, T. Hatziioannou, J.L. Smith, APOBEC3H

- structure reveals an unusual mechanism of interaction with duplex RNA, *Nat. Commun.* 8 (2017) 1021.
- [51] M. Ooms, B. Brayton, M. Letko, S.M. Maio, C.D. Pilcher, F.M. Hecht, J.D. Barbour, V. Simon, HIV-1 Vif adaptation to human APOBEC3H haplotypes, *Cell Host Microbe* 14 (2013) 411–421.
- [52] M.M. Li, L.I. Wu, M. Emerman, The range of human APOBEC3H sensitivity to lentiviral Vif proteins, *J. Virol.* 84 (2010) 88–95.
- [53] G.J. Starrett, E.M. Luengas, J.L. McCann, D. Ebrahimi, N.A. Temiz, R.P. Love, Y. Feng, M.B. Adolph, L. Chelico, E.K. Law, M.A. Carpenter, R.S. Harris, The DNA cytosine deaminase APOBEC3H haplotype I likely contributes to breast and lung cancer mutagenesis, *Nat. Commun.* 7 (2016) 12918.
- [54] M.G. Romei, S.G. Boxer, Split green fluorescent proteins: scope, limitations, and outlook, *Annu. Rev. Biophys.* 48 (2019) 19–44.
- [55] C. Foglieni, S. Papin, A. Salvade, T. Afroz, S. Pinton, G. Pedrioli, G. Ulrich, M. Polymenidou, P. Paganetti, Split GFP technologies to structurally characterize and quantify functional biomolecular interactions of FTD-related proteins, *Sci. Rep.* 7 (2017) 14013.
- [56] K. Shi, O. Demir, M.A. Carpenter, J. Wagner, K. Kurahashi, R.S. Harris, R.E. Amaro, H. Aihara, Conformational switch regulates the DNA cytosine deaminase activity of human APOBEC3B, *Sci. Rep.* 7 (2017) 17415.
- [57] J. Li, Y. Chen, M. Li, M.A. Carpenter, R.M. McDougale, E.M. Luengas, P.J. Macdonald, R.S. Harris, J.D. Mueller, APOBEC3 multimerization correlates with HIV-1 packaging and restriction activity in living cells, *J. Mol. Biol.* 426 (2014) 1296–1307.
- [58] I.J. Byeon, J. Ahn, M. Mitra, C.H. Byeon, K. Hercik, J. Hritz, L.M. Charlton, J.G. Levin, A.M. Gronenborn, NMR structure of human restriction factor APOBEC3A reveals substrate binding and enzyme specificity, *Nat. Commun.* 4 (2013) 1890.
- [59] T. Kouno, T.V. Silvas, B.J. Hilbert, S.M.D. Shandilya, M.F. Bohn, B.A. Kelch, W.E. Royer, M. Somasundaran, N. Kurt Yilmaz, H. Matsuo, C.A. Schiffer, Crystal structure of APOBEC3A bound to single-stranded DNA reveals structural basis for cytosine deamination and specificity, *Nat. Commun.* 8 (2017) 15024.
- [60] M. Ooms, M. Letko, M. Binka, V. Simon, The resistance of human APOBEC3H to HIV-1 NL4-3 molecular clone is determined by a single amino acid in Vif, *PLoS One* 8 (2013) e57744.
- [61] K. Zhao, J. Du, Y. Rui, W. Zheng, J. Kang, J. Hou, K. Wang, W. Zhang, V.A. Simon, X.F. Yu, Evolutionarily conserved pressure for the existence of distinct G2/M cell cycle arrest and A3H inactivation functions in HIV-1 Vif, *Cell Cycle* 14 (2015) 838–847.
- [62] M. Mitra, D. Singer, Y. Mano, J. Hritz, G. Nam, R.J. Gorelick, I.J. Byeon, A.M. Gronenborn, Y. Iwatani, J.G. Levin, Sequence and structural determinants of human APOBEC3H deaminase and anti-HIV-1 activities, *Retrovirology* 12 (2015) 3.
- [63] D. Kamiyama, S. Sekine, B. Barsi-Rhyne, J. Hu, B. Chen, L.A. Gilbert, H. Ishikawa, M.D. Leonetti, W.F. Marshall, J.S. Weissman, B. Huang, Versatile protein tagging in cells with split fluorescent protein, *Nat. Commun.* 7 (2016) 11046.
- [64] G. Haché, K. Shindo, J.S. Albin, R.S. Harris, Evolution of HIV-1 isolates that use a novel Vif-independent mechanism to resist restriction by human APOBEC3G, *Curr. Biol.* 18 (2008) 819–824.
- [65] J.T. Becker, N.M. Sherer, Subcellular localization of HIV-1 gag-pol mRNAs regulates sites of virion assembly, *J. Virol.* 91 (2017).
- [66] J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, A. Cardona, Fiji: an open-source platform for biological-image analysis, *Nat. Methods* 9 (2012) 676–682.