



ELSEVIER

Contents lists available at ScienceDirect

Microbiological Research

journal homepage: www.elsevier.com/locate/micres

Muscodor brasiliensis sp. nov. produces volatile organic compounds with activity against *Penicillium digitatum*

Lorena C. Pena^a, Gustavo H. Jungklaus^a, Daiani C. Savi^a, Lisandra Ferreira-Maba^a, André Servienski^a, Beatriz H.L.N.S. Maia^b, Vinicius Annies^b, Lygia V. Galli-Terasawa^a, Chirlei Glienke^a, Vanessa Kava^{a,*}

^a Departamento de Genética, Universidade Federal do Paraná, Cx. Postal 19071, 81531-980, Curitiba, PR, Brazil

^b Departamento de Química, Universidade Federal do Paraná, Av. Coronel Francisco Heráclito dos Santos, 210, 81531-980, Curitiba, PR, Brazil

ARTICLE INFO

Keywords:

Biocontrol
VOCs
Endophytic fungus
Plant diseases
Green mold

ABSTRACT

Endophytic fungi belonging to *Muscodor* genus are considered as promising alternatives to be used in biological control due to the production of volatile organic compounds (VOCs). The strains LGMF1255 and LGMF1256 were isolated from the medicinal plant *Schinus terebinthifolius* and, by morphological data and phylogenetic analysis, identified as belonging to *Muscodor* genus. Phylogenetic analysis suggests that strain LGMF1256 is a new species, which is herein introduced as *Muscodor brasiliensis* sp. nov. The analysis of VOCs production revealed that compounds phenylethyl alcohol, α -curcumene, and E (β) farnesene until now has been reported only from *M. brasiliensis*, data that supports the classification of strain LGMF1256 as a new species. *M. brasiliensis* completely inhibited the phytopathogen *P. digitatum* *in vitro*. We also evaluated the ability of VOCs from LGMF1256 to inhibit the development of green mold symptoms by inoculation of *P. digitatum* in detached oranges. *M. brasiliensis* reduced the severity of diseases in 77%, and showed potential to be used for fruits storage and transportation to prevent the green mold symptoms development, eventually reducing the use of fungicides.

1. Introduction

The *Muscodor* genus was described as a sterile endophytic fungus isolated from *Cinnamomum zeylanicum* in Honduras (Worapong et al., 2001), thereafter the isolation of strains belonging to this genus has been reported in tropical rainforests in Australia, Central and South America, and Southeastern and Northeastern Himalayas (Zhang et al., 2010, Suwannarach et al., 2013, Saxena et al., 2015, Pena et al., 2017). The *Muscodor* genus comprises 18 legitimate species deposited at MycoBank database (<http://www.mycobank.org>) that have been described based on phylogenetic analysis of the *ITS1-5.8S-ITS2* sequences and volatile organic composition analysis (Zhang et al., 2010, Suwannarach et al., 2013, Saxena et al., 2015). Morphologically, these endophytic fungi are differentiated by colony and mycelial characteristics, that appear differ in hyphal arrangement (Strobel et al., 2001, Zhang et al., 2010).

Muscodor species are used in biological control due to the production of volatile organic compounds (VOCs) with antimicrobial properties (Strobel et al., 2001, Worapong et al., 2002, Pena et al., 2017). Interestingly, fungi of the genus *Muscodor* can inhibit a large variety of

fungi, but do not inhibit fungi of the same genus (Ezra et al., 2004). The use of VOCs is especially effective against postharvest diseases in view of the ability to act and diffuse through heterogeneous environments, whereas water-soluble antimicrobial agents are limited to proximal interactions via aqueous-phase diffusion (Worapong et al., 2001, Fialho et al. 2011, Hutchings et al., 2017). Thus, VOCs have already been used to prevent postharvest disease during storage and transport of grapes (Gabler et al., 2010) and peaches (Schnabel and Mercier, 2006).

Among the postharvest diseases, the green mold caused by *Penicillium digitatum* (Macarasin et al., 2007, Marcet-Houben et al., 2012) has great impact in the citrus production worldwide, including in Brazil (FAOSTAT, 2015), accounting for half of the total fruit wastage (Milind, 2008). The green mold symptoms in oranges are characterized by slight fruit discoloration in the affected region, and at an advanced stage, the mycelium assumes a greenish color due to spore production. The lesion generally shows rapid development and can easily compromise the entire fruit (Timmer et al., 2002). Control of green mold is mainly based on application of several fungicides and maturation regulators (Agrios, 2005). However, the widespread use of fungicides for green mold control significantly increases the production costs and has also been

* Corresponding author.

E-mail addresses: vankava@ufpr.br, vanessagenetica@gmail.com (V. Kava).

<https://doi.org/10.1016/j.micres.2019.01.002>

Received 6 December 2018; Received in revised form 8 January 2019; Accepted 17 January 2019

Available online 18 January 2019

0944-5013/ © 2019 Elsevier GmbH. All rights reserved.

associated with negative environmental effects and selection of resistant pathogens (Agrios, 2005, Fischer et al., 2011, Talibi et al., 2014). In this scenario, promising results have been found for the biocontrol of this disease using yeasts from the Brazilian Cerrado (Sperandio et al., 2015). Therefore, the use of VOCs produced by *Muscodor* species for biocontrol (Worapong et al., 2001) could be an alternative treatment of postharvest diseases, such as green mold.

Based on the reasoning presented so far, a decision was made to study the Brazilian medicinal plant *Schinus terebinthifolius* as a source of *Muscodor* isolates able to act against the citrus green mold disease. Endophytes from *S. terebinthifolius* have shown high antimicrobial activity (Dos Santos et al., 2016, Tonial et al. 2016); however, *Muscodor* species have not been previously reported as part of the endophytic community of this medicinal plant. Hence, the following objectives were established for this work i) isolate *Muscodor* strains from *S. terebinthifolius*, ii) conduct morphological, VOCs, and phylogenetic analyses to check if the isolated strains belong to the *Muscodor* genus, and iii) investigate the ability of the isolated strains to inhibit *P. digitatum* development.

2. Materials and methods

2.1. Biological material

Muscodor sp. LGMF1254, previously obtained by our research group (Pena et al., 2017) was used as a selective factor to obtain new isolates of this genus, since it is reported in the literature that *Muscodor* isolates were resistant to *Muscodor* VOCs. *P. digitatum* LGMF1507 was obtained from contaminated market oranges and its identity was confirmed through ITS DNA sequence analysis (GenBank accession No.: KY978834). All the strains were deposited in the LabGeM Culture Collection, Federal University of Parana (UFPR), Curitiba, Brazil.

2.2. *Muscodor* isolation

Ten leaves of *Schinus terebinthifolius*, without scratches, or wounds, from healthy plants were collected in Curitiba, Brazil (25°26'52.5"S 49°14'00.3"W). The leaves surfaces were disinfected by immersion in 70% (v/v) ethanol for 1 min, followed by incubation in 2.5% NaClO (v/v available chlorine) for 4 min and washed in 70% ethanol for 30 s. The leaves were rinsed twice with sterilized distilled water for 6 min and cut into fragments of 7 × 7 mm and transferred to Petri dishes containing potato dextrose agar (PDA) medium (pH 5.8) and a previously cultivated inoculum (5 day old) of *Muscodor* sp. LGMF1254. The Petri dishes were incubated at 24 °C. Emerging hyphae for the next 2 weeks were transferred to a new PDA plate for further analyses. Two isolates, named as LGMF1255 and LGMF1256, were then isolated and deposited in the LabGeM Culture Collection, UFPR, Brazil.

2.3. Morphological characterization

2.3.1. Macroscopic analysis

To assess the differences in growth rates and colony morphology, strains LGMF1255 and LGMF1256 were grown in Petri dishes containing PDA, malt extract agar medium (MEA; malt extract 20.0 g/L, dextrose 20.0 g/L, peptone 1.0 g/L, agar 15.0 g/L), Sabouraud agar (SDA; peptone 15 g/L, dextrose 45 g/L, agar 15 g/L), or a minimal medium (MM; NaNO₃ 6 g/L, KH₂PO₄ 1.5 g/L, MgSO₄·7H₂O 0.5 g/L, KCl 0.5 g/L, FeSO₄ 0.01 g/L, ZnSO₄ 0.01 g/L, dextrose 10 g/L, agar 15.0 g/L), at 24 °C for 8 days. The characteristics of colonies were analyzed by visual inspection, and the growth rate was evaluated by measuring the colony area with the software Image J (Rasband, 2016).

2.3.2. Light microscopy analysis

Microcultures of isolates LGMF1255 and LGMF1256 were grown separately based on Kern and Blevins (1997). The isolates were grown

in PDA medium at 24 °C for 5 and 10 days, and the slides were stained with lactophenol cotton blue or analyzed directly with distilled water. The slides were examined on Axio Imager Z2 (Carl Zeiss, Jena, DE), with software Metafer 4/VSlide (Metasystems, Altlußheim, DE), by differential interference contrast (DIC) illumination. Micrometry was carried out with the Image J software (Rasband, 2016) (National Institutes of Health, Bethesda, MD) with 50 measurements per structure.

2.3.3. Scanning electron microscopy (SEM)

The isolates were grown on PDA at 24 °C for 10 days, and fixed in Karnovsky solution (2.5% glutaraldehyde and 2.5% paraformaldehyde in 0.05 M sodium cacodylate buffer, CaCl₂ 0.001 M, pH 7.2) for 24 h. The samples were dehydrated in ascending series of ethanol solutions (10%, 30%, 50%, and 70%) for 10 min each step, then ethanol 95% for 15 min, and finally maintained 72 h in 100% ethanol. The samples were brought to a critical point, dried, and coated with 80-nm gold. The observed images of structures and arrangements of mycelium were captured under low-vacuum conditions in a VEGA 3 TESCAN microscope.

2.4. DNA extraction, sequencing, and phylogenetic analysis

Genomic DNA extraction was performed using the UltraClean™ Microbial DNA Kit (MO Bio, Carlsbad, CA). For amplification of the ITS region of ribosomal DNA, primers ITS1 and ITS4 (White and Morrow, 1990; Gerrits Van Den Ende and De Hoog, 1999) were used. All the amplicons were purified using the enzymes Exo1 and FastAP™ (GE Healthcare, USA) and were sequenced using the BigDye Sequencing Kit (Applied Biosystems). The sequences were analyzed on an ABI3500 DNA Sequencer (Applied Biosystems). The BioEdit software was applied to edit the sequences, which were compared with sequences from the type species available in the GenBank database of NCBI (<http://www.ncbi.nlm.nih.gov/>) and Mycobank (<http://www.mycobank.org/>), and aligned in the CLUSTAL_X v.1.81 software. The best-fit model was identified in web-based MODELTEST 3.7: Kimura 2 parameters plus Gamma were selected. The Bayesian inference phylogenetic analysis was carried out with 10,000 generations by means of the MrBayes software, v.3.1.1, using *Xylaria berteri* (GenBank accession No. GU324749) as an outgroup. Sequences obtained in this study were deposited in GenBank under the following accession numbers ITS: KY924493 (LGMF1255) and KY924494 (LGMF1256). The final alignment of ITS was deposited in TreeBASE under accession number 21938.

2.5. Identification of volatile organic compounds

The identification of VOCs produced by isolates LGMF1255 and LGMF1256 was performed using solid-phase microextraction coupled gas chromatography with mass spectrometry (SPME-GC-MS). For trapping the VOCs produced, the cultures were set up with 5 mL of PDA in a 20 mL glass test tube closed with a rubber stopper and incubated at 24 °C; a syringe with a stable, flexible fiber made of 50/30 divinylbenzene (DVB)/carboxen on polydimethylsiloxane (PMDS) (Supelco, Sigma-Aldrich) was employed. The syringe was manually inserted into a small hole in the rubber stopper of the test tube, and the fiber was exposed to the air space above the 10 days older mycelium for 60 min and immediately desorbed into the hot injector of the gas chromatograph coupled to a mass spectrometer.

Volatile constituents were analyzed on a Shimadzu GC-2010 gas chromatographer coupled with a GCMS-QP2010 Plus mass spectrometer, fitted with a Shimadzu AOC-20i auto sampler and equipped with a Rtx-5MS capillary column (30 m × 0.25 mm × 0.25 μm). The device was operated at programmed temperatures from 60 °C to 250 °C changed at 3 °C min, and at an injector temperature of 250 °C, with an injection of 15 min. The interface ion source was at 300 °C, analyzing molecular fragments between 40 and 350 m/z, with helium as a carrier gas at a flow rate of 1.02 mL/min, in the ionization mode: electron




Fig. 1. Morphological characteristics of isolates LGMF1255 (box 1) and LGMF1256 (*Muscodor brasiliensis*) (box 2) after 8 days of growth. Macromorphology of isolate LGMF1255 (*Muscodor* sp.) (box 1) cultivated on PDA (1 A), MM (1B), MEA (1C), and SDA (1D). Micromorphological aspects of isolate LGMF1255 showing coiling formation (1E, F), cauliflower-like structures (1 G), vesicles (1 H), hyphae interlacing (1I), and mycelium arrangement (1 J). Macromorphology of isolate LGMF1256 on PDA (2 A), MM (2B), MEA (2C), and SDA (2D). Micromorphological aspects showing mycelium development (2E-F), cauliflower-like structures (2G) and mycelium arrangements (2H, 2I, 2J).

Notes: The scale bars on panels A–D are 4 cm. The scale bars on light microscopy images (E and F) are 10 μ m. In the SEM images, the scale is 5 μ m (G), 10 μ m (H and J), and 20 μ m (I).

impact 70 eV. The constituents of the volatiles were identified by calculating retention indexes (RI) and compared with mass spectra from the NIST08 library and with data from the literature (Adams, 2007). Compounds identified in control test tubes, and the silylated compounds from the column were subtracted from the analysis. Experimental RIs were determined by means of the *n*-alkane series (C_9 – C_{24}) standard, using the equations of Van der Dool and Kratz (Van Den Dool and Kratz, 1963). The components were quantified according to the peak area, without correction for response factors. Each condition (PDA and PDA plus *Muscodor* isolate) was analyzed in triplicate.

2.6. Activity against *Penicillium digitatum*

2.6.1. Inhibition of *P. digitatum* mycelial development

The production of VOCs with antifungal activity, against *Penicillium digitatum* (LGMF1507), by strains LGMF1255 and LGMF1256 were evaluated primarily in PDA media using plates with a divisor (Pena et al., 2017). The strain that showed the highest antifungal activity was selected for evaluation of VOCs production in autoclaved wheat or oatmeal grains as a nutrient source. Petri dishes with two compartments

were used, one compartment containing PDA (10 mL) for the phytopathogen development and in the other wheat or oatmeal. Mycelial discs (5-mm diameter) of 10-day-old colonies of selected strain were added to the compartments containing wheat, or oatmeal. After 7 days, an inoculum of *P. digitatum* LGMF1507 (1-mm diameter) was placed in the PDA compartment. As a positive control only the phytopathogen was inoculated. The Petri dishes were sealed with Parafilm[®] and Scotch[®] tape and incubated at 24 °C for 8 days. The inhibition analysis was performed comparing the growth of the pathogen (colony area: cm²) in comparison to control.

2.6.2. Biocontrol assay in fruits

The experiment was conducted using 40 organic sweet oranges (*Citrus sinensis* (L) Obsbeck). The fruits were disinfected, first immersed in 70% (v/v) ethanol for 1 min, followed by immersion in 2.5% NaClO (v/v available chlorine) for 3 min and washed with 70% ethanol for 30 s and finally the fruits were rinsed twice with sterilized distilled water. The *P. digitatum* LGMF1507 development in citrus fruits was performed as described by López-Pérez et al., (2015). To evaluate the ability of endophytic strain to inhibit the *P. digitatum* development, each orange

was placed in disinfected transparent plastic cylinders (volume of 500 mL, $r = 5.5$ cm, $h = 8$ cm) and the fruit was inoculated with $10 \mu\text{L}$ of phytopathogen spore solution (10^6 conidia/mL) deposited into 12 small injuries distributed in an area of 1 cm^2 . In the treatment, a culture plate without the lid containing the *Muscodora* strain, growth on PDA for 20 days was deposited inside of the plastic cylinder. In the positive control only the phytopathogen was inoculated, and the negative control consists in the inoculation of a saline solution (NaCl 0.85%), instead of the phytopathogen. As fungicide control, Magnate 500 EC® (2 mL/L), a systemic imidazole, was sprayed directly over the injuries. The experiment was conducted using 10 oranges for each condition. The cylinders were closed and sealed with the Scotch® tape and incubated at 24°C . The development of green mold in oranges were evaluated daily, until the oranges in the positive control group were fully covered by the green mold, but no later than 10 days after inoculation. The ability of selected strain to inhibit *P. digitatum* development in fruit was evaluated comparing the diameter of lesion in the treatment with the control.

2.7. Statistics

The results on growth in different media, the antifungal effect against *P. digitatum* on PDA, grains, and on the detached oranges were analyzed according to the last day of evaluation, and normality of the residual distribution was verified (Shapiro and Wilk, 1965). The obtained means were grouped by the Tukey test ($p \leq 0.05$) and the Student *t* test, or the mean pair comparison was conducted when there was a pair of responses due to variable function. For quantitative analysis, regression analyses were conducted, with adjustment of curves and functions by linear, quadratic, and cubic models (Supporting material S4). The R program (Core Team, 2013) was used to implement the functions and to generate the routines of the statistical tests, with the aid of the ExpDes package and Agricolae package (Mendiburu, 2010) for statistical functions implementations. All the experiments above were conducted at least in triplicate.

3. Results and discussion

3.1. Isolation and identification of *Muscodora* isolates

In this study, we present the prospection of two *Muscodora* isolates, LGMF1255 and LGMF1256, obtained from the medicinal plant *S. terebinthifolius* against the causal agent of citrus green-mold, *P. digitatum*.

Two endophytic isolates from leaves of *S. terebinthifolius*, LGMF1255 and LGMF1256, showed morphological characteristics consistent with the genus *Muscodora*, such as ivory mycelium, without reproductive structures (Fig. 1) and with a musty odor. Isolate LGMF1255 showed faster growth on PDA, MEA, and SDA, than MM medium (Fig. 1.1) after 8 days of incubation (Supporting material Figures S1, S2). Using light microscopy, hyaline thin-walled septate hyphae, diameter of 0.45 to $2.04 \mu\text{m}$ ($\bar{x} = 1.05 \mu\text{m}$, $\text{SD} = 0.355 \mu\text{m}$), with right angles between branches and coiling structures in the center (Fig. 1) were observed. SEM showed structures previously described for the *Muscodora* genus, called cauliflower-like or bloom structures (Fig. 1.1 G) and vesicles (Fig. 1.1H, Table S2). It was also possible to notice the hyphal growth interlacing on PDA (Fig. 1.1I) and no specific organization of the mycelium (Fig. 1.1J).

Isolate LGMF1256 had a dense mycelium, with appearance of multiple layers of growth in colonies and irregular edges on PDA. On MEA, its appearance was also dense but more uniform than on PDA (Fig. 1.2A and 1.2C). On SDA (Fig. 1.2D), the colonies showed a regular colony border, with prominent radial lines along the mycelium extension and a dense mycelium. The area of colonies on the eighth day was 4.79 , 4.29 , or 4.24 cm^2 on PDA, MEA, and SDA, respectively. The growth was slower on MM than on the other media, with colony area of 2.35 cm^2 on average, after 8 days (Fig. 1.2B and figure S2). Light

microscopy analysis of isolate LGMF1256 revealed hyaline septate hyphae, thin walls with diameter of 0.79 – $2.19 \mu\text{m}$ ($\bar{x} = 1.45 \mu\text{m}$, $\text{SD} = 0.34 \mu\text{m}$), and straight angles between branches (Fig. 1.2E and 1.2F, Table S2). The hyphae showed interlacing and formed ropelike strands. SEM revealed cauliflower-like structures (Fig. 1.2G), and no specific organization of the mycelium was observed (Figs. 1.2H, 1.2I and 1.2J). Strain LGMF1255 showed the presence of coiling hyphae; unlike in isolate LGMF1256, where these structures were not observed (Table S2). Coiling hyphae are common among *Muscodora* species; however, this characteristic is not associated with the phylogenetic analysis, since it is reported for different species belonging to different phylogenetic groups into *Muscodora* genus (Daisy et al., 2002, Kudalkar et al., 2012, Meshram et al., 2013, Meshram et al., 2014).

3.2. Phylogenetic analysis

ITS sequence analysis included 22 *Muscodora* sequences, with 18 from type strains recognized as legitimate species by the Mycobank database (Mycobank.org) and the outgroup *Xylaria berterii*. The alignment contained 371 characters, with 289 conserved sites, 82 variables and 42 parsimony-informative. Based on ITS sequence analysis, *Muscodora* species were grouped in two clades (Fig. 2), and isolates LGMF1255 and LGMF1256 were in Clade 2. Strain LGMF1255 was found to be close related to species *M. vitigenus*, *M. equiseti*, and *M. sutura* showing 100% sequence similarity with these strains, a cryptic group into *Muscodora* genus, in which it is not possible to perform species identification based on ITS phylogenetic analysis (Pena et al., 2017).

The position of isolate LGMF1256, as a sister species of *M. yucatanensis* and *M. coffeanum*, but located in a separate branch, suggests this strain as a new species in the *Muscodora* genus, named *Muscodora brasiliensis*. All the branches and clades in the phylogenetic analysis are supported by high posterior probability values (> 0.94). The designation of *M. brasiliensis* as a new species is also supported by micro-morphology analysis - hyphae diameter of $2.19 \mu\text{m}$, in comparison to *M. equiseti*, *M. yucatanensis*, and *M. coffeanum* (Table S2) that have greater diameters: 2.9 , 4.0 and $5.5 \mu\text{m}$, respectively (González et al., 2009; Suwannarach et al., 2013; Hongsanan et al., 2015). Names of *Muscodora* species are commonly related with its host (González et al., 2009; Suwannarach et al., 2013; Hongsanan et al., 2015), however we obtained two different species from the same host, consequently, we proposed the name of *M. brasiliensis*, based on the country where it was isolated.

3.3. Identification of volatile organic compounds

Once VOCs were used for *Muscodora* species identification, we analyzed this characteristic of both isolates LGMF1255 and LGMF1256. In *M. brasiliensis* LGMF1256 14 VOCs were detected, and *Muscodora* sp. LGMF1255 analysis yielded 12 chemical compounds overall. All the compounds present in the control were disregarded, as were silylated compounds belonging to the equipment column. Both strains produced 10 compounds in common (Table 1). The compounds produced by both strains were octylformate, 2-phenyl ethyl acetate, 2-undecanone, β -elemene, *n*-tetradecane, α -guaiene, β -selinene, aciphyllene, α -bulnesene, and pogostol. Among them, many already have been found to be produced by other microorganisms or plants, and their bioactive properties have not been described (Strobel et al., 2007, Swamy and Sinniah, 2015). The active compound, phenylethyl alcohol, detected in *M. brasiliensis* LGMF1256 was reported inhibiting the growth of gram-negative microorganisms by interfering with DNA synthesis, cell membranes stability (Corre et al., 1990), and may be related to the inhibition of *P. digitatum*, as observed in our experiments. This compound also is produced by *M. albus* (Strobel et al., 2007) and *M. fenyangensis* (Zhang et al., 2010), strains with antimicrobial activity. The most representative VOC produced by both isolates was pogostol,




Fig. 2. Bayesian phylogenetic tree of *Muscodor* species based on the partial sequences of the ITS region (ITS1-5.8S-ITS2).

Notes: *Xylaria berteri* (GU324749) as outgroup. Scale bar: 0.006 substitutions per nucleotide position. Values at the nodes indicate Bayesian posterior probabilities.

Table 1

Main volatile compounds produced by *Muscodor* sp. LGMF1255 and *Muscodor brasiliensis* LGMF1256 on potato dextrose agar (PDA) at 24 °C after 10 days by SPME-GC-MS analysis.

<i>Muscodor</i> sp. LGMF1255				
RT (min)	AI	Average area (%)	AI ^S -	Possible compound
13.01	1135	7.21	1131	Octylformate
16.807	1224	3.84**	1254	2-phenyl ethyl acetate
19.423	1284	1.48	1293	2-undecanone
22.81	1362	1.13	1359	neoio-dihydrocarveolacetate
23.68	1382	6.12	1390	β-elemene
24.457	1400	1.01	1400	n-tetradecane
24.81	1408	1.28	1409	Z- caryophyllene*
25.68	1429	4.59	1439	α-guaiene
27.603	1475	2.67	1489	β-selinene
28.07	1486	1.54**	1501	Aciphyllene
28.33	1492	6.53	1509	α-bulnesene
33.73	1628	62.60	1651	Pogostol

<i>Muscodor brasiliensis</i> LGMF1256				
RT (min)	AI	Average area (%)	AI ^S -	Possible compound
10.903	1083	16.98	1106	phenylethylalcohol*
13.020	1135	8.15	1131	octyl formate
16.803	1224	12.17	1254	2-phenylethylacetate*
18.950	1273	0.97**	1293	2-undecanone*
23.680	1382	2.73**	1390	β-elemene
24.453	1400	1.24	1400	n-tetradecane
25.677	1429	3.50	1439	α-guaiene*
27.213	1465	3.83	1479	α-curcumene
27.513	1473	3.57	1454	E-(β)-farnesene*
27.610	1475	2.46**	1489	β-selinene
28.073	1486	0.66	1501	Aciphyllene
28.330	1492	5.96	1509	α-bulnesene
28.643	1500	1.28**	1500	n-pentadecane
33.727	1628	36.51	1651	Pogostol

Notes: RTx retention time, AI arithmetic index; AI^S arithmetic index by Adams, 2007;

* Compounds identify by NIST and Adams, 2007.

** Average obtained by two measures; Bold compounds names are produced by *Muscodor* sp. LGMF1255 and *Muscodor brasiliensis* LGMF1256.

isolated first from *Pogostemon cablin*, plant from the family Lamiaceae, compound that has been already used as an antimicrobial agent (Swamy and Sinniah, 2015). Compound α-curcumene, produced by *Curcuma zedoaria*, causes apoptotic effect on ovarian tumor cells (Shin and Lee, 2013), and has been found in some fungi and bacteria (Korpi et al., 2009). However, it is the first report of α-curcumene production by a *Muscodor* species (Table 1), which can also be used to differentiate *M. brasiliensis* from the other species of the *Muscodor* genus.

3.4. Taxonomic description

Muscodor brasiliensis Pena, Serviński & Kava, sp. nov.

Mycobank No.: MB#820079

GenBank accession No.: ITS: KY924494; RPB2: MF510171

Etymology. Named after the country in which it was isolated, Brazil.

The fungus in nature is associated with the plant *Schinus terebinthifolius* and is an Ascomycete with a sterile mycelium. The colonies of *M. brasiliensis* grown on PDA, MEA, and SDA media, with diameters of 4.79, 4.79, and 4.24 cm respectively (Table S2), at 24 °C, 8 days after inoculation. Growth covered the Petri dishes of 9 cm PDA in 3–4 weeks and produced a characteristic odor of the genus that was also detected in the other culture media used. The light microscopy analysis revealed hyaline hyphae, with diameter of 0.79–2.19 μm (\bar{x} = 1.45 μm, SD = 0.34 μm) septate, branched at a right angle, thin-walled. The hyphae were intertwining and forming rope-like strands. No conidia or other sporulation structures were observed. *M. brasiliensis* produces, α-curcumene and n-pentadecane as exclusively VOCs, compared to others *Muscodor* species.

Specimen examined: Brazil, Paraná, Curitiba 25°26'52.5"S 49°14'00.3"W. *Muscodor brasiliensis* is an endophytic fungus from internal tissue of leaves of *Schinus terebinthifolius* (Anacardiaceae) isolated in September 2013 by Serviński, A. & Kava, V. (Holotype: 91050, ex-type culture: LGMF1256).

3.5. Antifungal activity of *Muscodor brasiliensis* against *Penicillium digitatum*

As the VOCs produced by *Muscodor* species are related to high biotechnological potential we decided to analyze the ability of the new




Fig. 3. Antifungal activity of VOCs produced by *Muscodor brasiliensis* LGMF1256, cultivated in different nutritional sources, against *Penicillium digitatum* LGMF1507.

Notes: A–C) positive control: only *P. digitatum* LGMF1507; Antifungal activity of VOCs produced by *M. brasiliensis* in PDA D), oatmeal E) and wheat F). The scale bars on panels represent 1, 2 and 4 cm.

Table 2

Percentage inhibition of radial growth (PIRG) of *Penicillium digitatum* LGMF1507 after 5 days of exposure to volatile organic compounds (VOCs) produced by *Muscodor brasiliensis* LGMF1256 on different nutrient sources.

Medium	PIRG of <i>P. digitatum</i>
Negative control	0 a *
PDA	99.5 b
Wheat	98.4 b
OatMeal	97.9 b
S-W =	0.60***
CV (%) =	2.5

Notes: * Original Medium Averages; ** Means followed by the same letter in the column do not differ by the Tukey test at 5% probability; *** Values S–W in bold mean normal distribution by Shapiro-Wilk at .05 significance level. Negative control represents the evaluation without inoculation of *M. brasiliensis*.

species, *M. brasiliensis* LGMF1256 to control an important post-harvest pathogen, *P. digitatum*. Strains LGMF1255 and LGMF1256, on PDA medium, inhibited *P. digitatum* growth by 70% and 100%, respectively. In view of the great inhibition caused by the VOCs produced by *M. brasiliensis* LGMF1256, this strain was selected for the analysis of VOCs production using wheat or oatmeal as a nutritional source. Similarly, on these substrates, the VOCs produced completely inhibited the phytopathogen growth (Fig. 3, Table 2), data that suggest a great potential to be used in industrial scale (Table 2).

The potential use of *M. brasiliensis* LGMF1256 to control the green mold during fruit transportation was demonstrated in the experiment with detached organic orange fruits artificially infected with *P. digitatum* LGMF1507. *M. brasiliensis* LGMF1256 was able to suppress the green-mold lesion size by approximately 77% in the detached organic oranges (Fig. 4), lesions reached 25.2 cm² in the positive control (Fig. 4C), and 5.9 cm² in the treatment with *M. brasiliensis* LGMF1256 (Fig. 4F; Table S1 - Tukey at the significance level of 0.05%). In the negative control, green-mold symptoms were not observed, and the fungicide treatment completely inhibited the pathogen development.

The reduction (77%) in the severity of green mold symptoms, suggests that LGMF1256 VOCs have great potential to disease control. Similar results were obtained by Suwannarach et al. (2013) with *M. suthpensis* for biofumigation-based to control tangerine rot, also caused

by *P. digitatum*. The use of VOCs to inhibit microbial growth has been studied for different purposes and there are already indications for the use of these compounds to combat citrus diseases. Fialho et al. (2011) reported a high inhibitory activity (87% inhibition) of the VOC ethyl acetate produced by *Saccharomyces cerevisiae* against *Phyllosticta citricarpa*, the causal agent of citrus black spot (CBS). In addition, previous results from our research group demonstrated the potential of one *Muscodor* strain, LGMF1254, suppressing *P. citricarpa* growth (Pena et al., 2017).

The presence and mixture of several types of VOCs (3-methyl-1-butanol, naphthalene and propanoic acid), is necessary for *M. albus* to inhibit the phytopathogens *S. sclerotiorum*, *Pythium ultimum*, and *Rhizoctoniasolani*, indicating the synergism among the different components of the mixture (Ezra et al., 2004). In this view, it is more probably that a mix of compounds, instead of a single compound, can be responsible for the biological activity of *M. brasiliensis*. The use of a mix of compounds is particularly interesting as a treatment of phytopathogens because the compounds may represent different mechanisms operating at the same time, preventing the selection of resistant strains (Tonial et al., 2017).

M. brasiliensis LGMF1256 was able to suppress the green-mold lesion size by approximately 77% in the detached organic oranges (Fig. 4), lesions reached 25.2 cm² in the positive control (Fig. 4C), and 5.9 cm² in the treatment with *M. brasiliensis* LGMF1256 (Fig. 4F; Table S1 - Tukey at the significance level of 0.05%). In the negative control, green-mold symptoms were not observed, and the fungicide treatment completely inhibited the pathogen development.

In conclusion, *Schinus terebinthifolius* is a host of two strains belonging to the *Muscodor* genus: *Muscodor* sp. LGMF1255 and the new species *Muscodor brasiliensis* LGMF1256. The VOCs produced by *M. brasiliensis* were able to significantly decrease the severity of green mold symptoms on oranges, suggesting the potential use of these compounds during the fruits storage and transportation to inhibit green mold development, mainly in organic fruits.

Funding

This research was supported by CNPq [486206/2012-2] and Fundação Araucária [39428-316/2014]. L.C.P. is also grateful to CAPES for the scholarship. V.K. would like to thank to CNPq for her research productivity grant [312316/2016-0].




Fig. 4. Inhibition of green-mold symptoms in detached organic oranges caused by *Muscodora brasiliensis* LGMF1256 VOCs produced in PDA. Notes: A, B, and C: positive control group with *Penicillium digitatum* on the 4th day, 5th day, and 6th day, respectively. D, E, and F: treatment with *M. brasiliensis* LGMF1256 on the 4th day, 5th day, and 6th day, respectively. G, H and I: treatment with the fungicide on the 4th day, 5th day, and 6th day, respectively

Acknowledgements

The authors would like to thank Dr. Gary Strobel for kindly providing the strain *M. sutura*, Breno Cezar Marinho Juliatti for the help with statistics analyses, Rodrigo Aluizio for the technical support, and Dr. Célia Regina Cavichiolo Franco for assistance with SEM analysis.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.micres.2019.01.002>.

References

- Adams, R.P., 2007. Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy. Carol Stream. Allured Publishing Corporation., Illinois, USA.
- Agrios, G.N., 2005. Plant Pathology, fifth ed. Elsevier Academic Press, Burlington.
- CORE TEAM Development, 2013. R: a Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Available at <http://www.R-project.org/>. Accessed in April 2016.
- Corre, J., Lucchini, J.J., Mercier, G.M., Cremieux, A., 1990. Antibacterial activity of phenethyl alcohol and resulting membrane alterations. Microbiol. Res. 141, 483–497.
- Daisy, B., Strobel, G., Ezra, D., Hess, W.M., 2002. *Muscodora vitigenus* sp. nov., an endophyte from *Paullinia paullinioides*. Mycotaxon 84, 39–50.
- Dos Santos, P.J.C., Savi, D.C., Gomes, R.R., Goulin, E.H., Senkiv, C.D.C., Tanaka, F.A., Almeida, A.M., Galli-Terasawa, L., Kava, V., Glienke, C., 2016. *Diaporthe endophytica* and *D. terebinthifolii* from medicinal plants for biological control of *Phyllosticta citricarpa*. Microbiol. Res. 186, 153–160.
- Ezra, D., Hess, W.M., Strobel, G., 2004. New endophytic isolates of *Muscodora albus*, a volatile-antibiotic-producing fungus. Microbiology 150, 4023–4031.
- FAOSTAT, 2015. Food and Agriculture Organization of the United Nations Database. Available at <http://faostat3.fao.org/home/E> Accessed in May 2015.
- Fialho, M.B., Moraes, M.H.D., Tremocoldi, A.M., Pascholati, S.F., 2011. Potential of antimicrobial volatile organic compounds to control *Sclerotinia sclerotiorum* in bean seeds. Pesqui. Agropecu. Bras. 46, 137–142.
- Fischer, I.H., Zanette, M.M., Spósito, M.B., Amorim, L., 2011. Doenças pós-colheita em laranja 'Valência' e caracterização da população fúngica em pomares orgânicos e convencionais. Trop. Plant Pathol. 36, 390–399.
- Gabler, F.M., Mercier, J., Jiménez, J.I., Smilanick, J.L., 2010. Interaction of continuous biofumigation with *Muscodora albus* with pre-cooling fumigation with ozone or sulfur dioxide to control postharvest gray mold of table grapes. Postharv. Biol. Technol. 55, 78–84.
- Gerrits Van Den Ende, A.H.G., De Hoog, G.S., 1999. Variability and molecular diagnostics of neurotropic species *Cladophialophora bantiana*. Stud. Mycol. 43, 151–162.
- González, M.C., Anaya, A.L., Glenn, A.E., Marcias-Rubiacava, M.L., Hernandez-Bautista, B., Hanlin, R.T., 2009. *Muscodora yucatanensis*, new endophytic ascomycete from Mexican chakah, *Bursera simaruba*. Mycotaxon 110, 363–372.
- Hongsanan, S., Hyde, K.D., Bahkali, A.H., Camporesi, E., Chomnunti, P., Ekanayaka, H., Gomes, A.A.M., Hofstetter, V., Gareth Jones, E.B., Pinho, D.B., Pereira, O.L., Tian, Q.,

- Wanasinghe, D.N., Xu, J.C., Buyck, B., 2015. Fungal biodiversity profiles 11–20. *Cryptogamie Mycol.* 36, 355–380.
- Hutchings, M.L., Alpha-Cobb, C.J., Hiller, D.A., Berro, J., Strobel, S.A., 2017. Mycofumigation through production of the volatile DNA methylating agent N-methyl-N-nitrosoisobutyramide by fungi in the genus *Muscodor*. *J. Appl. Biol. Chem.* 1–27.
- Kern, M.E., Blevins, K.S., 1997. *Medical Mycology: A Self-Instructional Text*, second ed. FA Davis, Philadelphia.
- Korpi, A., Jarnberg, J., Pasanen, A.L., 2009. Microbial volatile organic compounds. *Crit. Rev. Toxicol.* 39, 139–193.
- Kudalkar, P., Strobel, G., Ul-Hassan, S.R., Geary, B., 2012. *Muscodor sutura*, a novel endophytic fungus with volatile antibiotic activities. *Mycoscience* 53, 319–325.
- López-Pérez, M., Ballester, A.R., González-Candelas, L., 2015. Identification and functional analysis of *Penicillium digitatum* genes putatively involved in virulence towards citrus fruit. *Mol. Plant Pathol.* 16, 262–275.
- Macarasin, D., Cohen, L., Eick, A., Rafael, G., Belsa, V.E., Wisniewski, M., Droby, S., 2007. *Penicillium digitatum* suppresses production of hydrogen peroxide in host tissue during infection of citrus fruit. *Phytopathol.* 97, 1491–1500.
- Marcet-Houben, M., Ballester, A.R., De La Fuente, B., Harries, E., Marcos, J.F., González-Candelas, L., Gabaldón, T., 2012. Genome sequence of the necrotrophic fungus *Penicillium digitatum*, the main postharvest pathogen of citrus. *BMC Genomics* 13, 646–663.
- Mendiburu, F., 2010. *Agricolae: Statistical Procedures for Agricultural Research. R Package Version 1.0-9*. Available at <http://CRAN.R-project.org/package=agricolae> Accessed in April 2016.
- Meshram, V., Kapoor, N., Saxena, S., 2013. *Muscodor kashayum* sp. nov. – a new volatile antimicrobial producing endophytic fungus. *Mycology* 4, 196–204.
- Meshram, V., Saxena, S., Kapoor, N., 2014. *Muscodor strobilii*, a new endophytic species from South India. *Mycotaxon* 128, 93–104.
- Milind, S.L., 2008. *Citrus Fruit Biology, Technology and Evaluation*, first ed book. Copyright Elsevier Inc. Elsevier, USA Elsevier.
- Pena, L.C., Jung, L.F., Savi, D.C., Serviński, A., Aluizio, R., Galli-Terasawa, L.T., Sales Maia, B.H.L.N., Annes, V., Franco, C.R.C., Glienke, C., Kava, V., 2017. A *Muscodor* strain isolated from *Citrus sinensis* and its production of volatile organic compounds inhibiting *Phyllosticta citricarpa* growth. *J. Plant. Dis. Prot.* 123, 1–12.
- Rasband, W.S., 2016. *ImageJ*, U.S. National Institutes of Health, Bethesda, Maryland, USA. Available at <http://imagej.nih.gov/ij> Accessed in April 2016.
- Saxena, S., Meshram, V., Kapoor, N., 2015. *Muscodor tigerii* sp. nov. -Volatile antibiotic producing endophytic fungus the northeastern Himalayas. *Ann. Microbiol.* 65, 47–57.
- Schnabel, G., Mercier, J., 2006. Use of a *Muscodor albus* pad delivery system for the management of brown rot of peach in shipping cartons. *Postharv. Biol. Technol.* 42, 121–123.
- Shapiro, S.S., Wilk, M.B., 1965. An analysis of variance test for normality (Complete Samples). *Biometrika* 52, 591–611.
- Shin, Y., Lee, Y., 2013. Cytotoxic activity from *Curcuma zedoaria* through mitochondrial activation on ovarian cancer cells. *Toxicol. Res.* 29, 257–261.
- Sperandio, E.M., Martins do Vale, H.M., Moreira, G.A.M., 2015. Yeast from native Brazilian Cerrado plants: occurrence, diversity and use in the biocontrol of citrus green mold. *Fungal Biol.* 119, 984–993.
- Strobel, G.A., Dirske, E., Sears, J., Markworth, C., 2001. Volatile Antimicrobials from *Muscodor albus*, a novel endophytic fungus. *Microbiology* 147, 2943–2950.
- Strobel, G.A., Katreena, K., Hess, W.M., Sears, J., Ezra, D., Vargas, P.N., 2007. *Muscodor albus* E-6, an endophyte of *Guazuma ulmifolia* making volatile antibiotics: isolation, characterization and experimental establishment in host plant. *Microbiology* 1532, 613–620.
- Suwannarach, N., Kumla, J., Bussaban, B., Hyde, K., Matsui, K., Lumyong, S., 2013. Molecular and morphological evidence support four new species in the genus *Muscodor* from northern Thailand. *Ann. Microbiol.* 63, 1341–1351.
- Swamy, M.K., Sinniah, U.R., 2015. A comprehensive review on the phytochemical constituents and pharmacological activities of *Pogostemon cablin* Benth.: an aromatic medicinal plant of industrial importance. *Molecules* 20, 8521–8547.
- Talibi, I., Boubaker, H., Boudyach, E.H., Ben, Ait, Aoumar, A., 2014. Alternative methods for the control of postharvest citrus diseases. *J. Appl. Microbiol.* 117, 1–17.
- Timmer, L.W., Gernsey, S.M., Graham, J.H., 2002. *Plagas Y Enfermedades De Los Cítricos*. APS, USA.
- Tonial, F., Maia, B.H.L.N.S., Gomes-Figueiredo, J.A., Sobottka, A.M., Bertol, C.D., Nepel, A., Savi, D.C., Vicente, V.A., Gomes, R.R., Glienke, C., 2016. Influence of culturing conditions on bioprospecting and the antimicrobial potential of endophytic fungi from *Schinus terebinthifolius*. *Curr. Microbiol.* 72, 173–183.
- Tonial, F., Maia, B.H.L.N.S., Sobottka, A.M., Savi, D.C., Vicente, V.A., Gomes, R.R., Glienke, C., 2017. Biological activity of *Diaporthe terebinthifolii* extracts against *Phyllosticta citricarpa*. *FEMS Microbiol. Lett.* 364. <https://doi.org/10.1093/femsle/fnx026>.
- Van Den Dool, H., Kratz, P.D., 1963. A generalization of the retention index system including linear temperature programmed gas–liquid partition chromatography. *J. Chromatogr.* 11, 463–471.
- White Jr., J.F., Morrow, A.C., 1990. Endophyte-host associations in forage grasses. XII. A fungal endophyte of *Trichachne insular* is belonging to *Pseudocercospora*. *Mycologia* 82, 218–226.
- Worapong, J., Strobel, G., Hess, W.M., 2001. *Muscodor albus* anam. Gen. Et sp. nov., an endophyte from *Cinnamomum zeylanicum*. *Mycotaxon* 79, 67–79.
- Worapong, J., Strobel, G., Daisy, B., Castillo, U.F., Baird, G., Hess, W.M., 2002. *Muscodor roseus* anam. Sp. nov., an endophyte from *Grevillea pteridifolia*. *Mycotaxon* 81, 463–475.
- Zhang, C.L., Wang, G.P., Mao, L.J., Komon-Zelazowska, M., Yuan, Z.L., Lin, F.C., Druzhinina, I.S., Kubicek, C.P., 2010. *Muscodor fengyangensis* sp. nov. from southeast China: morphology, physiology and production of volatile compounds. *Fungal Biol.* 114, 797–808.