



# Molecular mechanisms of *Vibrio parahaemolyticus* pathogenesis

Lingzhi Li<sup>a,b,1</sup>, Hongmei Meng<sup>a,b,1</sup>, Dan Gu<sup>a,b,1,\*</sup>, Yang Li<sup>a,b</sup>, Mengdie Jia<sup>a,b</sup>

<sup>a</sup> Jiangsu Key Laboratory of Zoonosis/Jiangsu Co-Innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Yangzhou University, Yangzhou, 225009, China

<sup>b</sup> Joint International Research Laboratory of Agriculture and Agri-Product Safety/Key Laboratory of Prevention and Control of Biological Hazard Factors (Animal Origin) for Agrifood Safety and Quality, The Ministry of Education of China, Yangzhou University, Yangzhou, 225009, China

## ARTICLE INFO

### Keywords:

*Vibrio parahaemolyticus*  
Virulence factors  
Hemolysins  
T3SS  
T6SS

## ABSTRACT

*Vibrio parahaemolyticus* is a Gram-negative halophilic bacterium that is mainly distributed in the seafood such as fish, shrimps and shellfish throughout the world. *V. parahaemolyticus* can cause diseases in marine aquaculture, leading to huge economic losses to the aquaculture industry. More importantly, it is also the leading cause of seafood-borne diarrheal disease in humans worldwide. With the development of animal model, next-generation sequencing as well as biochemical and cell biological technologies, deeper understanding of the virulence factors and pathogenic mechanisms of *V. parahaemolyticus* has been gained. As a globally transmitted pathogen, the pathogenicity of *V. parahaemolyticus* is closely related to a variety of virulence factors. This article comprehensively reviewed the molecular mechanisms of eight types of virulence factors: hemolysin, type III secretion system, type VI secretion system, adhesion factor, iron uptake system, lipopolysaccharide, protease and outer membrane proteins. This review comprehensively summarized our current understanding of the virulence factors in *V. parahaemolyticus*, which are potentially new targets for the development of therapeutic and preventive strategies.

## 1. Introduction

*Vibrio parahaemolyticus* is the leading cause of diarrheal disease associated with seafood worldwide. Symptoms after ingestion include two to ten days of watery diarrhea, abdominal cramps, nausea, vomiting, and headaches (DePaola et al., 2003; Austin B, 2010). Septicemia following gastroenteritis and wound infections can lead to mortality, particularly for individuals with preexisting liver disease. Antibiotics have been used as the main treatment for *V. parahaemolyticus* infection (Letchumanan et al., 2014; Letchumanan et al., 2016). However, antibiotics may interfere with beneficial microbiota and promote post-antibiotic pathogen expansion (Langdon et al., 2016; Laniro et al., 2016). More importantly, recent studies have shown that majority of *V. parahaemolyticus* isolated from both environmental and clinical settings are multi-drug resistant, highlighting that alternative treatment and prevention strategies are needed (Freire-Moran et al., 2011; Yano et al. (2013); Obaidat et al., 2017; Lee and Raghunath, 2018). Previous studies have shown that the bacteriophage could be used to control *V. parahaemolyticus* (Letchumanan et al., 2016; Zhang et al., 2018a). Targeting virulence determinants has been increasingly

applied for the development of alternative antibacterial treatments. With the development of the first infant rabbit animal model that recapitulate many symptoms in humans, next generation sequencing and biochemical, cell biological technologies, significant advances have been made on the virulence factors and pathogenic mechanisms of *V. parahaemolyticus*. This review articles summarized eight types of virulence factors and the molecular mechanisms by which they contribute to the pathogenesis of *V. parahaemolyticus*.

## 2. Hemolysins

It is currently known that *V. parahaemolyticus* can produce three hemolysins: thermostable direct hemolysin (TDH), TDH-related hemolysin (TRH) and thermolabile hemolysin (TLH) encoded by the *tdh*, *trh* and *tlh* genes, respectively. TDH and TRH are considered to be the main pathogenic factors (Baker-Austin et al., 2010; Mahoney et al., 2010; Paranjpye et al., 2012). Epidemiological studies showed that most of the clinical isolates contained *tdh* and *trh* genes, while very few environmental isolates contained *tdh* and *trh* genes (Theethakaew et al., 2013). However, recently studies also showed that *tdh* and *trh* were

\* Corresponding author at: Jiangsu Key Laboratory of Zoonosis/Jiangsu Co-Innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Yangzhou University, Yangzhou, 225009, China.

E-mail address: [006491@yzu.edu.cn](mailto:006491@yzu.edu.cn) (D. Gu).

<sup>1</sup> These authors contributed equally to this manuscript.

<https://doi.org/10.1016/j.micres.2019.03.003>

Received 11 January 2019; Received in revised form 27 February 2019; Accepted 7 March 2019

Available online 08 March 2019

0944-5013/ © 2019 Elsevier GmbH. All rights reserved.

identified from 48% and 8.3% of the environmental isolates, respectively Gutierrez West et al. (2013), indicating that *tdh* and *trh* genes may also play a role in the bacterial fitness to the environment. TDH is a heat-resistant, pore-forming toxin composed of 156 amino acids (Sakurai et al., 1973; Hamada et al., 2007). Studies have shown that mannitol, high concentrations of sodium chloride, bile acids and amino acids can promote the production of TDH (Osawa et al., 2002). There is only one copy of the *tdh* gene in the *tdh*<sup>+</sup>*trh*<sup>+</sup> strain. In contrast, *tdh2*<sup>+</sup>*RH*<sup>+</sup> strain has two copies of *tdh*: *tdh1* and *tdh2*, both of which encode proteins with hemolytic function (Kishishita et al., 1992). However, *tdh2* has higher expression level than *tdh1* (Shalu et al., 2012). TDH acts directly on red blood cells and has hemolytic activity, enterotoxin activity, cardiotoxicity and cytotoxicity.

The hemolysis process of TDH is mediated by receptors. TDH tightly binds erythrocyte cell membranes and destroys cell membrane and lysosomal membrane in a temperature-dependent manner (Huntley et al., 1993). The transcriptional regulatory factor Hfq inhibits the expression of *tdh* gene (Nakano et al., 2008), while CalR inhibits *tdh2* transcription (Zhang et al., 2017a). The mechanism of cytotoxicity of TDH is different from its hemolytic activity (Naim et al., 2001). High concentration of TDH increases the concentration of calcium ions in intestinal epithelial cells, leading to the opening of the chloride channel, and increased secretion of chloride ions in the intestinal cells (Raimondi et al., 1995; Takahashi et al., 2000a, b; Takahashi et al., 2001). *trh* shares 54.8–68.8% homology with *tdh*, and is present in a small number of clinical isolates (Kishishita et al., 1992). TRH is composed of 189 amino acids, *trh1* and *trh2* are two encoding genes of TRH, and the similarity of *trh1* and *trh2* are 84% (Nishibuchi et al., 1989; Kishishita et al., 1992). Isolate carrying *trh2* does not contain the *tdh* and *trh1* genes, suggesting that *trh2* does not coexist with the *tdh* and *trh1* genes (Nishibuchi et al., 1989). TLH is an atypical phospholipase that can only exert hemolysis in the presence of lecithin (Zhang et al., 2005). Expression of *tlh* is up-regulated in the intestine, but the function and pathogenesis of TLH is still unclear (Gotoh et al., 2010; Broberg et al., 2011). Both clinical isolates and environmental isolates contain *tlh*. Due to the highly conserved of the TLH, and studies showed that *tlh* could be used for detection of *V. parahaemolyticus* (Park et al., 2013). The newly established method Propidium monoazide-Loop mediated isothermal amplification (PMA-LAMP) targeting *tlh*, *tdh* and *trh* gene can rapidly, sensitively and specifically screen the VBNC status of *V. parahaemolyticus* (Zhong et al., 2017). A green fluorescent protein-based TLH antibody has been developed, which provides the possibility to monitor TLH in real time (Chen et al., 2015). Overall, these studies suggested that TDH, TRH and TLH may play an important role in the pathogenesis of *V. parahaemolyticus* during infections in humans, and the combination of these genes could be potentially used for differentiating virulent strains from those that are less virulent.

The study showed that TDHs trigger NLRP3 inflammasome activation, while the activation of NLRP3 could promote the elimination of pathogens (Higa et al., 2013). The TDHs may be used in the evasion of inflammasome sensing at the interaction between host and microbe. Besides, development of strategies that specifically targeting other hemolysins could be potentially used to reduce the virulence of *V. parahaemolyticus*.

### 3. Type III secretion system (T3SS)


T3SS is usually encoded by a 30–40 kb genetic region present either in a plasmid or chromosome as a pathogenicity island (PAI) (Calder et al., 2014a). T3SS is a major virulence factor, which can inject a variety of effector proteins into host cells through a syringe-like transmembrane device. Effectors manipulate modify host's protein function or signaling pathways to exert virulence (Shames and Finlay et al., 2012; Portaliou et al., 2016). T3SS consists of over 30 proteins, which can be divided into four categories: structural proteins, translocators, effector proteins and molecular chaperones (Zhou et al., 2013).

*V. parahaemolyticus* has two sets of T3SSs, which are located on chromosome 1 (T3SS1) and chromosome 2 (T3SS2) (Calder et al., 2014a). T3SS1 mainly contributes to cytotoxicity, while T3SS2 mainly contributes to enterotoxicity (Hiyoshi et al., 2010). Both clinical and environmental isolates contain eight T3SS1 genes (VP1667, VP1690, VP1656, VP1680, VP1686, VP1696, VP1670 and vp1659) including the secretion apparatus, putative translocators or effectors, thus T3SS1 in combination with Kanagawa test (hemolysis test) can be used as a genetic marker for *V. parahaemolyticus* (Paranjpye et al., 2012). T3SS1 mainly affects the biofilm formation, motility and cytotoxicity, and contributes to the survival of *V. parahaemolyticus* in the environment. T3SS2 is involved in the negative regulation of cellular inflammatory response, which is conducive to the process of immune evasion of pathogenic bacteria in the host (Calder et al., 2014a). Analysis of the *V. parahaemolyticus* isolates obtained from the stool samples of 1900 diarrheal patients in Shanghai Hospital found that T3SS1 and T3SS2 were present in all isolates, and only the *trh*<sup>+</sup> strain contained the T3SS2β gene (Li et al., 2017a).

Regulation of T3SS1 in *V. parahaemolyticus*. T3SS1 is regulated by a cascade of regulatory systems, which composes the interaction of ExsA, ExsC, ExsD and ExsE proteins (Fig. 1) (Kodama et al., 2010a). Transcription of T3SS1 is initiated by ExsA (Liu and Thomas et al., 2015), a transcription factor belonging to the AraC family. ExsD is an anti-activator that binds directly to ExsA and inhibits the activity of ExsA. ExsC is an anti-anti-activator that can directly bind ExsD to release ExsA (Zhou et al., 2008, 2010a). ExsE contributes to the negative regulation of T3SS1 and modulation of the adherence phenotype while infection the HeLa cells (Erwin et al., 2012). *in vitro* studies have shown that growth under DMEM condition or contact with epithelial cells, e.g., HeLa or Caco-2 cells, stimulates the expression of T3SS1 via modulating the expression level of ExsC, ExsA, ExsD and their interaction (Zhou et al., 2010b). *in vivo* transcriptome studies also showed that *V. parahaemolyticus* in the small intestine has higher level of T3SS1 transcription than those grown *in vitro* under LB condition (Zhou et al., 2010b). In addition, the HlyU and H-NS can modulate the T3SS1 through regulating the expression of *exsA* (Kodama et al., 2010a; Getz et al., 2018). Besides, the CalR and ToxR protein also involved in the regulation of T3SS1 (Osei-Adjei et al., 2017). Interestingly, recent studies demonstrated that bacterial colonization of *exsD* mutant is reduced comparing to the wild-type, suggesting that the overexpression of T3SS1 impairs bacterial colonization in the small intestine (Hubbard et al., 2016). Both the NLRP3 and NLR4 inflammasomes are activated by T3SS1 in response to *V. parahaemolyticus* infection indicating that T3SS1 effectors appears to be one strategy for bacterial evasion of the host proinflammatory response (Higa et al., 2013). Thus, overexpression of T3SS1 may lead to high level of inflammatory response, which is harmful for bacterial colonization in the small intestine. It remains to be elucidated that what signals in the small intestine modulate T3SS1 expression and how overexpression of T3SS1 impairs its colonization.

Effectors of T3SS1. T3SS1 has four secreted proteins: VopQ, VopS, VPA0450 and VopR (VP1683). VopQ is the most important protein responsible for T3SS1-mediated eukaryotic cytotoxicity. VopQ binds to V-ATPase on lysosomal membrane and changes the proton concentration in the lysosome, which promotes lysosomal cleavage and autophagic vesicle formation, leading to host cell autophagy and cytotoxicity. VopQ can also promote IL-8 secretion by activating p38 in the MAPKs signaling pathway (Matlawska-Wasowska et al., 2010; Shimohata et al., 2011). VopS is a post-translational regulatory protein. It contains a Fic domain that mediates adenylation (AMPylation) of Rho GTPase (RhoA, Rac1 and Cdc42), resulting in its inability to bind to downstream substrate proteins and collapse of the actin cytoskeleton (Casselli et al., 2008).

VPA0450 is an inositol phosphatase that hydrolyzes the phosphate group of 4,5-diphosphosphatidylinositol on the plasma membrane (Broberg et al., 2010). Subsequently, the connection between actin



**Fig. 1.** Regulation of T3SS1 genes in *V. parahaemolyticus*. When culture at DMEM medium or contact the host cell, the ExsC binds ExsD and release the ExsA, then the ExsA protein can directly promote the expression of T3SS1. Besides, the HlyU and H-NS can regulate the expression of *exsA*, while the CalR can inhibit the ExsA to bind T3SS1 promoter.

cytoskeleton and the plasma membrane is disrupted, leading to the formation of bubble in the cytoplasmic membrane, cell rounding and lysis (Broberg et al., 2010). VopR is encoded by locus vp1683 and contains a domain similar to the phosphoinositide domain (BPD). N-terminal domain of VopR binds to phosphoinositide on the host cell membrane. VopR may promote the correct folding of T3SS effectors in the host cells (Broberg et al., 2010; Salomon et al., 2013a). The correct secretion of many effectors requires molecular chaperones. There are currently two T3SS1 specific chaperone proteins: VecA and VPA0451. VecA is a chaperone protein of VopQ. VecA can direct bind to VopQ, which is essential for its secretion and stability in the host cells (Akeda et al., 2009). VPA0451 is a chaperone protein of VPA0450 and required for VPA0450 translocation into the host cell membrane (Waddell et al., 2014).

**Regulation of T3SS2 in *Vibrio parahaemolyticus*.** T3SS2 is only present in the pathogenic *V. parahaemolyticus* isolated from patients with sepsis and other diseases and has a low similarity with other T3SS. Therefore, T3SS2 is a specific secretion system for pathogenic *V. parahaemolyticus* (Li et al., 2017b). The expression of T3SS2 is regulated by two transcriptional regulators (VtrA and VtrB). VtrA acts as a transmembrane protein and upregulates the expression of *vtrB*, and VtrB in turn activates the transcriptional expression of T3SS2 genes. VtrA and VtrC form a barrel-like structure on the cell membrane of *V. parahaemolyticus*. Such barrel-like structure acts as a receptor that can recognize and bind bile salt, leading to VtrB and subsequent T3SS2 gene expression (Kodama et al., 2010b; Li et al., 2016). ToxR protein plays essential roles in the pathogenicity of *V. parahaemolyticus* and contributes to the colonization of *V. parahaemolyticus* in the mammalian intestine (Hubbard et al., 2016). Studies showed that the over-expression of VtrA in the absence of ToxR can restore the expression of T3SS2 (Hubbard et al., 2016), indicating that ToxR controls the expression of T3SS2 through VtrA. It is possible that ToxR enhances the ability of VtrA or the protein-protein interactions between ToxR and VtrA, leading to the activation of VtrB expression and further increase the expression of T3SS2 (Fig. 2).

**Effectors of T3SS2.** The enterotoxicity of T3SS2 is mediated its effectors that destroy the cytoskeleton and manipulate cell signaling transduction. Currently, seven effectors of T3SS2 (VopA, VopT, VopL, VopV, VopC, VopZ and VPA1380) have been identified and characterized (Fig. 3) (Trosky et al., 2004; Kodama et al., 2007; Liverman et al.,

2007; Hiyoshi et al., 2011; Zhang et al., 2012; Zhou et al., 2013; Calder et al., 2014b). VopA and VopZ inhibit immune response by inhibiting different components of the MAPKs signaling pathway (Trosky et al., 2004). VopC, VopL and VopV can act on the actin cytoskeleton in the host cells. VopA is an acetyltransferase that acetylates Ser/Thr residues of MAPKs and blocks MAPKs signaling, thereby inhibiting cell growth and ultimately leading to cell death (Trosky et al., 2004). VopZ inhibits the activation of kinase (TAK1), which in turn inhibits the downstream kinases MAPK and NF- $\kappa$ B signaling pathways (Zhou et al., 2013). It is hypothesized that inhibition of TAK1 activation may lead to disruption of the intestinal epithelial cells and diarrheal disease as deletion of TAK1 can cause intestinal epithelial cell damage, death, and inflammatory response (Zhou et al., 2013). VopT is an ADP ribosyltransferase that modifies Ras GTPase by transferring ADP-ribose from NAD<sup>+</sup> to the substrates. VopT is one of the effectors responsible for T3SS2-mediated cytotoxicity (Kodama et al., 2007). VopC has deamidase/transglutaminase activity. VopC activates Rac1 and Cdc42 by deamidation, leading to the modification of actin cytoskeleton. Therefore, VopC can induce changes in actin cytoskeleton and promote *V. parahaemolyticus* invasion into host cells (Zhang et al., 2012). VopV is a key effector that can bind both actin and filamin with its repeat and C-terminus domains, respectively. More importantly, binding to either actin or filamin is sufficient to mediate its diarrheal disease, suggesting these two actin and filamin binding domains are functionally redundant (Hiyoshi et al., 2011). It remains to be elucidated how binding of VopV to either actin or filamin leads to diarrheal disease. VopL binds to microfilaments to replace nucleation promoting during in microfilament synthesis, leading to the acceleration of nucleation reactions (Liverman et al., 2007). VopL can also neutralize reactive oxygen species (ROS) produced by the nicotinamide dinucleotide phosphate (NADPH) oxidase complex in host cells to facilitate bacteria survival in host cells (de Souza Santos et al., 2017). VPA1380 has a structure similar to the cysteine protease domain in various bacterial toxins and requires inositol hexose phosphate (IP6) as an activator to mediate cytotoxicity (Calder et al., 2014b).

#### 4. Type 6 secretion system (T6SS)

**Structure and function of T6SS in *V. parahaemolyticus*.** T6SS is also a secretory device capable of secreting effectors into host mammalian




Fig. 2. Regulation of T3SS2 in *V. parahaemolyticus*. While there is no bile salt, VtrC binds VtrA, and inhibits expression of *vtrB*. When bile salt is present, it binds VtrA and releases VtrA protein, leading to the expression of *vtrB*, and T3SS2. ToxR regulates the ability of VtrA to activate the expression of *vtrB*.

cells as well as target bacterial cells. Structurally, T6SS is similar T1SS, T3SS and T4SS (Fig. 4). In *V. parahaemolyticus*, there are two T6SSs (T6SS1 and T6SS2 located on chromosome 1 and 2, respectively). Both T6SS1 and T6SS2 can be present in clinical and environmental isolates (Li et al., 2017a). T6SS2 is expressed and is most active under low salt conditions (Salomon et al., 2013b). T6SS consists of a series of components including structural proteins, translocators, secreted proteins, and some other proteins with auxiliary function (Bönemann et al., 2010). The core components of T6SS structural proteins, DotU and IcmF, are located in the cell membrane and stabilize the stability of the plasma membrane (Bönemann et al., 2010). The ClpV ATPase provides energy for the secretory function of T6SS. Typically, ClpV interacts with IcmF to initiate an energy supply system. The T6SS translocation protein Hcp (hemolysin co-regulatory protein) and VgrG (valine-glycine repeat protein G) together constitute a tube structure to secrete proteins, which Hcp is a six-membered ring protein that can be stacked in

the periplasm to form a long tube that transports secreted proteins (Sha et al., 2013). The VgrG protein forms a needle-like spike structure that penetrates the lipoprotein bilayer of the cell membrane and exposes the hidden effector region within the host cell membrane to the cytoplasm. VgrG binds to the target protein in the host cell, inducing physiological response. When the VgrG is detached from Hcp tube, the secreted protein of T6SS is released into the host cell through the Hcp tube (Durand et al., 2014). The T6SS2 in *V. parahaemolyticus* plays a role in bacterial invasion into host cells. VgrG2 of T3SS2 can also induce autophagy in macrophage RAW264.7 (Yu et al., 2015). STK (a serine/threonine kinase) and STP (a serine/threonine phosphatase) regulate the biological activity of T6SS by phosphorylation and dephosphorylation of Fha (Mougous et al., 2007).

Regulation of T6SS. T6SS in *V. parahaemolyticus* is regulated by quorum sensing (QS) system. OpaR and AphA are the core regulators of QS that are highly expressed at high bacterial density and low bacterial





Fig. 3. Function of T3SS2 effectors. Seven effectors of T3SS2 (VopA, VopV, VopL, VopT, VopC, VopZ, VPA1380) have been characterized. The signaling pathways that are hijacked or impaired by these effectors are shown.



**Fig. 4.** Structure of Type VI secretion system. IcmF and DotU is the core element of T6SS and located at the inner membrane to stabilize the stability of the plasma membrane. ClpV interacts with the IcmF to provide the energy, and the STK can regulate the T6SS by phosphorylation and dephosphorylation of Fha. VgrG can be secreted to the host cell through the Hcp tube.

density, respectively. At low cell densities, the regulatory factor LuxO is active, inhibiting the activity of OpaR and activating AphA. At high cell densities, the activity of LuxO is restricted, resulting in inhibition of AphA activity and activation of OpaR (Kalburge et al. (2017)). OpaR can inhibit the transcription of *hcp1* and negatively regulate T6SS1, but positively regulate T6SS2 (Ma et al., 2012). AphA has a negative regulatory effect on both T6SS1 and T6SS2 (Wang et al., 2013; Zhang et al., 2017c). H-NS is a nucleic acid binding protein and can bind to AT-rich DNA sequences, thereby inhibiting gene transcription. H-NS inhibits the expression of both T6SS1 and T6SS2 (Salomon et al., 2014; Sun et al., 2014). CalR (a member of LysR family) can positively regulate the expression of T6SS1 and T6SS2. CalR is an antagonistic factor of H-NS and can activate the transcription of the T6SS2 gene by competing for the binding site of the H-NS to the promoter region of the target genes (Zhang et al., 2017b). As a membrane-bound virulence-regulating protein of *V. parahaemolyticus*, ToxR works together with AphA and OpaR to inhibit the activity of T6SS1. The expression of *toxR* gene is closely related to QS. In addition, T6SS1 has the highest expression level between when OD600 is between 0.6 and 0.8, indicating that the virulence of *V. parahaemolyticus* is inextricably linked with the bacterial QS (Zhang et al., 2017c).

## 5. Adhesion factors


The adhesion of pathogens is closely related to their pathogenicity. Microbial adhesion to host tissues is the initial event during the process of infection (Navarre et al., 1999). The main factors that are associated with *V. parahaemolyticus* adhesion to host cells are hemagglutinin, enolase and T6SS2 (Yamamoto et al., 1989; Jiang et al., 2014; Zhang et al., 2017). Studies have shown that mannose-sensitive hemagglutinin (MSHA) is an important factor affecting bacterial-host cell adhesion and pathogenicity. MSHA has a high affinity for polysaccharides such as sialic acid and GM1 ganglioside, indicating that MSHA might be receptors for MSHA in the gut, promoting their colonization to intestinal epithelial cells (O'Boyle et al. (2013)). In *V. parahaemolyticus*, the

enolase is a plasminogen-binding, surface-exposed protein. Enolase could facilitate bacterial adherence to Hep-2 cells, which could promote pathogen-host interaction (Jiang et al., 2014). Besides, the CalR could regulate the *V. parahaemolyticus* adhesion by regulating the expression of T6SS2 (Zhang et al., 2017). The sigma factor RpoN is a regulatory protein that controls the synthesis of flagella. The *rpoN* mutant has a higher colonization in the intestine than the wild-type, indicating that RpoN may significantly affect the colonization of *V. parahaemolyticus* (Whitaker et al., 2014). The adhesion factor MAM-7, which is widely distributed in Gram-negative bacteria, binds to fibronectin and membrane phosphatidic acid on the host cell membrane to assemble a three-molecular polymer, which mediates pathogen adhesion to host cells. At the initial stage, the expression of MAM-7 in non-pathogenic lactic acid bacteria can make them adhere to host cells (Beltran et al., 2016). Inhibition of MAM-7 expression can reduce the adhesion and virulence of *V. parahaemolyticus* (Krachler et al., 2011). This phenomenon is also present in other Gram-negative bacteria containing MAM-7, suggesting that MAM-7 is a good vaccine or therapeutic target to prevent and treat infections caused by Gram-negative bacteria. Deeper comprehensive understanding of the adhesion mechanism could provide the new perspective to control the colonization of *V. parahaemolyticus* in the host and interrupt the first step of infection.

## 6. Iron uptake system

A recent study found that metal ions play an important role in the regulation of gene expression in *V. parahaemolyticus*. High calcium and low iron growth conditions stimulate group behavior and T3SS transcription (Gode-Potratz et al., 2010). Iron is an indispensable element in the growth and metabolism of some pathogens. The iron in the host cells is mainly found in red blood cells, lactoferrin and transferrin, and there are very few free iron ions, so it cannot meet the demand of iron for pathogens. There are two main ways for pathogenic *Vibrio* to obtain iron: one is to produce exotoxin to destroy red blood cells and release hemoglobin; the other is to produce a low molecular weight iron chelating agent, which has high affinity for iron ions in heme (Yamamoto et al., 1994; Kustusich et al., 2011). The iron uptake system in *V. parahaemolyticus* consists of an iron membrane-bound receptor and an iron chelate. *V. parahaemolyticus* can produce a carrier, called vibrioferrin, capable of chelation of irons, and the resulting iron-carrier complex can be transported to cells via the bacterial outer membrane protein receptor for iron assimilation (Yamamoto et al., 1995). Under the iron sufficient condition, Fur, as a DNA-binding protein, can inhibit the transcription of genes associated with the iron uptake system (Tanabe et al., 2013). Studies have shown that iron overload in humans increases the virulence of pathogens (Bullen et al., 1991; Berlutti et al., 2005; Pieracci et al., 2005). In humans, iron is found mainly in hemoglobin (Hb), heme, hemin, and iron chelators (Tf, Lf, or Ft). As an iron carrier, vibrioferrin binds to the iron-binding protein of the host cell and is recognized by the receptors PvuA1 and PvuA2 on the outer membrane of *V. parahaemolyticus* cells (Tanabe et al., 2003). Heme and ferritin (Hb) can also be recognized by these receptors. PvuA can bind the ABC transport system, the PvuBCDE system in the inner membrane, which transports iron-Vibrioferrin to the inner membrane (Tanabe et al., 2011). In addition, *V. parahaemolyticus* also contains the TonB system, which is composed of TonB, TonB1 and TonB2. Iron-Vibrioferrin is transported to the inner membrane by PvuA1 by the energy supplied by TonB1. Energy supply of PvuA2 system depends on TonB1 and TonB2 (Kuehl et al., 2010). *V. parahaemolyticus* can also utilize siderophores produced by other bacteria, such as exogenous aerobactin, desferri-ferrichrome, and enterobactin (Funahashi et al., 2003, 2009; Tanabe et al., 2012).

*V. parahaemolyticus* can secrete an iron-reducing protein, which reduces the ferric ion to divalent iron. Besides, *V. parahaemolyticus* can also degrade hemoglobin (Hb) by secreting a protease to obtain iron from the humans (León-Sicairos et al., 2015). In addition, some outer



**Fig. 5.** The molecular mechanism of iron uptake system. The iron uptake system contains an iron membrane-bound receptor and an iron chelate in the outer membrane. Iron-Vibrioferrin can be transported to the inner membrane by PvuBCDE system, TonB system or ABC system. The DNA-binding protein Fur can regulate the expression of genes associated with the iron uptake system.

membrane protein receptors are important for the acquisition of iron. VctA and IrgA are receptors for enterobactin, FhuA is a receptor for desferri-ferrichrome and aerobactin, and HutA and HutR are receptors for heme (Tanabe et al., 2012). This outer membrane protein receptor transports the iron-iron carrier into the periplasm and binds to the periplasmic binding protein. This process requires TonB, ExbB, and ExbD on the inner membrane to provide energy and ultimately transport it to the interior of the cell via the ABC system (Fig. 5) (O'Malley et al. (1999); Kustusch et al., 2011; Tanabe et al., 2011, 2012).

## 7. Lipopolysaccharide (LPS)

LPS is the main substance of bacterial endotoxin, and one of the main components of the cell wall in Gram-negative bacteria. LPS consists of three parts: core polysaccharide, lipid A and O antigen. The formation of extracellular polysaccharide (EPS) in biofilm plays a crucial role in bacterial adhesion to living organisms and non-living organisms. Transcription and translation of cpsA-J operon are controlled by ScrABC, extracellular c-di-GMP concentration, CpsS-CpsR-CpsQ regulator and quorum sensing. H-NS can positively regulate the transcription of cpsA-J to promote EPS production (Zhang et al., 2018b). Bandekar et al. have shown that LPS of *V. parahaemolyticus* has an effect on murine peritoneal macrophages (Bandekar and Nerkar (1988)). By increasing the dose of LPS, RNA content and lysosomal activity of peritoneal macrophages can be significantly increased. Studies have found that the recombinant protein LVLGBP can be used as a pattern recognition protein (PRP) to recognize and bind LPS and activate innate immune response in shrimp (Valli et al., 2012; Chen et al., 2016).

## 8. Proteases

Protease secreted to the outside the cell is closely related to the pathogenesis of *Vibrio*. For example, in *Vibrio cholerae*, hemagglutinin protease can not only activate the A subunit of cholera enterotoxin, but also separate the bacteria from the host cell membrane and infect other host cells (Finkelstein et al., 1992). The *V. parahaemolyticus* strain also secretes hemagglutinin protease, but its mechanism of action is unclear.

Lee et al obtained a protease A from a *V. parahaemolyticus* clinical isolate that does not contain *tdh* and *trh* genes and found that purified protease A has obvious toxic effects on experimental cells and has lytic activity on red blood cells (Lee et al., 2002). It can also cause tissue hemolysis, and death in mice in severe cases. Thus, protease A is a virulence factor of *V. parahaemolyticus*. Kim et al. cloned and expressed the metalloproteinase gene in *V. parahaemolyticus* and found that this protease has high homology with the collagenase of *Vibrio alginolyticus* (Kim et al., 2002). Since *V. alginolyticus* collagenase plays a role in infection, it is hypothesized that metalloproteinase may have a similar effect in *V. parahaemolyticus* infection. In addition, studies have found that in addition to metalloproteinases, extracellular serine proteases contribute to the pathogenicity of *Vibrio vulnificus* (Wang et al., 2008). In *V. parahaemolyticus*, PrtA acts as a serine protease with hemolytic activity and cytotoxicity (Osei-Adjei et al., 2018). The quorum sensing regulator (OprA) and general stress response sensor (RpoS) in *V. parahaemolyticus* can regulate PrtA synthesis pathway by binding to the promoter of *prtA* gene. Both OprA and RpoS can up-regulate PrtA (Chang et al., 2018).

## 9. Outer membrane proteins (OMPs)

*Vibrio* OMPs are located at the outer membrane surface and thus can interact extensively with the outside of the cell (Li et al., 2014). LptD is an important outer membrane protein that plays an important role in mediating the transport of LPS to the outer leaflet. Studies have shown that LptD is widely distributed, protective, and has surface antigenic activity in the major *Vibrio* genus, indicating that LptD is involved in immune responses and has a strong ability to stimulate antibody responses (Zha et al., 2016). OMP is considered to be a potential candidate vaccine. To facilitate the screening and identification of potential candidate vaccines, immunoproteomics techniques based on two-dimensional electrophoresis and immunoblotting can be used to screen for immunogenic OMP (Pan et al., 2016).

## 10. Conclusion and prospective

As a halophilic food-borne pathogen, *V. parahaemolyticus* have various virulence factors responsible for hemolytic activity, cytotoxicity, intestinal toxicity, gastroenteritis, diarrhea, sepsis and even death. At present, the main virulence factors include hemolytic toxin, adhesin, protease, outer membrane protein, lipopolysaccharide, urease, type III (T3SS) and type VI (T6SS) secretion systems. In addition, the pathogenicity of *V. parahaemolyticus* is the result of a combination of multiple virulence factors, and thus it is necessary to comprehensively consider various virulence factors and the regulatory networks. Further studies on the mechanisms by which the environmental signals integrate into the genetic circuit to regulate the expression of these virulence factors would provide novel strategies to prevent and treat infection caused by *V. parahaemolyticus*.

## References

- Akeda, Y., Okayama, K., Kimura, T., Dryselius, R., Kodama, T., Oishi, K., Iida, T., Honda, T., 2009. Identification and characterization of a type III secretion-associated chaperone in the type III secretion system 1 of *Vibrio parahaemolyticus*. *FEMS Microbiol. Lett.* 296 (1), 18–25.
- Austin, B., 2010. Vibrios as causal agents of zoonoses. *Vet. Microbiol.* 140, 310–317.
- Baker-Austin, C., Stockley, L., Rangdale, R., Martinez-Urtaza, J., 2010. Environmental occurrence and clinical impact of *Vibrio vulnificus* and *Vibrio parahaemolyticus*, a European perspective. *Environ. Microbiol. Rep.* 2 (1), 7–18.
- Bandekar, J.R., Nerkar, D.P., 1988. Biological activities of lipid A from *Vibrio parahaemolyticus*, stimulation of murine peritoneal macrophages. *Microbiol. Immunol.* 32 (3), 275–282.
- Beltran, S., Munoz-Bergmann, C.A., Elola-Lopez, A., Quintana, J., Segovia, C., Trombert, A.N., 2016. The expression of heterologous MAM-7 in *Lactobacillus rhamnosus* reduces its intrinsic capacity to inhibit colonization of pathogen *Vibrio parahaemolyticus* *in vitro*. *Biol. Res.* 49, 2.
- Berlutti, F., Morea, C., Battistoni, A., Sarli, S., Cipriani, P., Superti, F., Ammendolia, M.G., Valenti, P., 2005. Iron availability influences aggregation, biofilm, adhesion and invasion of *Pseudomonas aeruginosa* and *Burkholderia cenocepacia*. *Int. J. Immunopathol. Pharmacol.* 18 (4), 661–670.
- Bönemann, G., Pietrosiuk, A., Mogk, A., 2010. Tubules and donuts, a type VI secretion story. *Mol. Microbiol.* 76 (4), 815–821.
- Broberg, C.A., Zhang, L., Gonzalez, H., Laskowski-Arce, M.A., Orth, K., 2010. A *Vibrio* effector protein is an inositol phosphatase and disrupts host cell membrane integrity. *Science* 329, 1660–1662.
- Broberg, C.A., Calder, T.J., Orth, K., 2011. *Vibrio parahaemolyticus* cell biology and pathogenicity determinants. *Microbes Infect.* 13, 992–1001.
- Bullen, J.J., Spalding, P.B., Ward, C.G., Gutteridge, J.M., 1991. Hemochromatosis, iron and septicemia caused by *Vibrio vulnificus*. *Arch. Intern. Med.* 151 (8), 1606–1609.
- Calder, T., de Souza Santos, M., Attah, V., Klimko, J., Fernandez, J., Salomon, D., Krachler, A.M., Orth, K., 2014a. Structural and regulatory mutations in *Vibrio parahaemolyticus* type III secretion systems display variable effects on virulence. *FEMS Microbiol. Lett.* 361 (2), 107–114.
- Calder, T., Kinch, L.N., Fernandez, J., Salomon, D., Grishin, N.V., Orth, K., 2014b. *Vibrio* type III effector VPA1380 is related to the cysteine protease domain of large bacterial toxins. *PLoS One* 9 (8), e104387.
- Casselli, T., Lynch, T., Southward, C.M., Jones, B.W., Devinney, R., 2008. *Vibrio parahaemolyticus* inhibition of Rho family GTPase activation requires a functional chromosome I type III secretion system. *Infect. Immun.* 76, 2202–2211.
- Chang, S.C., Lee, C.Y., 2018. OpaR and RpoS are positive regulators of a virulence factor PrtA in *Vibrio parahaemolyticus*. *Microbiology* 164 (2), 221–231.
- Chen, Y., Huang, X., Wang, R., Wang, S., Shi, N., 2015. The structure of a GFP-based antibody (fluorobody) to TLH, a toxin from *Vibrio parahaemolyticus*. *Acta Crystallogr. F Struct. Biol. Commun.* 71 (7), 913–918.
- Chen, Y.Y., Chen, J.C., Kuo, Y.H., Lin, Y.C., Chang, Y.H., Gong, H.Y., Huang, C.L., 2016. Lipopolysaccharide and beta-1,3-glucan-binding protein (LGBP) bind to seaweed polysaccharides and activate the phenoloxidase system in white shrimp *Litopenaeus vannamei*. *Dev. Comp. Immunol.* 55, 144–151.
- De Souza Santos, M., Salomon, D., Orth, K., 2017. T3SS effector VopL inhibits the host ROS response, promoting the intracellular survival of *Vibrio parahaemolyticus*. *PLoS Pathog.* 13 (6), e1006438.
- DePaola, A., Ulaszek, J., Kaysner, C.A., Tenge, B.J., Nordstrom, J.L., Wells, J., Puh, N., Gendel, S.M., 2003. Molecular, serological and virulence characteristics of *Vibrio parahaemolyticus* isolated from environmental, food, and clinical sources in North America and Asia. *Appl. Environ. Microbiol.* 69, 3999–4005.
- Durand, E., Cambillau, C., Cascales, E., Journet, L., 2014. VgrG, Tae, Tle, and beyond, the versatile arsenal of Type VI secretion effectors. *Trends Microbiol.* 22 (9), 498–507.
- Finkelstein, R.A., Boesman-Finkelstein, M., Chang, Y., Häse, C.C., 1992. *Vibrio cholerae* hemagglutinin/protease, colonial variation, virulence, and detachment. *Infect. Immun.* 60 (2), 472–478.
- Freire-Moran, L., Aronsson, B., Manz, C., Gyssens, I.C., So, A.D., Monnet, D.L., Cars, O., 2011. Critical shortage of new antibiotics in development against multidrug-resistant bacteria—time to react is now. *Drug Resist. Updates* 14, 118–124.
- Funahashi, T., Tanabe, T., Aso, H., Nakao, H., Fujii, Y., Okamoto, K., Narimatsu, S., Yamamoto, S., 2003. An iron-regulated gene required for utilization of aerobactin as an exogenous siderophore in *Vibrio parahaemolyticus*. *Microbiology* 149 (5), 1217–1225.
- Funahashi, T., Tanabe, T., Shiuchi, K., Nakao, H., Yamamoto, S., 2009. Identification and characterization of genes required for utilization of desferri-ferrichrome and aerobactin in *Vibrio parahaemolyticus*. *Biol. Pharm. Bull.* 32 (3), 359–365.
- Getz, L.J., Thomas, N.A., 2018. The transcriptional regulator HlyU positively regulates expression of *exsA*, leading to type III secretion system 1 activation in *Vibrio parahaemolyticus*. *J. Bacteriol.* 200 (15), e00653–17.
- Gode-Potratz, C.J., Chodur, D.M., McCarter, L.L., 2010. Calcium and iron regulate swarming and type III secretion in *Vibrio parahaemolyticus*. *J. Bacteriol.* 192 (22), 6025–6038.
- Gotoh, K., Kodama, T., Hiyoshi, H., Izutsu, K., Park, K.S., Dryselius, R., Akeda, Y., Honda, T., Iida, T., 2010. Bile acid-induced virulence gene expression of *Vibrio parahaemolyticus* reveals a novel therapeutic potential for bile acid sequestrants. *PLoS One* 5, e13365.
- Gutierrez West, C.K., Klein, S.L., Lovell, C.K., 2013. High frequency of virulence factor genes *tdh*, *trh*, and *tli* in *Vibrio parahaemolyticus* strains isolated from a pristine estuary. *Appl. Environ. Microbiol.* 79 (7), 2247–2252.
- Hamada, D.T., Higurashi, K., Mayanagi, T., Miyata, T., Fukui, T., Iida, T., Honda, Yanagihara, I., 2007. Tetrameric structure of thermostable direct hemolysin from *Vibrio parahaemolyticus* revealed by ultracentrifugation, small-angle X-ray scattering and electron microscopy. *J. Mol. Biol.* 365, 187–195.
- Naomi Higa, N., Toma, C., Koizumi, Y., Nakasone, N., Nohara, T., Masumoto, J., Kodama, T., Iida, T., Suzuki, T., 2013. *Vibrio parahaemolyticus* effector proteins suppress inflammation activation by interfering with host autophagy signaling. *PLoS Pathog.* 9 (1), e1003142.
- Hiyoshi, H., Kodama, T., Iida, T., Honda, T., 2010. Contribution of *Vibrio parahaemolyticus* virulence factors to cytotoxicity, enterotoxicity, and lethality in mice. *Infect. Immun.* 78 (4), 1772–1780.
- Hiyoshi, H., Kodama, T., Saito, K., Gotoh, K., Matsuda, S., Akeda, Y., Honda, T., Iida, T., 2011. VopV, an F-actin-binding type III secretion effector, is required for *Vibrio parahaemolyticus* induced enterotoxicity. *Cell Host Microbe* 10, 401–409.
- Hubbard, T.P., Chao, M.C., Abel, S., Blondel, C.J., Abel Zur Wiesch, P., Zhou, X., Davis, B.M., Waldor, M.K., 2016. Genetic analysis of *Vibrio parahaemolyticus* intestinal colonization. *PNAS* 113 (22), 6283–6288.
- Huntley, J.S., Hall, A.C., 1993. The effects of membrane active agents on the leak induced in erythrocyte membranes by Kanagawa haemolysin. *Biochem. Soc. Trans.* 21, 2195.
- Jiang, W., Han, X., Wang, Q., Li, X., Yi, L., Liu, Y., Ding, C., 2014. *Vibrio parahaemolyticus* enolase is an adhesion-related factor that binds plasminogen and functions as a protective antigen. *Appl. Microbiol. Biotechnol.* 98 (11), 4937–4948.
- Kalburge, S.S., Carpenter, M.R., Rozovsky, S., Boyd, E.F., 2017. Quorum sensing regulators are required for metabolic fitness in *Vibrio parahaemolyticus*. *Infect. Immun.* 85 (3), e00930–16.
- Kim, S.K., Yang, J.Y., Cha, J., 2002. Cloning and sequence analysis of a novel metalloprotease gene from *Vibrio parahaemolyticus* 04. *Gene* 283 (1–2), 277–286.
- Kishishita, M., Matsuoka, N., Kumagai, K., Yamasaki, S., Takeda, Y., Nishibuchi, M., 1992. Sequence variation in the thermostable direct hemolysin-related hemolysin (*trh*) gene of *Vibrio parahaemolyticus*. *Appl. Environ. Microbiol.* 58, 2449–2457.
- Kodama, T., Rokuda, M., Park, K.S., Cantarelli, V.V., Matsuda, S., Iida, T., Honda, T., 2007. Identification and characterization of VopT, a novel ADP-ribosyltransferase effector protein secreted via the *Vibrio parahaemolyticus* type III secretion system 2. *Cell. Microbiol.* 9, 2598–2609.
- Kodama, T., Yamazaki, C., Park, K.S., Akeda, Y., Iida, T., Honda, T., 2010a. Transcription of *Vibrio parahaemolyticus* T3SS1 genes is regulated by a dual regulation system consisting of the ExsACDE regulatory cascade and H-NS. *FEMS Microbiol. Lett.* 311 (1), 10–17.
- Kodama, T., Gotoh, K., Hiyoshi, H., Morita, M., Izutsu, K., Akeda, Y., Park, K.S., Cantarelli, V.V., Dryselius, R., Iida, T., Honda, T., 2010b. Two regulators of *Vibrio parahaemolyticus* play important roles in enterotoxicity by controlling the expression of genes in the Vp-PAI region. *PLoS One* 5, e8678.
- Krachler, A.M., Orth, K., 2011. Functional characterization of the interaction between bacterial adhesion multivalent adhesion molecule 7 (MAM7) protein and its host cell ligands. *J. Biol. Chem.* 286 (45), 38939–38947.
- Kuehl, C.J., Crosa, J.H., 2010. The TonB energy transduction systems in *Vibrio* species. *Future Microbiol.* 5, 1403–1412.
- Kustusch, R.J., Kuehl, C.J., Crosa, J.H., 2011. Power plays, iron transport and energy transduction in pathogenic vibrios. *Biomaterials* 24, 559–566.
- Langdon, A., Crook, N., Dantas, G., 2016. The effects of antibiotics on the microbiome throughout development and alternative approaches for therapeutic modulation. *Genome Med.* 8, 39.
- Laniro, G., Tilg, H., Gasbarrini, A., 2016. Antibiotics as deep modulators of gut microbiota: between good and evil. *Gut* 65, 1906–1915.
- Lee, L.-H., Raghunath, P., 2018. Editorial: vibronaceae diversity, multidrug resistance and management. *Front. Microbiol.* 9, 563.
- Lee, C.Y., Cheng, M.F., Yu, M.S., Pan, M.J., 2002. Purification and characterization of a putative virulence factor, serine protease, from *Vibrio parahaemolyticus*. *FEMS Microbiol. Lett.* 209 (1), 31–37.
- León-Sicairos, N., Angulo-Zamudio, U.A., de la Garza, M., Velázquez-Román, J., Flores-Villaseñor, H.M., Canizalez-Román, A., 2015. Strategies of *Vibrio parahaemolyticus* to acquire nutritional iron during host colonization. *Front. Microbiol.* 6, 702.
- Letchumanan, V., Chan, K.-G., Lee, L.-H., 2014. *Vibrio parahaemolyticus*: a review on the pathogenesis, prevalence, and advance molecular identification techniques. *Front. Microbiol.* 5, 705.
- Letchumanan, V., Chan, K.G., Pusparajah, P., Saokaew, S., Duangjai, A., Goh, B.H.,

- Mutalib, N.-S., Lee, L.-H., 2016. Insights into bacteriophage application in controlling *Vibrio* species. *Front. Microbiol.* 7, 1114.
- Li, C., Ye, Z., Wen, L., Chen, R., Tian, L., Zhao, F., Pan, J., 2014. Identification of a novel vaccine candidate by immunogenic screening of *Vibrio parahaemolyticus* outer membrane proteins. *Vaccine* 32 (46), 6115–6121.
- Li, P., Rivera-Cancel, G., Kinch, L.N., Salomon, D., Tomchick, D.R., Grishin, N.V., Orth, K., 2016. Bile salt receptor complex activates a pathogenic type III secretion system. *Elife* 5 (5).
- Li, P., Kinch, L.N., Ray, A., Dalia, A.B., Cong, Q., Nunan, L.M., Camilli, A., Grishin, N.V., Salomon, D., Orth, K., 2017a. Acute hepatopancreatic necrosis disease-causing *Vibrio parahaemolyticus* strains maintain an antibacterial type VI secretion system with versatile effector repertoires. *Appl. Environ. Microbiol.* 83 (13) e00737-17.
- Li, H., Tang, R., Lou, Y., Cui, Z., Chen, W., Hong, Q., Zhang, Z., Malakar, P.K., Pan, Y., Zhao, Y., 2017b. A comprehensive epidemiological research for clinical *Vibrio parahaemolyticus* in Shanghai. *Front. Microbiol.* 8, 1043.
- Liu, A.C., Thomas, N.A., 2015. Transcriptional profiling of *Vibrio parahaemolyticus* *sexA* reveals a complex activation network for type III secretion. *Front. Microbiol.* 6, 1089.
- Liverman, A.D., Cheng, H.C., Trosky, J.E., Leung, D.W., Yarbrough, M.L., Burdette, D.L., Rosen, M.K., Orth, K., 2007. Arp2/3-independent assembly of actin by *Vibrio* type III effector VopL. *PNAS* 104, 17117–17122.
- Ma, L., Zhang, Y., Yan, X., Guo, L., Wang, L., Qiu, J., Yang, R., Zhou, D., 2012. Expression of the type VI secretion system 1 component Hcp1 is indirectly repressed by OpaR in *Vibrio parahaemolyticus*. *Sci. World J.* 2012, 982140.
- Mahoney, J.C., Gerding, M.J., Jones, S.H., Whistler, C.A., 2010. Comparison of the pathogenic potentials of environmental and clinical *Vibrio parahaemolyticus* strains indicates a role for temperature regulation in virulence. *Appl. Environ. Microbiol.* 76 (22), 7459–7465.
- Matlawska-Wasowska, K., Finn, R., Mustel, A., O'Byrne, C.P., Baird, A.W., Coffey, E.T., Boyd, A., 2010. The *Vibrio parahaemolyticus* type III secretion systems manipulate host cell MAPK for critical steps in pathogenesis. *BMC Microbiol.* 10, 329.
- Mougous, J.D., Gifford, C.A., Ramsdell, T.L., Mekalanos, J.J., 2007. Threonine phosphorylation post-translationally regulates protein secretion in *Pseudomonas aeruginosa*. *Nat. Cell Biol.* 9, 797–803.
- Naim, R., Iida, T., Takahashi, A., Honda, T., 2001. Monodansylcadaverine inhibits cytotoxicity of *Vibrio parahaemolyticus* thermostable direct hemolysin on cultured rat embryonic fibroblast cells. *FEMS Microbiol. Lett.* 196 (2), 99–105.
- Nakano, M., Takahashi, A., Su, Z., Harada, N., Mawatari, K., Nakaya, Y., 2008. Hfq regulates the expression of the thermostable direct hemolysin gene in *Vibrio parahaemolyticus*. *BMC Microbiol.* 8, 155.
- Navarre, W.W., Schneewind, O., 1999. Surface proteins of gram-positive bacteria and mechanisms of their targeting to the cell wall envelope. *Microbiol. Mol. Biol. Rev.* 63 (1), 174–229.
- Nishibuchi, M., Taniguchi, T., Misawa, T., Khaeomane-Iam, V., Honda, T., Miwatani, T., 1989. Cloning and nucleotide sequence of the gene (*trh*) encoding the hemolysin related to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. *Infect. Immun.* 57 (9), 2691–2697.
- O'Boyle, N., Houeix, B., Kilcoyne, M., Joshi, L., Boyd, A., 2013. The MSHA pilus of *Vibrio parahaemolyticus* has lectin functionality and enables TTSS-mediated pathogenicity. *Int. J. Med. Microbiol.* 303 (8), 563–573.
- O'Malley, S.M., Mouton, S.L., Occhino, D.A., Deanda, M.T., Rashidi, J.R., Fuson, K.L., Rashidi, C.E., Mora, M.Y., Payne, S.M., Henderson, D.P., 1999. Comparison of the heme iron utilization systems of pathogenic *Vibriosis*. *J. Bacteriol.* 181, 3594–3598.
- Obaidat, M.M., Salaman, A.E.B., Roess, A.A., 2017. Virulence and antibiotic Resistance of *Vibrio parahaemolyticus* isolates from seafood from three worldwide environmental, seafood, and clinical isolates from 2000 to 2017. *J. Food Prot.* 80 (12), 2060–2067.
- Osawa, R., Arakawa, E., Okitsu, T., Yamai, S., Watanabe, H., 2002. Levels of thermostable direct hemolysin produced by *Vibrio parahaemolyticus* O3,K6 and other serovars grown anaerobically with the presence of a bile acid. *Curr. Microbiol.* 44 (4), 302–305.
- Osei-Adjei, G., Gao, H., Zhang, Y., Zhang, L., Yang, W., Yang, H., Yin, Z., Huang, X., Zhang, Y., Zhou, D., 2017. Regulatory actions of ToxR and CalR on their own genes and type III secretion system 1 in *Vibrio parahaemolyticus*. *Oncotarget.* 8 (39), 65809–65822.
- Osei-Adjei, G., Huang, X., Zhang, Y., 2018. The extracellular proteases produced by *Vibrio parahaemolyticus*. *World J. Microbiol. Biotechnol.* 34 (5), 68.
- Pan, J., Li, C., Ye, Z., 2016. Immunoproteomic approach for screening vaccine candidates from bacterial outer membrane proteins. *Methods Mol. Biol.* 1404, 519–528.
- Paranjpye, R., Hamel, O.S., Stojanovski, A., Liermann, M., 2012. Genetic diversity of clinical and environmental *Vibrio parahaemolyticus* strains from the Pacific Northwest. *Appl. Environ. Microbiol.* 78 (24), 8631–8638.
- Park, J.Y., Jeon, S., Kim, J.Y., Park, M., Kim, S., 2013. Multiplex real-time polymerase chain reaction assays for simultaneous detection of *Vibrio cholerae*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*. *Osong Public Health Res. Perspect.* 4 (3), 133–139.
- Pieracci, F.M., Barie, P.S., 2005. Iron and the risk of infection. *Surg Infect. (Suppl)*, S41–46.
- Portaliou, A.G., Tsois, K.C., Loos, M.S., Zorzini, V., Economou, A., 2016. Type III secretion, building and operating a remarkable nanomachine. *Trends Biochem. Sci.* 41, 175–189.
- Raimondi, F., Kao, J.P., Kaper, J.B., Guandalini, S., Fasano, A., 1995. Calcium-dependent intestinal chloride secretion by *Vibrio parahaemolyticus* thermostable direct hemolysin in a rabbit model. *Gastroenterology* 109 (2), 381–386.
- Sakurai, J., Matsuzaki, A., Miwatani, T., 1973. Purification and characterization of the thermostable direct hemolysin of *Vibrio parahaemolyticus*. *Infect. Immun.* 8 (5), 775–780.
- Salomon, D., Guo, Y.R., Kinch, L.N., Grishin, N.V., Gardner, K.H., Orth, K., 2013a. Effectors of animal and plant pathogens use a common domain to bind host phosphoinositides. *Nat. Commun.* 4, 2973.
- Salomon, D., Gonzalez, H., Updegraff, B.L., Orth, K., 2013b. *Vibrio parahaemolyticus* type VI secretion system 1 is activated in marine conditions to target bacteria, and is differentially regulated from system 2. *PLoS One* 8, e61086.
- Salomon, D., Klimko, J.A., Orth, K., 2014. H-NS regulates the *Vibrio parahaemolyticus* type VI secretion system 1. *Microbiology* 160 (9), 1867–1873.
- Sha, J., Rosenzweig, J.A., Kozlova, E.V., Wang, S., Erova, T.E., Kirtley, M.L., van Lier, C.J., Chopra, A.K., 2013. Evaluation of the roles played by Hcp and VgrG type 6 secretion system effectors in *Aeromonas hydrophila* SSU pathogenesis. *Microbiology* 159 (6), 1120–1135.
- Shalu, O.A., Pisanov, R.V., Monakhova, E.V., 2012. Effectiveness of expression of *tdh* gene of *Vibrio parahaemolyticus* depends on two point mutations in promoter region. *Genetika* 48 (12), 1364–1371.
- Shames, S.R., Finlay, B.B., 2012. Bacterial effector interplay, a new way to view effector function. *Trends Microbiol.* 20, 214–219.
- Shimohata, T., Nakano, M., Lian, X., Shigeyama, T., Iba, H., Hamamoto, A., Yoshida, M., Harada, N., Yamamoto, H., Yamato, M., Mawatari, K., Tamaki, T., Nakaya, Y., Takahashi, A., 2011. *Vibrio parahaemolyticus* infection induces modulation of IL-8 secretion through dual pathway via VP1680 in Caco-2 cells. *J. Infect. Dis.* 203, 537–544.
- Sun, F., Zhang, Y., Qiu, Y., Yang, H., Yang, W., Yin, Z., Wang, J., Yang, R., Xia, P., Zhou, D., 2014. H-NS is a repressor of major virulence gene loci in *Vibrio parahaemolyticus*. *Front. Microbiol.* 5, 675.
- Takahashi, A., Kenjyo, N., Imura, K., Myonsun, Y., Honda, T., 2000a. Cl(-) secretion in colonic epithelial cells induced by the *Vibrio parahaemolyticus* hemolytic toxin related to thermostable direct hemolysin. *Infect. Immun.* 68 (9), 5435–5438.
- Takahashi, A., Sato, Y., Shiomi, Y., Cantarelli, V.V., Iida, T., Lee, M., Honda, T., 2000b. Mechanisms of chloride secretion induced by thermostable direct haemolysin of *Vibrio parahaemolyticus* in human colonic tissue and a human intestinal epithelial cell line. *J. Med. Microbiol.* 49 (9), 801–810.
- Takahashi, A., Iida, T., Naim, R., Naykaya, Y., Honda, T., 2001. Chloride secretion induced by thermostable direct haemolysin of *Vibrio parahaemolyticus* depends on colonic cell maturation. *J. Med. Microbiol.* 50 (10), 870–878.
- Tanabe, T., Funahashi, T., Nakao, H., Miyoshi, S., Shinoda, S., Yamamoto, S., 2003. Identification and characterization of genes required for biosynthesis and transport of the siderophore vibrioferrin in *Vibrio parahaemolyticus*. *J. Bacteriol.* 185, 6938–6949.
- Tanabe, T., Funahashi, T., Okajima, N., Nakao, H., Takeuchi, Y., Miyamoto, K., Tsujibo, H., Yamamoto, S., 2011. The *Vibrio parahaemolyticus* *pvuA1* gene (formerly termed *psuA*) encodes a second ferric vibrioferrin receptor that requires *tonB2*. *FEMS Microbiol. Lett.* 324, 73–79.
- Tanabe, T., Funahashi, T., Shiuchi, K., Okajima, N., Nakao, H., Miyamoto, K., Tsujibo, H., Yamamoto, S., 2012. Characterization of *Vibrio parahaemolyticus* genes encoding the systems for utilization of enterobactin as a xenosiderophore. *Microbiology* 158, 2039–2049.
- Tanabe, T., Funahashi, T., Nakao, H., Maki, J., Yamamoto, S., 2013. The *Vibrio parahaemolyticus* small RNA RyhB promotes production of the siderophore vibrioferrin by stabilizing the polycistronic mRNA. *J. Bacteriol.* 195 (16), 3692–3703.
- Theethakaew, C., Feil, E.J., Castillo-Ramírez, S., Aanensen, D.M., Suthienkul, O., Neil, D.M., Davies, R.L., 2013. Genetic relationships of *Vibrio parahaemolyticus* isolates from clinical, human carrier, and environmental sources in Thailand, determined by multilocus sequence analysis. *Appl. Environ. Microbiol.* 79 (7), 2358–2370.
- Trosky, J.E., Mukherjee, S., Burdette, D.L., Roberts, M., McCarter, L., Siegel, R.M., Orth, K., 2004. Inhibition of MAPK signaling pathways by VopA from *Vibrio parahaemolyticus*. *J. Biol. Chem.* 279, 51953–51957.
- Valli, J.S., Vaseeharan, B., 2012. cDNA cloning, characterization and expression of lipopolysaccharide and b-1,3-glucan binding protein (LGBP) gene from the Indian white shrimp *Fenneropenaeus indicus*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 163 (1), 74–81.
- Waddell, B., Southward, C.M., McKenna, N., Devinney, R., 2014. Identification of VPA0451 as the specific chaperone for the *Vibrio parahaemolyticus* chromosome 1 type III-secreted effector VPA0450. *FEMS Microbiol. Lett.* 353 (2), 141–150.
- Wang, J., Sasaki, T., Maehara, Y., Nakao, H., Tsuchiya, T., Miyoshi, S., 2008. Variation of extracellular proteases produced by *Vibrio vulnificus* clinical isolates, genetic diversity of the metalloprotease gene (*vvp*), and serine protease secretion by *vvp*-negative strains. *Microb. Pathog.* 44 (6), 494–500.
- Wang, L., Wang, L., Zhou, D., Mao, P., Zhang, Y., Hou, J., Hu, Y., Li, J., Hou, S., Yang, R., Wang, R., Qiu, J., 2013. Cell density- and quorum sensing-dependent expression of type VI secretion system 2 in *Vibrio parahaemolyticus*. *PLoS One* 8 (8), e73363.
- Whitaker, W.B., Richards, G.P., Boyd, E.F., 2014. Loss of sigma factor RpoN increases intestinal colonization of *Vibrio parahaemolyticus* in an adult mouse model. *Infect. Immun.* 82 (2), 544–556.
- Yamamoto, T., Yokota, T., 1989. Adherence targets of *Vibrio parahaemolyticus* human small intestines. *Infect. Immun.* 57 (8), 2410–2419.
- Yamamoto, S., Okujo, N., Matsuura, S., Fujiwara, I., Fujita, Y., Shinoda, S., 1994. Siderophore-mediated utilization of iron bound to transferrin by *Vibrio parahaemolyticus*. *Microbiol. Immunol.* 38, 687–693.
- Yamamoto, S., Hara, Y., Tomochika, K., Shinoda, S., 1995. Utilization of heme and hemoglobin as iron sources by *Vibrio parahaemolyticus* and identification of an iron-repressible heme-binding protein. *FEMS Microbiol. Lett.* 128, 195–200.
- Yano, Y., Hamano, K., Satomi, M., Tsutsui, I., Ban, M., Aue-umneoy, M., 2013. Prevalence and antimicrobial susceptibility of *Vibrio* species related to food safety isolated from shrimp cultured at inland ponds in Thailand. *Food Control* 38, 30–36.
- Yu, Y., Fang, L., Zhang, Y., Sheng, H., Fang, W., 2015. VgrG2 of type VI secretion system 2 of *Vibrio parahaemolyticus* induces autophagy in macrophages. *Front. Microbiol.* 6, 168.

- Zha, Z., Li, C., Li, W., Ye, Z., Pan, J., 2016. LptD is a promising vaccine antigen and potential immunotherapeutic target for protection against *Vibrio* species infection. *Sci. Rep.* 6, 38577.
- Zhang, H.Z., Austin, B., 2005. Haemolysins in *Vibrio* species. *J. Appl. Microbiol.* 98, 1011–1019.
- Zhang, L., Krachler, A.M., Broberg, C.A., Li, Y., Mirzaei, H., Gilpin, C.J., Orth, K., 2012. Type III effector VopC mediates invasion of *Vibrio* species. *Cell Rep.* 1, 453–460.
- Zhang, Y., Zhang, Y., Gao, H., Zhang, L., Yin, Z., Huang, X., Zhou, D., Yang, H., Yang, W., Wang, L., 2017a. *Vibrio parahaemolyticus* CalR down regulates the thermostable direct hemolysin (TDH) gene transcription and thereby inhibits hemolytic activity. *Gene* 613, 39–44.
- Zhang, L., Osei-Adjei, G., Zhang, Y., Gao, H., Yang, W., Zhou, D., Huang, X., Yang, H., Zhang, Y., 2017b. CalR is required for the expression of T6SS2 and the adhesion of *Vibrio parahaemolyticus* to HeLa cells. *Arch. Microbiol.* 199 (6), 931–938.
- Zhang, Y., Gao, H., Osei-Adjei, G., Zhang, Y., Yang, W., Yang, H., Yin, Z., Huang, X., Zhou, D., 2017c. Transcriptional regulation of the type VI secretion system 1 genes by quorum sensing and ToxR in *Vibrio parahaemolyticus*. *Front. Microbiol.* 8, 2005.
- Zhang, H., Yang, Z.Q., Zhou, Y., Bao, H.D., Wang, R., Li, T.W., Pang, M.D., Sun, L.C., Zhou, X.H., 2018a. Application of a phage in decontaminating *Vibrio parahaemolyticus* in oysters. *Int. J. Food Microbiol.* 275, 24–31.
- Zhang, L., Weng, Y., Wu, Y., Wang, X., Yin, Z., Yang, H., Yang, W., Zhang, Y., 2018b. H-NS is an activator of exopolysaccharide biosynthesis genes transcription in *Vibrio parahaemolyticus*. *Microb. Pathog.* 116, 164–167.
- Zhong, H., Zhong, Y., Deng, Q., Zhou, Z., Guan, X., Yan, M., Hu, T., Luo, M., 2017. Virulence of thermolabile haemolysin *tlh*, gastroenteritis related pathogenicity *tdh* and *trh* of the pathogens *Vibrio parahaemolyticus* in Viable but Non-Culturable (VBNC) state. *Microb. Pathog.* 111, 352–356.
- Zhou, X., Shah, D.H., Konkel, M.E., Call, D.R., 2008. Type III secretion system 1 genes in *Vibrio parahaemolyticus* are positively regulated by ExsA and negatively regulated by ExsD. *Mol. Microbiol.* 69 (3), 747–764.
- Zhou, X., Konkel, M.E., Call, D.R., 2010a. Regulation of type III secretion system 1 gene expression in *Vibrio parahaemolyticus* is dependent on interactions between ExsA, ExsC and ExsD. *Virulence* 1 (4), 260–272.
- Zhou, X., Konkel, M.E., Call, D.R., 2010b. Vp1659 is a *Vibrio parahaemolyticus* type III secretion system 1 protein that contributes to translocation of effector proteins needed to induce cytolysis, autophagy, and disruption of actin structure in HeLa cells. *J. Bacteriol.* 192 (13), 3491–3502.
- Zhou, X., Gewurz, B.E., Ritchie, J.M., Takasaki, K., Greenfeld, H., Kieff, E., Davis, B.M., Waldor, M.K., 2013. A *Vibrio parahaemolyticus* T3SS effector mediates pathogenesis by independently enabling intestinal colonization and inhibiting TAK1 activation. *Cell Rep.* 3, 1690–1702.