

- complicated chronic distal aortic dissections: intermediate outcomes and complications. *J Thorac Cardiovasc Surg* 2011;142:1074–83.
- 17 Thrumurthy SG, Karthikesalingam A, Patterson BO, Holt PJ, Hinchliffe RJ, Loftus IM, et al. A systematic review of mid-term outcomes of thoracic endovascular repair (TEVAR) of chronic type B aortic dissection. *Eur J Vasc Endovasc Surg* 2011;42:632–47.
  - 18 Rohlfs F, Tsilimparis N, Fiorucci B, Heidemann F, Debus ES, Kölbel T. The Candy-plug technique: technical aspects and early results of a new endovascular method for false lumen occlusion in chronic aortic dissection. *J Endovasc Ther* 2017;24:549–55.
  - 19 Etz CD, Weigang E, Hartert M, Lonn L, Mestres CA, Di Bartolomeo R, et al. Contemporary spinal cord protection during thoracic and thoracoabdominal aortic surgery and endovascular aortic repair: a position paper of the vascular domain of the European Association for Cardio-Thoracic Surgery. *Eur J Cardiothorac Surg* 2015;47:943–57.
  - 20 Katsargyris A, Oikonomou K, Kouvelos G, Renner H, Ritter W, Verhoeven EL. Spinal cord ischemia after endovascular repair of thoracoabdominal aortic aneurysms with fenestrated and branched stent grafts. *J Vasc Surg* 2015;62:1450–6.
  - 21 Maurel B, Delclaux N, Sobocinski J, Hertault A, Martin-Gonzalez T, Moussa M, et al. The impact of early pelvic and lower limb reperfusion and attentive peri-operative management on the incidence of spinal cord ischemia during thoracoabdominal aortic aneurysm endovascular repair. *Eur J Vasc Endovasc Surg* 2015;49:248–54.
  - 22 Spear R, Hertault A, Van Calster K, Settembre N, Delloye M, Azzaoui R, et al. Complex endovascular repair of postdissection arch and thoracoabdominal aneurysms. *J Vasc Surg* 2018;67:685–93.
  - 23 Mastracci TM, Carrell T, Constantinou J, Dias N, Martin-Gonzalez T, Katsargyris A, et al. Editor's choice - Effect of branch stent choice on branch-related outcomes in complex aortic repair. *Eur J Vasc Endovasc Surg* 2016;51:536–42.


*Eur J Vasc Endovasc Surg* (2019) 57, 109

## COUP D'OEIL

### 'If You Fall Down, You Have to Climb up Again'

Konstantinos Spanos\*, Athanasios D. Giannoukas

Vascular Surgery Department, University Hospital of Larissa, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa, Greece



A 69 year old male presented at the outpatient clinic concerned about being able to palpate his aneurysmal pulsation. He had had endovascular repair five years previously with a 34 mm Anaconda stent graft (Terumo Aortic, Inchinnan, UK) at another vascular centre, but failed to attend follow up appointments for personal reasons. Computed tomography angiography (CTA) revealed caudal migration of the stent graft towards the iliac bifurcation (panel A). Redo aneurysm exclusion was successfully achieved with a four fenestration stent graft (Zenith, Cook Aortic Intervention, Bloomington, IN, USA) with all splanchnic vessels patent with good fixation of the stent graft and no endoleak on follow up (panel B).

\* Corresponding author.

E-mail address: [spanos.kon@gmail.com](mailto:spanos.kon@gmail.com) (Konstantinos Spanos).

1078-5884/© 2018 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2018.10.027>