

Is Socioeconomic Status a Fundamental Cause of Racial Differences in End-of-Life Care Use? (TH321D)



Sarah Cross, LMSW MPH, Duke University, Durham, NC.

Objectives

- Articulate how social theories can help our understanding of inequities in EOLC use.
- Describe social characteristics associated with hospice use, life support use, and hospital death.

Original Research Background. Racial differences in end-of-life care (EOLC) are well documented. De-medicalized care is now the innovation at EOL. Sociological theories suggests those with more education should be the first to use palliative care/less aggressive care and this will lead to inequities in EOLC.

Research Objectives. To determine whether socioeconomic status (SES), as measured by level of education, mediates the relationship between race and three commonly accepted indicators of quality EOLC: hospice use, life support use, and place of death.

Methods. I used modified poisson regression to analyze Health and Retirement Study 2014 Core and Exit Interviews data. The HRS is a longitudinal study surveying a representative sample of Americans age 50 and older. The Exit Interview collects data for respondents who have died during the study. Separate analyses were run for each outcome (hospice use n=1,193; life support use n=893; place of death n=1,198).

Results. In unadjusted models, Blacks were less likely to use hospice (unadjusted risk ratio [RR] =.813; 95% CI=.679-.974; P=0.023), more likely to use life support use (unadjusted RR=1.48; 95% CI=1.18-1.86; P=0.001), and more likely to have a hospital death (uRR=1.29; 95% CI=1.06-1.57; P=0.013) than Whites. After adjusting for education and patient characteristics, Black race was no longer a significant predictor of hospice use (aRR=.993; 95% CI=.826-1.19; P=0.940) or hospital death (aRR=1.17; 95% CI=.942-1.46; P=0.151). Blacks were still significantly more likely to use life support (aRR=1.41; 95% CI=1.09-1.82; P=0.008) than Whites.

Conclusion. Race is significantly associated with indicators of poor quality EOLC. Higher levels of education are significantly associated with a greater risk of hospice use and a lesser likelihood of life support use and hospital death. SES appears to be a fundamental cause of racial inequities in hospice use, but not in life support use or hospital death.

Implications for Research, Policy, or Practice. EOLC quality indicators may ignore cultural heterogeneity of patient preferences. Efforts

should be made to better educate persons of lower SES groups about hospice.

Association of Timing and Type of Advance Care Planning Documentation on End-Of-Life Care for Patients with Serious Illness (TH322A)



Seelwan Sathitratanaheewin, MD, Chulalongkorn University, Bangkok, Thailand.

Objectives

- Describe the association between earlier advance care planning documentation and intensity of care at the end of life.
- Describe the association between late advance care planning documentation and intensity of care at the end of life.

Original Research Background. It is unclear how the timing of advance care planning (ACP) documentation and type of documentation influence the intensity of end-of-life care for patients with chronic illness.

Research Objectives. To examine associations between timing and type of ACP documentation with intensity of end-of-life care for patients with chronic illness receiving care in one healthcare system.

Methods. We developed a retrospective cohort using Washington State death certificates and electronic health records (EHR) to identify patients with chronic illnesses who died between 2010 and 2015. Association between timing and type of ACP documentation and end-of-life care were examined with multivariate probit regression analysis, adjusted for patient characteristics.

Results. Of 22,100 eligible decedents, 6,660 (30%) had an ACP document in the EHR and most (69%) were completed before the last six months of life, with 11% completed in the last 30 days of life. POLST forms (13%) were the least common ACP document, with increasingly higher documentation of health care directives (14%) and DPOA (20%). Earlier completion of ACP specifying treatment limitations were associated with reduced intensity of end-of-life care, whereas completion in the last 30 days of life was associated with increased intensity. Treatment-limiting POLST before the last 180 days had the strongest influence on end-of-life care, followed by treatment-limiting POLST between the last 30-180 days of life and healthcare directives before the last 180 days of life.

Conclusion. Treatment-limiting ACP documentation was associated with reduced intensity of care when completed more than 180 days prior to death.

However, treatment-limiting ACP documentation in the last 30 days of life was associated with higher intensity care, suggesting an inverse causal relationship that many of these documents are completed during hospitalization and that higher intensity care is leading to increased ACP documentation.

Implications for Research, Policy, or Practice. These findings are important for healthcare systems when exploring the association between ACP and intensity of care.

What Do Adolescents Want? Values, Goals, and Beliefs of Teens with Cancer (TH322B) 

Sarah Friebert, MD FAAP FAAHPM, Akron Children's Hospital, Akron, OH. Jessica Gaines, BSN RN CPN, Children's National Health System, Washington, D.C. Jennifer Needle, MD MPH, University of Minnesota, Minneapolis, MN. Justin Baker, MD FAAP FAAHPM, St. Jude Children's Hospital, Memphis, TN. Yao Cheng, MD, Children's National Health System, Washington, D.C. Maureen Lyon, PhD, Children's National Health System, Washington, D.C.

Objectives

- Describe patient-reported palliative care needs of teens with cancer.
- Discuss implications of patient-reported outcomes for family-centered advance care (ACP) planning.

Original Research Background. The National Cancer Institute's (NCI) 2016 recommendations advise parents to have open and honest communication about cancer with their children. Parents are interested in their adolescents-patient's voice. Best timing and strategies to structure and facilitate this communication is unknown.

Research Objectives. This study examined patient-identified palliative care needs, goals, and values of teens with cancer.

Methods. Surveyed adolescents with cancer randomized to a pediatric ACP intervention using the Lyon Advance Care Planning Survey-Adolescent Version-Revised.

Results. Adolescents' (N=45) mean age 17 years (range $\geq 14 < 21$ years); 39% male; 81% white. Though 91% felt that being able to complete an advance directive (AD) was important, most teens had never talked about EOL care wishes. 96% would want their family to be involved in EOL decisions. Problems rated as worse than death were: not being able to communicate wishes to family, 58%; living with great pain, 42%; and total physical dependency, 22%. At EOL, adolescents desire honest answers from physicians (100%), being at peace spiritually (98%), physical comfort (93%), feeling strongly about being able to stay in own home (73%), understanding treatment choices (98%), saying everything I want to say to people in my family (100%), not being a burden to loved ones (89%), and knowing how to say good bye (91%). 56% of teens want to die at home with or without hospice and 9% in hospital. 58% of teens preferred to have ACP conversations early (when healthy, first diagnosed).

Conclusion. Communicating EOL wishes to their family is very important to teens with cancer, consistent with ACS recommendations. Crucial information surrounding ADs and EOL wishes can be gained from teens with cancer.

Implications for Research, Policy, or Practice. Structured, adolescent/family-centered, evidence-based ACP interventions are one way to facilitate open communication about their cancer with their families.

Differences by Race, Religion, and Mental Health in Preferences for Life-Prolonging Treatment in Adverse Health States: Results from a National Sample of Medicare Beneficiaries (TH322C) 

Justin Sanders, MD MSC, Dana-Farber Cancer Institute, Boston, MA. Anna Berrier, BSPH (C), University of North Carolina at Chapel Hill, Chapel Hill, NC. Leonce Nshuti, MS, Vanderbilt University Medical Center, Nashville, TN. Charlotta Lindvall, MD PhD, Dana-Farber Cancer Institute, Boston, MA. James Tulsky, MD FACP FAAHPM, Dana-Farber Cancer Institute, Boston, MA.

Objectives

- Discuss factors that predict patient preferences for life-prolonging treatment in the setting of adverse health states that some would deem intolerable relative to death.
- Consider hypotheses to explain findings regarding preferences for life-prolonging treatment, including those from behavioral economics that might apply across multiple findings.
- Incorporate strategies to effectively elicit patient preferences.

Original Research Background. Goal-concordant care aligns patients' preferences with their medical treatments and is important for patients with serious illness, whose treatments may hasten death or prolong suffering. We lack population-level data on patient preferences, which can help prepare clinicians for advance care planning or goals of care discussions.

Research Objectives. To understand factors that underlie individual preferences regarding life-prolonging treatment in the setting of two adverse health scenarios using a nationally representative sample of Medicare beneficiaries.

Methods. Using the National Health and Aging Trends Study, we used descriptive statistics and multivariable logistic regression to compare sociodemographic and illness characteristics of patients who said they would accept or reject life-prolonging treatments in the setting of severe, constant pain and inability to walk, talk, or recognize others.

Results. Patients in all demographic groups were more likely than not to express a preference for