

- 26 Azria D, Magne N, Zouhair A, Castadot P, Culine S, Ychou M, et al. Radiation recall: a well recognized but neglected phenomenon. *Cancer Treat Rev* 2005;31:555–70.
- 27 Olszewski WL. Lymphovenous microsurgical shunts in treatment of lymphedema of lower limbs: a 45-year experience of one surgeon/one center. *Eur J Vasc Endovasc Surg* 2013;45:282–90.


Eur J Vasc Endovasc Surg (2019) 57, 867

## COUP D'OEIL

# A Roll for Guidewires in an Internal Iliac Artery Aneurysm

Mara Fanelli<sup>\*</sup>, Tiziano Tecchio

Vascular Surgery, Department of Medicine and Surgery, University of Parma, Parma, Italy



An 80 year old man underwent endovascular aneurysm repair of a 50 mm asymptomatic abdominal aortic aneurysm associated with a 73 mm asymptomatic left internal iliac artery aneurysm (LIIAA) with a bifurcated endograft (Anaconda Vascutek; Terumo, Inchinnan, UK). The LIIAA was embolised with 15 hydrophilic floppy guidewires (0.035" Glidewire 180 cm; Terumo). This is not a standard approach in the authors' centre, but was considered in this case as it was effective in providing long wire lengths, and thus the necessary coiling volume, and was also cheaper (€37/wire). This image represents the good treatment result on the two year computed tomography angiogram.

<sup>\*</sup> Corresponding author. Vascular Surgery, Department of Medicine and Surgery, University of Parma, 14 via Gramsci, 43126 Parma, Italy.

E-mail address: [mara.fanelli@libero.it](mailto:mara.fanelli@libero.it) (Mara Fanelli).

1078-5884/© 2019 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2019.02.013>