

A Clinical Validation Study of Anatomical Risk Scoring for Procedural Stroke in Patients Treated by Carotid Artery Stenting in the International Carotid Stenting Study

Djurre D. de Waard^{a,†}, Evelien E. de Vries^{a,†}, Anne E. Huibers^a, Mechteld M. Arnold^a, Paul J. Nederkoorn^b, Lukas C. van Dijk^c, Aad van der Lugt^d, Peter J. Koudstaal^e, Leo H. Bonati^f, Martin M. Brown^g, Gert J. de Borst^{a,*}

^a Department of Vascular Surgery, University Medical Centre Utrecht, Utrecht, the Netherlands

^b Department of Neurology, Academic Medical Centre, Amsterdam, the Netherlands

^c Department of Radiology, Haga Hospital, The Hague, the Netherlands

^d Department of Radiology & Nuclear Medicine, Erasmus MC, University Medical Centre Rotterdam, the Netherlands

^e Department of Neurology, Erasmus MC, University Medical Centre Rotterdam, the Netherlands

^f Department of Neurology and Stroke Centre, Department of Clinical Research, University Hospital Basel, Basel, Switzerland

^g Stroke Research Group, UCL Queen Square Institute of Neurology, London, UK

WHAT THIS PAPER ADDS

Although several anatomical risk factors have been suggested to influence procedural difficulty of carotid artery stenting (CAS), no validated risk score is available to help select suitable patients for CAS. The Delphi anatomical risk (DAR) score was developed to predict the difficulty of CAS in relation to procedural stroke risk, and hereby aid in patient selection. This is the first attempt to validate clinically the DAR score within this *post hoc* analysis of the International Carotid Stenting Study. No significant association between the DAR score and procedural stroke risk was found. However, the small sample size potentially rendered the study underpowered and the results should be replicated in the future.

Objectives: Vascular anatomy of the aortic arch and supra-aortic arteries has been suggested as influencing the risk of carotid artery stenting (CAS). The expert opinion based Delphi anatomical risk (DAR) score was developed to predict difficulty of CAS in relation to procedural stroke risk, and thereby aid patient selection. The aim was to validate the DAR score in the context of a randomised clinical trial.

Methods: In this *post hoc* analysis of the International Carotid Stenting Study (ICSS), only patients treated by CAS with available pre-procedural CT angiography (CTA) were included. Patients with tortuous anatomy unsuitable for stenting were excluded from ICSS. CTA based vascular anatomy was rated by two independent observers. Every possible combination of anatomy resulted in a risk score, divided in four categories of expected risk (low, < 5.0; low–intermediate, 5.0–5.9; high–intermediate, 6.0–6.9; high, ≥ 7.0). Binomial logistic regression was used to assess the relationship between anatomical risk score and procedural risk of any stroke. Differences between predefined age groups were also assessed.

Results: A total of 275 patients were included. Interobserver reliability for all anatomical risk factors was high ($\kappa = 0.76–0.84$). In total, 16 strokes (6%) occurred in the procedural period. No significant relationship was observed between the DAR score and risk of procedural stroke, with the risk of stroke being 9% in the high risk vs. 4% in the low risk categories ($p = .49$). A higher mean DAR score was observed in patients ≥70 years compared with younger patients (4.6 ± 1.5 vs. 3.9 ± 1.4 , $p < .001$), which was mainly explained by higher rates of arch atheroma (44% vs. 20%, $p < .001$). Prolonged intervention duration was significantly associated with increased stroke risk (11% vs. 4%, $p < .04$), but not with the DAR score.

Conclusions: No statistically significant association was found between anatomical difficulty, as defined in the DAR score, and procedural stroke risk. However, the small sample size potentially rendered the study underpowered to detect group differences, and confirmation with a larger sample is essential.

Keywords: Carotid artery stenting, Procedural stroke, Risk score, Vascular anatomy

Article history: Received 21 September 2018, Accepted 24 April 2019, Available online 25 September 2019

© 2019 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

[†] Both authors contributed equally.

* Corresponding author. Department of Vascular Surgery, University Medical Centre Utrecht, PO Box 85500, Utrecht, the Netherlands.

E-mail address: G.J.deBorst-2@umcutrecht.nl (Gert J. de Borst).

1078-5884/© 2019 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2019.04.035>

INTRODUCTION

Extracranial atherosclerotic carotid disease is an important risk factor for stroke. Carotid artery stenting (CAS) has gained popularity in the past decade as an alternative to carotid endarterectomy (CEA) for the treatment of significant carotid stenosis, and has been shown equally effective in long term stroke prevention.^{1,2} However, most large randomised trials comparing CEA with CAS have shown a higher risk of procedural stroke in patients treated by stenting.^{3–6} Thus, the reduction of these procedural strokes is pivotal in treatment optimisation. As the majority of procedural strokes occur on the day of stenting,^{7,8} a reduction is likely to be achieved through optimal selection of patients with favourable characteristics for CAS. As transfemoral CAS requires guidewire navigation through the aortic arch up to the carotid bifurcation, any difficulty with arterial navigation could increase the risks of arterial damage and subsequent thrombo-embolism. Consequently, several features of vascular anatomy have been suggested to influence the procedural risks of CAS.^{9–14}

The Delphi anatomical risk (DAR) score is an expert opinion based scoring system that grades expected difficulty of CAS based on seven individual anatomical features of the aortic arch and supra-aortic arteries, developed by Macdonald *et al.*¹⁵ as an aid to case selection. The assumption underlying the DAR score is that a higher score predicts the risk of procedural stroke during CAS. A recent substudy from the International Carotid Stenting Study (ICSS) showed that the DAR score predicted the risk of ischaemic lesions on magnetic resonance diffusion weighted imaging (DWI) following CAS, although not after correction for age.¹⁶ Up to now the DAR score has not been validated for the clinically relevant outcome of procedural stroke related to CAS.

The aim was to clinically validate the DAR score criteria within this *post hoc* analysis of the ICSS, a large randomised controlled trial that compared CAS and CEA in recently symptomatic patients. Individual anatomical features as well as the DAR score were related to the risk of any procedural stroke up to 30 days in patients undergoing CAS.

MATERIALS AND METHODS

Trial protocol and patient selection

The ICSS trial protocol (ISRCTN 25337470) has been published previously.¹⁷ In summary, patients with recently symptomatic moderate or severe carotid stenosis (>50%) who were eligible for both surgery and stenting were randomly assigned to CEA or CAS. Patients were excluded from ICSS if they had a stenosis known to be unsuitable for stenting because of tortuous anatomy proximal or distal to the stenosis, the presence of visible thrombus, proximal common carotid artery stenotic disease, or pseudo-occlusion (“string sign”). All procedures were performed by interventionists approved by a credentialing committee as having appropriate training and experience of carotid stenting. Individual interventionists who were not able to satisfy the credentialing requirements were identified as probationary investigators. Stenting procedures carried out

by probationary investigators were proctored by an experienced carotid interventionist, until the proctor was satisfied that the interventionist could carry out procedures without supervision.

For the purpose of this study, ICSS patients were included only if they had been randomised to CAS, the CAS procedure was completed, and pre-procedural computed tomography angiography (CTA) images were available for analysis (i.e. centrally stored in the ICSS archive). In order to calculate the DAR score, imaging had to include the aortic arch and supra-aortic arteries up to the base of the skull.

Anatomical risk scoring

The DAR score is a carotid anatomy scoring system, based on expert consensus, developed to categorise the expected difficulty of CAS. The final DAR score is calculated according to the flow chart in the original expert consensus paper.¹⁵ Table 1 shows seven anatomical risk factors and their definitions as included in the DAR score. Every possible combination of anatomy results in a risk score between 2.4 and 8.5, and has been divided in four categories of expected risk (low, < 5.0; low–intermediate, 5.0–5.9; high–intermediate, 6.0–6.9; high, ≥ 7.0).

Two trained independent observers (D.d.W., E.d.V.) assessed available pre-operative vascular CTAs in order to grade anatomical difficulty and the DAR score. For cases of disagreement, consensus was achieved through discussion with a third independent observer (A.H.).

Table 1. The seven anatomical risk factors that constitute the Delphi anatomical risk score, their definitions, and interobserver reliability per risk factor

Anatomical risk factor	Definition	Kappa (95% CI)	p
Arch atheroma	Severe arch atheroma or arch origin disease	0.80 (0.76–0.84)	<.001
Access		0.76 (0.72–0.80)	<.001
Diseased CCA	Atheroma CCA		
ECA problem	Diseased and/or occluded ECA		
Arch type		0.84 (0.80–0.87)	<.001
Bovine	Conjoint origin of left CCA and brachiocephalic trunk		
Type III	2 cm between highest point of arch and origin of brachiocephalic trunk		
Angulated distal ICA	Angulated distal ICA ≥90°	0.77 (0.73–0.81)	<.001
Pinpoint stenosis	90–99% stenosis ICA with flow beyond lesion	0.79 (0.75–0.83)	<.001

CCA = common carotid artery; CI = confidence interval; ECA = external carotid artery; ICA = internal carotid artery.

Procedural stroke (ipsi- and contralateral)

In ICSS, major events were adjudicated by an independent endpoint committee that was unaware of treatment allocation. Stroke was defined as a rapidly developing clinical syndrome of focal disturbance of cerebral function lasting more than 24 h or leading to death with no apparent cause other than that of vascular origin. Only procedural stroke was analysed, defined as any stroke occurring within 30 days of the stenting procedure, irrespective of the vascular territory of the event.

Duration of procedure

In ICSS, investigators were asked to complete technical data forms for all interventions, which allowed the exact duration of the stenting procedure to be studied. The prolonged duration of a stenting procedure was arbitrarily defined as any procedure lasting ≥ 75 min.

Statistical analysis

A power analysis was performed using Rstudio (version 1.0.143, RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA; <http://www.rstudio.com/>). The total 30 day stroke risk in ICSS was 7.0%. Assuming a procedural stroke risk of 10% in the highest DAR score group vs. 4% in the lowest DAR score group, 566 patients would be needed for the analysis (confirming to a power of 0.8 at an alpha level of 0.05 and a confidence interval of 95%).

The presence of anatomical risk factors was compared in two predefined age groups (<70 years and ≥ 70 years) using a chi-square test. Binomial logistic regression analysis was used to test procedural risk of stroke according to individual anatomical risk factors and the DAR score. Comparison between the reference group and risk factor under consideration was expressed in an odds ratio with Wald 95% confidence interval. Similarly, an association between anatomical difficulty and a prolonged intervention duration was analysed. All analyses were performed for the total cohort and divided by the predefined age groups.

An interobserver reliability analysis was performed for each anatomical risk factor in the DAR score. A kappa statistic (κ) was used to determine consistency among observers (D.d.W., E.d.V.). A κ value of less than 0.20 indicates poor; 0.20–0.39, fair; 0.40–0.59, moderate; 0.60–0.79, good; and 0.80–1.00, outstanding agreement between observers.¹⁸ Statistical analyses were conducted using SPSS 22.0 (IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY, USA). A p value < 0.05 was considered to be statistically significant for all analyses. Data are presented as mean \pm standard deviation, unless otherwise specified.

RESULTS

Study population

A total of 1713 patients were enrolled in ICSS, of whom 855 were randomly allocated to CAS. Pre-procedural CTA was

Table 2. Baseline patient characteristics of 275 patients included in the analyses, and of 853 patients included in ICSS⁵ as a whole

Baseline characteristics	CAS, current study (n = 275)	CAS, ICSS ⁵ (n = 853)
Sex		
Male	194 (71)	601 (70)
Age at intervention		
Age – y	70 \pm 9	70 \pm 9
<70	141 (51)	Unknown
≥ 70	134 (49)	Unknown
Vascular risk factors^a		
Treated hypertension	183 (66)	587 (69)
Treated hyperlipidaemia	173 (64)	522 (61)
Any diabetes mellitus	62 (23)	184 (24)
Cardiac failure	7 (3)	23 (3)
Atrial fibrillation	21 (8)	57 (7)
Angina in past six months	22 (8)	83 (10)
Previous myocardial infarction	48 (18)	151 (18)
Previous CABG	40 (15)	109 (13)
Current smoker	83 (31)	205 (24)
Ex-smoker	122 (45)	408 (48)
Peripheral arterial disease	48 (18)	139 (16)
Ipsilateral carotid stenosis		
50–69%	29 (11)	92 (11)
70–99%	246 (89)	761 (89)
Contralateral carotid stenosis		
<50%	166 (61)	565 (66)
50–69%	42 (15)	128 (15)
70–99%	45 (17)	105 (12)
Occlusion	18 (7)	49 (6)
Procedural outcome		
30 day stroke rate	16 (6)	58 (7)

Values are mean \pm standard deviation (SD), or n (%). CABG = coronary artery bypass graft; CAS = carotid artery stenting; ICSS = International Carotid Stenting Study.

^a For vascular risk factors information was available for 271 patients.

available for 275 of 855 patients (33%), of which 97% had been performed within three months of the procedure. Baseline characteristics are presented in Table 2. The mean age at intervention was 70 \pm 9 years and the majority were male (71%). Table 2 also shows the baseline characteristics of all patients included in ICSS as a whole, which confirms the representativeness of the selected sample.

Anatomical risk scoring

The overall interobserver reliability for the seven anatomical risk factors was high with all kappa values between 0.76 and 0.84 (Table 1). The third observer was consulted in 94 out of 275 patients (34%), in 69 out of 94 patients the disagreement concerned merely one factor. The presence of anatomical risk factors and the DAR score is shown in Table 3. The mean DAR score was 4.3 in the total cohort and only 11 (4%) patients were scored as high difficulty (DAR score ≥ 7.0). Patients aged 70 years and above had a significantly higher DAR score than younger patients (4.6 \pm 1.5 vs. 3.9 \pm 1.4, p < .001), which was mainly attributable to higher rates of arch atheroma (44% vs. 20%, p < .001).

Table 3. Number of patients with anatomical risk factors for the total cohort. Chi-square analysis and two sample *t* test comparing presence of risk factors and the mean DAR score between both age groups

Anatomical risk factors and the DAR score	Total cohort (n = 275)	<70 years (n = 141)	≥70 years (n = 134)	<i>p</i>
<i>Arch atheroma</i>				<.001
No atheroma	188 (68)	113 (80)	75 (56)	
Atheroma	87 (32)	28 (20)	59 (44)	
<i>Access</i>				.15
Normal	199 (72)	109 (77)	90 (67)	
Diseased CCA	50 (18)	20 (14)	30 (23)	
ECA problem	26 (9)	12 (9)	14 (11)	
<i>Arch type</i>				.77
Normal arch	190 (69)	99 (70)	91 (68)	
Bovine arch	31 (11)	17 (12)	14 (10)	
Type III arch	47 (17)	21 (15)	26 (20)	
Bovine and type III arch	7 (3)	4 (3)	3 (2)	
<i>Target vessel</i>				.11
Normal	201 (73)	109 (77)	92 (69)	
Angulated ICA	74 (27)	32 (23)	42 (31)	
<i>Lesion type</i>				.77
Normal	183 (67)	95 (67)	88 (66)	
Pinpoint stenosis	92 (33)	46 (33)	46 (34)	
<i>DAR score</i>				.003
<5.0	180 (65)	107 (76)	73 (55)	
5.0–5.9	58 (21)	20 (14)	38 (28)	
6.0–6.9	26 (9)	10 (7)	16 (12)	
≥7.0	11 (4)	4 (3)	7 (5)	
Mean	4.3 ± 1.4	3.9 ± 1.4	4.6 ± 1.5	<.001

Data are given as *n* (%) or mean ± standard deviation (SD). CCA = common carotid artery; DAR = Delphi anatomical risk; ECA = external carotid artery; ICA = internal carotid artery; SD = standard deviation.

Procedural stroke risk (ipsi- and contralateral)

A total of 16/275 (6%) patients had a stroke during the procedural period, with a higher risk of stroke in patients ≥ 70 than younger patients (9% vs. 3%, *p* = .03). All strokes were ischaemic and seven of 275 (3%) patients had a stroke that was fatal or disabling.

The risk of stroke according to the DAR score is presented in Table 4. Univariable logistic regression showed that the DAR score did not accurately predict procedural stroke, although the low risk group (DAR score < 5.0) had the lowest risk of stroke in the overall cohort. The risk of stroke in the high risk group (DAR score ≥ 7.0) was doubled when compared with the low risk group but this difference was not statistically significant (9% vs. 4%; OR 2.15, 95% CI 0.24–18.91, *p* = .49).

Several additional analyses were performed. A separate analysis for age groups <70 years vs. ≥70 years showed a significantly increased risk of stroke in the high–intermediate group (DAR score 6.0–6.9) when compared with the low risk group in patients < 70 years (OR 13.13, 95% CI 1.62–105.86, *p* = .02; data not shown). However, this was based on four strokes in the age group <70 years, which is reflected in the wide confidence interval. Furthermore, the DAR score was

Table 4. Procedural risk of any stroke following CAS according to the DAR score for the total cohort

DAR score	Patients (n = 275)	Strokes	Unadjusted OR (95% CI)	<i>p</i>
<5.0	180 (65)	8 (4)	1.00	
5.0–5.9	58 (21)	4 (7)	1.59 (0.46–5.50)	.46
6.0–6.9	26 (9)	3 (12)	2.80 (0.69–11.33)	.15
≥7.0	11 (4)	1 (9)	2.15 (0.24–18.91)	.49

Data are given as *n* (%) unless stated otherwise. Binomial logistic regression was used to give odds ratios (OR) per anatomical risk group where possible. “Low risk” anatomy (DAR score < 5.0) was used as reference category. CAS = carotid artery stenting; DAR = Delphi anatomical risk.

grouped into two categories of expected risk instead of four (low risk, < 6.0; high risk, ≥ 6.0), which did not result in any statistical significance (stroke risk 11% vs. 5%; OR 2.28, 95% CI 0.70–7.50, *p* = .17; data not shown).

Table 5 presents the risk of stroke according to the individual anatomical risk factors. None of the evaluated anatomical features showed a significant association with procedural stroke when analysed separately. Arch atheroma, the anatomical feature that was overexpressed in patients over 70 years of age, did not increase the risk of stroke in the total cohort. A separate analysis for the two predefined age groups showed a significantly higher risk of stroke in the patients <70 years with bovine and type III arch (25% vs. 1%; OR 32.67, 95% CI 1.63–656.41, *p* = .02; data not shown). However, this was based solely on one stroke in the four patients <70 years with bovine and type III arch, also reflected in the wide confidence interval.

Duration of intervention

Duration of intervention was available for 242 out of 275 (88%) patients and in these patients the median procedure duration was 65 (range 15–400) minutes. According to the predefined definition, a total of 76/242 (31%) CAS interventions were prolonged for 75 min or more. A prolonged intervention duration was significantly associated with increased risk of stroke (11% vs. 4%; OR 3.14, 95% CI 1.05–9.39, *p* = .04; data not shown). However, no association was found between higher DAR score and prolonged CAS intervention (data not shown).

DISCUSSION

Several features of vascular anatomy have been proposed to predict the difficulty of CAS, and are suggested to increase the risk of procedural stroke after CAS. In the current study, an attempt was made to validate the anatomical risk score as a substudy of ICSS, and demonstrate that the DAR score can be reliably assessed with good interobserver agreements. As the sample size was too small to reach adequate power, no definite conclusions could be drawn on the relationship between the DAR score and procedural risk of stroke after CAS. Of note, as the analysis was confined to CAS patients, no conclusions can be drawn regarding comparisons with carotid endarterectomy.

Table 5. Procedural risk of any stroke following CAS for the total cohort

Risk factor	Patients (n = 275)	Strokes	Unadjusted OR (95% CI)	p
<i>Arch atheroma</i>				
Not present	188 (68)	10 (5)	1.00	
Present	87 (32)	6 (7)	1.32 (0.46–3.75)	.61
<i>Access</i>				
Normal	199 (72)	12 (6)	1.00	
ECA problem	50 (18)	2 (4)	0.65 (0.14–3.00)	.58
Diseased CCA	26 (9)	2 (8)	1.30 (0.27–6.16)	.74
<i>Arch type</i>				
Normal arch	190 (69)	8 (4)	1.00	
Bovine arch	31 (11)	2 (7)	1.57 (0.32–7.76)	.58
Type III arch	47 (17)	5 (11)	2.71 (0.84–8.30)	.09
Bovine and type III arch	7 (3)	1 (14)	3.79 (0.41–35.34)	.24
<i>Target vessel</i>				
Normal	201 (73)	13 (7)	1.00	
Angulated ICA	74 (27)	3 (4)	0.61 (0.17–2.21)	.45
<i>Lesion type</i>				
Normal	183 (67)	9 (5)	1.00	
Pinhole stenosis	92 (33)	7 (8)	1.59 (0.57–4.42)	.37

Data are given as n (%) unless stated otherwise. Binomial logistic regression was used to give odds ratios (OR) per anatomical risk factor where possible. "Normal" anatomy was used as reference category. CAS = carotid artery stenting; CCA = common carotid artery; ECA = external carotid artery; ICA = internal carotid artery; CI = confidence interval; OR = odds ratio.

Perhaps now more than ever, patient selection for carotid interventions is a topic of broad interest, and guidelines pay careful attention to optimising the selection of suitable patients for CAS.¹⁹ If it had been shown that vascular anatomy predicted procedural stroke after CAS, assessment of anatomy could be used to identify high risk patients, and thereby reduce the peri-procedural complication risk. In a recent substudy of ICSS, it was suggested that the DAR score was able to predict the occurrence of new post-procedural DWI lesions after CAS, although not after correction for age.¹⁶ This is the first study to validate the DAR score for the clinically relevant outcome of procedural stroke after CAS. Unfortunately, with only 275 patients the study was insufficiently powered to allow final conclusions. However, some interesting findings could be observed. A doubling of stroke risk was seen in the high risk compared with the low risk DAR categories. Second, a prolonged duration of intervention was significantly associated with increased risk of stroke. This might reflect the degree of difficulty of the intervention, which is likely to be associated with anatomical difficulty, although not statistically significant in this study. Lastly, it was demonstrated that patients over 70 years of age had a significantly higher DAR score than younger patients, and strokes were over three times as common in this elderly patient group. The higher degree of anatomical difficulty in older patients was largely explained by a higher incidence of aortic arch atheroma. This is in line with previous studies in which octogenarians were shown to have increased incidence of aortic arch calcification and innominate

stenosis.^{20,21} Consequently, it has been suggested that the observation of increased procedural risk in elderly patients is related to increased anatomical complexity.²² In the current study, elderly patients had both a higher procedural risk and a higher DAR score. However, although twice as many strokes were seen in the high anatomical risk patients, this did not reach statistical significance.

The only relevant anatomical exclusion criteria from ICSS were the presence of a stenosis thought to be unsuitable for stenting because of tortuous anatomy proximal or distal to the stenosis and pseudo-occlusion. These criteria might be thought to correspond to the factors included in the DAR score of angulated distal ICA and pinpoint stenosis. However, it is notable that although 166 patients in the current study were included in ICSS with one or more of these two factors, they did not have a significantly higher rate of procedural stroke. Thus, the results support excluding such patients only if the anatomy is considered unfavourable for stenting by an experienced interventionist.

The findings need to be interpreted in the light of the following study limitations. Firstly, only a third of ICSS CAS patients had the required baseline CTA available, resulting in a representative but relatively small cohort. Patients in whom only digital subtraction angiography and/or magnetic resonance angiography had been performed were deliberately excluded, in order to achieve a dataset as homogenous as possible. It is notable that there were only small numbers of patients in the high risk groups. This might be a reflection of patient selection in ICSS, and the ICSS CAS population included probably reflected DAR intermediate to low risk patients in everyday practice. Second, although the stroke rate was high in this cohort, the absolute number of strokes was small. The limited power made it less likely to find more modest contributors to risk of procedural strokes. Thirdly, in accordance with the DAR score, only anatomical factors were studied. Plaque characteristics may also be an important predictor of stroke, especially in patients with challenging anatomy.^{23–25} In this study analysis was limited to anatomical risk factors included in the DAR score. A tortuous CCA was rated most difficult in the DAR score expert consensus study, and was excluded from the final risk score because it would rate too difficult in combination with other anatomical risk factors.¹⁵ In addition, several other anatomical factors like severe (circumferential) lesion calcification^{26,27} or lesion length,^{10,28} might also contribute to procedural hazard, but had not been included in the final DAR score.

Finally, it is notable that among the anatomical factors included in the DAR score, only pinpoint stenosis might be expected to influence risk during stent deployment. The others would be expected only to increase the risk of stroke resulting from carotid cannulation. The latter causes stroke in only a minority of stented cases. In ICSS as a whole, an analysis of the mechanisms of procedural stroke only identified four non-ipsilateral strokes out of 55 stented patients with procedural ischaemic stroke that could be related to cannulation and only one case that could be attributed to technical difficulties related to anatomical factors.⁷ Whether the DAR score would have predicted

stroke in these cases could not be determined because of the very small number of events.

CONCLUSIONS AND RECOMMENDATIONS

This is the first clinical validation analysis of the Delphi anatomical risk score for procedural stroke after CAS. No statistically significant association between anatomical difficulty, as defined in the DAR score, and procedural stroke risk was shown in this recently symptomatic patient group. However, the small sample size potentially rendered the study underpowered to give firm conclusions, and a trend towards higher stroke rate in high anatomical risk categories was noted. Nonetheless, it has been shown that the DAR score is a feasible and consistently reported instrument when based on CTA imaging. Furthermore, because only small numbers of strokes can be expected to be associated with anatomical risk factors, the study should be replicated in the future. In order to obtain a sufficient number of baseline CTAs and events, it is important that currently running trials, as well as trials still to be designed and registries, include these imaging work up in their study protocols.

CONFLICTS OF INTEREST

None.

FUNDING

None.

REFERENCES

- Bonati LH, Dobson J, Featherstone RL, Ederle J, Van Der Worp HB, De Borst GJ, et al. Long-term outcomes after stenting versus endarterectomy for treatment of symptomatic carotid stenosis: the International Carotid Stenting Study (ICSS) randomised trial. *Lancet* 2015;**385**:529–38.
- De Borst GJ, Naylor AR. In the end, it all comes down to the beginning! *Eur J Vasc Endovasc Surg* 2015;**50**:271–2.
- Mas J-L, Chatellier G, Beyssen B, Branchereau A, Moulin T, Becquemin J-P, et al. Endarterectomy versus stenting in patients with symptomatic severe carotid stenosis. *N Engl J Med* 2006;**355**:1660–71.
- Ringleb PA, Allenberg J, Brückmann H, Eckstein H-H, Fraedrich G, Hartmann M, et al. 30-day results from the SPACE trial of stent-protected angioplasty versus carotid endarterectomy in symptomatic patients: a randomised non-inferiority trial. *Lancet* 2006;**368**:1239–47.
- Ederle J, Dobson J, Featherstone RL, Bonati LH, van der Worp HB, de Borst GJ, et al. Carotid artery stenting compared with endarterectomy in patients with symptomatic carotid stenosis (International Carotid Stenting Study): an interim analysis of a randomised controlled trial. *Lancet* 2010;**375**:985–97.
- Rosenfield K, Matsumura JS, Chaturvedi S, Riles T, Ansel GM, Metzger DC, et al. Randomized trial of stent versus surgery for asymptomatic carotid stenosis. *N Engl J Med* 2016;**374**:1011–20.
- Huibers A, Calvet D, Kennedy F, Czuriga-kovács KR, Featherstone RL, Moll FL, et al. Mechanism of procedural stroke following carotid endarterectomy or carotid artery stenting within the international carotid stenting study (ICSS) randomised trial. *Eur J Vasc Endovasc Surg* 2015;**50**:281–8.
- Huibers A, De Borst GJ, Thomas DJ, Moll FL, Bulbulia R, Halliday A. The mechanism of procedural stroke following carotid endarterectomy within the asymptomatic carotid surgery trial 1. *Cerebrovasc Dis* 2016;**42**:178–85.
- Khan M, Qureshi AI. Factors associated with increased rates of post-procedural stroke or death following carotid artery stent placement: a systematic review. *J Vasc Interv Neurol* 2014;**7**:11–20.
- Naggara O, Touzé E, Beyssen B, Trinquart L, Chatellier G, Meder JF, et al. Anatomical and technical factors associated with stroke or death during carotid angioplasty and stenting: results from the endarterectomy versus angioplasty in patients with symptomatic severe carotid stenosis (EVA-3S) trial and systematic review. *Stroke* 2011;**42**:380–8.
- Werner M, Bausback Y, Bräunlich S, Ulrich M, Piorkowski M, Friedenberger J, et al. Anatomic variables contributing to a higher periprocedural incidence of stroke and TIA in carotid artery stenting: single Center Experience of 833 Consecutive Cases. *Catheter Cardiovasc Interv* 2012;**80**:321–8.
- Faggioli G, Ferri M, Gargiulo M, Freyrie A, Fratesi F, Manzoli L, et al. Measurement and impact of proximal and distal tortuosity in carotid stenting procedures. *J Vasc Surg* 2007;**46**:1119–24.
- Faggioli GL, Ferri M, Rapezzi C, Tonon C, Manzoli L, Stella A. Atherosclerotic aortic lesions increase the risk of cerebral embolism during carotid stenting in patients with complex aortic arch anatomy. *J Vasc Surg* 2009;**49**:80–5.
- Faggioli GL, Ferri M, Freyrie A, Gargiulo M, Fratesi F, Rossi C, et al. Aortic arch anomalies are associated with increased risk of neurological events in carotid stent procedures. *Eur J Vasc Endovasc Surg* 2007;**33**:436–41.
- Macdonald S, Lee R, Williams R, Stansby G. Towards safer carotid artery stenting: a scoring system for anatomic suitability. *Stroke* 2009;**40**:1698–703.
- Müller MD, Ahlhelm FJ, Von Hessling A, Doig D, Nederkoorn PJ, MacDonald S, et al. Vascular anatomy predicts the risk of cerebral ischemia in patients randomized to carotid stenting versus endarterectomy. *Stroke* 2017;**48**:1285–92.
- Featherstone RL, Brown MM, Coward LJ. International Carotid Stenting Study: protocol for a randomised clinical trial comparing carotid stenting with endarterectomy in symptomatic carotid artery stenosis. *Cerebrovasc Dis* 2004;**18**:69–74.
- Altman D. *Practical Statistics for Medical Research*. Taylor & Francis Ltd; 1990.
- Naylor AR, Ricco JB, de Borst GJ, Debus S, de Haro J, Halliday A, et al. Management of atherosclerotic carotid and vertebral artery disease: 2017 clinical practice guidelines of the European Society for Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg* 2018;**55**:3–81.
- Lam RC, Lin SC, DeRubertis B, Hyncek R, Kent KC, Faries PL. The impact of increasing age on anatomic factors affecting carotid angioplasty and stenting. *J Vasc Surg* 2007;**45**:875–80.
- Lin S, Trocciola S, Rhee J, Dayal R, Chaer R, Morrissey N, et al. Analysis of anatomic factors and age in patients undergoing carotid angioplasty and stenting. *Ann Vasc Surg* 2005;**19**:798–804.
- Howard G, Roubin GS, Jansen O, Hendrikse J, Halliday A, Fraedrich G, et al. Association between age and risk of stroke or death from carotid endarterectomy and carotid stenting: a meta-analysis of pooled patient data from four randomised trials. *Lancet* 2016;**387**:1305–11.
- Howard DPJ, Van Lammeren GW, Rothwell PM, Redgrave JN, Moll FL, De Vries JPPM, et al. Symptomatic carotid atherosclerotic disease: correlations between plaque composition and ipsilateral stroke risk. *Stroke* 2015;**46**:182–9.
- Van Lammeren GW, Reichmann BL, Moll FL, Bots ML, De Kleijn DPV, De Vries JPPM, et al. Atherosclerotic plaque vulnerability as an explanation for the increased risk of stroke in elderly undergoing carotid artery stenting. *Stroke* 2011;**42**:2550–5.
- Huibers A, De Borst GJ, Bulbulia R, Pan H, Halliday A. Plaque echolucency and the risk of ischaemic stroke in patients with asymptomatic carotid stenosis within the first asymptomatic carotid surgery trial (ACST-1). *Eur J Vasc Endovasc Surg* 2016;**51**:616–21.
- Setacci C, Chisci E, Setacci F, Iacoponi F, de Donato G, Rossi A. Siena carotid artery stenting score: a risk modelling study for individual patients. *Stroke* 2010;**41**:1259–65.

27 AbuRahma AF, DerDerian T, Hariri N, Adams E, AbuRahma J, Dean LS, et al. Anatomical and technical predictors of perioperative clinical outcomes after carotid artery stenting. *J Vasc Surg* 2017;66:423–32.

28 Bonati LH, Ederle J, Dobson J, Engelter S, Featherstone RL, Gaines PA, et al. Length of carotid stenosis predicts peri-procedural stroke or death and restenosis in patients randomized to endovascular treatment or endarterectomy. *Int J Stroke* 2014;9:297–305.


Eur J Vasc Endovasc Surg (2019) 58, 670

COUP D'OEIL

A Mycotic Saccular Aneurysm Diagnosed With ¹⁸F-Labelled Fluoro-2-Deoxyglucose Positron Emission Tomography/Computed Tomography Scanning

Spyros Papadoulas*, Stavros K. Kakkos

Department of Vascular Surgery, University of Patras Medical School, Patras, Greece



An 83 year old man presented with low grade fever, anorexia, and para-umbilical pain. C reactive protein levels and erythrocyte sedimentation rate were elevated, but the white cell count was normal. Abdominal computed tomography (CT) angiography revealed a 3.5 cm saccular aneurysm at the aortic bifurcation (A, arrow). Positron emission tomography with ¹⁸F-labelled fluoro-2-deoxyglucose integrated with CT revealed increased metabolic activity in the aneurysm sac. The peri-aortic and prevertebral fat showed higher density, suggesting a mycotic aneurysm (B, arrow). Blood cultures were negative. For family reasons, the patient was transferred to a centre in his home city, where he underwent endovascular repair.

* Corresponding author. Department of Vascular Surgery, University Hospital of Patras, Patras 26504, Greece.

E-mail address: spyros.papadoulas@gmail.com (Spyros Papadoulas).

1078-5884/© 2019 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2019.07.026>