

The Knee

British Patello-Femoral Society Special Edition

VOLUME 26 ISSUE 6

CONTENTS

DECEMBER 2019

EDITORIAL

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

Editorial

D. Clark, A. Metcalfe, J. Eldridge 1159

REVIEWS

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

Management of the first-time lateral patellar dislocation

D.S. Johnson, P.G. Turner (UK of Great Britain and Northern Ireland) 1161

Replacement for patellofemoral arthritis

D.S. Johnson, P.G. Turner (UK of Great Britain and Northern Ireland) 1166

Current concepts in the surgical management of patellar instability

P. Thompson, A.J. Metcalfe (UK) 1171

BASIC SCIENCE

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

The effect of surgery on patellar tendinopathy: Novel use of MRI questions the exploitability of the rat collagenase model to humans

M.J. Dan, R.A. Oliver, J.D. Crowley, V. Lovric, W.C.H. Parr, D. Broe, M. Cross, W.R. Walsh (Australia) 1182

Psychometric properties of the Norwich Patellar Instability Score in people with recurrent patellar dislocation

T.O. Smith, A. Choudhury, R. Navratil, C.B. Hing (UK) 1192

REGULAR ARTICLE

The relationship between lateral epicondyle morphology and iliotibial band friction syndrome: A matched case-control study

J.S. Everhart, J.C. Kirven, J. Higgins, A. Hair, A.A.W. Chaudhari, D.C. Flanigan (USA) 1198

BIOMECHANICS & KINEMATICS

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

Mid inter-epicondyle trochlea intersection (MIELTI): Proposal of a new index for identifying the deepest part of the trochlea

R. Keehan, A. Gill, L. Smith, R. Ahmad, J. Eldridge (UK) 1204

Influence of the Powers™ strap on pain and lower limb biomechanics in individuals with patellofemoral pain

H. Greuel, L. Herrington, A. Liu, R.K. Jones (UK) 1210

Mechanical characterization of the ligaments in subject-specific models of the patellofemoral joint using in vivo laxity tests

M. Akbar, F. Farahmand, N. Arjmand (Iran) 1220

Lateral patellar maltracking due to trochlear dysplasia: A computational study

S.C. Rezvanifar, B.L. Flesher, K.C. Jones, J.J. Elias (USA) 1234

REGULAR ARTICLE

Dual-pivot bearings improve ambulation and promote increased activity levels in Total knee arthroplasty: A match-controlled retrospective study

R. Sandberg, E.R. Deckard, M. Ziemba-Davis, S.A. Banks, R.M. Meneghini (USA) 1243

SOFT TISSUE

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

Traumatic injury of the knee extensor mechanism in skeletally immature patients: Outcome and classification

M.A. Yousef (USA) 1250

(continued on next page)

REGULAR ARTICLES

- Selective bundle reconstruction in partial ACL tears leads to excellent long-term functional outcomes and a low percentage of failures
S. Perelli, F. Ibañez, P.E. Gelber, J.I. Erquicia, X. Pelfort, J.C. Monllau (Spain) 1262
- Long-term clinical and radiological outcomes after multiligament knee injury using a delayed ligament reconstruction approach: A single-center experience
M. Hantes, A. Fylos, F. Papageorgiou, K. Alexiou, I. Antoniou (Greece) 1271

ARTHRITIS & OSTEOTOMIES

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

- Anteroposterior distance between the tibial tuberosity and trochlear groove in patients with patellar instability
M.J. Tanaka, T. D'Amore, J.J. Elias, G. Thawait, S. Demehri, A.J. Cosgarea (USA) 1278

REGULAR ARTICLES

- Association between pain catastrophizing, physical function and pain at first visit in the outpatient knee clinic
S. Birch, M. Stilling, I. Mechlenburg, M.B. Reinholdt, T.B. Hansen (Denmark) 1286
- The effect of a longitudinal tear of the medial meniscus on medial meniscal extrusion in anterior cruciate ligament injury patients
H. Katagiri, K. Miyatake, Y. Nakagawa, K. Otabe, T. Ohara, M. Shioda, I. Sekiya, H. Koga (Japan) 1292
- Proximal tibiofibular division in lateral closing wedge high tibial osteotomy does not increase varus instability of the knee
J.-Y. Park, J.-K. Kim, H.-S. Han, M.C. Lee (South Korea) 1299
- Return to sport following isolated opening wedge high tibial osteotomy
J.N. Liu, A. Agarwalla, G.H. Garcia, D.R. Christian, M.L. Redondo, A.B. Yanke, B.J. Cole (USA) 1306
- Factors affecting stability after medial opening wedge high tibial osteotomy using locking plate: A cadaveric study
Y.G. Na, D.-S. Kwak, S. Chong, T.K. Kim (Republic of Korea) 1313

REHABILITATION & ORTHOTICS

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

- A mathematical modelling study investigating the influence of knee joint flexion angle and extension moment on patellofemoral joint reaction force and stress
T. Tomoya, E. Mutsuaki, I. Takuma, T. Yuta, K. Masayoshi (Japan) 1323

ARTHROPLASTY

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

- Qualitative and quantitative analysis of patellar vascular anatomy by novel three-dimensional micro-computed-tomography: Implications for total knee arthroplasty
D. Wang, Z. Shen, D. Jiang, X. Li, X. Fang, H. Leng, W. Zhang (PR China) 1330
- Does total contact of the patella with the femoral trochlea during no thumb test significantly reduce anterior knee pain?
J. Manghwani, S.V. Vaidya, H. Patel, C.S. Vaidya (India) 1338
- The midterm results of a cohort study of patellofemoral arthroplasty from a non-designer centre using an asymmetric trochlear prosthesis
R. R., S. Gupta, P.Y.F. Lee, A. Chandratreya (UK) 1348
- The effect of patellar thickness on gait biomechanics following total knee arthroplasty
E.A. Parke, C.K. Nakasone, S.N. Andrews, A.R. Wright, C.D. Stickley (USA) 1354
- Early post-operative opioid consumption: A comparison between medial unicompartmental, patellofemoral, and total knee arthroplasty
C.M. Melnic, E.R. Kazarian, M.K. Dwyer, E.L. Domingo-Johnson, A.A. Freiberg, H.S. Bedair (USA) 1360

REGULAR ARTICLES

- A greater reduction in the distal femoral anterior condyle improves flexion after total knee arthroplasty in patients with osteoarthritis
K. Nishitani, R. Hatada, S. Kuriyama, S.L. Lyman, S. Nakamura, H. Ito, S. Matsuda (Japan, USA) 1364
- Inconsistent relationship between body weight/body mass index prior to total knee arthroplasty and the 12-year survival
J.-Y. Jenny, D. Saragaglia, M. Bercovy, A. Cazenave, T. Gaillard, F. Châtain, B. Jolles, J.-L. Rouvillain French Society for Hip and Knee Surgery (SFHG) (France, Switzerland) 1372
- Determining the stress biomarker profile in patients undergoing total knee replacement and the relationship with outcome at 12 months
A. Chawla, V. Paraoan, R. Rabiou, T. Clark, S. Powell, D. Grammatopoulos, C. Armitage, A. Penedo, A. Havard, A. Metcalfe, T. Barlow, C. Khatri (UK) 1379
- The fibular shaft axis and medial cortex of the proximal fibula are reliable landmarks for the mechanical axis of the tibia in patients with knee osteoarthritis
K. Xie, X. Han, X. Jiang, L. Wang, S. Ai, Z. Yu, Y. Hao, H. Wu, X. Qu, M. Yan (China) 1386

(continued on next page)

Different targets of mechanical alignment do not improve knee outcomes after TKA Y. Khatib, A. Xia, J.M. Naylor, I.A. Harris, R.M. Sorial (Australia)	1395
Fulfilment of preoperative expectations and postoperative patient satisfaction after total knee replacement. A prospective analysis of 200 patients A.H. Deakin, M.A. Smith, D.T. Wallace, E.J. Smith, M. Sarungi (UK)	1403
Simultaneous bilateral unicompartmental knee replacement improves gait parameters in patients with bilateral knee osteoarthritis M. Marullo, J.A. Vitale, E. Stucovitz, S. Romagnoli (Italy)	1413
A novel patient-specific instrument design can deliver robotic level accuracy in unicompartmental knee arthroplasty G.G. Jones, S. Clarke, S. Harris, M. Jaere, T. Aldalmani, P. de Klee, J.P. Cobb (UK, Ireland)	1421

TECHNICAL TIPS

SPECIAL ISSUE ON PATELLOFEMORAL JOINT

Referencing the trochlear groove based on three-dimensional computed tomography imaging improves the reliability of the measurement of the tibial tuberosity–trochlear groove distance in patients with higher grades of trochlea dysplasia K. Nha, Y.J. Nam, M.J. Shin, S.D. Sun, J.Y. Park, R. Debnath, B.H. Lee (Republic of Korea, India)	1429
Tibial tubercle proximalization as quadriceps lengthening in treating severe habitual patellar dislocation in adults G.-y. Song, H. Zhang, J. Zhang, Z.-j. Zhang, H. Feng (China)	1437

CASE REPORT

REGULAR ARTICLE

Cartilage regeneration in osteoarthritic knees treated with distal femoral osteotomy and intra-lesional implantation of allogenic human umbilical cord blood-derived mesenchymal stem cells: A report of two cases J.-S. Song, K.-T. Hong, N.-M. Kim, J.-Y. Jung, H.-S. Park, Y.S. Chun, S.J. Kim (Republic of Korea)	1445
--	------

LETTERS TO EDITOR & CORRECTIONS

REGULAR ARTICLES

Letter to Editor “Continuous adductor canal blockade facilitates increased home discharge and decreased opioid consumption after total knee arthroplasty.” M.C. Kendall (USA)	1451
Response to Editor “Continuous adductor canal blockade facilitates increased home discharge and decreased opioid consumption after total knee arthroplasty” M.R. Klement, W.M. Bullock, T.M. Seyler, M.P. Bolognesi, S.A. Grant	1453