


Amidation of aldehydes using mono-cationic half-sandwich rhodium(III) complexes with functionalized phenylhydrazone ligands

Neelakandan Devika, Subbiah Ananthalakshmi, Nandhagopal Raja, Gajendra Gupta and Bruno Therrien*




Supramolecular polymers derived from the Pt^{II} and Pd^{II} schiff base complexes via C(sp²)-H ... Hal hydrogen bonding: Combined experimental and theoretical study

Alexander G. Tskhovrebov*, Alexander S. Novikov, Olga V. Odintsova, Vladimir N. Mikhaylov, Viktor N. Sorokoumov, Tatiyana V. Serebryanskaya and Galina L. Starova




Supramolecular polymers via C(sp²)-H...Hal HB



NOTES

Heterotrimetallic complexes of iron and ruthenium based on vinyliminium dithiocarboxylate ligands

Lorenzo Mognon, Samuel Richardson, Gabriele Agonigi, Tamzin Bond, Fabio Marchetti** and James D.E.T. Wilton-Ely*



COMMUNICATIONS

Chiral salen Mn (III) immobilized on ZnPS-PVPA through alkoxy-triazole for superior performance catalyst in asymmetric epoxidation of unfunctionalized olefins

Jing Huang*, Sirui Liu, Yan Ma and Jiali Cai**

Chiral salen Mn (III) anchored on ZnPS-PVPA through alkoxy-triazole display superior catalytic ability either for the experimental scale reactions or for the large-scale reactions.

