

Comment

From theory to experimental evidence

Comment on “Chimera states in neuronal networks: A review” by S. Majhi, B.K. Bera, D. Ghosh and M. Perc

Nikita Frolov, Alexander Hramov *

Neuroscience and Cognitive Technology Laboratory, Innopolis University, 1 Universitetskaya str., Innopolis 420500, Russia

Received 18 February 2019; accepted 20 February 2019

Available online 25 February 2019

Communicated by J. Fontanari

Since the early 2000s, when *chimera state* has been firstly discovered by Kuramoto and Battogtokh in their pioneer work [1], this nonlinear phenomenon continues to arouse interest of scientific community. Despite the fact, that network models considered in numerous theoretical studies of chimera states represent simplified versions of real living systems, they surprisingly exhibit various types of unobvious emergent behavior [2]. Lately, much attention is paid to the analysis of more realistic models with complexly organized structure and different types of possible links between the elements [3–7]. In this sense, human brain represents “a very challenging complex system where neurons [...] form possibly the most complicated structure” as Majhi and colleagues expertly stated at the beginning of their review [8]. Indeed, self-organization, synchronization and chimera state as a particular case of pattern formation are inherent in normal brain functioning since these processes provide the mechanisms for neural communication and cognitive activity. In this context, the present article is extremely relevant. The Authors provide a comprehensive review on recent advances in theoretical analysis and state-of-the-art problems of chimera pattern formation with emphasis on neuronal networks. The paper is easy to follow and we have been pleased to learn that the Authors paid a special attention to biological background of the reviewed models.

However, one of the important problems of this field is a lack of focused experimental studies of the chimera states in neurobiological systems. This is due to the complexity of neural systems on the one hand and the non-triviality of their control schemes on the other hand. Further, we are going to propose a few possible directions for experimental investigation of chimera patterns in biological systems.

Recent advances in neurophysiological techniques for studying and control of single neuron behavior provide necessary tools for fundamental research in the area of self-organization of neural systems. In particular, the approach based on optogenetic neural stimulation and closed-loop control demonstrated by Pashae et al. [9] might be extremely helpful in this context. Using electrode array one may drive the dynamics in a whole neural population recreating interaction mechanisms from theoretical works. This technique opens a way for the development of methods for either generation or suppression of locally coherent patterns with given properties in brain neural network. Besides

DOI of original article: <https://doi.org/10.1016/j.plrev.2018.09.003>.

* Corresponding author.

E-mail address: hramovae@gmail.com (A. Hramov).

pure fundamental prospects, such direction of research have a clear practical application in the field of neural disease therapy and brain functioning improvement.

Another interesting research topic, in our opinion, is searching for the chimera patterns in normal human brain activity. This kind of behavior may not be as pronounced as pathological brain activity, but it reflects more fundamental aspects of brain dynamics. According to *coordination dynamics* theory [10,11] human cognitive functioning is associated with activation of neural groups of brain cortex, wherein the manner of interdependence among them switches in real-time due to performed task. This theory is closely related with concepts of multi- and metastability originated from nonlinear dynamics [12]. Generation of such self-organized dynamical patterns in cortical network may be referred to as chimera-like states. Recent experimental studies on perception of ambiguous visual stimuli show the multistability of cognitive brain activity in visual cortex, which manifests as emergence of different coherent patterns depending on image interpretation [13,14]. These findings could be considered as a starting point for a detailed experimental study of chimera-like behavior in normal brain activity.

Finally, we suggest that chimera-like patterns may be observed in experimental realization of theoretical concept of *network of networks* [15] as a mental interaction in a group of people. In recent work [6], the network of networks topology has been proved to promote chimera behavior. The mechanism of brain dynamics control through mental interaction in a pair of participants via brain-to-brain interface (BBI) during the performance of common cognitive task has been proposed in [16]. When sharing cognitive load, brain dynamics of both participants self-organizes into either anti-phase synchronization (optimal interaction) or desynchronization (non-optimal interaction) due to proper choice of coupling delay. Expanding this concept to a group of people one may expect to observe the emergence of various regimes of collective brain behavior, i.e. collective decision-making, including chimera-like patterns. It is remarkable that, in this case one may achieve chimera behavior on the different levels of such living network of networks: (i) on the macroscopic level, represented by group interactions; (ii) on the microscopic level manifested as neural interactions within a brain network of each participant.

To conclude, we consider that the problems reviewed by Majhi et al. are significant for multidisciplinary field of science and relevant for a wide audience including specialists from physics, network theory, biology and neuroscience. It has a potential to inspire future research in the area of theoretical and experimental studies of the chimera states in living systems. We strongly support the materials and the way they are presented in the Review paper. The Authors have accounted broad range of theoretical findings concerning chimera states in neural network, but they have paid sufficiently less attention to discussion of possibilities of chimera behavior experimental evidence. We believe that our brief comments partially make up this lack and complement the main Review paper.

Acknowledgements

The authors acknowledge the support of Russian Science Foundation (Grant 17-72-30003).

References

- [1] Kuramoto Y, Battogtokh D. Coexistence of coherence and incoherence in nonlocally coupled phase oscillators. *Nonlinear Phenom Complex Syst* 2002;5(4):380–5.
- [2] Panaggio MJ, Abrams DM. Chimera states: coexistence of coherence and incoherence in networks of coupled oscillators. *Nonlinearity* 2015;28(3):R67.
- [3] Hizanidis J, Kouvaris NE, Zamora-López G, Díaz-Guilera A, Antonopoulos CG. Chimera-like states in modular neural networks. *Sci Rep* 2016;6:19845.
- [4] Frolov NS, Maksimenko VA, Makarov VV, Kirsanov DV, Hramov AE, Kurths J. Macroscopic chimera-like behavior in a multiplex network. *Phys Rev E* 2018;98(2):022320.
- [5] Chouzouris T, Omelchenko I, Zakharova A, Hlinka J, Jiruska P, Schöll E. Chimera states in brain networks: empirical neural vs. modular fractal connectivity. *Chaos, Interdiscip J Nonlinear Sci* 2018;28(4):045112.
- [6] Makarov VV, Kundu S, Kirsanov DV, Frolov NS, Maksimenko VA, Ghosh D, et al. Multiscale interaction promotes chimera states in complex networks. *Commun Nonlinear Sci Numer Simul* 2019;71:118–29.
- [7] Kundu S, Bera BK, Ghosh D, Lakshmanan M. Chimera patterns in three-dimensional locally coupled systems. *Phys Rev E* 2019;99:022204.
- [8] Majhi S, Bera BK, Ghosh D, Perc M. Chimera states in neuronal networks: a review. *Phys Life Rev* 2019;28:100–21. <https://doi.org/10.1016/j.plrev.2018.09.003> [in this issue].
- [9] Pashaei R, Baumgartner R, Richner TJ, Brodnick SK, Azimipour M, Eliceiri KW, et al. Closed-loop optogenetic brain interface. *IEEE Trans Biomed Eng* 2015;62(10):2327–37.
- [10] Bressler SL, Kelso JS. Cortical coordination dynamics and cognition. *Trends Cogn Sci* 2001;5(1):26–36.

- [11] Kelso JS. Dynamic patterns: the self-organization of brain and behavior. Bradford Book; 1997.
- [12] Kelso JS. Multistability and metastability: understanding dynamic coordination in the brain. *Philos Trans R Soc B* 2012;367(1591):906–18.
- [13] Runnova AE, Hramov AE, Grubov VV, Koronovskii AA, Kurovskaya MK, Pisarchik AN. Theoretical background and experimental measurements of human brain noise intensity in perception of ambiguous images. *Chaos Solitons Fractals* 2016;93:201–6.
- [14] Hramov AE, Maksimenko VA, Pchelintseva SV, Runnova AE, Grubov VV, Musatov VY, et al. Classifying the perceptual interpretations of a bistable image using EEG and artificial neural networks. *Front Neurosci* 2017;11:674.
- [15] Gao J, Buldyrev SV, Havlin S, Stanley HE. Robustness of a network of networks. *Phys Rev Lett* 2011;107(19):195701.
- [16] Maksimenko VA, Hramov AE, Frolov NS, Lüttjohann A, Nedaivovov VO, Grubov VV, et al. Increasing human performance by sharing cognitive load using brain-to-brain interface. *Front Neurosci* 2018;12:949.