

Comment

One size fits all does not apply to brain lateralisation

Comment on “Phenotypes in hemispheric functional segregation? Perspectives and challenges” by Guy Vingerhoets

Michel Thiebaut de Schotten^{a,b,*}, Patrick Friedrich^{a,b}, Stephanie J. Forkel^{c,d}^a Brain Connectivity and Behaviour Laboratory (BCBLab), Sorbonne Universities, Paris, France^b Groupe d'Imagerie Neurofonctionnelle, Institut des Maladies Neurodégénératives-UMR 5293, CNRS, CEA University of Bordeaux, Bordeaux, France^c Centre for Neuroimaging Sciences, Department of Neuroimaging, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK^d Natbrainlab, Department of Forensics and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

Received 16 July 2019; accepted 18 July 2019

Available online 22 July 2019

Communicated by J. Fontanari

Our understanding of the functioning of the brain is primarily based on an average model of the brain's functional organisation, and any deviation from the standard is considered as random noise or a pathological appearance. Studying pathologies has, however, greatly contributed to our understanding of brain functions. For instance, the study of naturally-occurring or surgically-induced brain lesions revealed that language is predominantly lateralised to the left hemisphere [1] while perception/action and emotion are commonly lateralised to the right hemisphere [2,3]. The lateralisation of function was subsequently replicated by task-related functional neuroimaging in the healthy population [4–6]. Despite its high significance and reproducibility, this pattern of lateralisation of function is true for most, but not all participants. Bilateral and flipped representations of classically lateralised functions have been reported during development [7–10] and in the healthy adult population for language [11], perception/action [12] and emotion [13]. Understanding these different functional representations at an individual level is crucial to improve the sophistication of our models and account for the variance in developmental trajectories, cognitive performance differences and clinical recovery. With the availability of *in vivo* neuroimaging, it has become feasible to study large numbers of participants and reliably characterise individual differences, also referred to as phenotypes. Yet, we are at the beginning of inter-individual variability modelling, and new theories of brain function will have to account for these differences across participants [14].

Within this context, Guy Vingerhoets argues in his review [15] that three main phenotypes of functional lateralisation prevail: The *order* type corresponds to the common pattern of lateralisation, where language functions are dominant in the left, while perception/action and the processing of emotional stimuli are dominant in the right hemisphere. The *reverse* type corresponds to a “flipped” pattern, with language being dominant in the right and per-

DOI of original article: <https://doi.org/10.1016/j.plrev.2019.06.002>.

* Corresponding author at: Brain Connectivity and Behaviour Laboratory (BCBLab), Institut des Maladies Neurodégénératives - CNRS UMR 5293, 146 Rue Léo Saignat - CS61292, 33076 BORDEAUX.

E-mail address: michel.thiebaut@gmail.com (M. Thiebaut de Schotten).

ception/action and emotion in the left hemisphere. Lastly, in the *crowded* type, functions that are usually dissociated between the two hemispheres, are co-located within the same hemisphere. In healthy controls, the *order* and *reverse* types have a small general cognitive advantage on standard neuropsychological assessments [16,17], whereas the *crowded* type may be advantageous in some situations. For instance, a strong dominance of attention in the right hemisphere in left-handers combined with a right eye dominance has been suggested as a critical advantage for sporting duels [18].

Two hypotheses may explain the existence of a *crowded* type. The pathological hypothesis suggests that the *crowded* type is associated with an interrupted or disturbed development of functional lateralisation [19], which manifests with a small cognitive disadvantage. The evolutionary hypothesis, on the other hand, suggests that the *crowded* type is a normally developing phenotype that thrives in specific situations and hence survived through evolution. While genes, handedness, gender and *situs invertus* have been shown to have a minimal association with lateralisation types (see Vingerhoets et al. [15] for an extensive review), lateralisation phenotypes are critical in the presence of brain damage and may determine the potential for the recovery of function (e.g. language recovery [20,21], visuospatial neglect recovery [22]). *Crowded* or *reverse* functional lateralisation is evident in numerous neurodevelopmental and psychiatric conditions [23], including a stronger right lateralised attention network in attention deficit hyperactivity disorder [24] and a weaker left lateralisation of language networks in autism [25] and schizophrenia [26]. Whether this lateralisation causes or is contingent to psychiatric conditions remains to be clarified [19]. This body of literature may suggest that functional lateralisation has been finely tuned across millions of years of evolution, and characterising these phenotypes is critical to advancing personalised medicine.

How does one accurately decipher the phenotypical organisation of the lateralisation of brain functions? Studying brain lateralisation has been hampered by various limitations, including the sometimes shallow relationship between structural and functional asymmetries [27], the low replicability of behavioural, functional and anatomical measures and the often small group sizes. Guy Vingerhoets suggests that a comprehensive model of the lateralisation of brain functions will require a high-dimensional, rather than a uni- or bidimensional approach [28], to capture the complementary trade-off or independence between lateralisation in behavioural measures, functional brain activations, brain anatomy, and their relationship with genetic data across a large subsample of the population. At the dawn of the open science/open data revolution, this endeavour seems feasible and timely, as preliminary evidence recently demonstrated for structural lateralisation [29]. The high-dimensional analysis of brain lateralisation will allow for the demonstration of the existence and the origin of an entirely reverse type; indicate whether *order*, *reverse* and *crowded* phenotypes are truly segregated or part of the same continuum; potentially reveal other phenotypes (i.e., is partially crowded possible?), and identify predictive biomarkers for brain lateralisation that will subsequently be translated to the clinic. It will also allow enable the objective testing of whether separate and overlapping Gaussian distributions are hidden behind brain lateralisation [30] and will lead to new hypotheses on mechanisms supporting functional lateralisation and recovery after brain damage [31]. A preliminary hypothesis proposed by Guy Vingerhoets [15] amongst other authors [32,33], would be that function is lateralised in the brain in order to minimise functional overlap and redundancy, and thus enhance cognitive capacities. Accordingly, functional segregation across the hemispheres and the co-lateralisation of function can be advantageous or disadvantageous according to whether they rely on redundant processes. Such explorations will help one understand the functioning of the brain and elucidate some of the evolutionary mechanisms that support functional specialisation and segregation. This is in line with preliminary evidence that suggests that variability may be a significant landmark of recent evolutionary trends [34] and might impact limbic functions (e.g., emotion) differently than higher cognitive functions (e.g., language). Hence, examining the variability of brain lateralisation phenotypes across species may provide insights into the evolution of hemispheric asymmetries.

To conclude, we strongly agree with Guy Vingerhoets' perspective and challenges [15]. We believe that actions are needed to take the next critical steps towards multidimensional participant-centric analyses. Taking other successful initiatives [35] as an example, the initial step would be for researchers who investigate functional brain lateralisation to unite and pool efforts to create an extensive high-dimensional comprehensive database to harvest together as a consortium. This effort ensures a large database and productive collaborations within an open science framework to take the next steps together.

Acknowledgements

This work has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 818521 to MTS) and by the Ruhr University Research School PLUS, funded by Germany's Excellence Initiative [DFG GSC 98/3 to PF].

References

- [1] Broca P. Sur le siege de la faculte du langage articule. *Bull Soc Anthropol* 1865.
- [2] Sperry RW. Lateral specialization in the surgically separated hemispheres. New York: Rockefeller University Press; 1974.
- [3] Mills CK. The cerebral mechanisms of emotional expression. *Trans Coll Phys Phila* 1912;34:381–90.
- [4] Tzourio N, Crivello F, Mellet E, Nkanga-Ngila B, Mazoyer B. Functional anatomy of dominance for speech comprehension in left handers vs right handers. *NeuroImage* 1998;8:1–16.
- [5] Shulman GL, Pope DL, Astafiev SV, McAvoy MP, Snyder AZ, Corbetta M. Right hemisphere dominance during spatial selective attention and target detection occurs outside the dorsal frontoparietal network. *J Neurosci* 2010;30:3640–51.
- [6] Karolis VR, Corbetta M, Thiebaut de Schotten M. The architecture of functional lateralisation and its relationship to callosal connectivity in the human brain. *Nat Commun* 2019;10:1417.
- [7] Perani D, Saccuman MC, Scifo P, Anwander A, Spada D, Baldoli C, et al. Neural language networks at birth. *Proc Natl Acad Sci USA* 2011;108:16056–61.
- [8] Friederici AD, Brauer J, Lohmann G. Maturation of the language network: from inter- to intrahemispheric connectivities. *PLoS ONE* 2011;6:e20726.
- [9] Szaflarski JP, Holland SK, Schmithorst VJ, Byars AW. fMRI study of language lateralization in children and adults. *Hum Brain Mapp* 2006;27:202–12.
- [10] Vannest J, Karunanayaka PR, Schmithorst VJ, Szaflarski JP, Holland SK. Language networks in children: evidence from functional MRI studies. *Am J Roentgenol* 2009;192:1190–6.
- [11] Mazoyer B, Zago L, Jobard G, Crivello F, Joliot M, Percey G, et al. Gaussian mixture modeling of hemispheric lateralization for language in a large sample of healthy individuals balanced for handedness. *PLoS ONE* 2014;9:e101165.
- [12] Thiebaut de Schotten M, Dell'Acqua F, Forkel SJ, Simmons A, Vergani F, Murphy DG, et al. A lateralized brain network for visuospatial attention. *Nat Neurosci* 2011;14:1245–6.
- [13] Ioannucci S, George N, Cerliani L, Thiebaut de Schotten M. White matter correlates of hemi-face dominance in emotional expression. *bioRxiv*, 232926, 2017.
- [14] Thiebaut de Schotten M, Shallice T. Identical, similar or different? Is a single brain model sufficient?. *Cortex* 2017;86:172–5.
- [15] Vingerhoets G. Phenotypes in hemispheric functional segregation? Perspectives and challenges. *Phys Life Rev* 2019;(30):1–18. <https://doi.org/10.1016/j.plrev.2019.06.002> [in this issue].
- [16] Mellet E, Zago L, Jobard G, Crivello F, Petit L, Joliot M, et al. Weak language lateralization affects both verbal and spatial skills: an fMRI study in 297 subjects. *Neuropsychologia* 2014;65:56–62.
- [17] Groen MA, Whitehouse AJ, Badcock NA, Bishop DV. Does cerebral lateralization develop? A study using functional transcranial Doppler ultrasound assessing lateralization for language production and visuospatial memory. *Brain Behav* 2012;2:256–69.
- [18] Petit L, Zago L, Mellet E, Jobard G, Crivello F, Joliot M, et al. Strong rightward lateralization of the dorsal attentional network in left-handers with right sighting-eye: an evolutionary advantage. *Hum Brain Mapp* 2015;36:1151–64.
- [19] Bishop DV. Cerebral asymmetry and language development: cause, correlate, or consequence?. *Science* 2013;340:1230531.
- [20] Bartha-Doering L, Novak A, Kollndorfer K, Schuler AL, Kasprian G, Langs G, et al. Atypical language representation is unfavorable for language abilities following childhood stroke. *Eur J Paediatr Neurol* 2019;23:102–16.
- [21] Forkel SJ, Thiebaut de Schotten M, Dell'Acqua F, Kalra L, Murphy DG, Williams SC, et al. Anatomical predictors of aphasia recovery: a tractography study of bilateral perisylvian language networks. *Brain* 2014;137:2027–39.
- [22] Lunven M, Thiebaut De Schotten M, Bourslon C, Duret C, Migliaccio R, Rode G, et al. White matter lesional predictors of chronic visual neglect: a longitudinal study. *Brain* 2015;138:746–60.
- [23] Wexler BE. Cerebral laterality and psychiatry: a review of the literature. *Am J Psychiatr* 1980;137:279–91.
- [24] Sanefuji M, Craig M, Parlatini V, Mehta MA, Murphy DG, Catani M, et al. Double-dissociation between the mechanism leading to impulsivity and inattention in Attention Deficit Hyperactivity Disorder: a resting-state functional connectivity study. *Cortex* 2017;86:290–302.
- [25] Knaus TA, Silver AM, Kennedy KA, Lindgren KA, Dominick KC, Siegel J, et al. Language laterality in autism spectrum disorder and typical controls: a functional, volumetric, and diffusion tensor MRI study. *Brain Lang* 2010;112:113–20.
- [26] Oertel V, Knochel C, Rotarska-Jagiela A, Schonmeyer R, Lindner M, van de Ven V, et al. Reduced laterality as a trait marker of schizophrenia – evidence from structural and functional neuroimaging. *J Neurosci* 2010;30:2289–99.
- [27] Herve PY, Zago L, Petit L, Mazoyer B, Tzourio-Mazoyer N. Revisiting human hemispheric specialization with neuroimaging. *Trends Cogn Sci* 2013;17:69–80.
- [28] Nachev P, Rees G, Frackowiak R. Lost in translation. *F1000Res* 2018;7:620.
- [29] Kong XZ, Mathias SR, Guadalupe T, Glahn DC, Franke B, et al. ELW Group. Mapping cortical brain asymmetry in 17,141 healthy individuals worldwide via the ENIGMA Consortium. *Proc Natl Acad Sci USA* 2018;115:E5154–63.
- [30] McManus C. Right hand, left hand. London: Phoenix; 2003.

- [31] Bartolomeo P, Thiebaut de Schotten M. Let thy left brain know what thy right brain doeth: inter-hemispheric compensation of functional deficits after brain damage. *Neuropsychologia* 2016.
- [32] Vallortigara G, Rogers LJ. Survival with an asymmetrical brain: advantages and disadvantages of cerebral lateralization. *Behav Brain Sci* 2005;28:575–89. discussion 89–633.
- [33] Friston KJ, Price CJ. Degeneracy and redundancy in cognitive anatomy. *Trends Cogn Sci* 2003;7:151–2.
- [34] Croxson PL, Forkel SJ, Cerliani L, Thiebaut de Schotten M. Structural variability across the primate brain: a cross-species comparison. *Cereb Cortex* 2018;28:3829–41.
- [35] Milham MP, Ai L, Koo B, Xu T, Amiez C, Balezeau F, et al. An open resource for non-human primate imaging. *Neuron* 2018;100:61–74. e2.