

Review

Diverse Genome Topologies Characterize Dosage Compensation across Species

William Jordan III,¹ Leila E. Rieder,^{1,2} and Erica Larschan^{1,*}

Dosage compensation is the process by which transcript levels of the X chromosome are equalized with those of autosomes. Although diverse mechanisms of dosage compensation have evolved across species, these mechanisms all involve distinguishing the X chromosome from autosomes. Because one chromosome is singled out from other chromosomes for precise regulation, dosage compensation serves as an important model for understanding how specific *cis*-elements are identified within the highly compacted 3D genome to co-regulate thousands of genes. Recently, multiple genomic approaches have provided key insights into the mechanisms of dosage compensation, extending what we have learned from classical genetic studies. In the future, newer genomic approaches that require little starting material show great promise to provide an understanding of the heterogeneity of dosage compensation between cells and how it functions in nonmodel organisms.

Composition of Chromatin

Within the eukaryotic cell nucleus, each chromosome consists of a single molecule of double-stranded DNA. Frequently thought of in linear space, these DNA molecules are compacted within the confines of the nucleus, as much as 10 000-fold, by adopting diverse 3D conformations. Despite extensive compaction of the chromosomes within the nucleus, specific *cis*-elements need to be precisely identified to regulate genes spatially and temporally.

At the most basic level, chromatin consists of a repeating array of nucleosomes, which are octamers of histone proteins, around which DNA is wrapped. There are two copies of each of the four core histone proteins (H2A, H2B, H3, and H4) within each nucleosome, which is surrounded by 146 bp of DNA [1,2]. Chromatin serves as the substrate for essential cellular processes such as transcription, replication, recombination, and cell division [1]. The idea of crumpled polymer globules, or the beads on a string framework for chromatin, was initially introduced 40 years ago [3,4]. Surprisingly, chromatin only accounts for roughly half of the available nuclear space [5]; specialized nuclear bodies and diverse chromatin regulators reside in the remaining interchromatin space. Nuclear bodies are dynamic structures that often reflect the transcriptional state of a cell and serve as sites of transcription and/or RNA processing [5,6].

Chromatin can be generalized into two contrasting types: euchromatin, which generally resides in the interior of the nucleus, and heterochromatin, which generally resides close to the nuclear envelope. These types of chromatin differ in their accessibility: euchromatin is regarded as active as it contains gene-rich regions, and positively correlates with chromatin accessibility, and heterochromatin is regarded as inactive, is gene poor, and negatively correlates with chromatin accessibility [7–9]. Chromatin may vary dramatically between cell types as histone proteins, histone modifiers (or readers/writers), and other transcription

Highlights

Dosage compensation equalizes transcript levels on the X chromosome with those on autosomes.

Dosage compensation serves as a model for understanding how specific *cis*-elements are collectively targeted for coordinated regulation within the genome.

Genomic approaches to study chromatin and 3D genome organization have provided new insights into dosage compensation.

¹Department of Molecular Biology, Cellular Biology and Biochemistry, Brown University, Providence, RI, USA

²Department of Biology, Emory University, Atlanta, GA, USA

*Correspondence: erica_larschan@brown.edu (E. Larschan).

factors are able to recognize and modulate specific loci, further differentiating chromatin compartments and fine-tuning gene expression. Therefore, while chromatin broadly influences gene regulation, it does not fully explain the modulation in activity of individual genes; rather chromatin establishes local environments that are more or less favorable for gene-specific expression [10–13]. In contrast, topologically associated domains (TADs), are physical megabase partitions of the genome. Within an organism, TAD organization is believed to be largely invariant and conserved across cell types [2]. Smaller domains are known by various names, including sub-TADs, chromatin loops, and neighborhoods. These smaller domains may be highly variable between cell types [10,12] and thus play a fundamental role in genomic function and regulation.



Across species, TAD boundaries are enriched for insulator elements, which are bound by a specialized class of insulator proteins that regulate 3D genome organization [9,14]. TAD boundary regions are also enriched for active genes, though recent studies in *Drosophila* suggest 3D chromatin organization is established during zygotic genome activation, independent of transcription [15]. Few factors, including chromatin regulators and noncoding RNAs, modulate genome organization [9]. Therefore, much remains unknown regarding how specific nuclear domains are established and how domains are targeted to modulate gene regulation.

Overview of Dosage Compensation

Diverse models of sex chromosome dosage compensation, in which the sex chromosomes of one sex are specifically targeted and transcriptionally regulated, serve as model systems in which to study how 3D genome organization regulates the expression of large groups or co-regulated genes. Analyses of dosage compensation across species, described below, have revealed the mechanisms by which large groups of genes are co-regulated within the context of 3D genome organization.

Sex chromosome dosage compensation is an essential, yet diverse, process in many species. There exist many distinct dosage compensation strategies to equalize transcript levels of genes located on the sex chromosomes between the sexes [16,17]. A conserved first step in heterogametic XY species, including humans and *Drosophila*, is distinguishing the X chromosome from the autosomes. For example, in many mammals, one female X chromosome is singled out for inactivation. A conserved dosage compensation mechanism, shared across several species including mammals, *Caenorhabditis elegans* and *Drosophila*, is upregulation of transcription of genes on the active X chromosome to equalize their expression with that of autosomal genes [16–18] (Figure 1).

In contrast to the X chromosome, there are no known examples in which the Y chromosome is targeted for dosage compensation. The Y chromosome is small and heterochromatic, although in some species it does carry specialized genes that have male sex-specific functions [19]. X-chromosome upregulation was first proposed by Ohno, who hypothesized that transcription of the X chromosome in mammals was upregulated during early evolution of sex chromosomes in order to compensate for degeneration of the Y chromosome [20]. Furthermore, Ohno theorized X-chromosome upregulation was not limited to males but also occurred in females. Ohno proposed that X-chromosome downregulation evolved in mammals in order to counteract X upregulation in XX females, therefore maintaining equalized dosage with autosomes [20]. This idea is widely known as Ohno's hypothesis. Data exist that both support and oppose the existence of such a mechanism across species [18,20–22].

Organism	Sex Chr.	Dosage Compensation	3-D Chromosome Changes
(A) <i>C. elegans</i> 	XX, AA	1. X-upregulation in both sexes: XX , AA XO , AA	Dosage compensation results in long-range contacts on—and compaction of—the X-chromosome.
	XO, AA	2. DCC-mediated X-down-regulation in hermaphrodites: XX , AA XO , AA	
(B) <i>Drosophila</i> 	XX, AA	MSL-mediated X-upregulation in males: XX, AA XY , AA	Long-range contacts on the X-chromosome exist in both sexes.
	XY, AA		
(C) Mouse 	XX, AA	1. Xist-mediated X-inactivation in females: X •, AA XY, AA	Xist spreading leads to additional long-range contacts to the XIST locus. The X-chromosome is highly compacted into the Barr body.
	XY, AA	2. Upregulation of active X in both sexes, mechanism unknown: X •AA XY , AA	

Trends in Genetics

Figure 1. Dosage Compensation Strategies: The Dosage Compensated Sex Is Represented in Red.

Mechanisms of Dosage Compensation

In the nematode worm, *C. elegans*, males carry a single X chromosome and two copies of each autosome (XO and AA), while hermaphroditic females carry two copies of the X chromosome and each autosome (XX and AA). An unknown mechanism is thought to upregulate expression of the X chromosome in both sexes. A well-defined dosage compensation complex (DCC) then downregulates gene expression for both X chromosomes in hermaphrodites to equalize X-linked gene expression to that of males [16,17,23–25] (Figure 1A). The DCC initially localizes to a small number of X-linked recruitment element on X (*rex*) sites, which are located in regions of euchromatin and contain binding motifs that cluster in 2D and 3D space [25]. The process of *C. elegans* dosage compensation is believed to occur through X-chromosome compaction as compaction is lost in the absence of DCC activity and the X chromosome becomes enlarged. Enlargement of the X chromosome may contribute to upregulation of X-linked genes by a mechanism that likely involves the function of specific X-linked DNA sequences rather than global regulation of all genes on the X chromosome [22,26].

In the fruit fly *Drosophila melanogaster*, males carry a single X chromosome and two copies of each autosome (XY and AA), while females carry two copies of the X chromosome and each autosome (XX and AA) (Figure 1B). A dosage compensation complex known as the male specific lethal complex (MSL), is specifically assembled in males and upregulates transcription of the single male X chromosome to approach levels of female X-linked gene expression. Similar to *C. elegans*, the initial sites of MSL recruitment, chromatin entry sites (CESs), are located in euchromatin and contain binding motifs that cluster in 2D and 3D space [14,27,28]. Furthermore, H4K16ac is important for increasing transcript levels on the male X chromosome [29,30].

In mammals, such as mice and humans, males carry a single X chromosome and two copies of each autosome (XY and AA), while females carry two copies of the X chromosome and each autosome (XX and AA) (Figure 1C). In females, one copy of the X chromosome is randomly

inactivated in a process known as X inactivation (XI), which normalizes female X-linked gene expression with that of males [16]. XI is characterized by extensive DNA methylation that is thought to maintain the inactive state [31].

In mammals, the inactive female X chromosome forms two large domains. In *C. elegans*, compaction along the entire length of both X chromosomes in the hermaphrodite is associated with reduced transcription levels [26]. In *Drosophila*, the X chromosome exhibits enhanced accessibility in both males and females independent of MSL [32]. In the future, 3D techniques could be applied to reveal which species-specific factors are important for setting up and or maintaining the 3D chromosome conformation many dosage compensation complexes use to target an entire chromosome.

Techniques for Probing Chromatin Structural Dynamics




Dosage compensation requires the targeting and regulation of an entire chromosome. Therefore, sex chromosomes represent a model that provides great insights into how smaller chromatin domains are targeted. Many improved techniques have had a substantial impact on the study of chromatin structure and function. While most of our understanding of dosage compensation comes from classical genetics and microscopy, which have been powerful, these approaches do not address the heterogeneity of dosage compensation across cells and tissue types. Further application of the new approaches described below will allow for even greater insight into how this fundamental process functions at high resolution in single cells.

Chromosome Conformation Capture (3C)

The development of 3C and the subsequent development of the current 3C family of techniques – chromosome conformation capture on chip (4C), chromosome capture carbon copy (5C), and genome-wide chromosome conformation capture (Hi-C) [33–35] – have significantly contributed to our understanding of how 3D genome structure contributes to genomic function (Figure 2A). There have been many recent improvements in this family of techniques. For example, *in situ* Hi-C may be performed in intact nuclei, which reduces experimental noise due to random ligation events and enables the production of higher-resolution data from fewer reads compared to the original technique (dilution Hi-C) [36]. 4C has also been improved by the incorporation of unique molecular identifiers, which reduce the impact of PCR-inflated ligation read counts, allowing the generation of 4C profiles with as few as 100 000 reads per bait [37] compared with 500 000 to 3 million reads for traditional 4C-seq [38].

Several groups have used chromosome conformation capture approaches to study dosage compensation in mammals [39–42], *C. elegans* [25,43], and *Drosophila* [14,44]. Fluorescent *in situ* hybridization (FISH) and Hi-C on *C. elegans* embryos revealed that the long-range contacts seen between DCC recognition sites are also lost in the absence of dosage compensation [43]. This likely occurs because X-chromosome remodeling that occurs during active dosage compensation is dependent on a condensin I^{DC} subcomplex that is part of the DCC [26,43]. Furthermore, the histone modifiers SET-1, SET-4, and SIR-2.1 are also necessary for dosage compensation X-chromosome remodeling [26,45,46]. Hi-C was performed in several *Drosophila* cell lines and embryos and discovered that, in contrast to *C. elegans*, MSL likely takes advantage of a pre-existing 3D topology, as long-range contacts between CES are present in both sexes and are independent of dosage compensation [14,26,44]. However, the mechanism by which the long-range contacts between CES are established in both males and females remains unknown.

RNA antisense purification in mouse lung fibroblasts in conjunction with Hi-C analysis revealed that, upon initiation of XI, the long noncoding RNA Xist utilizes the existing 3D topology of the X chromosome in order to spread from the XIST locus to distant loci [39]. The spreading of Xist

Technique	Insights	DC advances	Future advances
(A) Chromosome conformation capture (3C, 4C, 5C, Hi-C)	<p>High-resolution 3-D contacts</p>  <p>Dosage compensated chromosome</p>	Determine the relationship between structure of the DC chromosome and the dosage compensation machinery.	Determine how the 3-dimensional structure of the dosage compensated X-chromosome differs from that of autosomes.
(B) Micrococcal nuclease (MNase)-seq Assay for transposon-Accessible chromatin (ATAC)-seq	<p>Chromatin accessibility and nucleosome positioning</p> 	Determine the relationship between accessibility of the DC chromosome and the dosage compensation machinery.	Determine how dosage compensated X-chromosome accessibility differs from that of autosomes.
(C) Cleavage under targets & release under nuclease (CUT&RUN)	<p>High-resolution interaction (direct vs. indirect) 3-D contact mapping</p> 	None so far.	Identify factor binding sites and 3-dimensional interactions of dosage compensation regulators.
(D) <i>In situ</i> single cell or fluorescence activated cell sorting (FACS)	<p>Cell-to-cell heterogeneity</p> 	None so far.	Define how dosage compensation functions differently between cell types and tissues.

Trends in Genetics

Figure 2. Emerging Technologies and Their Current Impacts within the Dosage Compensation Field.

leads to structural changes that cause new regions of the chromosome to come in closer proximity to the XIST locus; this then allows the XIST RNA to transfer to these newly proximal sites, eventually coating the entire chromosome [39]. The inactive X chromosome eventually takes on a unique, compacted structure known as a Barr body [47].

The inactive X is enriched for the histone marks H3K9me3 and H3K27me3, although it appears that H3K27me3 is not necessary for XI [31]. The compacted state of the inactive X requires the presence of two proteins, a heterochromatin protein 1 binding protein (HBP1) and structural maintenance of chromosomes hinge domain-containing protein (SMCHD1) [31]. Based on allele-specific Hi-C and RNA-seq experiments, SMCHD1 antagonizes TAD boundary formation and is required for gene silencing during a specific developmental window; it is not required for maintenance, which may occur through other redundant pathways [48].

Using allele-specific Hi-C analysis, it was demonstrated that both X chromosomes display classical TAD topology prior to XI [40]. However, after XI, the inactive X chromosome adopts a unique 3D structure, which consists of two large megabase domains, termed superdomains, separated by a boundary region [40]. Within these two superdomains, classical TAD structure is rare, but does occur at gene clusters that escape XI [40]. Hi-C data demonstrates that these structures are conserved in humans [36]. However, the genomic content of the two domains differs between humans and mice [41]. CRISPR deletions of the boundary locus in mice results in fusion of the two superdomains, but the boundary is not necessary for the formation of XI in either mice or humans [36,41,42]. Furthermore, XI has been studied more extensively but recent work suggests that X upregulation also occurs in mammals [18,49].

Micrococcal Nuclease Sequencing (MNase-seq) and Assay for Transposase-Accessible Chromatin Using Sequencing (ATAC-seq)

MNase-seq and ATAC-seq are two popular methods for probing chromatin accessibility genome-wide [50–52] (Figure 2B). ATAC-seq is a quick, cost-effective, two-step procedure

that is able to identify regions of open chromatin (highly accessible regions) and map nucleosome positioning and transcription factor binding sites [50]. The latest version of MNase-seq, which utilizes a titration series, is also able to probe nucleosome positioning as well as relative chromatin accessibility (lowest to highest regions of accessibility) genome-wide [53]. While both techniques offer the ability to determine regions of open chromatin, MNase-seq with titration series offers the added benefit of providing relative chromatin accessibility, allowing diverse regulatory regions throughout the genome to be directly compared.

MNase-seq in *Drosophila* cultured cells reveals that MSL regulates chromatin accessibility by modulating nucleosome positioning at CES [14]. Furthermore, a non-sex-specific dosage compensation protein, chromatin-linked adaptor for MSL proteins (CLAMP), regulates chromatin accessibility at CES as well as globally across the entire male X chromosome [32].

Cleavage Under Targets and Release Using Nuclease (CUT&RUN)

CUT&RUN is a modification of chromatin immunoprecipitation (ChIP) that allows for simultaneous high-resolution chromatin mapping of specific factor localization and probing of the local chromatin environment [54] (Figure 2C). CUT&RUN is performed *in situ* under native conditions. It requires only a tenth of the sequencing depth of traditional ChIP-seq and has the ability to map histone modifications, even within compacted chromatin [54]. Furthermore, CUT&RUN may be used to map contact sites at near-base-pair resolution and differentiate direct protein binding sites versus indirect sites resulting from long-range interactions [54]. This approach has not yet been applied to dosage compensation, however, it has potential to contribute to better understanding the targeting requirements of various dosage compensation complexes by more accurately identifying binding sites and 3D interactions between dosage compensation regulators across species.

Single-Cell Techniques

Until recently, performing many of the techniques discussed above required large pools of cells. Fluorescence-activated cell sorting allows specific marked cell populations to be assayed, but still requires a large number of input cells [55] (Figure 2D). More recently, the modernization of genomic approaches allows them to be performed *in situ*. This has led to the ability to assay the genomic landscape in single cells. Single cell approaches are powerful; no longer is an average reading recorded from a bulk sample. As a result, cell to cell heterogeneity is uncovered and cells may be clustered into a number of unique groups.

Single-cell techniques are currently being used to probe chromatin accessibility [56,57], DNA variation [58], transcription factor binding sites [59], methylation [60], and 3D genome structure [61]. Additionally, microscopy-based approaches such as single-molecule super-resolution microscopy combined with Oligopaints – specialized FISH probes that allow for super-resolution microscopy and visualization of chromatin organization [62] – and assay of transposase accessible chromatin with visualization (ATAC-see) – a modification of ATAC-seq that reveals open chromatin regions targeted by transposons [63] – allow for single-cell visualization of chromatin architecture. Single-cell methods have not yet been applied to study dosage compensation. However, these approaches have potential to define how it functions in different tissues and nonmodel organisms.

Concluding Remarks and Future Perspectives

Dosage compensation is an essential process across species that evolved rapidly to equalize transcript levels from the X chromosome and autosomes. Dosage compensation systems provide excellent models for defining how a specific part of the genome may be regulated in a

Outstanding Questions

How is the X chromosome specifically distinguished from autosomes across species?

How heterogeneous is dosage compensation across cell types and tissues?

How does dosage compensation function across a wide array of species including non-model organisms?

context-specific way. Moreover, although diverse dosage compensation mechanisms have evolved, they all share the common feature of specifically distinguishing the X chromosome from autosomes. In the future, applying the single-cell genomic and imaging techniques described above to study how the X chromosome is distinguished for specific regulation will reveal how heterogeneous this process is across cell types and organisms. Although little is understood about how dosage compensation works beyond classical genetic model organisms (see Outstanding Questions), this new frontier of techniques that require little starting material will allow us to study how sex chromosomes differ from autosomes beyond traditional model organisms, which is key to understanding such a rapidly evolving process.

References

- Widom, J. (1989) Toward a unified model of chromatin folding. *Annu. Rev. Biophys. Biophys. Chem.* 18, 365–395
- Dixon, J.R. *et al.* (2016) Chromatin domains: the unit of chromosome organization. *Mol. Cell* 62, 668–680
- Grosberg, A.Y. *et al.* (1988) The role of topological constraints in the kinetics of collapse of macromolecules. *J. Phys.* 49, 2095–2100
- Grosberg, A. *et al.* (1993) Crumpled globule model of the three-dimensional structure of DNA. *EPL (Europhys. Lett.)* 23, 373
- Rajapakse, I. and Groudine, M. (2011) On emerging nuclear order. *J. Cell Biol.* 192, 711–721
- Aumiller, W.M., Jr *et al.* (2014) Phase separation as a possible means of nuclear compartmentalization. *Int. Rev. Cell Mol. Biol.* 307, 109–149
- Gilbert, N. *et al.* (2004) Chromatin architecture of the human genome: gene-rich domains are enriched in open chromatin fibers. *Cell* 118, 555–566
- Solovei, I. *et al.* (2013) LBR and lamin A/C sequentially tether peripheral heterochromatin and inversely regulate differentiation. *Cell* 152, 584–598
- Bonev, B. and Cavalli, G. (2016) Organization and function of the 3D genome. *Nat. Rev. Genet.* 17, 661
- Dixon, J.R. *et al.* (2015) Chromatin architecture reorganization during stem cell differentiation. *Nature* 518, 331
- Li, L. *et al.* (2015) Widespread rearrangement of 3D chromatin organization underlies polycomb-mediated stress-induced silencing. *Mol. Cell* 58, 216–231
- Barutcu, A.R. *et al.* (2015) Chromatin interaction analysis reveals changes in small chromosome and telomere clustering between epithelial and breast cancer cells. *Genome Biol.* 16, 214
- Cavalli, G. and Misteli, T. (2013) Functional implications of genome topology. *Nat. Struct. Mol. Biol.* 20, 290
- Ramírez, F. *et al.* (2015) High-affinity sites form an interaction network to facilitate spreading of the MSL complex across the X chromosome in *Drosophila*. *Mol. Cell* 60, 146–162
- Hug, C.B. *et al.* (2017) Chromatin architecture emerges during zygotic genome activation independent of transcription. *Cell* 169, 216–228
- Ercan, S. (2015) Mechanisms of X chromosome dosage compensation. *J. Genomics* 3, 1–19
- Deng, X. *et al.* (2011) Evidence for compensatory upregulation of expressed X-linked genes in mammals, *Caenorhabditis elegans* and *Drosophila melanogaster*. *Nat. Genet.* 43, 1179–1185
- Deng, X. *et al.* (2014) X chromosome regulation: diverse patterns in development, tissues and disease. *Nat. Rev. Genet.* 15, 367
- Graves, J.A.M. (2006) Sex chromosome specialization and degeneration in mammals. *Cell* 124, 901–914
- Veitia, R.A. *et al.* (2015) X chromosome inactivation and active X upregulation in therian mammals: facts, questions, and hypotheses. *J. Mol. Cell Biol.* 7, 2–11
- Albritton, S.E. *et al.* (2014) Sex-biased gene expression and evolution of the X chromosome in nematodes. *Genetics* 197, 865–883
- Wheeler, B.S. *et al.* (2016) Chromosome-wide mechanisms to decouple gene expression from gene dose during sex-chromosome evolution. *eLife* 5, e17365
- Meyer, B. (2005) *X Chromosome Dosage Compensation*, WormBook
- Pferdehirt, R.R. *et al.* (2011) An MLL/COMPASS subunit functions in the *C. elegans* dosage compensation complex to target X chromosomes for transcriptional regulation of gene expression. *Genes Dev.* 25, 499–515
- Albritton, S.E. *et al.* (2017) Cooperation between a hierarchical set of recruitment sites targets the X chromosome for dosage compensation. *eLife* 6, e23645
- Lau, A.C. *et al.* (2014) The *C. elegans* dosage compensation complex mediates interphase X chromosome compaction. *Epigenetics Chromatin* 7, 31
- Alekseyenko, A.A. *et al.* (2008) A sequence motif within chromatin entry sites directs MSL establishment on the *Drosophila* X chromosome. *Cell* 134, 599–609
- Kuzu, G. *et al.* (2016) Expansion of GA dinucleotide repeats increases the density of CLAMP binding sites on the X-chromosome to promote *Drosophila* dosage compensation. *PLoS Genet.* 12, e1006120
- Armstrong, R.L. *et al.* (2018) Chromatin conformation and transcriptional activity are permissive regulators of DNA replication initiation in *Drosophila*. *Genome Res.* 28, 1688–1700
- Copur, Ö. *et al.* (2018) Sex-specific phenotypes of histone H4 point mutants establish dosage compensation as the critical function of H4K16 acetylation in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* 115, 13336–13341
- Nozawa, R.-S. *et al.* (2013) Human inactive X chromosome is compacted through a PRC2-independent SMCHD1–H3K9me3 pathway. *Nat. Struct. Mol. Biol.* 20, 566
- Urban, J. *et al.* (2017) Enhanced chromatin accessibility of the dosage compensated *Drosophila* male X-chromosome requires the CLAMP zinc finger protein. *PLoS One* 12, e0186855
- Dekker, J. *et al.* (2002) Capturing chromosome conformation. *Science* 295, 1306–1311
- Simonis, M. *et al.* (2006) Nuclear organization of active and inactive chromatin domains uncovered by chromosome conformation capture–on-chip (4C). *Nat. Genet.* 38, 1348
- Lieberman-Aiden, E. *et al.* (2009) Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326, 289–293
- Rao, S.S.P. *et al.* (2014) A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 159, 1665–1680
- Schwartzman, O. *et al.* (2016) UMI-4C for quantitative and targeted chromosomal contact profiling. *Nat. Methods* 13, 685
- Van De Werken, H.J.G. *et al.* (2012) Robust 4C-seq data analysis to screen for regulatory DNA interactions. *Nat. Methods* 9, 969
- Engreitz, J.M. *et al.* (2013) The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science* 341, 1237973
- Giorgetti, L. *et al.* (2016) Structural organization of the inactive X chromosome in the mouse. *Nature* 535, 575
- Deng, X. *et al.* (2015) Bipartite structure of the inactive mouse X chromosome. *Genome Biol.* 16, 152

42. Darrow, E.M. *et al.* (2016) Deletion of DXZ4 on the human inactive X chromosome alters higher-order genome architecture. *Proc. Natl. Acad. Sci.* 113, E4504–E4512
43. Crane, E. *et al.* (2015) Condensin-driven remodelling of X chromosome topology during dosage compensation. *Nature* 523, 240–244
44. Schauer, T. *et al.* (2017) Chromosome topology guides the *Drosophila* Dosage Compensation Complex for target gene activation. *EMBO Rep.* 18, 1854–1868
45. Vielle, A. *et al.* (2012) H4K20me1 contributes to downregulation of X-linked genes for *C. elegans* dosage compensation. *PLoS Genet.* 8, e1002933
46. Wells, M.B. *et al.* (2012) *Caenorhabditis elegans* dosage compensation regulates histone H4 chromatin state on X chromosomes. *Mol. Cell. Biol.* 32, 1710–1719
47. Barr, M.L. and Bertram, E.G. (1949) A morphological distinction between neurones of the male and female, and the behaviour of the nucleolar satellite during accelerated nucleoprotein synthesis. *Problems Birth Defects* 101–102 Springer
48. Gdula, M.R. *et al.* (2019) The non-canonical SMC protein SmcHD1 antagonises TAD formation and compartmentalisation on the inactive X chromosome. *Nat. Commun.* 10, 30
49. Deng, X. *et al.* (2013) Mammalian X upregulation is associated with enhanced transcription initiation, RNA half-life, and MOF-mediated H4K16 acetylation. *Dev. Cell* 25, 55–68
50. Buenrostro, J.D. *et al.* (2013) Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat. Methods* 10, 1213
51. Zhang, Z. and Pugh, B.F. (2011) High-resolution genome-wide mapping of the primary structure of chromatin. *Cell* 144, 175–186
52. Tolstorukov, M.Y. *et al.* (2010) Analysis of the primary structure of chromatin with next-generation sequencing. *Epigenomics* 2, 187–197
53. Mieczkowski, J. *et al.* (2016) MNase titration reveals differences between nucleosome occupancy and chromatin accessibility. *Nat. Commun.* 7, 11485
54. Skene, P.J. and Henikoff, S. (2017) An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites. *eLife* 6, e21856
55. Griffiths, J.A. *et al.* (2018) Using single-cell genomics to understand developmental processes and cell fate decisions. *Mol. Syst. Biol.* 14, e8046
56. Cusanovich, D.A. *et al.* (2015) Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* 348, 910–914
57. Buenrostro, J.D. *et al.* (2015) Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* 523, 486
58. Zong, C. *et al.* (2012) Genome-wide detection of single-nucleotide and copy-number variations of a single human cell. *Science* 338, 1622–1626
59. Rotem, A. *et al.* (2015) Single-cell ChIP-seq reveals cell subpopulations defined by chromatin state. *Nat. Biotechnol.* 33, 1165
60. Farlik, M. *et al.* (2015) Single-cell DNA methylome sequencing and bioinformatic inference of epigenomic cell-state dynamics. *Cell Rep.* 10, 1386–1397
61. Nagano, T. *et al.* (2013) Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature* 502, 59
62. Beliveau, B.J. *et al.* (2015) Single-molecule super-resolution imaging of chromosomes and *in situ* haplotype visualization using Oligopaint FISH probes. *Nat. Commun.* 6, 7147
63. Chen, X. *et al.* (2016) ATAC-seq reveals the accessible genome by transposase-mediated imaging and sequencing. *Nat. Methods* 13, 1013