

PE81 3D real time image reproduction of the prostate: Can it be used on virtual reality (VR) headsets and/or tilepro of Da Vinci surgical system as a guide during robotic radical prostatectomy?

EUR Urol Suppl 2019;18(6):e2660


Aksoy S.F.¹, Altinmakas E.², Canda A.E.³, Koseoglu E.³, Esen T.⁴


¹Sabancı University, Collaboration Space, Istanbul, Turkey, ²Koç University, School of Medicine, Dept. of Radiology, Istanbul, Turkey, ³Koç University, School of Medicine, Dept. of Urology, Istanbul, Turkey, ⁴Koç University, School of Medicine and VKF American Hospital, Dept. of Urology, Istanbul, Turkey

Introduction & Objectives: To create 3D prostate images that can be transferred to virtual reality (VR) headsets & tilepro of Da Vinci surgical system.

Materials & Methods: Axial T2W 3D-TSE sequence was used in multiparametric prostate magnetic resonance imaging (MRI). Borders of the tumor(s) and anatomical structures were marked by the uro-radiologist. In order to obtain the accurate segmentation of the tumor(s) and surrounding anatomical structures, interactive medical image segmentation methods were used such as, fast growcut algorithm and morphological contour interpolation. Simple region growing techniques were used after primary applications for improving the results of the initial methods. Urinary bladder, prostate (peripheral zone was separated) and urethra were segmented to give a better understanding.

Results: Tumor 1, Fig 1. A 2cm markedly hypointense mass arising from the right peripheral zone of the midgland is seen on axial T2W images with restricted diffusion on DWI and ADC map. Mean ADC value is $570 \times 10^{-6} \text{ mm}^2/\text{s}$, PIRADS5. Tumor 2, Fig 2. A 0.8cm lesion is noted at the left base of prostate gland with markedly hypointense signal on T2W images and restricted diffusion on DWI and ADC map. Mean ADC value is $760 \times 10^{-6} \text{ mm}^2/\text{s}$, PIRADS4. 3D images were constructed and were transferred to VR headsets. Tumor 1 had 851.781 mm^2 surface area, 1762.45 mm^3 and 1.76 cm^3 volumes. Tumor 2 had 129.258 mm^2 surface area, 123.931 mm^3 and 0.12 cm^3 volumes.





Conclusions: 3D image representation of the prostate and transfer to VR headsets and/or tilepro of Da Vinci surgical system might be used as a promising tool to help the console surgeon to better understand the size, location and characteristics of the tumor(s) and the prostate. It might be used as a guide during robotic radical prostatectomy and might have an impact on the learning curve and oncologic outcomes.