

$p = 0.5$ ), cT2a (3.2 vs. 6.5%,  $p = 0.4$ ), and use of MRI during AS (55.8 vs. 55.2%,  $p = 0.9$ ) were seen. Overall, 45 (47.4%) and 76 (37.8%) patients aged  $\leq 60$ yr and  $> 60$ yr, respectively progressed during AS. Unadjusted progression-free rates at 2 years of followup were 29.9 vs. 30.2% for younger vs. older patients, respectively ( $p = 0.54$ ). At MVA analyses, younger age was not associated with increased risk of progression (HR: 1.27; 95%CI: 0.71–2.31;  $p = 0.4$ ). Finally, RP was performed in 30 (66.7%) younger and in 36 (47%) older patients, respectively. Pathological evaluation of RP patients showed adverse characteristics in 8 (24.2%) younger vs. 21 (51.2%) older patients, respectively ( $p = 0.03$ ).

**Discussion:** Even applying less stringent criteria for inclusion in AS, younger age was not associated with increased risk of progression in AS or with more adverse pathological characteristics at RP after AS. Young patients are those who may gain the highest benefit from an AS approach without being exposed to increasing risk of progression or misclassification.

## SC37

### Do we really need detailed biopsy assessment of patients with low-risk prostate cancer candidate to active surveillance? A prospective validation of the ISUP recommendation

M. Bandini, N. Suardi, S. Scarcella, L. Nocera, G. Gandaglia, N. Fossati, V. Cucchiara, E. Zaffuto, S. Shariat, N. Longo, V. Mirone, S. Scuderi, A. Larcher, D. Robesti, P. Karakiewicz, A. Rizzo, D. Cannoletta, A. Pellegrino, F. Barletta, F. Montorsi, A. Briganti (Milano)

**Aim of the study:** A recent report of the International Society of Urological Pathology (ISUP) recommend accurate histopathologic assessment at biopsy to properly select low-risk prostate cancer (PCa) patients suitable for active surveillance (AS). However, the outcome of AS is excellent even when all men with low risk PCa are considered regardless of the extent of cancer involvement at biopsy. Therefore, we examined whether such detailed biopsy assessment is really needed and capable to predict the risk of progression during AS in low risk patients.

**Materials and methods:** Between 2010–18, 248 patients with low risk PCa (PSA  $< 10$  ng/ml, cT  $\leq 2$ a and biopsy Gleason Score [GS] 6) were prospectively enrolled in our AS program. Patients referred from other institutions underwent central biopsy review. Only patients with all available ISUP criteria ( $n = 174$ ) were included. Uni- and multi-variable (UVA and MVA) Cox regression models tested the effect of each ISUP criteria (number of cores taken, number of positive cores per side and percentage of cancer in the most involved core) on the risk of progression during AS. Covariates included baseline PSA, prostate volume, cT stage, age and use of mp-MRI. Progression was defined as upgrade in GS ( $\geq 3 + 4$ ). Finally, C-index assessed and compared the discrimination of the models with and without ISUP criteria.

**Results:** Median number of cores examined, median % of positive cores, mean % of positive cores in dominant side and median % of PCa in dominant core were 17.5, 7.1%, 96.3% and 5%, respectively. 96 men (55.2%) had mp-MRI performed before confirmatory biopsy. Within a median follow-up of 27.5 months, 63 patients experienced progression. At UVA, only % of PCa in dominant core (HR 1.02;  $p = 0.03$ ), baseline PSA (HR 1.10;  $p = 0.03$ ) and mpMRI (HR 0.55;  $p = 0.03$ ) were associated with the risk of AS progression. At MVA, none of the examined ISUP criteria reached independent prediction status (all  $p > 0.2$ ). Only PSA and mp-MRI represented independent predictors (HR 1.17 and 0.3, respectively; all  $p < 0.005$ ). The same C-Index was obtained for the predictive models with and without ISUP criteria (0.79 for both models).

**Discussion:** Our results showed that none of the detailed biopsy features recommended by ISUP represented independent predictors of AS progression after accounting for the use of mp-MRI as well as

clinical variables. Therefore, although such variables might be considered reliable proxies of tumor volume and PCa multi-focality, their use is at least questionable in clinical practice for patients candidate to AS.

## SC38


### A novel nomogram to identify candidates for active surveillance among patients with International Society of Urological Pathology Grade Group 1 prostate cancer diagnosed with systematic biopsies and submitted to multiparametric magnetic resonance imaging

S. Luzzago, G. Cozzi, M. Catellani, A. Conti, F. Mistretta, I. Sabatini, E. Di Trapani, P. Pricolo, S. Alessi, A. Brescia, G. Cordima, D. Matei, G. Petralia, G. Musi, O. de Cobelli (Milano)

**Aim of the study:** To create a novel nomogram for including a greater proportion of men in active surveillance (AS), compared to Prostate Cancer Research International: Active Surveillance (PRIAS) criteria, without compromising AS oncologic safety.

**Materials and methods:** Overall, 1,103 patients with ISUP GG 1 prostate cancer submitted to radical prostatectomy and to multiparametric magnetic resonance imaging (mpMRI) were included. The outcome of interest was the presence of unfavourable disease (csPCa): ISUP GG  $\geq 3$  and/or pT  $\geq 3a$  and/or pN1. Logistic regression models including PRIAS (Model 0), and a multivariable model including PSA, cT, PSA-D, and the percentage of positive cores (PCP) (Model 1), were fitted; subsequently PI-RADS (Model 2), extracapsular (ECE) score (Model 3), and PI-RADS+ECE score (Model 4) were added to Model 1. Discrimination was measured by the area under the ROC curve (AUC). Decision-curve analyses (DCA) determined the clinical net benefit associated with the best performing model.

**Results:** Overall, 371 (33.6%) men presented csPCa. Moreover, 348 (31.6%) patients respected PRIAS criteria and, of them, 89 (25%) harboured csPCa. Model 4 had the greatest AUC (0.83 vs 0.56 of Model 0;  $p < 0.001$ ). Significant predictors (all  $p < 0.001$ ) at multivariate analysis were PSA-D (OR:1.2), ECE score (OR:8.2 for ECE score  $\geq 4$ ) and PI-RADS (OR:2.5 for PI-RADS 5). Moreover, PSA (OR:1.01), cT (OR:1.4) and PCP (OR:1.1) were included. The adoption of a risk score based on Model 4 improved clinical risk prediction against threshold probabilities of csPCa approximately between 21% and 75%. Using a 25% threshold would increase the proportion of AS eligible patients from 31.6% (PRIAS) to 62.3% with an estimated risk of csPCa of 13.4%



**Discussion:** A novel nomogram that includes clinical, biopsic and mpMRI findings, is able to increase of roughly  $\sim 30\%$  the absolute number of patients suitable for AS, compared to PRIAS, with an

estimated 13.4% risk of csPCa. External validation of our model is needed.

SC39

**Development of novel criteria for active surveillance based on multiparametric MRI alone in men with Gleason 3 + 4 prostate cancer: Use of imaging to safely expand the eligibility for active surveillance**

A. Briganti, G. Gandaglia, G. Ploussard, M. Valerio, A. Mattei, C. Fiori, N. Fossati, A. Stabile, J. Beauval, B. Malavaud, M. Roumiguié, P. Dell'Oglio, N. Suardi, M. Moschini, S. Zamboni, A. Rakauskas, V. Mirone, F. De Cobelli, F. Porpiglia, F. Montorsi (*Milano*)

**Aim of the study:** Clinical guidelines recommend active Surveillance (AS) in men with low-risk prostate cancer (PCa) and in some highly selected low-volume Gleason 3 + 4 within a prospective protocol. However, the latter recommendation is based on results obtained in the pre-MRI era. We hypothesized that mpMRI characteristics might assist physicians in the identification of Gleason 3 + 4 patients suitable for AS regardless of the number of positive cores.

**Materials and methods:** Overall, 309 patients with Gleason 3 + 4 PCa at biopsy who underwent MRI-targeted biopsy plus systematic cores and radical prostatectomy (RP) between 2016 and 2018 were identified. Prostate biopsy and prostatectomy specimens were evaluated by dedicated uro-pathologists. Misclassification was dened as non-organ coned or ISUP grade group  $\geq 3$  disease at RP. Multivariable logistic regression analyses assessed the association between mpMRI parameters (i.e., extracapsular extension [ECE] and maximum lesion diameter) and misclassification after adjusting for confounders. The diameter of the lesion at mpMRI was dichotomized according to the most informative cut-off predicting misclassification. A novel risk classification was proposed based clinical characteristics and mpMRI findings. A comparison of the proportion of patients eligible for AS and the rates of misclassification was performed between men selected for AS according to the new imaging criteria vs. established criteria of low volume Gleason 3 + 4 ( $\leq 2$  positive cores).

**Results:** Median lesion diameter was 8 mm and 39 (13.2%) patients had suspicious ECE at mpMRI. Overall, 185 (59.9%) patients experienced misclassification. At multivariable analyses, ECE at mpMRI ( $p=0.03$ ) and lesion diameter ( $p=0.01$ ) were associated with the risk of misclassification. The ROC-derived area under the curve of the model was 75%. The most informative cut-off of lesion diameter for misclassification was 12 mm. Patients without ECE at mpMRI and with a lesion  $<12$  mm were classed as low risk regardless of the number of positive cores. The proportion of patients potentially eligible for AS increased from 18.1 to 22.5% when adopting the new mpMRI criteria compared to the inclusion of only men with 1–2 positive cores. The rate of misclassification was lower in intermediate-risk patients selected according to the new mpMRI criteria vs. those selected according to the number of positive cores (44.6 vs. 53.6%).

**Discussion:** Information obtained at mpMRI can assist physicians in the identification of men with Gleason 3 + 4 more likely to experience misclassification. Men with biopsy Gleason 3 + 4 without ECE and with a lesion  $<12$  mm at mpMRI biopsy could be included in AS protocols regardless of the number of positive cores at biopsy thus expanding AS indications while decreasing the rates of misclassification at the same time.

SC40

**The role of magnetic resonance in active surveillance for prostate cancer. The romas project: Four years experience in a randomized, prospective study**

M. Guerra, R. Schiavina, M. Borghesi, L. Bianchi, P. Verze, A. Angiolini, F. Mineo Bianchi, S. Di Fresco, P. Piazza, R. La Rocca, L. Lambertini, V. Mirone, E. Brunocilla (*Bologna*)

**Aim of the study:** Multiparametric Magnetic Resonance Imaging (mpMRI) has joined increasing enthusiasm as a diagnostic tool able to identify clinically significant Pca (csPCa). However, the role of mpMRI for patients who are included in AS protocols is still a matter of debate. The aim of our study was to evaluate the impact of mpMRI in the early identification (within 3 months) of csPCa in patients fulfilling AS criteria.

**Materials and methods:** From May 2015 to March 2019, 93 very low and low risk PCa patients were included in our AS protocol. Inclusion criteria for AS were: 35–75 year-old men; diagnosis of PCa within 8 weeks; PSA  $\leq 10$  ng/ml; Clinical T1c or T2 assessed with DRE and TRUS; Gleason Score  $< 7$ ;  $< 3$  positive cores; at least 12 biopsy cores taken; PSA density (PSAd) less or equal to 0.2; ASA score  $\leq 3$ ; central revision of the biopsy core confirming the presence of indolent Pca. Patients were randomized (1:1) in two groups: group 1 ( $n=38$ ) were given an mpMRI at 3 months from the beginning of AS; group 2 (control group,  $n=33$ ) did not receive the mpMRI within 12 months of surveillance. Patients in both Groups had a PSA dosage every 3 months and underwent digital rectal examination (DRE) every 6 months. In group 1, patients with at least one PIRADS  $\geq 3$  lesion at mpMRI underwent a Fusion biopsy (FB) plus systematic random biopsy; patients with negative mpMRI (namely PIRADS  $\leq 2$  lesion at mpMRI) in group 1 and patients included in group 2 underwent a confirmation systematic biopsy 12 months after the beginning of the study and subsequent prostate biopsy at 48 and 72 months. In case of reclassification (both due to upgrading and to upsizing) patients were scheduled for radical treatments (including radical prostatectomy or primary radiotherapy).

**Results:** Table 1 shows the clinical characteristics of the entire patient population. The two groups of patients were comparable in terms of initial Age, PSA, PSA density, clinical stage, number of positive and total cores. Table 2 shows the percentage of reclassified patients at 3 and 12 months. Of the 48 patients who underwent mpMRI at 3 months, 14 (26.9%) had at least one significant lesions nine of these patients (18.7%) were reclassified: 8 (87.5%) men due to upgrading and 1 patient due to upsizing. At 12-month random re-biopsy, only 2 out of 25 patients in group 1 (8%) and 11 out of 33 patients in group 2 (33%) were reclassified ( $p=0.03$ ). Kaplan-Meier plots (figure 3) estimate active treatment free survival among the 2 groups.

**Discussion:** The early implementation of a mpMRI into an AS protocol was able to obtain earlier reclassification rates and to significantly reduce the percentage of patients who would be reclassified at 12 months with a systematic biopsy. Therefore, our data suggest to include a mpMRI in AS protocols in order to offer timely radical treatments in patients with csPCa. A larger study population and longer follow up may help to achieve more accurate results.