

outcomes and postoperative success rate. Our study, first in literature, proves the interest in assessing the safety of no drainage placement also in the pediatric field.

SC90

Does the body weight influence the outcome in children treated with robotic pyeloplasty?

A. Grosso, S. Sforza, A. Mari, R. Tellini, F. Di Maida, L. Viola, E. Rosi, M. Carini, A. Minervini, L. Masieri (Firenze)

Aim of the study: To investigate the influence of the body weight on peri and postoperative outcome in a series of pediatric patients with a diagnosis of ureteropelvic junction obstruction (UPJO) treated with robot-assisted laparoscopic pyeloplasty (RALP) at a single tertiary referral centre.

Materials and methods: Fifty consecutive patients treated with RALP from January 2016 to October 2018 were divided according to their weight: group A < 15 Kg, group B ≥ 15 Kg and were included in the present study. Eligible criteria for surgery were symptomatic UPJO, worsening of hydronephrosis or obstructive pattern at renogram. Success criteria were resolution of the hydronephrosis at imaging and absence of flank pain. All procedures were performed by one expert robotic surgeon.

Results: 14 patients were included in group A (median weight 12 Kg) and 36 patients in group B (median weight 36 Kg). We registered one (7.1%) Clavien 3b complication (omental hernia after removal of the drainage requiring surgical correction) and one (2.8%) Clavien 2 complication in group B (urinary infection). No difference has been found in length of hospital stay, length of catheterization, duration of procedure between the groups ($p > 0,05$). At a median follow up of 21.5 months, overall success rate was 98%. One patient needed nephrostomy positioning for persistent hydronephrosis.

Preoperative data		Kg = 15 (n ; %)			
		Kg < 15 (14; 28%)		Kg > 15 (36; 72%)	
Gender, n. %	Male	6	42,9%	21	58,3%
	Female	8	57,1%	15	41,7%
Side, n. %	Right	4	28,6%	16	44,4%
	Left	10	71,4%	20	55,6%
Age (months), median IQR		25	18-36	118	89-171
Weight (kg), median IQR		12	11-13,8	36	27,4-51
Symptoms at diagnosis, n. %		9	64,3%	32	88,9%
Anatomic variant, n. %		0	0%	9	25%
Crossing vessels, n. %		3	21,4%	15	41,7%
Preoperative Nephrostomy, n. %		1	7,1%	2	5,6%
APD preoperative, median IQR		27	18-36	30	24,5-41,5
Surgical and perioperative data		Kg < 15 (14; 28%)		Kg > 15 (36; 72%)	
		Kg < 15 (14; 28%)		Kg > 15 (36; 72%)	
Overall Operative time, median IQR		95	90-110	90	85-108
Console time, median IQR		75	70-85	73	65-80
Set-up time, median IQR		20	15-25	20	15-28
Drainage, n. %	yes	10	71,4%	14	38,9%
	No	4	28,6%	22	61,1%
Catheter, median IQR		3	3-3	3	3-3
APD postoperative, median IQR		12	9-18	12	10-14,5
LOS, median IQR		4	3-4	4	3-4
Follow-up months, median IQR		20	14-25	25	14-30
Readmission, n. %		1	7,1%	1	2,8%
Relapse, n. %		1	7,1%	0	0%
Complication, n. %		1	7,1%	1	2,8%

Table 1. APD, anterior-posterior pelvic diameter; LOS, length of stay

Differences in symptoms and anatomic variant are found statistically significant

There were no significant differences found in any of the parameter

SC91

Results and complications of retrograde approach (URS/RIRS) in pediatric urolithiasis: A multicentric experience

S. Ferretti, D. Campobasso, T. Bocchialini, L. Bevilacqua, G. Bianchi, C. Di Pietro, P. Granelli, S. Micali, B. Rocco, P. Salsi, M. Sighinolfi, U. Maestroni, A. Frattini (Parma (PR) – Italia)

Aim of the study: Incidence of pediatric urolithiasis is increasing in industrialized countries. Treatment of this disease follows the same surgical procedures as in adults.

Materials and methods: We have retrospectively reviewed all the kidney units (RUs) which underwent to URS/RIRS at 3 centers from January 2009 to December 2018 up to 16 years of age. Data reported include: lithiasic volume, radiological exposure, operative time, complications according to Clavien classification at 3 months and stone free rate.

Results: We have performed 74 procedures (44 RIRS and 30 URS) in 66 RUs with an average age of 7.24 years (range: 2–16 yy). RUs treated in pre-school age (0–4 yy) were 19; 34 in pre-puberty (5–11 yy) and 13 in puberty (12–16 yy). In 39 cases a pre-operative urethral stent was applied (17 cases with age <4, 18 between 5–11 and 4 between 12–16). Average lithiasic volume was 1.165 cm². Mean operative time was 77.3 ± 41.64 minutes (range: 15–140 min) with a radiological exposure of 25 “± 28” (range: 5 “–1’20”). At the end of the procedure a Double J (DJ) was applied in 45 cases (8 cases with age <4, 27 between 5–11 and 10 between 12–16) and a Mono J (MJ) in 29 cases (13 cases with age <4, 11 between 5–11 and 5 between 12–16). Stone-free rate has been 81.8% (54/66, 13 cases with age <4, 31 between 5–11 and 10 between 12–16). Nine patients required a second intervention to achieve a complete stone clearance. Complications recorded at 90 days include a case (1.5%) of migration of a fragment in the urethra causing hydronephrosis and renal colics at 49 days to the operation (Clavien 3b) that was treated with URS; 8 cases (12.1%) which required the administration of painkillers during the first 48 hours postoperative (Clavien I); 5 cases of urinary tract infections during hospitalization (7.5%) treated with antibiotic therapy (Clavien II).

Discussion: Miniaturization of the instruments with the increased incidence of pediatric urolithiasis have led the endourologist to treat more patients in pediatric age. However, particular attention should be paid to treatment planning in patients under 6 years of age, where, based on our experience it is useful to apply a urethral stent for kidney stones or proximal and mid-lumbar ureteral stones before surgery. In our cases, in only three patients under 6 years of age, no pre-operative DJ stent has been required, since it were two patients with stones located in the distal part of the ureter. Another consideration should be the choice of the urethral stent to be applied at the end of the intervention (DJ vs MJ) to avoid excessive use of painkillers or access to the hospital for intolerance, bearing in mind the need to submit the patient to an additional anesthesia for DJ removal. In 60% of patients under the age of 5 years we preferred to apply a MJ, leaving a DJ alone in cases requiring a second intervention, where the procedure lasted more than an hour or in presence of lithiasic sand related to the dusting of the stone.