

GUA-52	Results of application of radio-frequency ablation in treatment of malignant and beneficial new formations of parichmatous organs and bones
GUA-53	The role of minimally invasive treatment methods in patients with neoplasms of the bladder with macrohematuria
GUA-54	X-ray vascular treatment of prostate cancer
GUA-55	The quality of life in renal transplant recipients and dialysis patients
GUA-56	Tubeless mini PCNL with antegrade stent tether in preschool children
GUA-57	Transurethral enucleation of prostate by one-step en-bloc technique
GUA-58	Implementation of optimized ERAS protocols after radical cystectomy
GUA-59	Salvage cystectomy in patients with muscle-invasive bladder cancer
GUA-60	Intraoperative complications of radical cystectomy with different types of urinary diversion

#### GUA-01 Laparoscopic Radical Cystectomy with Intracorporeal Neobladder, Report of First Case in Georgia

Giorgi Khvadagiani, Anna Chikovani, George Khvadagiani  
Caucasus Medical Center, Tbilis, Georgia; Tbilisi Medical State University, Tbilis, Georgia

**Background:** Radical cystectomy remains the gold standard for muscle-invasive, organ-confined transitional cell carcinoma of the bladder. Laparoscopic and more recently robotic surgery has been used increasingly by urologists as a minimally invasive procedure to treat bladder cancer. The present study reports technical aspects of, to our knowledge, the first case of radical cystectomy with bilateral pelvic lymphadenectomy and intracorporeal ileal neobladder performed in Georgia.

**Case report:** The patient was a male, 69 years old with high grade muscle invasive urothelial carcinoma of the bladder with negative metastatic workout. He had undergone open cystolithotomy 8 years ago and has coronary artery stents. Laparoscopic radical cystectomy and pelvic lymphadenectomy were performed using 5 abdominal ports. Bipolar and ultrasonic energies were used for vascular control. 40 cm of ileum was taken. Neobladder formation, urethral and both ureteral anastomosis were done using the completely intracorporeal laparoscopic technique. Operative time was 420 minutes. Estimated blood loss was 340 ml. Lymphorea occurred for 10 days. Patient was discharged on day 5. Ureteric stents and urethral catheter were removed on day 21. The post-operative pathologic report indicated pT2b N0, 25 lymph nodes and free resection margins. On the 6 month of follow-up patients has no recurrence.

**Conclusion:** Our groups initial experience in Laparoscopic Radical cystectomy with intracorporeal ileal neobladder demonstrated that it was a feasible treatment modality. However, more cases are required in the future to confirm its efficacy and improve laparoscopic technical performance in the treatment of organ-confined urothelial cancer.

#### GUA-02 Anatomic endoscopic enucleation of prostate using bipolar energy and mushroom technique

Giorgi Khvadagiani, Anna Chikovani, George Khvadagiani  
Caucasus Medical Center, Tbilis, Georgia; Tbilisi Medical State University, Tbilis, Georgia

**Background:** Transurethral resection of the **prostate** has been considered as the gold standard for benign prostatic hyperplasia (BPH). Anatomic endoscopic enucleation of prostate, mainly using LASER technology, is becoming a new standard, as it causes less bleeding and theoretically has no upper limit of prostate volume. The downside of LASER is initial cost of equipment, morcellation and steep learning curve. Transurethral **enucleation** with **bipolar** has emerged as an alternative prostatic **enucleation** procedure. We present our initial experience in bipolar enucleation and mushroom resection technique.

**Materials and methods:** Thirty patients with BPH and indications for surgery underwent bipolar enucleation from May 2018 to December 2018. Patients with **prostate** size >40 g were selected. All surgeries were done by a single urologist. Preoperative and postoperative International **Prostate** Symptom Score (IPSS) scores, Qmax, duration of surgery, duration of **enucleation**, drop in hemoglobin, weight of resected tissue, and the incidence of stress urinary incontinence were measured.

**Results and limitations:** The mean age was 67 years and mean prostatic size was 70 gr. Sixteen patients had urinary retention. The mean IPSS score in remaining patients was 23.5. The mean preoperative maximal flow 10.2 mL/s. Mean duration of surgery was 94 min. Mean duration of enucleation was 45 min. Mean drop in hemoglobin was 0.9 g/dl. Mean weight of resected tissue was 52 g. 15 patients had transient stress urinary incontinence after surgery. All of them resolved after 3 weeks. Mean IPSS score after bipolar enucleation was 7.5 showing significant improvement in all aspects of IPSS. Mean postoperative Qmax was 25 mL/s.

Main limitations of the study are its small cohort size and non-randomized nature.

**Conclusion:** Bipolar enucleation is an effective surgical management of BPH. It allows enucleation of large adenomas in a single sitting, mimicking conventional open enucleation of the prostate while having all the advantages of a minimally invasive surgery. Mushroom technique is more time consuming than morcellation, but feasible in case of absence of equipment.

#### GUA-03 Percutaneous nephrolithotomy in Galdakao-modified supine Valdivia position in comparison to prone position: our initial experience

Giorgi Khvadagiani, Anna Chikovani, George Khvadagiani  
Caucasus Medical Center, Tbilis, Georgia; Tbilisi Medical State University, Tbilis, Georgia

**Background:** Despite the fact that it is almost two decades since the description of supine percutaneous nephrolithotomy (PCNL) technique, it is still underutilized. Aim of this study was to compare retrospectively the safety, efficacy, and complications of percutaneous nephrolithotomy in Galdakao-modified supine Valdivia position vs prone position.

**Patients and methods:** The study included all patients who underwent PCNL from June 2018 to January 2019. 41 procedures were performed in supine and 22 in prone position. All kidney punctures were performed by urologist under ultrasound control. Length of stay, OR time, complication rate, rate of transfusion and stone free rates were compared.

**Results:** Both the groups were comparable regarding the male/female ratio, stone size, and site. No significant differences were found in terms of the stone-free rate (84% and 82%), blood transfusion rate (no blood transfusion) and complication rates (1% and 1%). Significant difference was reported in mean operative time between supine and prone positions (84 and 115 minutes, respectively). Mean hospital stay was similar for both groups.

**Conclusions:** PCNL in both positions was equally successful with no significant differences in stone free rate and complications. PCNL in Galdakao-modified supine Valdivia position was superior to PCNL in the prone position regarding operative time. As rotating the patient to prone position increases some anesthesiologic risks, supine is also more comfortable for anesthesiologists.

#### GUA-04 Our treatment experience in urolithiasis

Stepanenko Gennady Anatolyevich<sup>1</sup>, Pankov Viktor Sergeevich<sup>2</sup>  
<sup>1</sup>Associate Professor, NAO MUK, Karaganda, Gogol 40, Kazakhstan;  
<sup>2</sup>Urologist, NAO MUK, Karaganda, Gogol 40, Kazakhstan

**Background:** Urolithiasis has a high medical and social significance, which is due to a fairly high incidence rate, reaching 10% in the world and its annual growth in many countries. In recent decades, indisputable progress has been made in the development of high-tech instrumental and operational methods for removing urinary stones. The introduction of Extracorporeal Shock Wave Lithotripsy, percutaneous, pro-urethral nefroureterolithotripsy and lithoextraction into clinical practice has once again significantly changed the approach to treatment of ICD. Since January 2017, Laser lithotripsy of urinary stones has been performed in City Hospital N1 of Karaganda.

**Purpose:** To assess the quantity and quality of laser lithotripsy in urolithiasis on the basis of the City Hospital №1 of the city of Karaganda.

**Materials and methods:** A retrospective analysis of inpatient cards of patients in the surgical ward of hospital No. 1 from January 2017 to

May 2019 was conducted. An Olympus 70 6.4/7.8Fr ureteroscope and a DMHSolvo holmium laser lithotripter were used.

**Results and discussion:** According to the data we received from January 2017 to May 2019, the number of operated patients was 646, of which males made up (64%), females (36%). Age groups: from 18 to 59 years old (73%), 60–69 years old (17%) and over 70 years old (10%). According to the localization of calculi in the urinary system in 92% of the patients, the stones were in the ureter and only in 8% of the patients in the bladder. By localization of calculus in the ureter: in 77% the stone was in the lower third of the ureter, in the middle third of the ureter in 14%, in the lower third of the ureter in 9% of patients. The operation was performed under spinal anesthesia. The average duration is 45 minutes. In 90% of patients, the operation ended with the installation of a ureteral stent 5–6Fr for a period of 2 weeks to 4 weeks. In the postoperative period, a Foley catheter was placed in the bladder for 1 day (until the patient was activated). All patients underwent surgery satisfactorily. In the postoperative period, such complications as exacerbation of pyelonephritis were observed in 5 patients, which was resolved by conservative measures. In 30 patients, on the background of ureteral stenting, pain syndrome was observed, which was stopped in 15 patients with non-narcotic analgesics, in 3 patients, severe pain syndrome was not stopped by narcotic analgesics, which required removal of the stent; also in 5 patients it was necessary to remove the stent in connection with severe dysuria and signs of gross hematuria. After surgery, patients were discharged from the hospital for 4–7 days.

**Conclusions:** The use of laser lithotripsy with stones in the ureter showed high efficiency and small invasiveness. The method allows to reduce the timing of the patient's stay in the hospital and reduce the number of postoperative complications.

#### GUA-05 Antegrade revascularization of the penis with retroperitoneoscopic artery fence

P.S. Kyzlasov, A.T. Mustafayev, D.I. Volodin, V.P. Sergeev, A.G. Martov  
The Department of Urology and Andrology, Biomedical University of Innovation and Continuing Education of State Research Center A.I. Burnasyan Federal Medical Biophysical Center FMBA of Russia, Moscow, Russian Federation

**Background:** Despite a variety of methods of correction of erectile dysfunction, today the only method meeting criterion of physiology is the penis revascularization. The modern methods of revascularization of a penis based both on the principle arterio-arterial and on the principle of an arteriovenous anastomosis, are united by the general source of additional perfusion – the lower epigastric artery which owing to the anatomo-physiological features is an ideal donor vessel. We present a new method of arterio-venous anastomosis with retroperitoneoscopic method of artery intake. In addition, the risk of developing one of the main complications, anastomotic thrombosis, is reduced.

**Objective:** To estimate short-term results of revascularization of a penis with a retroperitoneoscopy harvesting of an artery and imposing of an antegrade anastomosis between the lower epigastric artery and a deep dorsal vein.

**Materials and methods:** On the basis of State Research Center A.I. Burnasyan Federal Medical Biophysical Center 14 patients with vasculogen erectile dysfunction during the period from 2018 to 2019 were operated.

Patients were uniform in clinical laboratory data and on age (middle age at patients 33 years (22–35 years), according to a penile ultrasonography with doppler at 10 patients is revealed only the ED arterial component, 5 patients have a mixed form of vasculogen erectile dysfunction. Operations it was carried out by the technique developed by us. The Retroperitoneoscopy stage of allocation of the