



Review

Is visceral manipulation beneficial for patients with low back pain? A systematic review of the literature



Jacob Marten Switters^{a,*}, Stefan Podar^b, Luke Perraton^c, Zuzana Machotka^d

^a JMS-Praxis, Markgraf-Rüdiger-Straße 25/5, 1150, Vienna, Austria

^b Top-Physio, Liniengasse 53, 1060, Vienna, Austria

^c Department of Physiotherapy, Monash University, Frankston, VIC, Australia

^d La Trobe Sport and Exercise Medicine Research Centre, La Trobe University, Bundoora, Australia

ARTICLE INFO

Keywords:

Visceral manipulation
Visceral range of motion
Low back pain
Systematic review

ABSTRACT

Objectives: Visceral dysfunction (e.g., mobility or motility restriction) may be an underlying cause or contributing factor for some non-specific LBP and can be treated by osteopathic manipulative treatment (OMT). The aim of this registered systematic review (CRD42018100633) is to determine the effectiveness of visceral mobilization for non-specific LBP and explore associations between changes in range of motion of the viscera and LBP symptoms.

Data sources: In November 2018 peer-reviewed studies published in English or German where retrieved from the following databases from inception: Medline, Cochrane library, Science Direct, PEDro, OSTMED.RD and Osteo web res.

Study selection: Articles identified during searching were screened using the eligibility criteria based on title and abstract. Studies were included following independent review of full-text versions.

Data extraction: Study quality appraisal (risk of bias tool and the PEDro score) and data extraction (means and standard deviations for patient-reported outcome measures and impairments - pain, function and ultrasound measurements of change in visceral mobility) were completed/extracted by two independent authors.

Data synthesis: Cohen's effect sizes were calculated. Meta-analysis was not performed due to heterogeneity in study populations and methods. A total of four RCT's were included with a moderate to good methodological quality. Two studies reported a significant short-term (< 3 days) improvement in pain and visceral range of motion, although the clinical significance of these differences were unclear. Only one study reported significant differences in the long term (52 weeks) for pain and one for the medium term (6 weeks) for quality of life. Adverse events were poorly reported.

Introduction

Low back pain (LBP) is a highly prevalent [1] and costly [2] condition. Musculoskeletal discomfort associated with LBP is one of the key reasons that people seek osteopathic consultation [3]. The average recovery time of acute low back pain symptoms is around a month, in which time 85% of people return to work [4]. Nearly 90% of people with LBP are diagnosed with non-specific low back pain, meaning with no known cause [5,6].

The definition of a visceral dysfunction is a restriction in the mobility or motility of the viscera and connected fascial, neural, skeletal, vascular and lymphatic components [7]. A problem in the viscera can cause a diffuse and poorly localized pain which can be referred to other somatic structures [8]. For example, renal/uterine colics [9] and

gastrointestinal tract problems [10,11] can cause referred pain to the back. People who have a combination of anterior trunk pain and LBP report more pain and disability in comparison to people with only LBP [12].

The theory behind the mechanism and effect of visceral manipulation on low back is not fully understood, but may be explained in part by the neurological 'structure function' model, which is one of the five structure-function models of osteopathy [13]. This model aims to explain the link between visceral and somatic systems based on their similar innervation. This so called viscerosomatic reflex was first described in the osteopathic literature 1907 by Louise Burns [14]. A viscerosomatic reflex arises from the afferent (i.e., sensory) fibers from the viscera to the somatic structures [15]. Nociceptive input from the viscera may lead to altered sensation in the autonomic innervated

* Corresponding author. Markgraf-Rüdiger-Straße 25/5, 1150, Vienna, Austria.
E-mail address: info@jms-praxis.at (J.M. Switters).

segments of the viscera [16] and causes hyperalgesia [17–20], trophic changes in the muscle (atrophy) and subcutaneous tissue (increased thickness) in the referred pain area [17,21]. Nevertheless, it is not clear if a visceral dysfunction (e.g., a mobility restriction) causes similar symptoms as a visceral pathology/disease.

Visceral manipulation is one of many techniques used in osteopathic manipulative treatment (OMT) for LBP and other peripheral conditions [7,13]. The aim of visceral manipulation is to reduce pain in the somatic structure that shares innervation with the viscera that is manipulated [22], decrease adhesion and therefore improve mobility of the viscera [23], activation of viscerosomatic reflexes and increase of the intestinal pressure by putting pressure on the abdomen which may stimulate bowel movements [24]. In the literature visceral manipulation has been described as a treatment in the management of female infertility [23], irritable bowel syndrome [25] and chronic constipation [26,27]. Manipulation of the sigmoid colon has previously demonstrated hypoalgesia effects in the lumbar spine [22] which suggests it may be useful in the treatment of LBP. However, no systematic reviews have been conducted to determine the effectiveness of visceral mobilization for people with LBP.

Aims

The aim of this study is to systematically review the available literature to provide clinicians and patients with greater knowledge of the strength and quality of the research evidence underpinning this commonly used treatment technique. This systematic review focuses on the following two questions: 1) what is the effectiveness of visceral manipulation on pain, function and disability in individuals with low back pain and 2) is there a change in visceral range of motion after visceral manipulation?

Methods

This systematic review was registered on PROSPERO (CRD42018100633)[28] and was undertaken according to PRISMA guidelines [29]. The Eligibility criteria are reported in Table 1. Only randomized controlled trials (RCT's) of OMT interventions for LBP published in English or German language were included.

The search was performed in November 2018. The following single or combination of search terms were used: Low Back pain, Lumbar spine, Visceral Mobilization, Visceral manipulation, Osteopathy, Osteopathic, Physiotherapy and Manipulation. The following electronic medical databases were searched from inception: Medline, Cochrane library, Science Direct, PEDro, OSTMED.RD and Osteo web res. In addition, the reference lists of retrieved articles were searched individually for articles which were possibly overlooked during the primary search.

Data collection and selection

Two authors independently performed the search strategy, study

Table 1
Eligibility criteria.

Inclusions	Exclusions	Restrictions
Intervention	Spinal manipulations	Studies published in English and German
Low back pain	Test on animals	
Visceral manipulation	Visceral diseases	
Visceral mobilization	Pregnancy	No year restrictions
Osteopathy	Post pregnancy	
Physiotherapy	Constipation	Study design
	Irritable bowel syndrome	Audits, qualitative studies and mixed methods research
Study design		
Only RCT's		

selection and the data extraction as recommended by the Cochrane Collaboration [30]. Disagreements were resolved with discussion. If no agreement could be reached the third reviewer was consulted.

Study characteristics

For each included study, data relating to the study characteristics, population, intervention, comparison and outcome (PICO) were extracted. The National Health and Medical Research Council (NHMRC) Hierarchy of Evidence was used to rate the different levels of research evidence included in the review [31]. The following information was extracted: country of origin, where and how subjects were recruited, number of therapists and their education (e.g. physiotherapist, osteopath or other health professional), inclusions and exclusion criteria reported, outcome measurements used, follow up duration, characteristics of the intervention and control group, treatment duration, and report of adverse effects.

Methodological quality

To identify the methodological strengths and weaknesses of the included studies the 10-item Physiotherapy Evidence Database (PEDro) scale was used (Table 2). The PEDro scale is validated [32] and has an acceptable reliability [33].

Assessment of risk of bias

In order to assess the risk of bias it is important to determine the strength of the available evidence. For example, articles with a high risk of bias such as errors in design, insufficient sample sizes or inappropriate allocation of subjects, subjective outcome measurements, may lead to an underestimation or overestimation of the available evidence [34]. To evaluate the risk of bias of randomized trials the Cochrane risk of bias tool was used [30]. The Cochrane risk of bias tool provides thirteen questions and is answered with low, unclear or high risk of bias [30]. Two reviewers made the assessment of the bias, if there are disagreement a third reviewer is consulted.

Data analysis

The effect size was calculated using the methods of Cohen [35] i.e., calculating standardized mean differences (SMD) and was calculated post treatment and for the follow up. The effect size was defined as small (0.2–0.5), medium (> 0.5 to 0.8) or large as (> 0.8)³⁶.

Results

Four RCTs [36–39] met the inclusion criteria (Fig. 1).The search identified 621 studies; 598 studies were excluded based on title and abstract and eleven duplicates were removed. After reading the full text of twelve studies, eight articles did not meet the inclusion criteria

Table 2
PEDro scale.

1. Eligibility criteria mentioned ^a
2. Randomization
3. Allocation Concealment
4. Groups at baseline similar
5. Blinding of subjects
6. Blinding of therapist
7. Blinding of assessors
8 .Measurement of at least > 85% of the follow up outcome
9 Intention-to-treat analysis
10 Between-group statistical comparison
11 Point and variability for the outcome measurement.

^a Item 1 does not contribute to the total score.




Fig. 1. Flowchart of the search Strategy.

because of the following reasons; one was a thesis and not peer-reviewed [40], one was not an RCT [41], one was a review paper [42], and three were narrative literature reviews [43–45]. One RCT included an asymptomatic subject [22], and one focused mainly on the effectiveness of a diaphragm manipulation intervention on low back pain and did not focus on visceral manipulation [46].

Overall, 303 subjects were included across the four studies. The number of visceral manipulation treatment sessions ranged from one to 12. Adverse events after visceral manipulation were either not reported or discussed in three studies [37–39] (Table 3). All subjects were recruited from clinical populations, three out of four articles were conducted in Europe [37–39] and the follow up duration ranged from immediately post-intervention to one year (Table 4).

The statistically significant differences between the intervention and control groups are summarized in Table 5. In two articles, treatment was performed by an osteopath and reported significant differences in pain reduction and increases in visceral range of motion for the visceral intervention groups compared to sham treatment groups [38,39]. One article found that standard care and standard care plus visceral manipulation both led to significant improvements in pain and function [37]. However, no significant difference was found between the two groups. There was a statistically significant improvement in quality of life for the visceral intervention group in the following subgroups; energy, emotional role limitations, mental health and total mental health score [38]. Panagopoulos et al. (2015) found no significant differences in disability and function when multi-modal physiotherapy techniques

were compared to visceral manipulation at 2, 6 and 52 weeks [36], with the exception that there was a significant difference found in favor of visceral manipulation for pain at 52 weeks post treatment, despite no differences found at 2 and 6 weeks post intervention.

Methodological quality

PEDro scores for the four RCT articles ranged from 5/10³⁸ to highest 9/10³⁷ (Table 6). As the therapist delivered the intervention, blinding of the therapist (criterion 6) was not possible for this study design.

Risk of bias

The evaluation of the risk of bias (Table 7) revealed more than 50% of the criteria were considered as low risk (33/52) and 25% (13/52) were considered as a high risk across all four studies. All included studies scored a high risk on blinding of the therapist and personnel. Blinding of the therapist is difficult when a manual intervention is performed. One article demonstrated high risk of bias by not explicitly mentioning if the subjects or outcome assessor were blinded, and did not describe the dropouts rates [37]. Moreover, the description of the intervention lacked sufficient detail to allow for reproducibility, with multiple possible combinations of the intervention described. The number of sessions each participant attended was similarly not reported and therefore individual patient exposure and compliance considered as high risk of bias.

Table 3
Intervention characteristics.

First Author	Treatment Duration	Adverse events		Intervention group	Control group
Panagopoulos 2015	12 Treatments for 6 weeks, one to two session a week	No	Subjects	32	32
			Age range	18–80	18–80
			Age mean	N/M	N/M
			Sex	13 (M)	12 (M)
Tamer 2017	10 Treatments for 5 weeks at two sessions a week	Not mentioned	Sex	19 (F)	20 (F)
			Subjects	20	19
			Age median	42	36
			Age IQR	34.2–51.5	29–47
Tozzi 2011	1 Treatment	Not mentioned	Sex	8 (M)	10 (M)
			Subjects	12 (F)	9 (F)
			Age range	30 (LBP)	30
			Age mean	21–58	28–52
Tozzi 2012	1 Treatment	Not mentioned	Age mean	39.1	39
			Sex	18 (M)	22 (M)
			Subjects	12 (F)	8 (F)
			Age Range	20–59	23–55
			Age mean	39.8	37.6
			Sex	54 (M)	20 (M)
				55(F)	11 (F)

M = Male F = Female N/M = Not mentioned, IQR = Interquartile range.

Two studies [38,39] reported that the subjects were randomized; however, the authors did not describe the randomization process, e.g. computer-generated random sequence. The two studies that used ultrasound as a primary outcome measure tool did not demonstrate a valid and reliable method of ultrasound assessment [38,39]. The compliance criteria is deemed irrelevant for single session interventions as was the case for two studies [38,39].

Effect size

All four articles had detailed information to calculate effect size (Table 8). Two articles showed large effect sizes for pain (1.1³⁹ and 1.2⁴⁰), large/moderate effect sizes for visceral range of motion - kidney (2.6³⁹ and 0.8⁴⁰) and bladder (2.3)³⁹. Another article showed a moderate effect size for quality of life (0.7), pain (0.5) and function (0.4)³⁸. Panagopoulos et al. (2015) reported small effect sizes for pain, disability and patient-reported function at six weeks post-intervention. A large effect size was found two weeks post intervention for disability (0.9) and a moderate effect size was maintained one year post intervention for pain (0.7)³⁷.

Pain

All four articles used reliable and validated outcome measurement for pain including the visual analogue scale (VAS) [37], numeric pain rating scale (NPRS) [36], and the short form of the McGill Pain Questionnaire (SF-MPQ) [38,39]. In the short term (< 3 days) visceral manipulation showed in two studies a significant improvement for pain in comparison with the control group [38,39]. One article found no significant difference at 2 and 6 weeks post intervention in comparison with the control group [36]. However, at 52 weeks post intervention there was a significant difference between groups in reduction of pain [36]. The other article that compared standard care with standard care plus visceral manipulation showed that both interventions were effective (P < 0.001) in reducing pain at 6 weeks from the start of the intervention [37]. No significant differences were found between the two groups (p = 0.154).

Disability and Function

Four reliable and validated questionnaires were used in two articles to measure disability and function [36,37]; Patient specific function

scale (PSFS) [36], Roland Morris disability questionnaire (RMDQ) [36], Quality of life score (SF-36)³⁸ and Oswestry function scale (OFS) [37]. Panagopoulos et al. found no significant differences for function and disability at the following outcome measurements: 2 weeks (P = 0.158), 6 weeks (P = 0.332) and 52 weeks (p = 0.882) after the intervention [36].

For the subgroups of quality of life score there was a difference between groups in favor of the physiotherapy with added visceral manipulation [37]. There was a significant difference (P < 0.05) in all subgroups for 1) physical function (p = 0.028), 2) energy (p = 0.034) and 3) the total physical score (p = 0.025)³⁸. Tamer et al. found that both groups had significant improvements in Oswestry function scores post-intervention for physiotherapy (P < 0.001) and physiotherapy with visceral manipulation added (P < 0.001)³⁸. No significant difference was found between the two interventions (P = 0.243).

Visceral range of motion

Two articles examined the range of motion of the viscera with ultrasound and the effectiveness on somatic complaints [38,39]. The following viscera were examined: kidney [38,39] and bladder [38].

To compare pre-intervention to post-intervention range of motion of the kidney the Kidney Renal-Diaphragmatic distance (RD) was measured. In both articles the control group had no significant pre-post differences in RD distance. One article reported a statistically significant difference in RD distance from pre-intervention (10.33 ± 4.70) to post-intervention (21.60 ± 7.06), units not reported [38]. The other article reported a statistical significant difference (p < 0.0001) between the pre-intervention distance of 5.79 mm (St. Dev. 8.55) against post-intervention distance of 11.34 mm (Std. Dev. 8.96)⁴⁰. The kidney mobility score (KMS) was also significantly different between asymptomatic subjects KMS (1.92 mm, Std. Dev. 1.14) and LBP patients (KMS 1.52 mm, Std. Dev. 0.79)⁴⁰. Hence, the patients with nonspecific low back pain included in this study had significantly reduced kidney mobility (P < 0.05) in comparison with asymptomatic subjects [39].

The range of motion of the bladder was the measurement of the distance of the neck of the bladder and the anterior vesical wall on the perpendicular line [38]. For the intervention group the pre-intervention distance was 12.70 mm (St. Dev. 4.18) compared to a post-intervention distance of 22.73 mm (St. Dev. 3.73) which was a significant difference (p < 0.0001)³⁹. None of the articles examined the reliability or

Table 4
Study and participants characteristic.

First Author	Level of research	Country of origin	Recruitment	Therapist Number	Inclusions and exclusions criteria reported	Outcome measurement	Follow up
Panagopoulos 2015	RCT (II)	Australia	Private physiotherapy clinic	Physiotherapist 2	Yes	Pain (NPRS) Disability (RMDQ) Function (PSFP) Pain (VAS) Disability (SF-36) Function (Oswestry function scale) Pain (SF-MPQ) Ultrasound on Kidney and Bladder Pain (SF-MPQ) Ultrasound on Kidney	2,6 and 52 weeks 6 weeks after the beginning of the intervention Pain: 3 days post-intervention Ultrasound: post-intervention Pain: 3 days post-intervention Ultrasound: post-intervention
Tamer 2017	RCT (II)	Turkey	University physical therapy department	Physiotherapist Not Mentioned	Yes		
Tozzi 2011	RCT (II)	Italy	Clinical Centre	Osteopath 1	Yes		
Tozzi 2012	RCT (II)	Italy	Clinical Centre	Osteopath 1	Yes		

Table 5
P Value between intervention and control group.

First author	Intervention	Control	P value*
Panagopoulos 2015	Physiotherapy plus visceral manipulation	Physiotherapy plus placebo visceral manipulation	Pain (NPRS) ● 2/52: P = 0.362 ● 6/52: P = 0.858 ● 52/52: P = 0.015 Disability (RMDQ) ● 2/52: P = 0.479 ● 6/52: P = 0.418 ● 52/52: P = 0.942 Function (PSFPS) ● 2/52: P = 0.158 ● 6/52: P = 0.332 ● 52/52: P = 0.882
Tamer 2017	Osteopathy plus physiotherapy (vOMT)	Physiotherapy (OMT)	Pain (VAS): ● 6/52: P = 0.154 Function (OFS): ● 6/52: P = 0.243 Quality of life (SF-36) ● 6/52: P = < 0.028
Tozzi 2011	Osteopathy	Sham	Kidneys ROM: ● P < 0.0001* Bladder ROM: ● P < 0.0001* Pain (SF-MPQ): ● 3/7 (P < 0.0001* Kidney ROM: ● P < 0.0001* Pain (SF MPQ) ● 3/7: P < 0.0001*
Tozzi 2012	Osteopathy	Sham	

*P value is from Post Intervention Both = Intervention and control.
Osteopathy: include MET, Visceral techniques, Still techniques, Fascial techniques, HVLA, lymphatic pump techniques.
Physiotherapy: include Manual therapy, Exercise therapy, Stretching Lumbar re-training, Functional exercise prescription, Massage.

Table 6
PEDro score.

First Author	1 ^a	2	3	4	5	6	7	8	9	10	11	Score
Panagopoulos et al., 2014	1	1	1	1	1	0	1	1	1	1	1	9/10
Tamer et al., 2016	1	1	1	1	0	0	0	0	0	1	1	5/10
Tozzi et al., 2011	1	1	0	0	1	0	1	1	0	1	1	6/10
Tozzi et al., 2012	0	1	0	1	1	0	1	0	0	1	1	6/10

^a Item 1 does not contribute to the total score.

validity of ultrasound variables [38,39].

Two articles measured changes in visceral range of motion and in pain [38,39]. Both reported significant differences between pre- and post-measurement range of motion and a significant difference in the short term (< 3 days) for pain.

Discussion

The aim of this systematic review was to investigate the effectiveness of visceral manipulation on low back pain and to explore associations between changes in range of motion of the viscera and LBP symptoms. The main finding was that visceral manipulation may be beneficial in the short term for low back pain symptoms and may be associated with increased range of motion of the viscera. However, due to insufficient follow up periods it is not clear how long this effect lasted. Therefore, further research to explore its effectiveness is needed.

Caution should be taken when generalizing the findings of this systematic review (303 participants) to the wider population of people with LBP. Of the 303 subjects, 191 received visceral manipulation as an isolated intervention or as part of a multi-modal intervention such as manual therapy, muscle energy techniques (MET), Still Technique or

Table 7
Risk of Bias of the included studies assessed by the Cochran Risk of bias.[30].

Criteria	First Author			
	Panagopoulos 2015	Tamer 2017	Tozzi 2011	Tozzi 2012
Randomized	Low	Low	Unclear	Unclear
Concealment	Low	Low	High	High
Blinding of the Patient	Low	High	Low	Low
Blinding of the care provider	High	High	High	High
Outcome assessor blinding	Low	High	Low	Low
Drop-outs reported and acceptable	Low	High	Low	Low
Were the randomized participant analyzed	Low	Low	Low	Low
Free of selective outcome	Low	Low	High	High
Groups similar at baseline	Low	Low	Low	Low
Co-intervention avoided or similar	Low	High	Low	Low
Compliance acceptable	Low	High	Unclear	Unclear
Similar Timing of outcome measurement	Low	Low	Low	Low
Other Bias	Low	Low	Unclear	Unclear

exercise. In terms of generalizing the findings of this review, it is important to consider that the osteopathic intervention only had one treatment [38,39], whereas the physiotherapy intervention had ten to twelve treatments [36,37].

A major concern of this systematic review is that more than half of the included articles did not report or measure adverse events related to visceral manipulation [37–39]. Therefore, it is hard to conclude if visceral manipulation is safe because there is no sufficient data available. Less than a quarter of the subjects (64/303) participating in the included studies were asked or checked for adverse effects after the visceral manipulation and no adverse effect was reported [36]. It has been estimated that the incidence rate for adverse events is 2%–6% after OMT intervention [47,48]. The most common complaints were pain and discomfort after the intervention and no serious adverse effect was reported [48]. This is relatively low in comparison with the minor and moderate side effects after manual therapy, e.g. mobilization and manipulation, which has been reported to be 16–41% [49,50]. The incidence for a serious side effect after manual therapy is considered low [50,51].

Serious adverse effects after visceral manipulation were only reported in three studies. These included hepatic hematoma, perforation of the sigmoid colon, ureteral stent displacement, embolization of the kidney, ruptured uterus [52,53] and bladder rupture [54]. The person who performed the treatment ranged from traditional healer, rolfing

therapist, family member or was not specifically mentioned [52–54]. The risk of serious adverse events should therefore be considered when making clinical decisions about the use of visceral manipulation.

Ultrasound was used in two articles as an outcome measurement to quantify whether visceral manipulation results in changes in the anatomical positions of the viscera after being treated [38,39]. Both articles did not mention if there were any reliable or validated studies to measure the visceral range of motion. To the knowledge of the authors there have been no recent reliability studies conducted for the renal length measured with ultrasound. Based on one older study, renal length has good intra-rater reliability [55]. Whilst these findings of these two articles are interesting, it is more clinically important to know whether these changes in visceral mobility are related to the viscera function and to the progression of LBP symptoms and function. Therefore, research is first needed to determine reliability of the diagnostic measurement of the viscera organ range of motion.

The range of motion of the following viscera, liver, spleen and kidneys is estimated from 0 mm to more than 27mm[56, 57]. However, the mobility of the kidney is highly dependent on both the intensity and type of breathing. For instance the cranio-caudal displacement of the kidney is threefold during forced aspiration compared to quiet breathing [58]. Standardizing breathing is a known problem described in a review on oncologic radiotherapy [59]. While the deepness of breathing compromises the reliability of kidney motion measurements,

Table 8
Effect size.

First Author	Outcome	Week	Intervention (SD)	Control (SD)	Effect size
Panagopoulos 2015	Pain (NPRS)	2/52	3.06 (2.08)	3.74 (2.25)	Small (0.3)
		6/52	2.31 (1.99)	2.33 (2.22)	Small (< 0.01)
		52/52	1.52 (1.65)	3.21 (2.27)	Medium (0.7)
	Disability (RMDQ)	2/52	5.78 (5.40)	6.26 (5.35)	Large (0.9)
		6/52	3.00 (2.96)	3.10 (3.98)	Small (0.02)
		52/52	2.06 (3.56)	3.50 (3.61)	Small (0.4)
	Function (PSFS)	2/52	6.10 (2.13)	6.15 (1.95)	Small (0.02)
		6/52	7.70 (1.81)	7.51 (1.86)	Small (0.1)
		52/52	8.43 (1.76)	7.55 (1.82)	Small (< 0.5)
Tamer 2017	Pain (VAS)	6/52	5.65 (1.64)	4.77 (1.91)	Small (< 0.5)
		6/52	42.95 (23.94)	22 (28.54)	Medium (0.7)
	Oswestry function scale	6/52	28.40 (18.19)	22.60 (14.09)	Small (0.4)
Tozzi 2011	Mc Gill (SF-MPQ)	3/7	15.517 (9.839)	25.05 (8.867)	Large (1.1)
	US Bladder	Post	22.73 (3.73)	12.90 (4.23)	Large (2.3)
	US Kidney	Post	21.60 (7.06)	10.10 (4.49)	Large (2.6)
Tozzi 2012	Mc Gill (SF-MPQ)	3/7	9.30 (5.55)	15.41 (5.24)	Large (1.2)
	US Kidney	Post	11.34 (8.96)	4.90 (8.15)	Medium (< 0.8)

Post = Post intervention 52 = weeks 7 = days.

the breathing pattern seems to be important as well. Due to increased diaphragmatic movement, abdominal breathers displayed more kidney motion than subject who used a thoracic breathing strategy [60]. Therefore, a standardization of the breathing should be reported in the article to receive consistent, dependable, repeatable, meaningful and trustworthy results.

The evidence for the effectiveness of visceral manipulation for low back pain is limited to a small number of randomized trials, of variable methodological quality with small sample sizes and limited to mainly short term (< 3 days) effects. Future randomized trials should be conducted with sufficient samples sizes as determined by power analysis and should focus on accurately reporting the treatment parameters (e.g. time, duration, treatment positions). Contra-indications to treatment, risks of adverse events and dropouts should also be reported clearly. Treatment reported in future studies should be standardized when possible and the long term (> 6 Months) effectiveness of visceral manipulation should be investigated.

Conclusion

In this systematic review it was shown that visceral manipulation for low back pain demonstrates beneficial effects for reducing pain and may improve visceral range of motion. However, these findings are based on a small number of studies with a high risk of bias. The lack of reporting of adverse effects in the articles is a major concern in relation to recommendations made for the safe use of visceral manipulation in the clinic.

Visceral manipulation is used to treat low back pain by clinicians; however, this systematic review highlights limitations to the reporting, quantity and quality of evidence underpinning its use. To justify ongoing use of this type of manual therapy it is of great importance that clinicians use reliable and validated outcome measurements to understand and reflect on the effectiveness of their treatment.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

Declarations of interest

All authors declare they have no competing interests.

References

- [1] Hoy D, Brooks P, Blyth F, Buchbinder R. The Epidemiology of low back pain. Best practice & research. Clin Rheumatol 2010;24(6):769–81. <https://doi.org/10.1016/j.berh.2010.10.002>.
- [2] Dagenais S, Caro J, Haldeman S. A systematic review of low back pain cost of illness studies in the United States and internationally. Spine J Off J North Am Spine Soc 2008;8(1):8–20. <https://doi.org/10.1016/j.spinee.2007.10.005>.
- [3] Morin C, Aubin A. Primary reasons for osteopathic consultation: a prospective survey in Quebec. PLoS One 2014;9(9):e106259 <https://doi.org/10.1371/journal.pone.0121180>.
- [4] Pengel LHM, Herbert RD, Maher CG, Refshauge KM. Acute low back pain: systematic review of its prognosis. Br Med J 2003;327(7410):323. <https://doi.org/10.1136/bmj.327.7410.323>.
- [5] Koes BW, van Tulder MW, Thomas S. Diagnosis and treatment of low back pain. Br Med J 2006;332(7555):1430–4. <https://doi.org/10.1136/bmj.332.7555.1430>.
- [6] Krismar M, van Tulder M. Strategies for prevention and management of musculoskeletal conditions. Low back pain (non-specific). Best practice & research. Clin Rheumatol 2007;21(1):77–91. <https://doi.org/10.1016/j.berh.2006.08.004>.
- [7] Educational Council on osteopathic principles (E.C.O.P.). Glossary of Osteopathic Terminology Usage Guide; 2011.
- [8] Cervero F, Laird JMA. Visceral pain. The Lancet 1999;353(9170):2145–8. [https://doi.org/10.1016/S0140-6736\(99\)01306-9](https://doi.org/10.1016/S0140-6736(99)01306-9).
- [9] Giamberardino MA, Affaitati G, Costantini R. Chapter 24 Referred pain from internal organs. Pain. Elsevier; 2006. p. 343–61.
- [10] Smith MD, Russell A, Hodges PW. How common is back pain in women with gastrointestinal problems? Clin J Pain 2008;24(3):199–203. <https://doi.org/10.1097/AJP.0b013e31815d3601>.
- [11] Smith MD, Russell A, Hodges PW. Do incontinence, breathing difficulties, and gastrointestinal symptoms increase the risk of future back pain? J Pain Off J Am Pain Soc 2009;10(8):876–86. <https://doi.org/10.1016/j.jpain.2009.03.003>.
- [12] Panagopoulos J, Hancock MJ, Kongsted A, Hush J, Kent P. Does anterior trunk pain predict a different course of recovery in chronic low back pain? Pain 2014;155(5):977–82. <https://doi.org/10.1016/j.pain.2014.01.023>.
- [13] World Health Organization. Benchmarks for training in traditional/complementary and alternative medicine.: benchmarks for Training in Osteopathy. Switzerland. 2010 Available from: <http://www.who.int/medicines/areas/traditional/BenchmarksforTraininginOsteopathy.pdf>, Accessed date: 26 June 2018.
- [14] Burns L. Viscero-somatic and somato-visceral spinal reflexes. J Am Osteopath Assoc 1907;100(4):249–58. 2000.
- [15] Beal MC. Viscerosomatic reflexes: a review. J Am Osteopath Assoc 1985;85(12):786–801.
- [16] van Buskirk RL. Nociceptive reflexes and the somatic dysfunction: a model. J Am Osteopath Assoc 1990;90(9):797–809. 792–4.
- [17] Giamberardino MA, Affaitati G, Lerza R, Lapenna D, Costantini R, Vecchiet L. Relationship between pain symptoms and referred sensory and trophic changes in patients with gallbladder pathology. Pain 2005;114(1–2):239–49. <https://doi.org/10.1016/j.pain.2004.12.024>.
- [18] Stawowy M, Rössel P, Bluhme C, Funch-Jensen P, Arendt-Nielsen L, Drewes AM. Somatosensory changes in the referred pain area following acute inflammation of the appendix. Eur J Gastroenterol Hepatol 2002;14(10):1079–84.
- [19] Stawowy M, Drewes AM, Arendt-Nielsen L, Funch-Jensen P. Somatosensory changes in the referred pain area before and after cholecystectomy in patients with uncomplicated gallstone disease. Scand J Gastroenterol 2006;41(7):833–7. <https://doi.org/10.1080/0036520500463332>.
- [20] Vecchiet L, Giamberardino MA, Dragani L, Albe-Fessard D. Pain from renal/ureteral calculosis: evaluation of sensory thresholds in the lumbar area. Pain 1989;36(3):289–95. [https://doi.org/10.1016/0304-3959\(89\)90087-0](https://doi.org/10.1016/0304-3959(89)90087-0).
- [21] Wesselmann U, Lai J. Mechanisms of referred visceral pain: uterine inflammation in the adult virgin rat results in neurogenic plasma extravasation in the skin. Pain 1997;73(3):309–17. [https://doi.org/10.1016/S0304-3959\(97\)00112-7](https://doi.org/10.1016/S0304-3959(97)00112-7).
- [22] McSweeney TP, Thomson OP, Johnston R. The immediate effects of sigmoid colon manipulation on pressure pain thresholds in the lumbar spine. J Bodyw Mov Ther 2012;16(4):416–23. <https://doi.org/10.1016/j.jbmt.2012.02.004>.
- [23] Rice AD, Patterson K, Wakefield LB, Reed ED, Breder KP, Wurn BF, et al. Ten-year retrospective study on the efficacy of a manual physical therapy to treat female infertility. Alternative therapies in health and medicine. 2015.
- [24] Liu Z, Sakakibara R, Odaka T, Uchiyama T, Yamamoto T, Ito T, et al. Mechanism of abdominal massage for difficult defecation in a patient with myelopathy (HAM/TSP). J Neurol 2005;252(10):1280–2. <https://doi.org/10.1007/s00415-005-0825-9>.
- [25] Müller A, Franke H, Resch K-L, Fryer G. Effectiveness of osteopathic manipulative therapy for managing symptoms of irritable bowel syndrome: a systematic review. J Am Osteopath Assoc 2014;114(6):470–9. <https://doi.org/10.7556/jaoa.2014.098>.
- [26] Ernst E. Abdominal massage therapy for chronic constipation: a systematic review of controlled clinical trials. Forschende Komplementärmed 1999;6(3):149–51. <https://doi.org/10.1159/000021240>.
- [27] Sinclair M. The use of abdominal massage to treat chronic constipation. J Bodyw Mov Ther 2011;15(4):436–45. <https://doi.org/10.1016/j.jbmt.2010.07.007>.
- [28] PROSPERO Available from: <https://www.crd.york.ac.uk/prospero/>, Accessed date: 17 August 2018.
- [29] Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Syst Rev 2015;4:1. <https://doi.org/10.1186/2046-4053-4-1>.
- [30] Furlan AD, Malmivaara A, Chou R, Maher CG, Deyo RA, Schoene M, et al. Updated method guideline for systematic reviews in the Cochrane back and neck group. Spine 2015;40(21):1660–73. <https://doi.org/10.1097/BRS.0000000000001061>.
- [31] Merlin T, Weston A, Tooher R. Extending an evidence hierarchy to include topics other than treatment: revising the Australian 'levels of evidence. BMC Med Res Methodol 2009;9:34. <https://doi.org/10.1186/1471-2288-9-34>.
- [32] Morton NA de. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. Aust J Physiother 2009;55(2):129–33.
- [33] Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. Phys Ther 2003;83(8):713–21.
- [34] Wood L, Egger M, Gluud LL, Schulz KF, Juni P, Altman DG, et al. Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study. Br Med J 2008;336(7644):601–5. <https://doi.org/10.1136/bmj.39465.451748.AD>.
- [35] Cohen J. A power primer. Psychol Bull 1992;112(1):155–9. <https://doi.org/10.1037/0033-2909.112.1.155>.
- [36] Panagopoulos J, Hancock MJ, Ferreira P, Hush J, Petocz P. Does the addition of visceral manipulation alter outcomes for patients with low back pain? A randomized placebo controlled trial. Eur J Pain 2015;19(7):899–907. <https://doi.org/10.1002/ejp.614>.
- [37] Tamer S, Öz M, Ülger Ö. The effect of visceral osteopathic manual therapy applications on pain, quality of life and function in patients with chronic nonspecific low back pain. J Back Musculoskelet Rehabil 2017;30(3):419–25. <https://doi.org/10.3233/BMR-150424>.
- [38] Tozzi P, Bongiorno D, Vitturini C. Fascial release effects on patients with non-specific cervical or lumbar pain. J Bodyw Mov Ther 2011;15(4):405–16. <https://doi.org/10.1016/j.jbmt.2010.11.003>.
- [39] Tozzi P, Bongiorno D, Vitturini C. Low back pain and kidney mobility: local osteopathic fascial manipulation decreases pain perception and improves renal

- mobility. *J Bodyw Mov Ther* 2012;16(3):381–91. <https://doi.org/10.1016/j.jbmt.2012.02.001>.
- [40] van der Linden S. Osteopathic treatment of lower back pain through mobilisation of the mesenterium. Research as to the added value of visceral manipulation in patients with non specific low back pain.: a randomized controlled clinical investigation Available from: http://www.osteopathicresearch.com/index.php?option=com_jresearch&view=publication&task=show&id=16182&lang=en, Accessed date: 12 November 2018.
- [41] Wilson D, Gorham J, Lamb T, Lui S, Todd D. Osteopathic manipulation improves functional status in patients with non-specific chronic back pain in a rural outpatient setting. *J Am Osteopath Assoc* 2018;28(2):29–33.
- [42] Bolton PS, Budgell B. Visceral responses to spinal manipulation. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 2012;22(5):777–84. <https://doi.org/10.1016/j.jelekin.2012.02.016>.
- [43] King HH. Visceral manipulation may affect pain perception. *J Am Osteopath Assoc* 2013;113(1):97–8.
- [44] King HH. Addition of osteopathic visceral manipulation to OMT for low back pain decreases pain and increases quality of life. *J Am Osteopath Assoc* 2017;117(5):333–4. <https://doi.org/10.7556/jaoa.2017.062>.
- [45] Seffinger M. Osteopathic fascial manipulation reduces low back pain and increases kidney mobility. *J Am Osteopath Assoc* 2013;113(1):102.
- [46] Martí-Salvador M, Hidalgo-Moreno L, Doménech-Fernández J, Lisón JF, Arguisuelas MD. Osteopathic manipulative treatment including specific diaphragm techniques improves pain and disability in chronic nonspecific low back pain: a randomized trial. *Arch Phys Med Rehabil* 2018;99(9):1720–9. <https://doi.org/10.1016/j.apmr.2018.04.022>.
- [47] Licciardone JC, Minotti DE, Gatchel RJ, Kearns CM, Singh KP. Osteopathic manual treatment and ultrasound therapy for chronic low back pain: a randomized controlled trial. *Ann Fam Med* 2013;11(2):122–9. <https://doi.org/10.1370/afm.1468>.
- [48] Degenhardt BF, Johnson JC, Brooks WJ, Norman L. Characterizing adverse events reported immediately after osteopathic manipulative treatment. *J Am Osteopath Assoc* 2018;118(3):141–9. <https://doi.org/10.7556/jaoa.2018.033>.
- [49] Carlesso LC, Gross AR, Santaguída PL, Burnie S, Voth S, Sadi J. Adverse events associated with the use of cervical manipulation and mobilization for the treatment of neck pain in adults: a systematic review. *Man Ther* 2010;15(5):434–44. <https://doi.org/10.1016/j.math.2010.02.006>.
- [50] Carnes D, Mars TS, Mullinger B, Froud R, Underwood M. Adverse events and manual therapy: a systematic review. *Man Ther* 2010;15(4):355–63. <https://doi.org/10.1016/j.math.2009.12.006>.
- [51] Nielsen SM, Tarp S, Christensen R, Bliddal H, Klokke L, Henriksen M. The risk associated with spinal manipulation: an overview of reviews. *Syst Rev* 2017;6(1):64. <https://doi.org/10.1186/s13643-017-0458-y>.
- [52] Ernst E. The safety of massage therapy. *Rheumatology* 2003;42(9):1101–6. <https://doi.org/10.1093/rheumatology/keg306>.
- [53] Grant KE. Massage safety: injuries reported in Medline relating to the practice of therapeutic massage—1965–2003. *J Bodyw Mov Ther* 2003;7(4):207–12. [https://doi.org/10.1016/S1360-8592\(03\)00043-3](https://doi.org/10.1016/S1360-8592(03)00043-3).
- [54] Posadzki P, Ernst E. The safety of massage therapy: an update of a systematic review. *Focus Altern Complement Ther* 2013;18(1):27–32. <https://doi.org/10.1111/ftc.12007>.
- [55] Ablett MJ, Coulthard A, Lee RE, Richardson DL, Bellas T, Owen JP, et al. How reliable are ultrasound measurements of renal length in adults? *Br J Radiol* 1995;68(814):1087–9. <https://doi.org/10.1259/0007-1285-68-814-1087>.
- [56] Brandner ED, Wu A, Chen H, Heron D, Kalnicki S, Komanduri K, et al. Abdominal organ motion measured using 4D CT. *Int J Radiat Oncol Biol Phys* 2006;65(2):554–60. <https://doi.org/10.1016/j.ijrobp.2005.12.042>.
- [57] Langen KM, Jones DT. Organ motion and its management. *Int J Radiat Oncol Biol Phys* 2001;50(1):265–78. [https://doi.org/10.1016/S0360-3016\(01\)01453-5](https://doi.org/10.1016/S0360-3016(01)01453-5).
- [58] Alt S, Alt B, Loudovici-Krug D, Knie R, Smolenski U. Respiratorische verschieblichkeit der Nieren. *Man Med* 2015;53(5):338–44. <https://doi.org/10.1007/s00337-015-0028-0>.
- [59] Pham D, Kron T, Foroudi F, Schneider M, Siva S. A review of kidney motion under free, deep and forced-shallow breathing conditions: implications for stereotactic ablative body radiotherapy treatment. *Technol Cancer Res Treat* 2014;13(4):315–23. <https://doi.org/10.7785/tcrt.2012.500387>.
- [60] Davies SC, Hill AL, Holmes RB, Halliwell M, Jackson PC. Ultrasound quantitation of respiratory organ motion in the upper abdomen. *Br J Radiol* 1994;67(803):1096–102. <https://doi.org/10.1259/0007-1285-67-803-1096>.