



ELSEVIER

www.obstetanesthesia.com

ORIGINAL ARTICLE

Design errors in vital sign charts used in consultant-led maternity units in the United Kingdom

R. Isaacs,^{a,b} G. Smith,^a L. Gale-Andrews,^c M. Wee,^{a,d} E. van Teijlingen,^e D. Bick,^{e,f} V. Hundley^e, on behalf of the Modified Obstetric Early Warning Systems (MObs) Research Group

^aCentre of Postgraduate Medical Research & Education (CoPMRE), Faculty of Health and Social Sciences, Bournemouth University, Bournemouth, UK

^bDepartment of Anaesthesia, University Hospital Southampton NHS Foundation Trust, Southampton, UK

^cBournemouth University Clinical Research Unit (BUCRU), Faculty of Health and Social Sciences, Royal London House, Christchurch Road, Bournemouth, UK

^dDepartment of Anaesthesia, Poole Hospital NHS Foundation Trust, Poole, UK

^eCentre for Midwifery, Maternal & Perinatal Health, Bournemouth University, Bournemouth, UK

^fDepartment of Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College, London, UK

ABSTRACT

Background: Paper-based charts remain the principal means of documenting the vital signs of hospitalised pregnant and postnatal women. However, poor chart design may contribute to both incorrect charting of data and clinical responses. We decided to identify design faults that might have an adverse clinical impact.

Methods: One hundred and twenty obstetric early warning charts and escalation protocols from consultant-led maternity units in the United Kingdom and the Channel Islands were analysed using an objective and systematic approach. We identified design errors that might impede their successful use (e.g. generate confusion regarding vital sign documentation, hamper the recognition of maternal deterioration, cause a failure of the early warning system or of any clinical response).

Results: We found 30% (n=36/120) of charts contained at least one design error with the potential to confuse staff, render the charts difficult to use or compromise patient safety. Amongst the most common areas were inadequate patient identification, poor use of colour, illogical weighting, poor alignment and labelling of axes, and the opportunity for staff to 'game' the escalation.

Conclusions: We recommend the urgent development of an evidence-based, standardised obstetric observation chart, which integrates 'human factors' and user experience. It should have a clear layout and style, appropriate colour scheme, correct language and labelling, and the ability for vital signs to be documented accurately and quickly. It should incorporate a suitable early warning score to guide clinical management.

Crown Copyright © 2019 Published by Elsevier Ltd. All rights reserved.

Keywords: Women's health; Patient safety; Standards of care; Patient record; Design; Error

Introduction

Maternal clinical deterioration can usually be identified by monitoring and charting key vital signs (e.g. respiratory rate, oxygen saturation (S_pO₂), blood pressure (BP)).¹ Paper-based charts remain the principal means of documenting vital signs, despite a drive to introduce

electronic automated systems.² These charts are intended to track maternal physiology temporally, highlight vital sign deviations and 'signpost' the need for a prompt clinical response. They employ a range of approaches to improve the recognition of abnormal physiology.³ Some charts use colour-coded shading, often yellow and red, to highlight 'mildly' and 'markedly' abnormal parameter ranges.⁴ Others incorporate an aggregate early warning system (EWS) score that allocates points to each vital sign measurement based on the extent of their derangement from predefined 'normal' ranges.⁵

Accepted January 2019

Correspondence to: Dr R. Isaacs, Department of Anaesthesia, University Hospital Southampton NHS Foundation Trust, Tremona Road, Southampton SO16 6YD, UK.

E-mail address: richard.isaacs@uhs.nhs.uk

However, observations are not always charted correctly, and appropriate and timely clinical responses to abnormal physiology do not always occur.^{1,4,6} Poor chart design may contribute to such failures.^{7–9} In 2007, the United Kingdom (UK) Confidential Enquiry into Maternal and Child Health (CEMACH) proposed that a single maternal observation chart should be used for all obstetric admissions in all settings.⁴ At the time, the ‘CEMACH’ chart was no more than an example of good practice, being neither evidence-based nor validated. It has not been universally used in UK midwifery or obstetric units.^{10,11} Recently, researchers in Australia have used an evidence-based approach, and an emphasis on ‘human factors’ and user experience, to design improved, ‘standardised’ vital signs charts for general patient use.^{8,9,12–15}

During an analysis of paper vital sign charts in consultant-led UK maternity units,³ design errors were commonly noted. Therefore, we decided to investigate this by focusing on common design faults that might hamper the charts’ safe and successful use.

Methods

In July 2014, as part of a survey of obstetric EWSs used in consultant-led maternity units in the UK and the Channel Islands,³ we wrote to 194 lead consultant anaesthetists registered with the Obstetric Anaesthetists’ Association (OAA). We requested a copy of the obstetric early warning chart and associated escalation protocol used in their unit. Where necessary, non-responders were contacted again between July 2014 and October 2014 via telephone (once), follow-up letter (three times) and email (three times). We used existing recommendations for chart design^{14,15} to identify faults that might lead to undesirable clinical impacts. Whilst acknowledging the importance of avoiding over-medicalisation of the management of pregnant and postnatal women, this article will often use the term ‘patient’ to refer to women.

Two members of the research team (GS and RI) analysed each chart independently. Charts were scrutinised for design errors that might impede their successful use by maternity staff (e.g. generate confusion regarding vital sign documentation, hamper recognition of maternal deterioration, cause a failure of the EWS or of any clinical response). All errors were recorded on a single spreadsheet. The following design aspects of the charts were evaluated:

- A. Identification of the patient
- B. Chart style and layout
 - B1. Use of colour
 - B2. Graphical versus tabular layouts/displays
 - B3. Integration of EWSs in the observation chart

- B4. Alignment of axes and their labelling; row and column shift
- B5. Illogical ranges
- B6. Opportunities for ‘gaming’ the escalation of care

In line with guidance from the National Health Service (NHS) Health Research Authority, this study did not require ethical review by an NHS Research Ethics Committee or management permission through the NHS Research and Development office. Approval for the initial survey was obtained from the OAA Audit Subcommittee.

Results

We received 120 usable charts from the 194 consultants approached. We identified many examples of good design, however 30% contained at least one design error with the potential to confuse staff, render the charts difficult to use or compromise patient safety. Examples of some errors have been re-created (for clarity and unit anonymity) and are presented in Figs. 1–10 and Supplementary Figs. 1–12.

A. Identification of the patient

All patient records should contain patient identifiers (i.e. first name, surname, date of birth, unique identifier such as hospital number).¹⁶ However, only 55/120 (46%) charts included space for a patient identification label, with a further 62/120 (52%) providing space for handwritten patient details. Three charts (2%) contained no such allocated space.

B. Chart style and layout

An observation chart should display data to facilitate the prompt detection of clinical deterioration. Charts did not always display critical information in order of importance, or use colour and lines thoughtfully. There was often data duplication and the display of unnecessary information. Additionally, some charts failed to display the measurement units of some vital signs. We also noted charts where vital sign documentation and the scoring guide/escalation instructions spanned two sides of paper, thereby requiring temporary data memorisation and potentially risking error. The design of other charts was such that considerable training seemed necessary for their correct use.

B1. Use of colour

Colour can be a powerful tool for improving chart usability.^{14,15} However, several charts were overcomplicated by its use, or demonstrated that poor colour choice or use can lead to errors. One chart (Fig. 1) used white, yellow, pink, red and purple for the weighting of




Fig. 1 Chart using white, yellow, pink, red and purple for the weighting of respiratory rate, with a value of two being applied to respiratory rates in both the red and pink zones of the chart




Fig. 2 Chart using the following scheme for the weighting of respiratory rate values: white=0, blue=1 and 2, and red=3, whilst a different scheme is used on the same chart for weighting temperature values, i.e. white=0, blue=1, 2 and 3




Fig. 3 Chart that uses yellow shading to identify abnormal values. Without appropriate training, staff might simply tick the ‘>38’ or ‘<36’ box if the patient’s temperature exceeds the normal ranges, rather than documenting actual values

respiratory rate, with a weighting of two being applied to respiratory rates in both the red and pink zones. Another chart (Fig. 2) used colours inconsistently, with the following scheme for the weighting of respiratory rate: white=0, blue=1 and 2, and red=3, but a different scheme for temperature: white=0, blue=1, 2 and 3. Supplementary Fig. 1 shows a typical chart employing a 0–3 aggregate EWS system and a pink shaded area to highlight abnormal physiological values. However, the whole of the row used to document oxygen saturation

(S_pO_2) values is shaded pink, i.e. all documented S_pO_2 values appear abnormal. The chart shown in Supplementary Fig. 2 uses unnecessary graduated grey shading.

The lack of colour standardisation is highlighted when considering the numerous schemes used to represent ‘mildly’ and ‘markedly’ abnormal respiratory rate ranges (e.g. red and orange, red and yellow, red and amber, orange and yellow, and pink and yellow, with the exact shades used differing to varying degrees).

Supplementary Fig. 3 demonstrates that coloured EWS weightings superimposed with good intention on vital sign ranges could distract users. Similarly, Supplementary Fig. 4 demonstrates how a lack of attention to detail can lead to errors in colour shading and weighting values (here for pulse rates of 81–90). We also found examples of charts using red and green concurrently – a potential hazard for staff with red-green colour-blindness.




Fig. 4 Chart where several vital signs are documented on the same section, cluttering the display or potentially obscuring deterioration

B2. Graphical versus tabular layouts/displays

Displaying data in a graphical or pictorial form facilitates its understanding and aids detection of abnormality.^{7,17} In one chart, the physiological trend would be undetectable because of the lack of granularity of the ordinate scale. In Fig. 3, although yellow shading is used to identify abnormal values, without appropriate training staff might simply tick the '>38' or '<36' box if the

patient's temperature exceeds the normal ranges, rather than documenting actual values.

We also found examples of several vital signs being plotted on the same chart section. This cluttered the display and could slow data interpretation, obscure deterioration and/or lead to vital signs being mixed up (Fig. 4 and Supplementary Fig. 2). In addition to the use of graduated grey shading, Supplementary Fig. 2 includes a thick horizontal black line that relates to a temperature of 37°C and/or a BP of 180 mmHg. However, both design aspects have the potential to conceal that a systolic BP >160 mmHg is abnormal and scores three points, as the value lies within the chart's unshaded zone and below the thick horizontal black line.

B3. Integration of early warning systems into the observation chart

Disparities between the physiological values used to reflect normal physiology in different areas of the chart and policy occurred frequently. On one chart (Fig. 5), the normal ranges for respiratory rate and S_pO₂ were identified as 11–20 breaths/min and 95–100%, respectively, whereas the chart's escalation plan used different values. Moreover, a S_pO₂ value of 91% lies within a red (i.e. markedly abnormal) zone on the chart but, according to the escalation plan, would not require a referral to the obstetric team. In contrast, a respiratory rate of 27 breaths/min lies within a yellow (i.e. mildly abnormal) zone on the chart, yet on its own does warrant a referral. A similar chart (Supplementary Fig. 5), using a yellow/red scheme to highlight 'mildly'/'markedly' abnormal ranges, used different 'normal' heart rate values on the chart and accompanying weighting table.

RESP rate (write rate in corresp. box)	>30		
	21-30		
	11-20		
	0-10		
Saturations	95-100%		
	92-95%		
	<92%		

Guidelines for Referring to Obstetric Team			
	Less Than	More Than	Only ONE Trigger required
Respiratory Rate	10	25	
Oxygen Saturation	90		
Heart Rate	50	120	
Systolic Blood Pressure	80	160	
Core Temperature	35	39	
Urine Output	30	200	for 2 consecutive hours
Other Significant Concerns e.g. unresponsiveness, abnormal blood gases			

Fig. 5 Chart using a yellow/red colour scheme where the normal ranges for respiratory rate and oxygen saturation (S_pO₂) are identified as 11–20 breaths/min and 95–100%, respectively, but the chart's escalation plan uses different values




Fig. 6 Chart where normal physiological ranges used on the chart differ from those in the early warning system (EWS) table. The shaded white, yellow and red zones on the chart bear little relationship to the weightings. For example, a temperature of 37.6°C lies in a ‘white’ (i.e. normal) zone on the chart but scores one point in the EWS weighting table, whereas a value of 38.4°C lies in the chart’s ‘red’ (i.e. markedly abnormal) zone and also scores one point




Fig. 7 Chart with a poorly aligned axis for heart rate values

Disparities become more prominent when charts attempt to incorporate both the yellow/red approach to escalation and an EWS weighting table. For example, in Fig. 6, the shaded white, yellow and red zones on the chart bear little relationship to the weightings in the EWS table. A temperature of 37.6°C lies in a ‘white’ (i.e. normal) zone on the chart but scores one point in the EWS table, whereas a value of 38.4°C also scores one point but lies in the chart’s ‘red’ (i.e. markedly abnormal) zone. Another error is illustrated in Supple-




Fig. 8 Chart with poor use of the ‘less than’ (<) and ‘more than’ (>) symbols within scoring ranges, making it impossible for staff to know which weighting value to apply to respiratory rates of 20 breaths/min and 30 breaths/min




Fig. 9 Chart with poor use of ranges, making it impossible for staff to know how to handle systolic blood pressure values of 90 mmHg, 100 mmHg and 150 mmHg

mentary Fig. 6, where the scoring system assigns two and four points to vital signs in the ‘yellow’ and ‘red’ ranges, respectively, making a score of five in the escalation algorithm impossible.

B4. Alignment and labelling of axes; row and column shift

To record physiological measurements accurately and quickly, a chart’s axes and their labelling must be clear and precise. We found numerous examples of poorly

B P	v ^	210	4	4
		200	4	4
		190	4	4
		180	4	4
		170	4	4
		160	4	4
		150	3	3
		140	0	0
		130	0	0
		120	0	0
		110	0	0
		100	1	1
		90	1	1
		80	2	2
		70	3	3
		60	3	3
		50	3	3

Fig. 10 Chart that facilitates ‘gaming’, in which staff can ‘choose’ a ‘desired’ escalation pathway by minor ‘manipulation’ of the documented physiological value. In this example, the small change in documented systolic blood pressure between 149 mmHg and 150 mmHg results in large changes in vital sign weighting and clinical actions

aligned axes, which could delay chart completion and introduce error when tracking values temporally (Fig. 7). There were also charts with different weightings on the left and right vertical axes (Supplementary Fig. 7); different pain score ranges on the left (0–4) and right (0–5) (Supplementary Fig. 8); and where a range referred to the wrong parameter (i.e. respiratory rate instead of S_pO_2) (Supplementary Fig. 9). Another chart (Supplementary Fig. 10) doubled the vertical scale width for the temperature range from 36°C to 38°C, which magnified the visual impact of temperature changes in the ‘normal’ range.

‘Row shift’ and ‘column-shift’ occur when users accidentally ‘jump’ into the wrong row or column when recording or reading vital signs data.^{12,14,15} Traditionally, vertical chart axes are labelled on the left. Good practice suggests that the vertical axis should be labelled on both sides of the chart to reduce the risk of ‘row shift’ and that placing thicker vertical black lines every three columns reduces the risk of ‘column shift’.¹⁵ Labelling vertical chart axes on both sides also assists left-handed staff, whose left hand may cover the left-hand scale during documentation. Many charts did not incorporate such features.

Labelling graphs may also prove problematic, ranging from the obvious omission of a legend for systolic BP (Supplementary Fig. 11) to the use of language for, specifically, abbreviations and measurement units. For example, we observed the use of several terms for the

same physiological parameter: ‘maternal heart rate’, ‘heart rate’, ‘pulse rate’, ‘pulse’, ‘HR’, ‘PR’ and ‘P’. Sometimes charts used two different abbreviations (e.g. ‘pulse rate’ and ‘HR’), for the same parameter, even though HR (heart rate) and pulse rate are different parameters and may not have the same value.

B5. Illogical ranges

A common error involved the use or omission of ‘less than’ (<) and ‘more than’ (>) symbols within scoring ranges, making it impossible for staff to know which weighting value to apply to a parameter, e.g. how to handle respiratory rates of 20 breaths/min and 30 breaths/min (Fig. 8), systolic BP values of 90 mmHg, 100 mmHg and 150 mmHg (Fig. 9), and S_pO_2 values of 85% (Supplementary Fig. 12).

B6. Opportunities for ‘gaming’ the escalation of care

Illogical vital signs ranges have the potential to provide staff with the opportunity to ‘choose’ a ‘desired’ escalation pathway by ‘manipulating’ the measured value that is documented. We describe this as ‘gaming’. One factor that encourages ‘gaming’ is the use of EWS weighting boundaries where minimal variation in a measured physiological value results in a large step in vital sign weighting. For example, in Fig. 10, a systolic BP value of 149 mmHg is considered ‘normal’, scoring zero, whereas a value of 150 mmHg scores three points and requires staff to summon immediate help. Therefore, staff can ‘elect’ whether to escalate care or not simply by manipulating the systolic BP value that they record.

Discussion

Whilst it is beyond the scope of this article to describe them, we found many examples where chart design adhered to the ‘good design practice’ outlined in the referenced documents.^{14,15} However, 30% of maternal observation charts contained at least one significant design error. The most common faults included inadequate patient identification, poor use of colour, illogical weighting and escalation ranges, poor alignment and labelling of axes, and the opportunity for staff to ‘game’ the escalation. Overall, the usability of some charts appeared poor. Such errors could lead to delays or difficulties in chart completion, inaccurate scoring and incorrect use of the clinical escalation protocol, resulting in the summoning of immediate support when not required, or manifestly worse, failure to escalate care when an urgent clinical response is necessary.

These findings imply a lack of attention to detail in the development of charts. Chart design is often undertaken by ‘interested’ individuals or by those perceived as having some experience in this regard. However, constructing a chart on the subjective judgements of health professionals is not ideal,^{7,13} especially if they lack

knowledge of design, ergonomics and human factors. Many of the charts surveyed had used the CEMACH-endorsed chart as their foundation,⁴ implementing it in an unchanged or slightly modified way.^{3,10} Consequently, we found that charts often contained similar design errors, probably arising from a practice of copying charts from one organisation to another.

For over a decade, there have been calls to introduce a national obstetric EWS and chart for the UK,^{1,4,18} similar to those used for non-pregnant adults. Given our findings, it seems time for a single body to show leadership by developing a properly designed, standardised obstetric observation chart, as has been done for non-pregnant adults by the Australian Commission on Safety and Quality in Health Care (ACSQHC).^{14,15,19}

Standardisation in healthcare brings many benefits, including reduced error rates, consistent clinical decision-making, opportunities for uniform staff training, better quality of care and improved patient safety.²⁰ However, standardised systems may be perceived as a challenge to professional autonomy and jurisdiction.²¹ Therefore, allowing focused, directed variation of a basic 'standard' chart, as is done by the ACSQHC, may be a way of engaging staff and facilitating improved care.²²

Our study has several strengths and weaknesses. It is the first to consider design errors in obstetric vital signs charts and to highlight their possible impact. Two researchers used a common, objective, systematic approach to interrogate the charts independently. However, not all maternity units in the UK and the Channel Islands are represented by the OAA audit database and this may have led to selection bias. In addition, our data may be subject to non-response and volunteer bias since only 65.5% of invited lead anaesthetists returned charts. Nevertheless, our findings would probably be unchanged (other than in magnitude) by charts from additional units.

Our results mirror the ACSQHC findings that general observation charts frequently have substantial usability problems which may affect the ability of clinical staff to record observations accurately or recognise patient deterioration. We believe that there must be a multi-pronged response to the issues that our work raises. We recommend that individual units urgently and carefully scrutinise their own obstetric observation charts and remedy any design errors. In addition, we recommend that a UK body urgently develops an evidence-based standard obstetric observation chart, which integrates the lessons learned from the scientific study of 'human factors' with user experience. This should be underpinned by research and consensus regarding: (a) 'normal' vital signs ranges in pregnancy and postnatally, (b) a preferred aggregate-weighted EWS, and (c) the EWS value at which care escalation occurs. Such a chart must have a clear layout and style,

an appropriate colour scheme, correct language and labelling, and the ability to record vital signs accurately and quickly. It should incorporate a suitable escalation score to guide clinical management.

Acknowledgements

We would like to acknowledge the OAA Audit Subcommittee for providing the contact details of the lead consultant anaesthetists. We would also like to thank those anaesthetists and other hospital staff who returned their unit's early warning system charts and associated escalation protocols.

Disclosure of interests

GBS was a member of the following groups: Royal College of Physicians of London's National Early Warning Score Development and Implementation Group; National Institute for Health and Care Excellence Guideline Development Group on 'Acutely ill patients in hospital. Recognition of and response to acute illness in adults in hospital'; National Patient Safety Agency Observatory group considering 'Deterioration not recognised or not acted on'; Department of Health Emergency Care Strategy Team's 'Competencies for Recognising and Responding to Acutely Ill Patients in Hospital'.

DB is part-funded by NIHR CLAHRC South London.

Funding

An OAA small project grant, supported by internal resourcing from Bournemouth University, was used to fund the project.

References

1. The Maternal Critical Care Working Group. Providing equity of critical and maternity care for the critically ill pregnant or recently pregnant woman. London: Royal College of Anaesthetists; 2011. Available at: <http://www.rcog.org.uk/womens-health/clinical-guideline/providing-equity-critical-and-maternal-care-critically-ill-pregnant>. Accessed December 2018.
2. Klumpner TT, Kountanis JA, Langen ES, Smith RD, Tremper KK. Use of a novel electronic maternal surveillance system to generate automated alerts on the labor and delivery unit. *BMC Anesthesiol* 2018;**18**:78.
3. Smith GB, Isaacs R, Andrews L, et al. Modified Obstetric Early Warning Systems (MObs) Research Group. Vital signs and other observations used to detect deterioration in pregnant women: an analysis of vital sign charts in consultant-led UK maternity units. *Int J Obstet Anesth* 2017;**30**:44–51.
4. Lewis, G (ed). The Confidential Enquiry into Maternal and Child Health (CEMACH). Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer – 2003–2005. The Seventh Report on Confidential Enquiries into Maternal Deaths in the United Kingdom. London: CEMACH, 2007.
5. Carle C, Alexander P, Columb M, Johal J. Design and internal validation of an obstetric early warning score: secondary analysis

- of the Intensive Care National Audit and Research Centre Case Mix Programme database. *Anaesthesia* 2013;**68**:354–67.
6. Knight M, Kenyon S, Brocklehurst P, Neilson J, Shakespeare J, Kurinczuk JJ (Eds.) on behalf of MBRRACE-UK. Saving Lives, Improving Mothers' Care – Lessons learned to inform future maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2009–12. Oxford: National Perinatal Epidemiology Unit, University of Oxford 2014.
 7. Chatterjee MT, Moon JC, Murphy R, et al. The “OBS” chart: an evidence based approach to re-design of the patient observation chart in a district general hospital setting. *Postgrad Med J* 2005;**81**:663–6.
 8. Christofidis MJ, Hill A, Horswill MS, Watson MO. Observation chart design features affect the detection of patient deterioration: a systematic experimental evaluation. *J Adv Nurs* 2016;**72**:158–72.
 9. Preece MHW, Hill A, Horswill MS, Watson MO. Supporting the detection of patient deterioration: observation chart design affects the recognition of abnormal vital signs. *Resuscitation* 2012;**83**:1111–8.
 10. Isaacs RA, Wee MYK, Bick DE, et al. A national survey of obstetric early warning systems in the United Kingdom: five years on. *Anaesthesia* 2014;**69**:687–92.
 11. Bick DE, Sandall J, Furuta M, et al. A national cross-sectional survey of heads of midwifery services of uptake, benefits and barriers to use of obstetric early warning systems (EWS) by midwives. *Midwifery* 2014;**30**:1140–6.
 12. Christofidis MJ, Hill A, Horswill MS, Watson MO. A human factors approach to observation chart design can trump health professionals' prior chart experience. *Resuscitation* 2013;**84**:657–65.
 13. Preece MH, Hill A, Horswill MS, et al. Designing observation charts to optimise the detection of patient deterioration: reliance on the subjective preferences of healthcare professionals is not enough. *Aust Crit Care* 2012;**25**:238–52.
 14. Horswill MS, Preece MHW, Hill A, Watson MO. Paper-based patient chart design information sheet. Sydney, New South Wales, Australia: Australian Commission on Safety and Quality in Health Care 2010. Available at: <https://espace.library.uq.edu.au/view/UQ:220755>. Accessed July 22, 2018.
 15. Preece MHW, Hill A, Horswill MS, et al. Developer's guide for observation and response charts. Report prepared for the Australian Commission on Safety and Quality in Health Care's program for Recognising and Responding to Clinical Deterioration. Available at: www.safetyandquality.gov.au/sites/default/files/migrated/ORC-Developers-Guide-4-Oct-2010.pdf. Accessed December 2018.
 16. Health and Social Care Information Centre, Academy of Medical Royal Colleges. Standards for the clinical structure and content of patient records. London: HSCIC; 2013.
 17. Horswill MS, Preece MHW, Hill A, et al. Human factors research regarding observation charts: research project overview. Report prepared for the Australian Commission on Safety and Quality in Health Care's program for Recognising and Responding to Clinical Deterioration. School of Psychology, The University of Queensland, May 2010.
 18. Quinn AC, Meek T, Waldmann C. Obstetric early warning systems to prevent bad outcome. *Curr Opin Anaesthesiol* 2016;**29**:268–72.
 19. Observation and response charts. Australian Commission on Safety and Quality in Health Care, Sydney. 2012. Available at: <https://www.safetyandquality.gov.au/our-work/recognising-and-responding-to-clinical-deterioration/observation-and-response-charts/> Accessed July 22, 2018.
 20. Leotsakos A, Zheng H, Croteau R, et al. Standardization in patient safety: the WHO High 5s project. *Int J Qual Health Care* 2014;**26**:109–16.
 21. Mackintosh N, Watson K, Rance S, Sandall J. Value of a modified early obstetric warning system (MEOWS) in managing maternal complications in the peripartum period: an ethnographic study. *BMJ Qual Saf* 2014;**23**:26–34.
 22. EE1 ORC2 Fact Sheet. Modifying the observation and response chart for local use. Australian Commission on Safety and Quality in Health Care. Sydney. 2012. Available at: <https://www.safetyandquality.gov.au/wp-content/uploads/2012/08/ee1-orc-2-fact-sheet.pdf>. Accessed July 22, 2018.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jjoa.2019.01.001>.