



ELSEVIER

[www.obstetanaesthesia.com](http://www.obstetanaesthesia.com)

CORRIGENDUM

**Corrigendum to “Ephedrine versus phenylephrine as a vasopressor for spinal anaesthesia-induced hypotension in parturients undergoing high-risk caesarean section: meta-analysis, meta-regression and trial sequential analysis” [*Int J Obstet Anesth* 2019;37:16–28]**

M. Heesen,<sup>a</sup> K. Rijs,<sup>b</sup> N. Hilber,<sup>a</sup> W.D. Ngan Kee,<sup>c</sup> R. Rossaint,<sup>d</sup> C. van der Marel,<sup>b</sup> M. Klimek<sup>b</sup>

<sup>a</sup>Department of Anaesthesia, Kantonsspital Baden, Baden, Switzerland

<sup>b</sup>Department of Anaesthesiology, Erasmus University Medical Centre, Rotterdam, The Netherlands

<sup>c</sup>Department of Anaesthesiology, Sidra Medicine, Doha, Qatar

<sup>d</sup>Department of Anaesthesia, University Hospital RWTH Aachen, Aachen, Germany

---

The authors regret that in Table 1 the ‘Definition for hypotension’ was incorrectly listed for the authors Dyer et al. IJOA and Dyer et al. ANAE (row 2 and 3). The ‘Definition for hypotension’ for these authors should have been listed as follows:

MAP >20% decrease from baseline and MAP <110 mmHg (rather than MAP >20% decrease from baseline or MAP <110 mmHg).

The authors would like to apologise for any inconvenience caused.