

May 2019, Volume 80, Number 5

CONTENTS

Continuing Medical Education

- CME 1179** Hair disorders in patients with cancer
Azael Freites-Martinez, MD, Jerry Shapiro, MD, Shari Goldfarb, MD, Julie Nangia, MD, Joaquin J. Jimenez, MD, Ralf Paus, MD, FRBS, and Mario E. Lacouture, MD
- 1197** CME examination
- 1198** Answers to CME examination (Identification No. JA0519), May 2019 issue of the *Journal of the American Academy of Dermatology*
- CME 1199** Hair disorders in cancer survivors
Azael Freites-Martinez, MD, Jerry Shapiro, MD, Corina van den Hurk, PhD, Shari Goldfarb, MD, Joaquin J. Jimenez, MD, Anthony M. Rossi, MD, Ralf Paus, MD, FRBS, and Mario E. Lacouture, MD
- 1213** Answers to CME examination (Identification No. JB0519), May 2019 issue of the *Journal of the American Academy of Dermatology*
- 1214** CME examination
- CME 1215** Acquired disorders with depigmentation: A systematic approach to vitiliginoid conditions
Mohammed D. Saleem, MD, MPH, Elias Oussedik, MD, Jennifer J. Schoch, MD, Adam C. Berger, MD, and Mauro Picardo, MD, PhD
- 1231** Answers to CME examination (Identification No. JC0519), May 2019 issue of the *Journal of the American Academy of Dermatology*
- 1232** CME examination
- CME 1233** Acquired disorders with hypopigmentation: A clinical approach to diagnosis and treatment
Mohammed D. Saleem, MD, MPH, Elias Oussedik, MD, Mauro Picardo, MD, PhD, and Jennifer J. Schoch, MD

	1251	CME examination
	1252	Answers to CME examination (Identification No. JD0519), May 2019 issue of the <i>Journal of the American Academy of Dermatology</i>
Letter from the Editor	1253	Planning better research projects: Advice for young researchers <i>Dirk M. Elston, MD</i>
A Clinician's Perspective	1254	Coming into focus: Dermoscopy for basal cell carcinomas <i>Warren R. Heymann, MD</i>
Original Articles	 1256	Melanoma prognosis in the United States: Identifying barriers for improved care Bottom line: States with increased melanoma incidence and dermatologist density had better melanoma survival. States with more active physicians and more non-Hispanic whites had worse melanoma survival. Efforts to improve survival should occur locally, as local differences explain most survival variations. <i>Zachary H. Hopkins, MD, Christopher Moreno, BS, Ryan Carlisle, BS, and Aaron M. Secrest, MD, PhD</i>
	1263	<i>CDKN2A</i> germline mutations are not associated with poor survival in an Italian cohort of melanoma patients Bottom line: Follow-up differences may be implicated in regional differences concerning the impact of <i>CDKN2A</i> on the survival of patients with melanoma. <i>Bruna Dalmasso, MD, Lorenza Pastorino, PhD, Giulia Ciccarese, MD, Virginia Andreotti, PhD, Federica Grillo, MD, Luca Mastracci, MD, Francesco Spagnolo, MD, Alberto Ballestrero, MD, PhD, Paola Queirolo, MD, William Bruno, MD, PhD, and Paola Ghiorzo, PhD</i>
	1272	Retrospective single-center study evaluating clinical and dermoscopic features of longitudinal melanonychia, ABCDEF criteria, and risk of malignancy Bottom line: Greater width and width percentage may signify subungual melanoma; ABCDEF criteria should be re-examined. <i>Dayoung Ko, BS, Clara Oromendia, MS, Richard Scher, MD, and Shari R. Lipner, MD, PhD</i>
	1284	Cutaneous nevi and risk of melanoma death in women and men: A prospective study Bottom line: Number of nevi might potentially serve as an independent prognostic factor of melanoma in clinical practices, particularly for men. <i>Wen-Qing Li, PhD, Eunyoung Cho, ScD, Martin A. Weinstock, MD, Suyun Li, PhD, Meir J. Stampfer, MD, DrPH, and Abrar A. Qureshi, MD, MPH</i>

- 1292** Clinicopathologic, misdiagnosis, and survival differences between clinically amelanotic melanomas and pigmented melanomas
Bottom line: Patients with amelanotic melanomas may be more likely to be older and have nonmelanoma skin cancer and red hair and less likely to have many nevi or a family history of melanoma. Clinicians should be alert to amelanotic lesions in patients with these factors.
Lauren C. Strazzulla, BA, Xiaoxue Li, PhD, Kathleen Zhu, BA, Jean-Phillip Okhovat, MD, MPH, Sandra J. Lee, ScD, and Caroline C. Kim, MD
- 1299** Age as key factor for pattern, timing, and extent of distant metastasis in patients with cutaneous melanoma: A study of the German Central Malignant Melanoma Registry
Bottom line: This study found an inverse relationship between age at diagnosis and the number of metastatic sites; these findings may have implications for optimizing outcomes of patients with melanoma.
Maximilian Gassenmaier, MD, Ulrike Keim, PHD, Ulrike Leiter, MD, Thomas K. Eigentler, MD, Martin Röcken, MD, Anja Gesierich, MD, Rose K. C. Moritz, MD, Lucie Heinzerling, MD, MPH, Thomas Tüting, MD, Uwe Wollina, MD, and Claus Garbe, MD
- 1308** Overall and subgroup prevalence of acne vulgaris among patients with hidradenitis suppurativa
Bottom line: Understanding the burden of acne vulgaris among patients with hidradenitis suppurativa may support evaluation and optimization of management for both conditions.
Sara Wertenteil, BA, Andrew Strunk, MA, and Amit Garg, MD
- 1314** Clindamycin versus clindamycin plus rifampicin in hidradenitis suppurativa treatment: Clinical and ultrasound observations
Bottom line: The study results suggest that clindamycin as monotherapy may be a useful and safe alternative to the combination of clindamycin and rifampicin regardless of the clinical stage of hidradenitis suppurativa.
Raffaele Dante Caposiena Caro, MD, Maria Vittoria Cannizzaro, MD, Elisabetta Botti, MD, Cosimo Di Raimondo, MD, Eleonora Di Matteo, MD, Roberta Gaziano, MD, and Luca Bianchi, MD
- 1322** Ustekinumab treatment is associated with decreased systemic and vascular inflammation in patients with moderate-to-severe psoriasis: Feasibility study using ¹⁸F-fluorodeoxyglucose PET/CT
Bottom line: Ustekinumab treatment was significantly associated with decreased systemic and vascular inflammation related to metabolic syndrome and cardiovascular disease among patients with psoriasis.
Byung-Soo Kim, MD, PhD, Won-Ku Lee, MD, Kyoungjune Pak, MD, PhD, Junhee Han, PhD, Gun-Wook Kim, MD, Hoon-Soo Kim, MD, PhD, Hyun-Chang Ko, MD, PhD, Moon-Bum Kim, MD, PhD, and Seong-Jang Kim, MD, PhD
- 1332** All-cause and cause-specific mortality in psoriasis: A systematic review and meta-analysis
Bottom line: Psoriasis is associated with increased mortality risk from all-causes in a dose-response manner with disease severity and from several specific causes, such as cardiovascular, liver, kidney, and infectious disease. Thus, psoriasis patients should receive appropriate screening and preventative interventions.
Ashar Dhana, MBBCh, MPH, Hsi Yen, MD, MPH, Hsuan Yen, MD, MPH, and Eunyoung Cho, ScD

- 1344** Secukinumab for moderate-to-severe palmoplantar pustular psoriasis: Results of the 2PRECISE study
Bottom line: The 2PRECISE randomized controlled trial shows potential benefits of secukinumab in the treatment of palmoplantar pustular psoriasis over 52 weeks, with improved quality of life and no unexpected adverse events.
Ulrich Mrowietz, MD, Hervé Bachelez, MD, PhD, A. David Burden, MD, Michael Rissler, PhD, Christian Sieder, PhD, Roberto Orsenigo, MD, and Kamel Chaouche-Teyara, PhD
- 1353** Increased topical generic prices by manufacturers
Bottom line: Price increases of topical generic drugs reflect price increases at the manufacturing level.
Mehul D. Bhatt, MD, MBA, Birju D. Bhatt, MD, James T. Dorrian, MD, and Beth N. McLellan, MD
- 1358** A healthy diet in women is associated with less facial wrinkles in a large Dutch population-based cohort
Bottom line: Dietary recommendations for skin aging preventing strategies could in addition help to improve overall health.
Selma Mekić, MD, Leonie C. Jacobs, MD, PhD, Merel A. Hamer, MD, M. Arfan Ikram, MD, PhD, Josje D. Schoufour, PhD, David A. Gunn, PhD, Jessica C. Kiefte-de Jong, PhD, and Tamar Nijsten, MD, PhD
- 1364** Association of antinuclear antibody status with clinical features and malignancy risk in adult-onset dermatomyositis
Bottom line: There is a strong association between negative antinuclear antibody status and increased risk of underlying malignancy in adults with dermatomyositis. More frequent follow-up and malignancy screening may be warranted in adults with dermatomyositis and a negative antinuclear antibody testing result.
Paul M. Hoesly, MD, Jason C. Sluzevich, MD, Anokhi Jambusaria-Pahlajani, MD, MSCE, Elizabeth R. Lesser, MS, Michael G. Heckman, MS, and Andy Abril, MD
- 1371** Vitiligo and major depressive disorder: A bidirectional population-based cohort study
Bottom line: Clinicians treating vitiligo patients should be aware of this bidirectional relationship and refer these patients for mental health services accordingly.
Isabelle A. Vallerand, PhD, Ryan T. Lewinson, MD, PhD, Laurie M. Parsons, MD, Jori Hardin, MD, MSc, Richard M. Haber, MD, Mark W. Lowerison, MSc, Cheryl Barnabe, MD, MSc, and Scott B. Patten, MD, PhD
- 1380** The diagnostic accuracy of dermoscopy for basal cell carcinoma: A systematic review and meta-analysis
Bottom line: Dermoscopy training should be included in dermatology residency programs because the level of expertise significantly affects diagnostic accuracy.
Ofer Reiter, MD, Ilit Mimouni, BMedSc, Michael Gdalevich, MD, MPH, Ashfaq A. Marghoob, MD, Assi Levi, MD, Emmilia Hodak, MD, and Yael Anne Leshem, MD, MCR

1389 Severity of disease and quality of life in parents of children with alopecia areata, totalis, and universalis: A prospective, cross-sectional study

Bottom line: Awareness of the impact of alopecia areata on the emotional health of parents should lead to more family-centered care for this disease. Parents need support systems as much as children do.

Elana Putterman, BS, Deepa P. Patel, MD, Gabriela Andrade, MD, Katya L. Harfmann, MD, Marcia Hogeling, MD, Carol E. Cheng, MD, Carolyn Goh, MD, Rachel S. Rogers, MS, and Leslie Castelo-Soccio, MD, PhD

1395 Serlopitant reduced pruritus in patients with prurigo nodularis in a phase 2, randomized, placebo-controlled trial

Bottom line: Serlopitant has potential as a therapeutic option for treatment of pruritus associated with prurigo nodularis.

Sonja Ständer, MD, Paul Kwon, MD, Joe Hirman, PhD, Andrew J. Perlman, MD, PhD, Elke Weisshaar, MD, Martin Metz, MD, and Thomas A. Luger, MD, for the TCP-102 Study Group

Dermatologic Surgery

1403 Melanoma and melanoma in-situ diagnosis after excision of atypical intraepidermal melanocytic proliferation: A retrospective cross-sectional analysis

Bottom line: Ambiguous melanocytic biopsy designations should be avoided, but, when present, warrant clinicopathologic correlation, with malignancy risk factors prompting consideration to excise as melanoma in situ.

Nina R. Blank, MD, Brian P. Hibler, MD, Ian W. Tattersall, MD, PhD, Courtney J. Ensslin, MD, Erica H. Lee, MD, Stephen W. Dusza, DrPH, Kishwer S. Nehal, MD, Klaus J. Busam, MD, and Anthony M. Rossi, MD

Reviews

1410 Screening of thyroid function and autoantibodies in patients with alopecia areata: A systematic review and meta-analysis

Bottom line: This meta-analysis suggests that it might be beneficial to screen for the signs and symptoms associated with thyroid dysfunction and autoimmune thyroid diseases in patients with alopecia areata.

Solam Lee, MD, Young Bin Lee, MD, Beom Jun Kim, MD, and Won-Soo Lee, MD, PhD

1414 Reflectance confocal microscopy terminology glossary for nonmelanocytic skin lesions: A systematic review

Bottom line: We built a glossary of reflectance confocal microscopy terms for diagnosing nonmelanocytic lesions. We identified terms with similar definition and histopathologic correlates and reduced the total number of terms by 52.5%. This systematic review could be used to attain terminology consensus.

Cristian Navarrete-Dechent, MD, Antonio P. DeRosa, MS, MLIS, AHIP, Caterina Longo, MD, Konstantinos Liopyris, MD, Margaret Oliviero, ARNP, Harold Rabinovitz, MD, Ashfaq A. Marghoob, MD, Allan C. Halpern, MD, Giovanni Pellacani, MD, Alon Scope, MD, and Manu Jain, MD

1428 Diagnosis and management of delusional parasitosis

Bottom line: Based on the published data on efficacy, the incidence of side effects, and attributable risk, risperidone (0.5-4 mg/day) is a reasonable first-line choice for pharmacotherapy.

Elliott H. Campbell, BSc, Dirk M. Elston, MD, James D. Hawthorne, MD, and David R. Beckert, MD

- 1435** Efficacy and safety of antiscabietic agents: A systematic review and network meta-analysis of randomized controlled trials
Bottom line: There is no one treatment that ranked highest in all aspects. Physicians should consider an easy-to-use treatment with a balance between efficacy and safety.
Kunlawat Thadanipon, MD, Thunyarat Anothaisintawee, MD, PhD, Sasivimol Rattanasiri, PhD, Ammarin Thakkinstian, PhD, and John Attia, MD, PhD

Research Letters

- 1445** Validated patient-reported outcome measurements for psoriasis may not reflect patients' current preferences
Oluwatobi Ogbechie-Godec, MD, MBA, Sarah Azarchi, BS, Jasmine Lee, BS, David E. Cohen, MD, MPH, Andrea Neimann, MD, and Arielle R. Nagler, MD
- 1447** What do patients undergoing Mohs micrographic surgery want? Results of a patient values survey based on the Outpatient and Ambulatory Surgery Consumer Assessments of Healthcare Providers and Systems
Paul R. Massey, MD, Ruibin Wang, MHS, Simi D. Cadmus, MS, Katherine R. Sebastian, RN, MPH, and Matthew C. Fox, MD
- 1449** Morphea and systemic sclerosis are associated with an increased risk for melanoma and nonmelanoma skin cancer
Emily Boozalis, BA, Ami A. Shah, MD, MHS, Fredrick Wigley, MD, Sewon Kang, MD, and Shawn G. Kwatra, MD
- 1451** A retrospective study: Application site pain with the use of crisaborole, a topical phosphodiesterase 4 inhibitor
Christine Pao-Ling Lin, BA, Samantha Gordon, MD, Min Ji Her, PharmD, and David Rosmarin, MD
- 1453** Idiopathic Stevens-Johnson syndrome and toxic epidermal necrolysis: Prevalence and patients' characteristics
Guillaume Chaby, MD, Saskia Ingen-Housz-Oro, MD, Nicolas De Prost, MD, PhD, Pierre Wolkenstein, MD, PhD, Olivier Chosidow, MD, PhD, and Laurence Fardet, MD, PhD
- 1455** Interruption of oral clindamycin plus rifampicin therapy in patients with hidradenitis suppurativa: An observational study to assess prevalence and causes
Luca Schneller-Pavelescu, MD, Eduardo Vergara-de Caso, MD, Antonio Martorell, MD, Jorge Romani, MD, PhD, Mireya Lázaro, MD, Eva Vilarrasa, MD, Blanca Díaz-Ley, MD, Igor Vázquez-Osorio, MD, Juan Manuel Segura Palacios, MD, José Manuel Azaña, MD, Marcos A. González-López, MD, Javier Cañueto, MD, Alejandro Molina-Leyva, MD, Maria Leiva-Salinas, MD, Francisco José Navarro-Triviño, MD, José Sánchez-Payá, MD, and José C. Pascual, MD
- 1458** Effects of new keratinocyte carcinomas on skin-related quality of life: Results from the Veterans Affairs Keratinocyte Carcinoma Chemoprevention Trial
Julia A. Siegel, MD, Meghan Beatson, BS, Mary-Margaret Chren, MD, and Martin A. Weinstock, MD, PhD; Veterans Affairs Keratinocyte Carcinoma Chemoprevention Trial group

- 1460** Screening for depression and suicidality in psoriasis patients:
A survey of US dermatologists
Sydney E. Liang, BS, Jeffrey M. Cohen, MD, and Roger S. Ho, MD, MS, MPH

- Dermatoethics Consultation** **1463** The ethical issue of cherry picking patients
Travis J. Morrell, MD, MPH, Sailesh Konda, MD, and Jane M. Grant-Kels, MD

JAAD ONLINE

Available at <http://www.jaad.org>

Clinical Pearls

- e99** Simple methods to enhance the diagnosis of scabies
Jiaoping Tang, BS, Zimeng You, BS, and Yuping Ran, MD, PhD
- e101** Prevention of thermal burns from magnetic resonance imaging in patients with tattoos
Jason F Wang, BA, and Nicole M. Hindman, MD
- e103** Hydrocolloid dressing application for the treatment of pediatric onychodystrophies
Solomon Geizhals, BA, Christine T. Lauren, MD, and Shari R. Lipner, MD, PhD
- e105** Mobile phone reminders for onychomycosis medication adherence
Krysta Lin, BS, and Shari R. Lipner, MD, PhD

Therapeutic Pearls

- e109** Platelet-rich plasma as an adjunctive treatment in lichen planopilaris
Abhijeet Kumar Jha, MD
- e111** Intralesional vitamin D3 in periungual warts
Deepak Jakhar, MD, Ishmeet Kaur, MD, and Rachita Misri, MD
- e113** An office-based evaluation of various itraconazole brands
Raj Kumar Mehta, MD, Deepak Jakhar, MD, and Ishmeet Kaur, MD

Surgical Pearls

- e115** Split earlobe repair with piercing preservation and optimal earlobe aesthetics
C. Helen Malone, MD, Vlad Codrea, MD, PhD, and Richard F. Wagner, Jr, MD
- e117** Use of QuikClot Combat Gauze during Mohs stages for intraoperative hemostasis
Valerie Fisher Shiu, MD, and Richard Keller, MD



- e119** Augmentation and eversion of lips without injections:
The lip peel
Carlos G. Wambier, MD, PhD, Ideli C. Neitzke, MD, Kachiu C. Lee, MD, MPH, Seaver L. Soon, MD, Peter P. Rullan, MD, Marina Landau, MD, João Carlos L. Simão, MD, PhD, J. Barton Sterling, MD, and Harold J. Brody, MD, for the International Peeling Society

	e121	The use of a tympanoplasty blade for tumor extirpation of the auricle <i>Lindsey E. Richards, MD, Faramarz Samie, MD, PhD, Scott Troob, MD, and Jesse M. Lewin, MD</i>
Technology Pearl	e123	A simple technique to increase field of view of a universal serial bus dermatoscope <i>Deepak Jakhar, MD, and Ishmeet Kaur, MD</i>
Notes & Comments	e125	Reply to: “The ethical issue of cherry picking patients” <i>Thomas Vazquez, BS</i>
	e127	Response to a letter to the editor regarding “The ethical issue of cherry picking patients” <i>Travis J. Morrell, MD, MPH, Sailesh Konda, MD, and Jane M. Grant-Kels, MD</i>
	e129	Acne vulgaris in patients with hidradenitis suppurativa <i>Astrid-Helene Ravn Jørgensen, MD, Hans Christian Ring, MD, PhD, and Simon Francis Thomsen, MD, PhD</i>
	e131	Overall and subgroup prevalence of acne vulgaris among patients with hidradenitis suppurativa <i>Sara Wertenteil, BA, Andrew Strunk, MA, and Amit Garg, MD</i>
Book Review	e133	Biodynamic excisional skin tension lines for cutaneous surgery <i>Derek W. Nickerson, MD</i>
Iotaderma	e135	Iotaderma (#303) <i>Robert I. Rudolph, MD, FACP</i>

JAAD Game Changers	1198	JAAD Game Changers: Acquired cold urticaria: Clinical features, particular phenotypes, and disease course in a tertiary care center cohort <i>Eden Lake, MD</i>
	1255	JAAD Game Changers: Multivariate analysis of potential risk factors for lymph node metastasis in patients with cutaneous squamous cell carcinoma of the head and neck <i>Eden Lake, MD</i>

To access the current version of the Information for Authors, Transfer of Copyright Form, Authorship Declaration Form, Conflict of Interest Disclosure, and Authors’ Rights Statement, as well as many other pertinent documents, please visit our website at www.jaad.org/content/auth_page.

Full text access to *Journal of the American Academy of Dermatology Online* is now available for Academy members and all print subscribers. To activate your online subscription, please visit *Journal of the American Academy of Dermatology Online* by pointing your browser to <http://www.jaad.org>.

Icon Key

	=	CME article
	=	Video content available at http://www.jaad.org
	=	Podcast available at http://www.aad.org/olc
	=	MOC credits available

Early online publication: Articles in press

The *Journal of the American Academy of Dermatology* posts in-press articles online before they appear in the print edition of the *Journal*. These Articles in Press are available from both the JAAD website (www.jaad.org) and Science Direct (www.sciencedirect.com). To access Articles in Press, simply click the Articles in Press link.

JAAD Case Reports

The *Journal of the American Academy of Dermatology* no longer publishes Case Reports. These can now be found in our new open access journal, *JAAD Case Reports*, at <http://www.jaadcasereports.org>.