

August 2019, Volume 81, Number 2

CONTENTS

Continuing Medical Education



Basic chemical peeling: Superficial and medium-depth peels
*Kachiu C. Lee, MD, MPH, Carlos G. Wambier, MD, PhD, Seaver L. Soon, MD,
J. Barton Sterling, MD, Marina Landau, MD, Peter Rullan, MD, and
Harold J. Brody, MD, on behalf of the International Peeling Society*

325 CME examination

326 Answers to CME examination (Identification No. JA0819),
August 2019 issue of the *Journal of the American Academy of
Dermatology*



Advanced chemical peels: Phenol-croton oil peel
*Carlos G. Wambier, MD, PhD, Kachiu C. Lee, MD, MPH, Seaver L. Soon, MD,
J. Barton Sterling, MD, Peter P. Rullan, MD, Marina Landau, MD, and
Harold J. Brody, MD, for the International Peeling Society*

337 CME examination

338 Answers to CME examination (Identification No. JB0819),
August 2019 issue of the *Journal of the American Academy of
Dermatology*

Letter from the Editor

339 Duplicate publication
Dirk M. Elston, MD

A Clinician's Perspective

340 The need to comprehend erosive pustular dermatosis of the scalp
is coming to a head
Warren R. Heymann, MD

- Original Articles** **342** Incidence of bloodstream infections and predictive value of qualitative and quantitative skin cultures of patients with overlap syndrome or toxic epidermal necrolysis: A retrospective observational cohort study of 98 cases
- Bottom line:** Skin cultures may help predicting the pathogens involved in bloodstream infections.
- Aude Lecadet, MD, Paul-Louis Woerther, MD, PhD, Camille Hua, MD, Audrey Colin, Camille Gomart, PharmD, MSc, Jean-Winoc Decousser, PharmD, PhD, Armand Mekontso Dessap, MD, PhD, Pierre Wolkenstein, MD, PhD, Olivier Chosidow, MD, PhD, Nicolas de Prost, MD, PhD, and Saskia Ingen-Housz-Oro, MD*
- 348** Vascular endothelial growth factor gene polymorphisms in patients with rosacea: A case-control study
- Bottom line:** Vascular endothelial growth factor may be a new target for future rosacea treatments.
- Yildiz Hayran, MD, Incilay Lay, MD, Mehmet Cem Mocan, MD, Tuba Bozduman, MD, and Sibel Ersoy-Evans, MD*
- 355** Nonbullous pemphigoid: Insights in clinical and diagnostic findings, treatment responses, and prognosis
- Bottom line:** Clinicians should realize that histopathology is not useful for diagnosing nonbullous pemphigoid; direct and indirect immunofluorescence microscopy are required. Patients have an increased all-cause mortality risk.
- Aniek Lamberts, MD, Joost M. Meijer, MD, PhD, Hendri H. Pas, PhD, Gilles F. H. Diercks, MD, PhD, Barbara Horváth, MD, PhD, and Marcel F. Jonkman, MD, PhD*
- 364** Sentinel lymph node biopsy in Merkel cell carcinoma: Predictors of sentinel lymph node positivity and association with overall survival
- Bottom line:** Following National Comprehensive Cancer Network guidelines and performing sentinel lymph node biopsy on patients with Merkel cell carcinoma without clinically evident nodal metastases could improve prognostic accuracy.
- Rosalynn R. Z. Conic, MD, PhD, Jennifer Ko, MD, Stephanie Saridakis, BS, Giovanni Damiani, MD, Pauline Funchain, MD, Allison Vidimos, MD, and Brian R. Gastman, MD*
- 373** Risk factors for keratinocyte carcinoma skin cancer in nonwhite individuals: A retrospective analysis
- Bottom line:** In nonwhite patients, there should be increased vigilance in sun-protected areas, particularly the anogenital region, during skin examinations, and all patients should be counselled on smoking cessation.
- Kumar S. Nadhan, MD, Christina L. Chung, MD, Erin M. Buchanan, PhD, Christine Shaver, MD, Scott Shipman, MS, Rina M. Allawh, MD, Melissa L. Hoffman, BS, Geoffrey Lim, MD, Mark Abdelmalek, MD, and Carrie Ann Cusack, MD*
- 379** Pigments in American tattoo inks and their propensity to elicit allergic contact dermatitis
- Bottom line:** Older metallic pigments have been replaced with organic dyes or less allergenic metals, such as zinc and barium. Cases of contact dermatitis to these organic dyes have been reported.
- Walter Liszewski, MD, and Erin M. Warshaw, MD, MS*

- 386** Estimating CDKN2A mutation carrier probability among global familial melanoma cases using GenoMELPREDICT
Bottom line: GenoMELPREDICT can aid in appropriate patient management, whether that is by recommending genetic testing or cancer risk counselling.
Nicholas J. Taylor, PhD, Nandita Mitra, PhD, Lu Qian, MS, Marie-Françoise Avril, MD, D. Timothy Bishop, PhD, Brigitte Bressac-de Paillerets, PhD, William Bruno, MD, PhD, Donato Calista, MD, Francisco Cuellar, MD, Anne E. Cust, PhD, Florence Demenais, MD, David E. Elder, MD, Anne-Marie Gerdes, PhD, Paola Ghiorzo, PhD, Alisa M. Goldstein, PhD, Thais C. Grazziotin, MD, Nelleke A. Gruis, PhD, Johan Hansson, MD, PhD, Mark Harland, PhD, Nicholas K. Hayward, PhD, Marko Hocevar, MD, PhD, Veronica Höiom, PhD, Elizabeth A. Holland, BSc, Christian Ingvar, MD, PhD, Maria Teresa Landi, MD, PhD, Gilles Landman, MD, PhD, Alejandra Larre-Borges, MD, Graham J. Mann, PhD, Eduardo Nagore, MD, PhD, Håkan Olsson, MD, PhD, Jane M. Palmer, RN, Barbara Perić, MD, PhD, Dace Pjanova, PhD, Antonia L. Pritchard, PhD, Susana Puig, MD, PhD, Helen Schmid, MPH, Nienke van der Stoep, PhD, Margaret A. Tucker, MD, Karin A. W. Wadt, MD, PhD, Xiaohong R. Yang, PhD, Julia A. Newton-Bishop, MD, and Peter A. Kanetsky, PhD, on behalf of the GenoMEL Study Group
- 395** Coffee, tea, caffeine, and risk of nonmelanoma skin cancer in a Chinese population: The Singapore Chinese Health Study
Bottom line: We showed that intake of coffee or caffeine may reduce the risk of squamous cell and basal cell carcinomas among Chinese in Singapore.
Choon Chiat Oh, MD, MRCP, Aizhen Jin, MSc, Jian-Min Yuan, MD, PhD, and Woon-Puay Koh, MD, PhD
- 403** Are there distinct clinical and pathological features distinguishing idiopathic from drug-induced subacute cutaneous lupus erythematosus? A European retrospective multicenter study
Bottom line: Our results show that older age at onset and leukocytoclastic vasculitis are more commonly seen in drug-induced cases, and mucin deposition and positive immunofluorescence are clues to the idiopathic form.
Federica Guicciardi, MD, Laura Atzori, MD, Angelo Valerio Marzano, MD, Simona Tavecchio, MD, Giampiero Girolomoni, MD, Chiara Colato, MD, Axel Patrice Villani, MD, Jean Kanitakis, MD, Christina Mitteldorf, MD, Rosanna Satta, MD, Bernard Cribier, MD, Laurence Gusdorf, MD, Maria Teresa Rossi, MD, Piergiacomo Calzavara-Pinton, MD, Isabel Bielsa, MD, Maria Teresa Fernandez-Figueras, MD, Werner Kempf, MD, Giorgio Filosa, MD, Luca Pilloni, MD, and Franco Rongioletti, MD
- 412** Improving nonattendance rates among pediatric patients with Medicaid or private insurance
Bottom line: Shortened time intervals between scheduling and appointment dates resulted in lower nonattendance among all patients. We hope that optimized scheduling might encourage increased acceptance of Medicaid-insured patients into dermatology practices.
Sofia B. Chaudhry, MD, Elaine Siegfried, MD, Umar A. Sheikh, BA, Cassandra Simonetta, MD, Niraj Butala, MD, and Eric Armbrecht, PhD

- 417** Reflectance confocal microscopy confirms residual basal cell carcinoma on clinically negative biopsy sites before Mohs micrographic surgery: A prospective study
Bottom line: Reflectance confocal microscopy may improve triage of patients with clinically negative biopsy sites, allowing selection of patients who should undergo surgery and those who can undergo observation only.
Cristian Navarrete-Dechent, MD, Miguel Cordova, MD, Saud Aleissa, MD, Konstantinos Liopyris, MD, Stephen W. Dusza, DrPH, William Phillips, BS, Anthony M. Rossi, MD, Erica H. Lee, MD, Ashfaq A. Marghoob, MD, and Kishwer S. Nehal, MD
- 427** A prospective multicenter study assessing effectiveness and safety of secukinumab in a real-life setting in 158 patients
Bottom line: Patients were more heterogeneous than in clinical trials but effectiveness rates were similar. Main predictors of treatment failure were high body mass index (>30 kg/cm²) and multiple previous treatment failures.
José-María Ortiz-Salvador, MD, Martina Saneleuterio-Temporal, MD, Jorge Magdaleno-Tapial, MD, Manel Velasco-Pastor, MD, Conrad Pujol-Marco, MD, Antonio Sahuquillo-Torralba, MD, Almudena Mateu-Puchades, MD, Gerard Pitarch-Bort, MD, Juan-Ignacio Mari-Ruiz, MD, Javier Mataix-Díaz, MD, Encarna Montesinos-Villaescusa, MD, Julia Miralles-Botella, MD, Laura García-Fernández, MD, Antonio Martorell-Calatayud, MD, Isabel Belinchón-Romero, MD, José-Luis Sánchez-Carazo, MD, and Amparo Pérez-Ferriols, MD
- 433** Dermatologic care of uninsured patients managed at free clinics
Bottom line: Free clinics can be made aware of the uninsured dermatologic patients they are most likely to see so that they can provide patients with higher-quality dermatology management.
Noura Ayoubi, BS, Abu-Sayef Mirza, MD, MPH, Justin Swanson, MPH, Rema Hamoui, BS, and Rahul Mhaskar, MPH, PhD
- 438** Issues in transgender dermatology: A systematic review of the literature
Bottom line: We aim to show dermatologists how to best provide routine dermatologic care, treat complications of common diseases, and provide safe and noninvasive treatment options for transgender patients.
Patrick Sullivan, MD, John Trinidad, MD MPH, and Dathan Hamann, MD
- 448** Management of Kaposi sarcoma after solid organ transplantation: A European retrospective study
Bottom line: Therapeutic management of post-transplant Kaposi sarcoma is mostly based on reduction of immunosuppression and conversion to mammalian target of rapamycin inhibitor, inducing response in more than 80% of patients.
Julie Delyon, MD, PhD, Clementine Rabate, MD, Sylvie Euvrard, MD, Catherine A. Harwood, MD, PhD, Charlotte Proby, FRCP, A. Tülin Güleç, MD, Deniz Seçkin, MD, Veronique Del Marmol, MD, PhD, Jan Nico Bouwes-Bavinck, MD, PhD, Carla Ferrándiz-Pulido, MD, PhD, Maria Andrea Ocampo, MD, Stephane Barete, MD, PhD, Christophe Legendre, MD, PhD, Camille Francès, MD, PhD, Raphael Porcher, MD, PhD, and Celeste Lebbe, MD, PhD, the Skin Care in Organ Transplant Patients Europe (SCOPE) group

- 456** Association between cycline antibiotic and development of pseudotumor cerebri syndrome
Bottom line: Cycline antibiotics may increase the risk of pseudotumor cerebri syndrome, but this association was not statistically significant in our models after accounting for confounding factors.
Luai Eldweik, MD, Collin McClelland, MD, Joshua D. Stein, MD, MS, Taylor S. Blachley, MS, and Michael S. Lee, MD
- 463** High accuracy of recognition of common forms of folliculitis by dermoscopy: An observational study
Bottom line: Dermoscopic evaluation of folliculitis has the potential to prevent unnecessary diagnostic tests and unnecessary treatments.
Murat Durdu, MD, Enzo Errichetti, MD, MSc, DVD, Ali Haydar Eskiocak, MD, and Macit Ilkit, MD
-  **472** Circulating bullous pemphigoid autoantibodies in the setting of negative direct immunofluorescence findings for bullous pemphigoid: A single-center retrospective review
Bottom line: Low-positive BP180 or BP230 autoantibody levels might be present in various clinical presentations and should not be overinterpreted as BP in the setting of a negative direct immunofluorescence.
Michael Wang, MD, Julia S. Lehman, MD, Michael J. Camilleri, MD, Lisa A. Drage, MD, and Carilyn N. Wieland, MD
- 480** Atrophic and hypertrophic photoaging: Clinical, histologic, and molecular features of 2 distinct phenotypes of photoaged skin
Bottom line: These clinically distinct forms of photoaging might benefit from specific types of therapies.
Dana L. Sachs, MD, James Varani, PhD, Heather Chubb, MS, Suzanne E. G. Fligiel, MD, Yilei Cui, PhD, Ken Calderone, BS, Yolanda Helfrich, MD, Gary J. Fisher, PhD, and John J. Voorhees, MD
- 489** Having a first-degree relative with melanoma increases lifetime risk of melanoma, squamous cell carcinoma, and basal cell carcinoma
Bottom line: This study revealed that the trunk and extremities are both melanoma-prone sites in individuals with a family history of melanoma; these individuals are also at an increased risk of basal cell carcinoma and extremity squamous cell carcinoma.
Erin X. Wei, MD, Xin Li, ScD, and Hongmei Nan, PhD
- 500** Anatomic location of primary melanoma: Survival differences and sun exposure
Bottom line: Results from this study identify independent associations of subsite with survival and indicate that previous models do not discriminate variability in survival and that more chronically sun-exposed sites have favorable survival.
Matthew D. Howard, MBBS (Hons), BPharm (Hons), Edmund Wee, MBBS (Hons), Rory Wolfe, PhD, Catriona A. McLean, FRCPA, MD, John W. Kelly, FACD, MD, and Yan Pan, MBBS, FACD

- 510** The blood proteomic signature of early-onset pediatric atopic dermatitis shows systemic inflammation and is distinct from adult long-standing disease

Bottom line: These data suggest the need for early intervention in pediatric atopic dermatitis to prevent disease chronicity and perhaps development of atopic comorbidities (“atopic march”) and other associated disorders.

Patrick M. Brunner, MD, MSc, Helen He, BS, Ana B. Pavel, PhD, Tali Czarnowicki, MD, MSc, Rachel Lefferdink, MD, Taylor Erickson, BS, Talia Canter, BS, Neha Puar, MD, Stephanie M. Rangel, PhD, Kunal Malik, MD, Yeriel Estrada, BS, James G. Krueger, MD, PhD, Emma Guttman-Yassky, MD, PhD, and Amy S. Paller, MS, MD

- 520** Reflectance confocal microscopy made easy: The 4 must-know key features for the diagnosis of melanoma and nonmelanoma skin cancers

Bottom line: Novice users of reflectance confocal microscopy may diagnose common skin cancers with reasonable sensitivity and specificity by recognizing a short list of quintessential confocal features. This simplified approach may facilitate dissemination of confocal among dermatologists.

Giovanni Pellacani, MD, Alon Scope, MD, Salvador Gonzalez, MD, PhD, Pascale Guitera, MD, PhD, Francesca Farnetani, MD, Josep Malvehy, MD, Alexander Witkowski, MD, PhD, Nathalie De Carvalho, MD, Omar Lupi, MD, and Caterina Longo, MD, PhD

- 527** Erosive pustular dermatosis of the scalp: A neutrophilic folliculitis within the spectrum of neutrophilic dermatoses

A clinicopathologic study of 30 cases

Bottom line: Biopsy specimens of intact follicular pustules reveal spongiotic vesiculopustules affecting the follicular infundibula. Erosive pustular dermatosis of the scalp is a neutrophilic superficial folliculitis, with some clinicopathologic similarities with other pathergic neutrophilic dermatoses, such as pyoderma gangrenosum.

Carlo Tomasini, MD, and Andrea Michelerio, MD

- 534** Hydroa vacciniforme—like lymphoproliferative disorder: Clinicopathologic study of 41 cases

Bottom line: Aggressive treatments, such as chemotherapy, should not be encouraged in the majority of Chinese patients.

Yanghe Liu, MD, Cuiling Ma, MD, PhD, Gang Wang, MD, PhD, and Lei Wang, MD, PhD

Dermatologic Surgery

- 541** Histopathologic upgrading of nonmelanoma skin cancer at the time of Mohs micrographic surgery: A prospective review

Bottom line: Tumors upgraded to more aggressive subtypes during Mohs surgery are larger and require more extensive surgery. Dermatologists should consider the limitations of superficial biopsies for detecting aggressive histology when weighing treatment options.

Rachel L. Kylo, MD, Karl W. Staser, MD, PhD, Ilana Rosman, MD, M. Laurin Council, MD, and Eva A. Hurst, MD

-
- Reviews**
- 548** Nodal staging of high-risk cutaneous squamous cell carcinoma
Bottom line: Nodal staging may be considered with high-risk cutaneous tumors without palpable lymphadenopathy via imaging and, if negative, sentinel lymph node biopsy.
Matthew Fox, MD, Marc Brown, MD, Nicholas Golda, MD, Dori Goldberg, MD, Christopher Miller, MD, Melissa Pugliano-Mauro, MD, Chrysalynne Schmults, MD, MSCE, Thuzar Shin, MD, PhD, Thomas Stasko, MD, Yaohui G. Xu, MD, PhD, and Kishwer Nehal, MD, from the High Risk Squamous Cell Carcinoma Workgroup, Dermatologic Surgery Section of the Association of Professors of Dermatology
- 558** Management of acute radiation dermatitis: A review of the literature and proposal for treatment algorithm
Bottom line: Washing with mild soap and water and use of topical corticosteroids and silver nylon dressings should be recommended by clinicians for the management of acute radiation dermatitis.
Amanda Rosenthal, BA, Rachel Israilevich, BS, and Ronald Moy, MD
- 568** Medical therapy for frontal fibrosing alopecia: A review and clinical approach
Bottom line: Intralesional steroids and 5 α -reductase inhibitors might be effective at preventing the progression of hair loss.
Anthony Ho, BA, and Jerry Shapiro, MD
- 581** Defining skin cancer local recurrence
Bottom line: We comprehensively review the literature to elucidate inconsistencies and propose a standardized algorithm for determination of tumor recurrence status across disciplines.
Ashley Wysong, MD, MS, Shauna Higgins, MD, Travis W. Blalock, MD, Dominic Ricci, MD, Rich Nichols, MD, Franki Lambert Smith, MD, and Iren Kossintseva, MD
-
- Research Letters**
- 600** Cutaneous post-transplantation lymphoma: Clinical features and survival outcomes
Seung Hwan Paik, MD, Young Jae Kim, MD, Chong Hyun Won, MD, PhD, Sung Eun Chang, MD, PhD, Mi Woo Lee, MD, PhD, Jee Ho Choi, MD, PhD, and Woo Jin Lee, MD, PhD
- 602** The association among thyroid dysfunction, thyroid autoimmunity, and clinical features of alopecia areata: A retrospective study
Sang-Myung Park, MD, Young-Jun Oh, MD, Bark-Lynn Lew, MD, PhD, and Woo-Young Sim, MD, PhD
- 605** Host characteristics and dynamics of *Staphylococcus aureus* colonization in patients with moderate-to-severe psoriasis before and after treatment: A prospective cohort study
Su-Hsun Liu, MD, PhD, Hsiao-Yun Yu, MS, Ya-Ching Chang, MD, Rosaline Chung-Yee Hui, MD, PhD, Yhu-Chering Huang, MD, PhD, and Yu-Huei Huang, MD, MS
- 607** Tattooing: A national survey in the general population of France
Nicolas Kluger, PhD, Laurent Misery, PhD, Sophie Seit , PhD, and Charles Taieb, MD

- 610** Nipple and areola lesions: Dermoscopy and reflectance confocal microscopy features
Elisa Cinotti, MD, PhD, Danila Galluccio, MD, Marco Ardigo', MD, PhD, Salvador Gonzalez, MD, PhD, Ausila Maria Manganoni, MD, Marina Venturini, MD, PhD, Paolo Broganelli, MD, Simone Ribero, MD, PhD, Francesca Farnetani, MD, Victor Desmond Mandel, MD, Giovanni Pellacani, MD, PhD, Linda Tognetti, MD, Francesco Lacarrubba, MD, PhD, Pascale Guitera, MD, PhD, Ignazio Stanganelli, MD, PhD, Iris Zalaudek, MD, PhD, Edith Johanna Arzberger, MD, Philippe Bahadoran, MD, PhD, Caterina Longo, MD, PhD, Giuseppe Spataro, MD, Jean-Luc Perrot, MD, PhD, and Pietro Rubegni, MD, PhD
- 613** A review of smartphone applications for promoting sun protection practices
Chelsea Moran, MA, and Ella Zetler, BAsC
- 615** Patient comfort and expectations for total body skin examinations: A cross-sectional study
Trissa McClatchey Connors, MD, MPH, Pooja Reddy, Emma Weiss, MD, Jaden Kohn, MD, MPH, Melinda Liu, MD, and Harry Dao, Jr, MD
- 617** Risk factors for acne development in the first 2 years after initiating masculinizing testosterone therapy among transgender men
Jason Andrew Park, SM, Erin Elizabeth Carter, MS, and Allison Ruth Larson, MD
- 618** Dermatologic care and sun protection practices need improvement in childhood cancer survivors
Edward Li, PhD, Patricia Todd, MD, Hannah Song, MD, and Jennifer T. Huang, MD
- 620** Assessment of biotin supplementation among patients in an outpatient dermatology clinic
Jason J. John, BS, Victoria Cooley, MS, and Shari R. Lipner, MD, PhD
- 621** Risk for neurodegenerative disorders in Korean patients with vitiligo: A nationwide retrospective cohort study
Tae-Gyun Kim, MD, PhD, Jaeyong Shin, MD, Shinwon Hwang, MD, Si-Hyung Lee, MD, PhD, Hyeon Chang Kim, MD, MPH, PhD, and Sang Ho Oh, MD, PhD
- 623** Rosacea: Relative risk versus absolute risk of malignant comorbidities
Leonardo A. Tjahjono, BS, Abigail Cline, MD, PhD, William W. Huang, MD, MPH, Alan B. Fleischer, Jr, MD, and Steven R. Feldman, MD, PhD
- 624** The association between obesity and hyperhidrosis: A nationwide, cross-sectional study of 2.77 million Israeli adolescents
Nadav Astman, MD, PhD, Itay Friedberg, MD, Jakob D. Wikstrom, MD, PhD, Estela Derazne, MSc, Orit Pinhas-Hamiel, MD, Arnon Afek, MD, MHA, Michal Freireich-Astman, MD, Aviv Barzilai, MD, Tarif Bader, MD, MHA, and Gilad Twig, MD, PhD

627 Impact of dermatology consultation on the care of children with chronic graft-versus-host disease of the skin

Anna Cristina Garza-Mayers, MD, PhD, L. Elizabeth Anderson, MD, JiaDe Yu, MD, and Jennifer T. Huang, MD



630 Rule of thumb: A simple tool to estimate 1% scalp surface area

Carlos Gustavo Wambier, MD, PhD, and Brett A. King, MD, PhD

631 Impact of the Patient Protection and Affordable Care Act on dermatologic health care utilization

Raghav Tripathi, MPH, Konrad D. Knusel, MS, Harib H. Ezaldeen, MD, Jeremy S. Bordeaux, MD, MPH, and Jeffrey F. Scott, MD

634 Effect of testosterone on chests and abdomens of transgender men

Catherine C. Motosko, BS, George A. Zakhem, BS, Miriam Keltz Pomeranz, MD, Rhonda Pomerantz, MD, Pierre B. Saadeh, MD, M. David Gothard, MS, and Alexes Hazen, MD

636 A comparison of the efficacy of ablative fractional laser–assisted photodynamic therapy according to ablative depth for actinic keratosis: A single-blinded, randomized, comparative, prospective study

Jeong-Wan Seo, MD, Ho-Jin Kim, MD, and Ki-Hoon Song, PhD

638 Pityriasis rubra pilaris: A study evaluating patient quality of life in 2 populations

A. Brooke Eastham, MD, Elizabeth Y. Tkachenko, BS, Alisa N. Femia, MD, Lisa K. Pappas-Taffer, MD, Misha Rosenbach, MD, Cara J. Joyce, PhD, Stephanie Liu, MD, and Ruth Ann Vleugels, MD, MPH

640 Factors associated with the utilization of Mohs micrographic surgery in the treatment of microcystic adnexal carcinoma

Walter Liszewski, MD, Derek R. Blanchette, MS, and Ian A. Maher, MD

642 Association of acetaminophen exposure with increased risk of eczema in children: A meta-analysis

Dongqiong Xiao, MD, Xiaojuan Su, PhD, Yi Qu, PhD, Yan Zhou, MD, Lingli Pan, MD, Xihong Li, MD, and Dezhi Mu, MD, PhD

644 Low-dose naltrexone therapy in benign chronic pemphigus (Hailey-Hailey disease): A case series

Constanza Riquelme-Mc Loughlin, MD, José Riera-Monroig, MD, Daniel Morgado-Carrasco, MD, Priscila Giavedoni, MD, Sebastian Podlipnik, MD, Pilar Iranzo, MD, and José M. Mascaró, Jr, MD

646 A pilot study of 2% tofacitinib cream with narrowband ultraviolet B for the treatment of facial vitiligo

Jacqueline McKesey, MD, and Amit G. Pandya, MD

648 Effectiveness and safety of low-dose oral minoxidil in male androgenetic alopecia

Juan Jimenez-Cauhe, MD, David Saceda-Corralo, MD, PhD, Rita Rodrigues-Barata, MD, Angela Hermosa-Gelbard, MD, Oscar M. Moreno-Arrones, MD, Diego Fernandez-Nieto, MD, and Sergio Vaño-Galvan, MD, PhD

- Commentary** **650** Highlights and implications of the 2019 proposed rule on sunscreens by the US Food and Drug Administration
Steven Q. Wang, MD, and Henry W. Lim, MD
- 652** Thirty years (and more) of hair research societies
Gillian E. Westgate, PhD, Wilma F. Bergfeld, MD, Victoria Ceh, MPA, Maria Hordinsky, MD, Rodney Sinclair, MBBS, MD, FACD, Bianca Maria Piraccini, MD, Hoon Kang, MD, PhD, and Manabu Ohyama, MD, PhD
- 654** Aligning research core outcome set development with clinical care performance measurement
Oliver J. Wisco, DO, Paul Imbriano, BA, MS, Matthew Zullo, BS, and David Adelson, MD

- Correction** **656** Lipner SR, Scher RK. Onychomycosis: Treatment and prevention of recurrence. *J Am Acad Dermatol.* 2019;80(4):853-67. Exam: 868

JAAD ONLINE

Available at <http://www.jaad.org>

- Clinical Pearls** **e23** Multiple lesions of erythema migrans in a “Christmas tree” pattern as a presenting sign of Lyme disease
Joseph C. Pierson, MD, Andrew R. Tegeder, MD, and Kendra Lesiak, MD
- e25** Postexposure prevention of *Toxicodendron* dermatitis by early forceful unidirectional washing with liquid dishwashing soap
Brett C. Neill, MD, John A. Neill, BA, Jim Brauker, PhD, Anand Rajpara, MD, and Daniel J. Aires, MD, JD
- e27** Telementoring and smartphone-based answering systems to optimize dermatology resident dermoscopy education
Kelly C. Nelson, MD, Jeffrey E. Gershenwald, MD, Stephanie A. Savory, MD, Janice M. Wilson, MD, Julie M. Simon, MHA, Elizabeth M. Burton, MBA, and Lauren E. Haydu, PhD
- e29** Scraping test for the diagnosis of acral subcorneal hemorrhage
Francesco Savoia, MD, Giulia Maria Ravaioli, MD, Michela Tabanelli, MD, Emi Dika, MD, and Annalisa Patrizi, MD
- Surgical Pearls** **e31** A noninvasive method for treating ingrown nail recurrence due to incomplete matricectomy
June-Ho Won, MD, Ji-Sun Chun, MD, and Young-Ho Won, MD, PhD
- e33** Treatment of oral mucosal neuromas with carbon dioxide laser
Emily A. Weig, Gretchen M. Roth, MD, Karolyn A. Wanat, MD, and Nkanyenzi N. Ferguson, MD
-  **e35** Use of disposable blade for harvesting epidermal skin graft
Rajsmita Bhattacharjee, MD, DNB, Keshavamurthy Vinay, MD, DNB, MNAMS, and Davinder Parsad, MD

| | | |
|-----------------------------|--|--|
| | e37 | Achieving eversion utilizing topical tissue adhesive <i>Marjon Vatanchi, MD, and Shang I. Brian Jiang, MD</i> |
| | e39 | Galeatomy: A useful technique aiding high-tension scalp closures <i>Marjon Vatanchi, MD, and Roy C. Grekin, MD</i> |
| Technology Pearl |  e41 | Live, remote control of an in vivo reflectance confocal microscope for diagnosis of basal cell carcinoma at the bedside of a patient 2500 miles away: A novel tele-reflectance confocal microscope approach <i>Gennady Rubinstein, MD, Jason Garfinkel, MS, and Manu Jain, MD</i> |
| Notes & Comments | e43 | Androgenetic alopecia and microneedling: Every needling is not microneedling <i>Abhijeet Kumar Jha, MD, and Keshavamurthy Vinay, MD, DNB</i> |
| | e45 | Comment on “Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics” <i>Andrew Greenspan, MD, and Sharon Popik, MD</i> |
| | e47 | Comment on “Incidence of pneumocystosis among patients exposed to immunosuppression” <i>Eric H. Kowalski, BS, Diana Kneiber, BS, Khalaf Kridin, MD, PhD, and Kyle T. Amber, MD</i> |
| | e49 | Reply to: “Comment on ‘Incidence of pneumocystosis among patients exposed to immunosuppression’” <i>Sergey Rekhman, MD, PharmD, MPH, Andrew Strunk, MA, and Amit Garg, MD</i> |
| | e51 | Frontal fibrosing alopecia, just lichen planopilaris? <i>Steven Kossard, MBBS, FACD, PhD</i> |
| | e53 | Frontal fibrosing alopecia should be renamed to lichen planopilaris of Kossard <i>Christos Tziotzios, MA, MB BChir (Cantab), MRCP(UK), FHEA, Catherine M. Stefanato, MD, FRCPath, John A. McGrath, MD, FRCP, FMedSci, and David A. Fenton, MBE, FRCP</i> |
| Iotaderma | e55 | Iotaderma (#306) <i>Robert I. Rudolph, MD, FACP</i> |

To access the current version of the Information for Authors, Transfer of Copyright Form, Authorship Declaration Form, Conflict of Interest Disclosure, and Authors’ Rights Statement, as well as many other pertinent documents, please visit our website at www.jaad.org/content/auth_page.

Full text access to *Journal of the American Academy of Dermatology Online* is now available for Academy members and all print subscribers. To activate your online subscription, please visit *Journal of the American Academy of Dermatology Online* by pointing your browser to <http://www.jaad.org>.

Icon Key

| | | |
|---|---|---|
|  | = | CME article |
|  | = | Video content available at http://www.jaad.org |
|  | = | Podcast available at http://www.aad.org/olc |
|  | = | MOC credits available |

Early online publication: Articles in press

The *Journal of the American Academy of Dermatology* posts in-press articles online before they appear in the print edition of the *Journal*. These Articles in Press are available from both the JAAD website (www.jaad.org) and Science Direct (www.sciencedirect.com). To access Articles in Press, simply click the Articles in Press link.

JAAD Case Reports

The *Journal of the American Academy of Dermatology* no longer publishes Case Reports. These can now be found in our new open access journal, *JAAD Case Reports*, at <http://www.jaadcasereports.org>.