

Retinal nerve fiber layer thickness in children with primary congenital glaucoma measured by spectral domain optical coherence tomography



Lucía Perucho-González, MD, PhD,^{a,c} José María Martínez de la Casa, MD, PhD,^{a,b,c,d}
Federico Sáenz-Francés, MD, PhD,^{a,c} Laura Morales-Fernandez, MD, PhD,^{a,c}
Carmen Dora Méndez-Hernández, MD, PhD,^{a,b,c,d} Rubén Sánchez-Jean,^{b,c,d} and
Julián García-Feijóo, MD, PhD^{a,b,c,d}

PURPOSE	To evaluate retinal nerve fiber layer (RNFL) thickness using spectral domain optical coherence tomography (SD-OCT) in a population of children diagnosed with primary congenital glaucoma (PCG).
METHODS	In this cross-sectional study, 59 eyes of 59 children diagnosed with PCG and 87 eyes of 87 healthy children were evaluated by SD-OCT to measure the RNFL. The global average peripapillary RNFL thickness and sectional RNFL thickness were evaluated in both groups. Differences in global average and sectional thickness were analyzed.
RESULTS	Mean age in the PCG group was 9.61 ± 3.23 years; in the control group, 8.47 ± 2.99 years ($P = 0.0516$). There were statistically significant differences ($P < 0.007$) in all sectors between both groups.
CONCLUSIONS	SD-OCT is a promising tool for evaluating the eyes of children diagnosed with PCG. Future research should examine the test–retest variability of SD-OCT parameters and their ability to diagnose progression in these children. (J AAPOS 2019;23:94.e1-4)

Glaucoma in children is characterized by the presence of elevated intraocular pressure (IOP) and characteristic optic disk cupping. Currently the most accepted methods for establishing the glaucoma-related optic nerve changes in children are optic nerve head examination and stereo photography. Some studies have reported differences in the retinal nerve fiber layer (RNFL) thickness between normal and glaucomatous eyes in children evaluated with optical coherence tomography (OCT) using both time domain¹ and spectral domain (SD-OCT)² systems. The current study investigated the role of ST-OCT in diagnosing glaucoma in children.

Subjects and Methods

This observational cross-sectional study included healthy volunteers and patients diagnosed with primary congenital glaucoma (PCG) treated in the glaucoma department of San Carlos Clinical

Author affiliations: ^aOphthalmology Service, Institute of Sanitary Research of the San Carlos Clinical Hospital, Madrid, Spain; ^bOcular Pathology, National Net OFTARED, Institute of Health Carlos III, University Hospital, Madrid, Spain; ^cGlaucoma Department, Institution of Sanitary Research of the San Carlos Clinical Hospital, Madrid, Spain; ^dUniversidad Complutense, Madrid, Spain

Submitted August 17, 2016.

Revision accepted March 3, 2017.

Published online April 2, 2019.

Correspondence: Lucía Perucho-González, MD, PhD, Ocular Pathology National Net OFTARED of the Institute of Health Carlos III University Hospital, Profesor Martín Lagos Street, 28040 Madrid, Spain (email: luciaperuchogonz@gmail.com).

Copyright © 2019, American Association for Pediatric Ophthalmology and Strabismus. Published by Elsevier Inc. All rights reserved.

1091-8531/\$36.00

<https://doi.org/10.1016/j.jaapos.2017.03.023>

Hospital, Madrid, a tertiary eye care center. The study was approved by the San Carlos University Hospital Institutional Review Board and complied with the guidelines of the Declaration of Helsinki. Written informed consent was obtained from the legal guardians of minors before enrollment, with the assent of children >12 years of age.

Children with known PCG—IOP of >22 mm Hg at diagnosis together with clinical evidence of glaucoma, such as enlarged corneal diameter, Haab's striae, glaucomatous optic disk head appearance, and glaucomatous visual fields (when available) were enrolled. Only PCG patients with IOP of <21 mm Hg after either medical therapy or surgery performed at least 6 months earlier and patients able to fixate on a target were included. Patients with other childhood glaucoma, presence of corneal edema precluding optical coherence tomography, nystagmus, uveitis, any concomitant retinal disease, or significant media opacity, such as cataract or pseudophakia, were excluded.

A total of 87 healthy volunteers were recruited from children referred for refraction who had normal examinations, children accompanying adult patients, and children of staff members who agreed to participate in the study. They had a normal eye examination, including a normal optic disk, IOP of <21 mm Hg, and no history of ocular surgery or trauma. Only one eye was randomly chosen in each participant unless only one eye met inclusion criteria (www.randomization.com).

All subjects underwent full ophthalmic examination, including dilated fundus examination, in one session. IOP was measured by Perkins handheld applanation tonometer (Haag-Streit, Koniz, Switzerland) when possible. Icare rebound tonometry (TA01I; Icare, Finland) was used if corneal irregularities were present or

when cooperation was poor. All OCT scans were obtained by the same trained ophthalmologist.

SD-OCT was used to obtain the RNFL (Spectralis; Heidelberg Engineering GmbH, Heidelberg, Germany; software version 5.4) thickness maps. A full RNFL circle scan containing 768 A-scans along a peripapillary circle of 360° was performed. The Spectralis software divides this circle into six regions: temporal quadrant thickness (90° sector), superotemporal quadrant thickness (45° sector), superonasal quadrant thickness (45° sector), nasal quadrant thickness (90° sector), inferonasal quadrant thickness (45° sector), inferotemporal quadrant thickness (45° sector), and average thickness (average of all 6 sectors or 360°), provided as a number in the center.³

Scans were included for analysis if the quality score was >20 dB. Three images were automatically obtained for each eye during the same session. If none of the three images met the quality criteria the eye was excluded from the analysis.

Statistical Analysis

All statistical tests were performed using the SPSS package version 18.0 for Windows (SPSS Inc, Chicago, IL). The Kolmogorov-Smirnov test was used to check for a normal distribution of quantitative data. Quantitative variables were expressed as their corresponding means and standard deviations. Medians and interquartile ranges were used to describe variables showing a non-normal distribution. Correlation between different treatments in PCG and RNFL thickness was assessed using the Pearson correlation coefficient. Bonferroni correction of the *P* value was performed for multiple comparisons and the level of significance was set at *P* value of <0.007. The diagnostic capacity of each variable to differentiate between normal and glaucoma eyes was determined by calculating the area under the receiver operating characteristics (ROC) curve (AUC). Differences between the ROC curves were tested to compare AUCs using the Hanley-McNeil method.⁴

Results

A total of 71 eyes of 71 children diagnosed with PCG and 98 eyes of 98 healthy children were eligible for inclusion. In the PCG group, 5 eyes with poor fixation, 4 eyes with low signal strength index on OCT, and 3 eyes with corneal opacities were excluded from the final analysis. In the control group, 6 eyes with poor fixation and 5 eyes with low signal strength index were excluded. For statistical analysis, 59 eyes of 59 children formed the PCG group; 87 eyes of 87 healthy children, the control group.

In the PCG group, mean age was 9.61 ± 3.23 years (range, 3-18); in the control group, 8.47 ± 2.99 years (range, 6-18; *P* = 0.0516). There were no significant differences in terms of sex distribution between groups. In the PCG group, the average age of diagnosis was 23.24 ± 9.28 months. Control eyes showed a significantly higher quality score (29.79 dB) compared with PCG eyes (25.63 dB; *P* < 0.0001). Clinical measurements of cup:disk ratio were 0.4 ± 0.1 in the control group and 0.6 ± 0.2 in PCG patients. Peripapillary RNFL measurements ob-

tained in both groups are summarized in Table 1. There were statistically significant differences (*P* < 0.007) in all sectors between groups, except for the nasal (*P* = 0.015) and inferonasal (*P* = 0.009) sectors. Mean IOP was 19.11 ± 4.23 mm Hg in the PCG group and 13.95 ± 1.98 mm Hg (*P* < 0.001) in the control group.

AUCs for the mean peripapillary RNFL and sectional RNFL thickness measurements are shown in Table 2 and Figure 1. The best AUC parameters were global or average RNFL (0.831 μ m), temporal superior RNFL (0.811 μ m), and nasal superior RNFL (0.770 μ m).

The likelihood ratio for a positive test was 6, meaning that a positive test is 6 times more likely to come from a PCG patient rather than control. The positive predictive value was 0.8, and the negative predictive value was 0.6.

Discussion

Expert clinical examination of the optic nerve remains the gold standard in diagnosing glaucomatous optic nerve damage in children as well as in adults. Morales and Brown⁵ demonstrated that ultra-short visual field programs that can be completed in <3 minutes offer the possibility of testing children successfully. However, they reported that, at <8 years of age, variability in individual maturity, concentration, and effort result in cause greater variability in accuracy.

SD-OCT can provide clinicians with objective measurements of RNFL thickness in both adult and pediatric patients. OCT scans have the advantages of being noncontact, rapid, and relatively patient friendly, making them ideally suited for testing children. Our study was a preliminary study to measure the RNFL thicknesses in children with PCG using SD-OCT.

The diagnostic accuracy of SD-OCT RNFL parameters for glaucoma in adults has been investigated extensively.⁶ In general, the parameters with the best diagnostic accuracy are the average peripapillary RNFL thickness and thicknesses in the inferior and superior quadrants.⁷⁻¹² Our results in PCG agree with those reported in glaucoma in adults. We found that the variables featuring the larger AUCs were global or average (0.831 μ m), temporal superior (0.811 μ m), nasal superior (0.770 microns), temporal inferior (0.753 μ m), and nasal inferior (0.748 μ m) RNFL thicknesses.

A well-known phenomenon to take into account is the reversal of optic disk cupping following IOP reduction in congenital and juvenile glaucoma, especially before 1 year of age.¹³⁻¹⁹ This change in optic disk cupping has also been recorded in adults by measuring RNFL thickness,¹⁸ but the results of these studies seem to be conflicting. For this reason, we decided to exclude PCG patients who had undergone surgery or experienced uncontrolled IOP in the previous 6 months.

Yanni and colleagues¹⁹ reported that the normal mean peripapillary RNFL thickness measured by SD-OCT in 83 healthy children 5-15 years of age was thicker (107.6 ± 1.2 μ m) than mean RNFL thickness in adults.

Table 1. Mean values of sectorial RNFL thickness in control group and PCG group

OCT parameter thickness, μm	Control group, mean \pm SD	PCG group, mean \pm SD	<i>P</i> value ^a
RNFL global	102.20 \pm 10.54	76.83 \pm 23.43	0.000
RNFL temporal	72.70 \pm 12.17	61.66 \pm 19.45	0.001
RNFL temporal superior	136.83 \pm 19.49	97.85 \pm 35.01	0.000
RNFL temporal inferior	148.09 \pm 23.91	114.25 \pm 45.65	0.000
RNFL nasal	77.43 \pm 18.72	57.63 \pm 23.37	0.015
RNFL nasal superior	111.29 \pm 24.14	76.73 \pm 32.43	0.005
RNFL nasal inferior	121.21 \pm 27.63	87.34 \pm 33.99	0.009

OCT, optical coherence tomography; PCG, primary congenital glaucoma; RNFL, retinal nerve fiber layer; SD, standard deviation.

^aCorrected significant difference ($P < 0.007$); *t* test (Bonferroni post hoc).

Table 2. Results of diagnostic capacity of each variable determined by calculating area under curve (AUC)

OCT parameter thickness, μm	AUC
RNFL global	0.831
RNFL temporal	0.666
RNFL temporal superior	0.811
RNFL temporal inferior	0.753
RNFL nasal	0.721
RNFL nasal superior	0.770
RNFL nasal inferior	0.748

OCT, optical coherence tomography; RNFL, retinal nerve fiber layer.

In central China a school-based cohort study designed to longitudinally observe the development of myopia and other ocular disease among children provided the baseline results in RNFL thickness in a cohort of 7-year-olds²⁰ as well as the RNFL thickness association with refractive error, axial length, and optic disk parameters.²¹

Another recent case-control study of 45 eyes of 37 children who underwent surgery for PCG and 72 eyes of 41 normal children, conducted at a tertiary eye care center in South India, found significant differences in all SD-OCT parameters between both groups.²² Our study found that all SD-OCT RNFL thickness measurements were significantly different ($P < 0.05$) in PCG eyes compared with normal healthy controls, suggesting that SD-OCT may be a valuable clinical tool to help monitor glaucoma in the pediatric population. The present study has several limitations. RNFL thickness parameters of the most severe cases are very difficult to obtain, and this could result in a selection bias. An important subgroup of patients of PCG associated with mutations in the *CYP11B1* could have been excluded; these patients have more severe PCG, an earlier age of onset, greater corneal alterations, and comparatively more surgical procedures.^{23,24} Our PCG cohort probably consisted of mild-to-moderate glaucoma cases. The average age at diagnosis in the PCG group was almost 2 years (23.24 ± 9.28 months); it is possible that most severe cases, which tend to be diagnosed at younger ages, were excluded,




FIG 1. Receiver operating characteristic curves for the variables showing the best discriminant capacity between the two subject groups. AUC: 0.831, RNFL global (G); 0.666, RNFL temporal (T); 0.811, RNFL superotemporal (TS); 0.753, RNFL inferotemporal (TI); 0.721, RNFL nasal (N); 0.770, RNFL superonasal (NS); 0.748, RNFL inferonasal (NI).

resulting in an underestimation of the difference in RNFL thickness between normal and PCG eyes. Also, it is known that axial length affects RNFL measurement in the direction of thinner RNFL and that glaucoma in infancy is associated with ocular enlargement (buphtalmos), corneal enlargement (usually before three years of age), increased measurements in axial length, and progressive myopia. Thus, we should consider that glaucomatous eyes included probably presented with higher axial length than eyes from control subjects. Future studies should obtain a random distribution of myopic and hyperopic subjects in both normal and glaucomatous subgroups.

Obtaining images of adequate quality using SD-OCT in children can be challenging—more so, perhaps, in children with PCG, who often have low visual acuity or refractive errors. For this reason, in the present study the cut-off of signal-to-noise ratio was fixed at 20 dB; for adults this value tends to be fixed at 40 dB. We found lower quality scores in patients with PCG than in controls, which may affect accuracy of measurement. Future research should also investigate the test-retest variability of SD-OCT parameters and their ability to diagnose progression in these children.

References

- Hess DB, Asrani SG, Bhide MG, Enyedi LB, Stinnett SS, Freedman SF. Macular and retinal nerve fiber layer analysis of normal and glaucomatous eyes in children using optical coherence tomography. *Am J Ophthalmol* 2005;139:509-17.
- Ghasia FF, El-Dairi M, Freedman SF, Rajani A, Asrani S. Reproducibility of spectral-domain optical coherence tomography measurements in adult and pediatric glaucoma. *J Glaucoma* 2015;24:55-63.

3. Bendschneider D, Tornow RP, Horn FK, et al. Retinal nerve fiber layer thickness in normals measured by spectral domain OCT. *J Glaucoma* 2010;19:475-82.
4. Hanley JA, McNeil BJ. A method of comparing the areas under receiver operating characteristic curves derived from the same cases. *Radiology* 1983;148:839-43.
5. Morales J, Brown SM. The feasibility of short automated static perimetry in children. *Ophthalmology* 2001;108:157-62.
6. Gracitelli CP, Abe RY, Medeiros FA. Spectral-domain optical coherence tomography for glaucoma diagnosis. *Open Ophthalmol J* 2015; 9:68-77.
7. Park SB, Sung KR, Kang SY, Kim KR, Kook MS. Comparison of glaucoma diagnostic capabilities of cirrus HD and stratus optical coherence tomography. *Arch Ophthalmol* 2009;127:1603-9.
8. Kim JS, Ishikawa H, Gabriele ML, et al. Retinal nerve fiber layer thickness measurement comparability between time domain optical coherence tomography (OCT) and spectral domain OCT. *Invest Ophthalmol Vis Sci* 2010;51:896-902.
9. Rao HL, Zangwill LM, Weinreb RN, Sample PA, Alencar LM, Medeiros FA. Comparison of different spectral domain optical coherence tomography scanning areas for glaucoma diagnosis. *Ophthalmology* 2010;117:1692-9.
10. Leite MT, Rao HL, Zangwill LM, Weinreb RN, Medeiros FA. Comparison of the diagnostic accuracies of the Spectralis, Cirrus, and RTVue optical coherence tomography devices in glaucoma. *Ophthalmology* 2011;118:1334-9.
11. Leung CK, Lam S, Weinreb RN, et al. Retinal nerve fiber layer imaging with spectral-domain optical coherence tomography. *Ophthalmology* 2010;117:1684-91.
12. Wang X, Li S, Fu J, et al. Comparative study of retinal nerve fibre layer measurement by RTVue OCT and GDx VCC. *Br J Ophthalmol* 2011;95:509-13.
13. Meirelles SH, Mathias CR, Bloise RR, et al. Evaluation of the factors associated with the reversal of the disc cupping after surgical treatment of childhood glaucoma. *J Glaucoma* 2008;17:470-73.
14. Mandal AK, Netland PA. *The Pediatric Glaucomas*. Philadelphia: Elsevier Inc; 2006:27-39.
15. Quigley HA. The pathogenesis of reversible cupping in congenital glaucoma. *Am J Ophthalmol* 1977;84:358-70.
16. Quigley HA. Childhood glaucoma: results with trabeculectomy and study of reversible cupping. *Ophthalmology* 1982;89:219-26.
17. Ely AL, El- Dairi MA, Freedman SF. Cupping reversal in pediatric glaucoma-evaluation of the retinal nerve fiber layer and visual field. *Am J Ophthalmol* 2014;158:905-15.
18. Raghu N, Pandav SS, Kaushik S, Ichhpujani P, Gupta A. Effect of trabeculectomy on RNFL thickness and optic disc parameters using optical coherence tomography. *Eye* 2012;26:1131-7.
19. Yanni SE, Wang J, Cheng CS, et al. Normative reference ranges for retinal nerve fiber layer, macula, and retinal layer thicknesses in children. *Am J Ophthalmol* 2013;155:354-60.
20. Zhu BD, Li SM, Li H, et al. Anyang Childhood Eye Study Group. Retinal nerve fiber layer thickness in a population of 12-year-old children in central China measured by iVue-100 spectral-domain optical coherence tomography: the Anyang Childhood Eye Study. *Invest Ophthalmol Vis Sci* 2013;54:8104-11.
21. Kang MT, Li SM, Li H, et al. Peripapillary retinal nerve fiber layer thickness and its association with refractive error in Chinese children: the Anyang Childhood Eye Study. *Clin Experiment Ophthalmol* 2016;44:701-9.
22. Srinivasan S, Addepalli UK, Rao HL, Garudadri CS, Mandal AK. Spectral domain optical coherence tomography in children operated for primary congenital glaucoma. *Br J Ophthalmol* 2014;98: 162-5.
23. Zenteno JC, Hernandez-Merino E, Mejia-Lopez H, et al. Contribution of *CYP11B1* mutations and founder effect to primary congenital glaucoma in Mexico. *J Glaucoma* 2008;17:189-92.
24. Morales- Fernandez L, Martinez-de-la-Casa JM, Garcia-Feijoo J, et al. Clinical variability of primary congenital glaucoma in a Spanish family with *Cyp11b1* gene mutations. *J Glaucoma* 2015;24: 630-34.