

ASSOCIATION BETWEEN CONGENITAL NASOLACRIMAL DUCT OBSTRUCTION AND MODE OF DELIVERY AT BIRTH

To the Editor: I read with interest the recent report by Tavakoli and colleagues¹ that concludes that cesarean section is an independent risk factor for the development of congenital nasolacrimal duct obstruction. Although statistical manipulations can produce *P* value significance and twinkling eyelids, the study's significant biases cannot support the authors' conclusions.

The authors report on 104 children with congenital nasolacrimal duct obstruction (CNLDO) from a population in which 48% of pregnancies are delivered by cesarean section. In such a setting, the finding that 61% of 104 CNLDO patients had a C-section (compared with 39% vaginal deliveries) may be due to chance alone, given the small number of patients. Moreover, the study suffers from ascertainment bias (tertiary referral center–observed patients) and the exclusion of those without a birth history (more likely to favor those having a vaginal delivery). The low spontaneous resolution rate (52%) and high percentage of bilateral disease (47.5%) in these cases illustrate the significant biases of this cohort.

As the authors point out, other studies have reported conflicting findings regarding the association between cesarean section and CNLDO, with some investigators² finding an association and others not.^{3–5} However, these studies are also relatively small in size, ranging from 40³ to 386⁵ cases for those that found no association and 107 infants in the sole prior study that reported an association.² In a recently published report of 1,998 children diagnosed with CNLDO by primary care providers in a medically isolated region, 404 (20.2%) were delivered via C-section compared with 19.2% for the general population (*P* = 0.26).⁶ Although this population-based study of 1998 afflicted children indicates that cesarean birth is not associated with the development of CNLDO, we may have to wait for big data to provide the final answer. From the cohort assembled by Tavakoli and colleagues, few if any risk factors can be named, certainly not cesarean birth, as an independent risk factor for congenital nasolacrimal duct obstruction.

Brian G. Mohny, MD
Department of Ophthalmology
Mayo Clinic
Rochester, Minnesota

References

1. Tavakoli M, Osigian CJ, Saksiriwutto P, et al. Association between congenital nasolacrimal duct obstruction and mode of delivery at birth. *JAAPOS* 2018;22:381-5.

2. Kuhli-Hattenbach C, Luchtenberg M, Hofmann C, Kohnen T. Increased prevalence of congenital dacryostenosis following cesarean section. *Ophthalmologie* 2016;113:675-83.
3. Yildiz A. Congenital nasolacrimal duct obstruction: cesarean section vs. vaginal delivery. *Med Glas (Zenica)* 2018;15:164-7.
4. Palo M, Gupta S, Naik MN, Ali MJ. Congenital nasolacrimal duct obstruction and its association with the mode of birth. *J Pediatr Ophthalmol Strabismus* 2018;55:266-8.
5. Spaniol K, Stupp T, Melcher C, Beheiri N, Eter N, Prokosch V. Association between congenital nasolacrimal duct obstruction and delivery by cesarean section. *Am J Perinatol* 2015;32:271-6.
6. Sathiamoorthi S, Frank RD, Mohny BG. Incidence and clinical characteristics of congenital nasolacrimal duct obstruction. *Br J Ophthalmol* 2018; <https://doi.org/10.1136/bjophthalmol-2018-312074>. pii: bjophthalmol-2018-312074.

<https://doi.org/10.1016/j.jaaapos.2019.01.002>

J AAPOS 2019;23:125.

Copyright © 2019, American Association for Pediatric Ophthalmology and Strabismus. Published by Elsevier Inc. All rights reserved.
1091-8531/\$36.00

REPLY

We thank Dr. Mohny for his interest in our publication and scholarly commentary. The following provides our response to his concerns. Given the differences between the demographics of our patients and those in other geographic locations, it is possible for something to be a risk factor for a condition in our population and not in another. For instance, there may be no association between cesarean section (CS) and congenital nasolacrimal duct obstruction (CNLDO) when the population rate of CS is 19.2% (in Olmsted County) and, at the same time, there may be an association between CS and CNLDO when the population rate of CS is 48% (Miami-Dade County). Similarly, there may be differences in race and ethnicity. In their recent study of residents of Olmsted County, Dr. Mohny's group concluded that the "Caucasian" race is associated with development of CNLDO.¹ The population census data in 2010 for this area was 86% white, compared to 2018 US census data of 60.7% white (not including Latino).^{2,3} Thus, if the same study was performed in a different population, a different result might be possible.

In our paper we discussed the theory that the pressure produced during vaginal delivery can potentially affect the opening of Hasner's membrane. We excluded patients without any known birth history; however, we suspect that this did not contribute largely to bias, because only 6 of 126 patients (4%) did not have a documented birth history. Also, in our study, we excluded babies who had an emergent CS delivery following a failed normal vaginal delivery