

September 2019, Volume 81, Number 3

CONTENTS

Continuing Medical Education

CME 657

The etiology, diagnosis, and management of hyperhidrosis:
A comprehensive review

Etiology and clinical work-up
Shiri Nawrocki, BA, and Jisun Cha, MD

667 CME examination

668 Answers to CME examination (Identification No. JA0919),
September 2019 issue of the *Journal of the American Academy of
Dermatology*

CME 669

The etiology, diagnosis, and management of hyperhidrosis:
A comprehensive review

Therapeutic options
Shiri Nawrocki, BA, and Jisun Cha, MD

681 CME examination

682 Answers to CME examination (Identification No. JB0919),
September 2019 issue of the *Journal of the American Academy of
Dermatology*

Letter from the Editor

683

Correcting the literature
Dirk M. Elston, MD

A Clinician's Perspective

684

Predicting sepsis in Stevens-Johnson syndrome—toxic epidermal
necrolysis: Should dermatologists be pro-procalcitonin?

Warren R. Heymann, MD

- Original Articles** **686** Risk factors and diagnostic markers of bacteremia in Stevens-Johnson syndrome and toxic epidermal necrolysis: A cohort study of 176 patients
- Bottom line:** The Bacteremia Risk Score, which assesses the presence of hemoglobin ≤ 10 g/dL, cardiovascular disease, and $\geq 10\%$ body surface area involvement on admission, is useful in predicting the development of bacteremia. Hypothermia and procalcitonin ≥ 1 $\mu\text{g/L}$ help in the timely detection of bacteremia during hospitalization.
- Hui Kai Koh, MBBS, Zi Teng Chai, MD, MRCP, Hui Wen Tay, BS, Stephanie Fook-Chong, MS, CStat, Karen J. L. Choo, MBChB, MRCP, Choon Chiat Oh, MBBS, MRCP, FAMS (Dermatology), Yi Wei Yeo, MBBS, MRCP, MMed (Int Med), FAMS (Dermatology), Hong Yi Koh, MBBS, MRCP, MMed (Int Med), FAMS (Dermatology), Shiu Ming Pang, MBBS, FRCP, Dip Dermatology, FAMS (Dermatology), and Haur Yueh Lee, MBBS, MRCP, MMed (Int Med), FAMS (Dermatology)
- 694** Cyclosporine for moderate-to-severe alopecia areata: A double-blind, randomized, placebo-controlled clinical trial of efficacy and safety
- Bottom line:** These results will guide clinicians in their choice of second-line agent for patients with alopecia areata who are steroid responsive but steroid dependent.
- Vivien Wai Yun Lai, MBBS (Hons), BMedSc (Hons), Gang Chen, PhD, MMSc, BMed, Douglas Gin, MBBS, FACD, and Rodney Sinclair, MBBS, MD, FACD
- 702** Determining the optimal dose of infliximab for treatment of hidradenitis suppurativa
- Bottom line:** For patients with hidradenitis suppurativa beginning treatment with infliximab, 10 mg/kg every 6 or 8 weeks may be a reasonable initial dosing regimen.
- Ashley N. Oskardmay, BS, J. Alex Miles, BA, MPH, and Christopher J. Sayed, MD
- 709** Lack of association of biologic therapy for psoriasis with psychiatric illness: An electronic medical records cohort study
- Bottom line:** The prescription of biologic therapies to treat psoriasis does not appear to be associated with psychiatric illnesses. The lack of an association is likely due to the prescribing behaviors of physicians (selection bias), who appear to be mediating these potential adverse psychiatric events by carefully selecting the patients receiving the treatment.
- David J. Margolis, MD, PhD, Daniel Shin, PhD, Megan H. Noe, MD, MPH, MSCE, Marilyn T. Wan, MBChB, MPH, Ketaki Bhate, MBBS, MSc, Shiyu Wang, MS, Ole Hoffstad, MS, Junko Takeshita, MD, PhD, MSCE, and Joel M. Gelfand, MD, MSCE
- 717** Modified shave surgery combined with nail window technique for the treatment of longitudinal melanonychia: Evaluation of the method on a series of 67 cases
- Bottom line:** To our knowledge, this is the largest case series to evaluate the postoperative outcome of shave surgery.
- Yuan Zhou, MD, Wei Chen, MD, Zhen-ru Liu, MD, Juan Liu, MD, Fei-ran Huang, MD, and Da-guang Wang, MD

- 723** Acne and obesity: A nationwide study of 600,404 adolescents
Bottom line: In youths, overweight and obesity are inversely associated with acne in a dose-dependent manner.
Igor Snast, MD, Adam Dalal, MD, Gilad Twig, MD, PhD, Nadav Astman, MD, PhD, Ron Kedem, PhD, Dan Levin, MD, Yifat Erlich, MD, Yael Anne Leshem, MD, MCR, Moshe Lapidot, MD, Emmilia Hodak, MD, and Assi Levi, MD
- 730** Cutaneous nontuberculous mycobacteria infections: A retrospective case series of 78 patients from the Texas Gulf Coast region
Bottom line: The diagnosis should be considered in treatment-resistant lesions, particularly with immunosuppression, trauma, invasive procedure, or environmental exposure. Simultaneous skin biopsy specimen and tissue culture can increase diagnostic yield. Multiple antibiotics combined with surgery may improve treatment success.
Rebecca C. Philips, MD, Paige E. Hoyer, MD, Skyler M. White, MD, Katherine T. Tinkey, MD, Michael Loeffelholz, PhD, Clark R. Andersen, MS, Michael G. Wilkerson, MD, Bernard R. Gibson, MD, and Brent C. Kelly, MD
- 740** Dermatology-specific and all-cause 30-day and calendar-year readmissions and costs for dermatologic diseases from 2010 to 2014
Bottom line: Identification of diseases with a high risk of readmissions should guide further efforts to reduce readmissions and costs.
Myron Zhang, MD, Alina Markova, MD, Joanna Harp, MD, Stephen Dusza, DrPH, Misha Rosenbach, MD, and Benjamin H. Kaffenberger, MD
- 749** Distinguishing Stevens-Johnson syndrome/toxic epidermal necrolysis from clinical mimickers during inpatient dermatologic consultation—A retrospective chart review
Bottom line: Nikolsky sign, atypical targets, fever, and lymphopenia aid in predicting the probability of true Stevens-Johnsons syndrome/toxic epidermal necrolysis.
Allison Weinkle, MD, Cory Pettit, BS, Aditi Jani, BS, Jesse Keller, MD, Yuanyuan Lu, MS, Stephen Malachowski, MD, MS, John C. Trinidad, MD, Benjamin H. Kaffenberger, MD, Elizabeth N. Ergen, MD, Lauren C. Hughey, MD, David Smith, MD, and Lucia Seminario-Vidal, MD, PhD
- 758** Cost analysis of a store-and-forward teledermatology consult system in Philadelphia
Bottom line: With engagement of stakeholders and proper planning, patients can obtain improved access to high-quality and sustainable dermatologic care in the setting of teledermatology reimbursement.
Xiaoshi Yang, PhD, John S. Barbieri, MD, MBA, and Carrie L. Kovarik, MD

- 765** Evaluating the cost-effectiveness of teledermatology
John S. Barbieri, MD, MBA, Xiaoshi Yang, PhD, and Carrie L. Kovarik, MD

Dermatologic Surgery

767

Mohs micrographic surgery for melanoma: A prospective multicenter study

Bottom line: With the goal of achieving negative margins 97% of the time, surgical margins of 10 mm for trunk/extremity and 12 mm for head/neck are recommended for primary noninvasive melanoma and invasive melanoma when 100% histologic margin evaluation is not used.

Patrick M. Ellison, MD, John A. Zitelli, MD, and David G. Brodland, MD

From the Academy



775

Joint American Academy of Dermatology—National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy

Craig A. Elmets, MD (Co-Chair), Henry W. Lim, MD, Benjamin Stoff, MD, MA, Cody Connor, MD, Kelly M. Cordoro, MD, Mark Lebwohl, MD, April W. Armstrong, MD, MPH, Dawn M. R. Davis, MD, Boni E. Elewski, MD, Joel M. Gelfand, MD, MSCE, Kenneth B. Gordon, MD, Alice B. Gottlieb, MD, PhD, Daniel H. Kaplan, MD, PhD, Arthur Kavanaugh, MD, Matthew Kiselica, BA/BS, Dario Kivelevitch, MD, Neil J. Korman, MD, PhD, Daniela Kroshinsky, MD, MPH, Craig L. Leonardi, MD, Jason Lichten, MD, Nehal N. Mehta, MD, MSCE, Amy S. Paller, MD, Sylvia L. Parra, MD, Arun L. Pathy, MD, Elizabeth A. Farley Prater, MD, Reena N. Rupani, MD, Michael Siegel, PhD, Bruce E. Strober, MD, PhD, Emily B. Wong, MD, Jashin J. Wu, MD, Vidhya Hariharan, PhD, and Alan Menter, MD (Co-Chair)

Reviews

805

Early detection of acral melanoma: A review of clinical, dermoscopic, histopathologic, and molecular characteristics

Bottom line: We have summarized current clinical guidelines on acral lentiginous melanoma, dermoscopic findings, histopathologic clues, and molecular findings. Integrating this information is required to diagnose this disease at an early, curable stage.

Claudia C. Darmawan, MBBS, Gwanghyun Jo, MD, Sara E. Montenegro, MD, Yoonjin Kwak, MD, PhD, Lee Cheol, MD, PhD, Kwang Hyun Cho, MD, PhD, and Je-Ho Mun, MD, PhD

813

Triggers, clinical manifestations, and management of pediatric erythema multiforme: A systematic review

Bottom line: Diagnosis of pediatric erythema multiforme is usually clinical, with biopsy most frequently required in infants. Management includes supportive care and, if severe or recurrent, corticosteroids or antimicrobial drugs.

Samer Zoghaib, MD, Elio Kechichian, MD, Karim Souaid, MD, Boutros Soutou, MD, Josiane Helou, MD, and Roland Tomb, MD, PhD

823

Melanoma on chronically sun-damaged skin: Lentigo maligna and desmoplastic melanoma

Bottom line: These melanomas often present subtly and can behave aggressively; thus, they require clinical acumen, the use of all available diagnostic tools, and appropriately aggressive treatment.

Madeline E. DeWane, BA, Andrew Kelsey, MD, Margaret Oliviero, ARNP, Harold Rabinovitz, MD, and Jane M. Grant-Kels, MD

- 834** Platelet-rich plasma and its utility in medical dermatology:
A systematic review
Bottom line: Activated platelet-rich plasma should be considered as adjunctive therapy to optimize outcomes in vitiligo and skin ulcers refractory to standard wound care.
Michael J. Hessler, MD, and Nikhil Shyam, MD

Research Letters

- 847** Allergenic ingredients in commercial topical cannabinoid preparations
Brandon L. Adler, MD, and Vincent A. DeLeo, MD
- 848** Real-world drug usage survival of spironolactone versus oral antibiotics for the management of female patients with acne
John S. Barbieri, MD, MBA, Juliana K. Choi, MD, PhD, William D. James, MD, and David J. Margolis, MD, PhD
- 851** Adult acne in women is not associated with a specific type of *Cutibacterium acnes*
Mélanie Saint-Jean, MD, Stéphane Corvec, PharmD, PhD, Jean-Michel Nguyen, MD, PhD, Marie Le Moigne, MD, Aurélie Boisrobert, BN, Amir Khammari, PhD, and Brigitte Dréno, MD, PhD
- 852** The proportion of nevus-associated invasive melanoma differs with Breslow thickness: A cross-sectional study of 1087 cutaneous melanomas
Esperanza Manrique-Silva, MD, Diana Reyes-García, MD, Belen Folgado, MD, Alejandro Martin-Gorgojo, PhD, Victor Traves, MD, Celia Requena, PhD, and Eduardo Nagore, PhD
- 854** Retrospective outcome analysis of 25 alopecia areata patients treated with simvastatin/ezetimibe
Carlos Morillo-Hernandez, BS, Jonathan J. Lee, MD, and Joseph C. English III, MD
- 857** Brodalumab in the treatment of moderate to severe psoriasis in patients when previous anti-interleukin 17A therapies have failed
Grace Kimmel, MD, Margot Chima, MD, Hee Jin Kim, MD, Jennifer Bares, MD, Christopher J. Yao, MPH, Giselle Singer, BS, Soo Jung Kim, MD, Jerry Bagel, MD, and Mark Lebwohl, MD
- 859** Adverse cutaneous drug reactions with antimalarials in cutaneous lupus and dermatomyositis: A retrospective cohort study
Cristian D. Gonzalez, MD, Christopher Hansen, MD, and Jennie T. Clarke, MD
- 860** Post—nail procedure analgesia: A randomized control pilot study
Nilton Di Chiacchio, MD, PhD, Jorge Ocampo-Garza, MD, César D. Villarreal-Villarreal, MD, Jesus Ancer-Arellano, MD, Leandro Fonseca Noriega, MD, and Nilton Gioia Di Chiacchio, MD, PhD

862 Atopic dermatitis is associated with increased hospitalization in US children

Tammy Hua, BA, and Jonathan I. Silverberg, MD, PhD, MPH

865 Hair transplant in frontal fibrosing alopecia: A multicenter review of 51 patients

Sergio Vañó-Galván, MD, PhD, Emilio Villodres, MD, PhD, Ramón Pigem, MD, M. R. Navarro-Belmonte, MD, Manuel Asín-Llorca, MD, Teresa Meyer-González, MD, Rita Rodrigues-Barata, MD, Óscar M. Moreno-Arrones, MD, PhD, David Saceda-Corralo, MD, PhD, Pierre Bouhanna, MD, PhD, and Alejandro Camps, MD, PhD

Health Policy & Practice

867 The impact of the Orphan Drug Act on Food and Drug Administration-approved therapies for rare skin diseases and skin-related cancers

Bottom line: We provide an analysis of dermatologic orphan drug approvals and forecast future drug development trends.

Laura Karas, MD, MPH, Christine Y. Lu, MSc, PhD, Pankaj B. Agrawal, MD, MMSc, and Maryam M. Asgari, MD, MPH

JAAD Game Changer

878 JAAD Game Changers: Variability in mitotic figures in serial sections of thin melanomas

Robert T. Brodell, MD

Correction

879 Erratum to “JAAD Game Changers: Multivariate analysis of potential risk factors for lymph node metastasis in patients with cutaneous squamous cell carcinoma of the head and neck.” *J Am Acad Dermatol.* 80 (2019) 1255-1255

JAAD ONLINE

Available at <http://www.jaad.org>

In Memoriam

e57 In memoriam: Richard L. Dobson, MD
Bruce H. Thiers, MD, and John C. Maize, Sr, MD

Clinical Pearl

e59 Application of cooled hydrogel dressing to minimize dyspigmentation from laser tattoo removal
Hao Feng, MD, MHS, Mitalee P. Christman, MD, and Roy G. Geronemus, MD

Surgical Pearls

e61 Use of multifilament nylon floss in onychocryptosis
Anuradha Bishnoi, MD, DNB, Keshavamurthy Vinay, MD, DNB, MNAMS, and Sunil Dogra, MD, DNB, FRCP



e63 Reverse beveling to improve wound edge apposition
Kourosh Beroukhim, MD, Lindsay R. Sklar, MD, and Daniel B. Eisen, MD

e65 Retrograde punch biopsy technique for removal of enlarged earlobe piercings
Jacques Pierre du Toit, MBCHB, MMED, FC DERM, Johann de Wet, MBCHB, MMED, and Willem Visser, MBCHB, MFAMMED, MMED

	e67	Microneedling with Dermaroller 192 needles along with 5-fluorouracil solution in the treatment of stable vitiligo <i>Akshy Kumar, MD, Rakesh Bharti, MD, and Shail Agarwal, MD</i>
Technology Pearl	e71	Absence of lesional features on reflectance confocal microscopy: Quality control steps to avoid false-negative results <i>Melissa Gill, MD, Jane M. Grant-Kels, MD, and Christi Alessi Fox, MS</i>
Notes & Comments	e75	Dermoscopy in folliculotropic mycosis fungoides—A possible mimicker of follicle-based inflammatory and infectious disorders <i>Shamir Geller, MD, Ayelet Rishpon, MD, and Patricia L. Myskowski, MD</i>
	e77	Reply: Application of dermoscopy in folliculotropic mycosis fungoides <i>Enzo Errichetti, MD, MSc, and Murat Durdu, MD, MSc</i>
	e79	Renaming frontal fibrosing alopecia is a step too far <i>Lachlan A. Byth, MD</i>
	e81	Comment on “Secukinumab drug survival in patients with psoriasis: A multicenter, real-world, retrospective study” <i>Laura Palacios-García, MD, Celia Gómez-de Castro, MD, Marc Mir-Bonafé, MD, Cristina Calzón, PharmD, Cristina Galache, MD, PhD, and Jorge Santos-Juanes, MD, PhD</i>
	e83	Reply to: “Comment on ‘Secukinumab drug survival in patients with psoriasis: A multicenter, real-world, retrospective study’” <i>Andrea Chiricozzi, MD, and Tiago Torres, MD</i>
	e85	Comment on “Just a quick pic: Ethics of medical photography:” Generative adversarial networks could be a solution <i>Soo Ick Cho, MD, Jung-Im Na, MD, PhD, Seung Seog Han, MD, PhD, and Jin Ho Chung, MD, PhD</i>
Iotaderma	e87	Iotaderma (#307) <i>Robert I. Rudolph, MD, FACP</i>

To access the current version of the Information for Authors, Transfer of Copyright Form, Authorship Declaration Form, Conflict of Interest Disclosure, and Authors’ Rights Statement, as well as many other pertinent documents, please visit our website at www.jaad.org/content/auth_page.

Full text access to *Journal of the American Academy of Dermatology Online* is now available for Academy members and all print subscribers. To activate your online subscription, please visit *Journal of the American Academy of Dermatology Online* by pointing your browser to <http://www.jaad.org>.

Icon Key

- | | | |
|---|---|---|
|  | = | CME article |
|  | = | Video content available at http://www.jaad.org |
|  | = | An interview with the author is available by podcast at: https://account.aad.org/Library/Podcasts |
|  | = | MOC credits available |

Early online publication: Articles in press

The *Journal of the American Academy of Dermatology* posts in-press articles online before they appear in the print edition of the *Journal*. These Articles in Press are available from both the JAAD website (www.jaad.org) and Science Direct (www.sciencedirect.com). To access Articles in Press, simply click the Articles in Press link.

JAAD Case Reports

The *Journal of the American Academy of Dermatology* no longer publishes Case Reports. These can now be found in our new open access journal, *JAAD Case Reports*, at <http://www.jaadcasereports.org>.