

punctate hyperreflective vitreous opacities seen on handheld spectral domain optical coherence tomography (SD-OCT).

Methods: This is a prospective observational study of infants requiring ROP screening between July 2015 and December 2017. Infants were imaged using handheld SD-OCT at the time of routine examinations. Trained graders masked to the clinical assessment analyzed each OCT scan of the right eye for vitreoretinal findings. Disagreement was mediated by a third trained grader. Punctate hyperreflective vitreous opacities seen on OCT were correlated with clinical ROP severity and other OCT vitreoretinal pathologies.

Results: Among 93 infants studied (51% male, mean gestational age 28.3 ± 2.9 weeks, mean birthweight $1008.2 \text{ kg} \pm 287.8 \text{ g}$), 22/93 (38%) developed ROP (14/93 [15%] stage 3). Agreement for OCT graders was 91% ($\kappa = 0.8$; $P < 0.001$). Punctate hyperreflective vitreous opacities developed in 31/93 (33%) of infants and were associated with the presence of ROP ($P = 0.005$), maximum ROP stage ($P = 0.005$), and pre-plus or plus disease ($P = 0.002$).

Discussion: Punctate hyperreflective vitreous opacities seen on handheld SD-OCT were strongly correlated with presence of ROP, maximum stage, and pre-plus or plus disease among premature infants screened for ROP. The opacities may represent cellular proliferation, protein or hemoglobin associated with advanced ROP.

Conclusions: Punctate hyperreflective vitreous opacities on OCT are a marker for advanced ROP. Further study should explore handheld SD-OCT as a noninvasive ROP screening tool.

Posters

041 Biomechanical analysis of ligatures and technique for managing drainage tubes in pediatric glaucoma. Carolina Adams, Steven Kane, Steven Brooks

Introduction: Absorbable ligatures are often used with glaucoma drainage tubes to avoid early postoperative hypotony. We sought to measure the force required to ligate a drainage tube and to develop a modified technique to promote earlier release in pediatric patients, where plate encapsulation occurs more quickly than in adults.

Methods: A precision digital force gauge was used to measure the tensile strength of several common ophthalmic sutures, and the necessary tensile force required to achieve tube ligation. A novel technique for tube ligation was devised to allow sutures as small as 10-0 to be effectively used.

Results: The mean tensile strengths of unknotted sutures varied from $211 \pm 127 \text{ g}$ for 10-0 Vicryl to $477 \pm 69 \text{ g}$ for 6-0 chromic gut. The mean tensile force required to ligate a Baerveldt or Ahmed tube was $35.9 \pm 0.9 \text{ g}$. However, 9-0 or 10-0 Vicryl could not be reliably used for ligation, due to breakage, unless a modified technique was employed, wherein the tube was first stretched to reduce its thickness and diameter.

Discussion: Frictional forces inherent to knot tying make it unfeasible to reliably use 9-0 or 10-0 Vicryl to ligate a drainage tube, despite the unknotted threads possessing apparently sufficient tensile strength. Our modified ligation technique overcomes this issue, allowing a wider range, of suture choices, and the potential for achieving more rapid release in pediatric cases.

Conclusions: A novel modification in technique allows the use of very small gauge sutures to ligate tubes in glaucoma drainage implants, allowing for potentially faster release times that may be favorable in managing pediatric glaucomas.

042 Determination of the strabismus surgery dosage accuracy using a new technology. Satenik G. Agagulyan, Elena A. Kudryashova, Victoria O. Balasanyan, Igor E. Aznauryan

Introduction: Accurate dosage of strabismus surgery is one of the most important issues in strabismology. The aim of this study was to show the accuracy of mathematical computer program 'Strabo' with the use of Gazelab technology. Gazelab technology shows the motility of eyes and the angle of deviation with one eye closed and both eyes open.

Methods: All patients passed standard pre- and postoperative ophthalmologic examinations. Additionally all patients undergone Gazelab Free or 5-point test before and on the 7th day after surgery.

Results: Eleven patients were operated on (mean age, 5.8 ± 1.5 years). The average value of the deviation with one eye covered was $34.7^\circ \pm 12.2^\circ$; with two eyes open, $27.7^\circ \pm 13.9^\circ$. Evaluation of the effectiveness of surgery dosage was measured by postoperative angle of strabismus measured by Gazelab technology. The average value of the deviation after surgery with one eye covered was $12.45^\circ \pm 7.1^\circ$; with two eyes open, $10.3^\circ \pm 6.1^\circ$. Mathematical surgery simulation differed from the actual surgical results by only $\pm 4.7^\circ$.

Discussion: Results obtained after the strabismus surgery correlated with the expected results that was received by "Strabo" program.

Conclusions: A mathematical model of the surgery allows to distribute the surgical effect on both eyes with a high cosmetic and functional result.

043 Digital biometry in children undergoing cataract surgery in vitrectomized eyes. Sumita S. Agarkar, Swathi Mailankody

Introduction: To report the accuracy of biometry in children undergoing cataract surgery in vitrectomized eyes using SRK-II formula.

Methods: A retrospective review of medical records of children undergoing cataract surgery between 2008 and 2017 with a prior history of pars plana vitrectomy was done. Biometry was performed using Ocuscan Rxp (Alcon) and keratometry was performed using a handheld keratometer IOL power calculation was performed under general anesthesia in uncooperative children. Adjustment in speed was made in silicon oil filled eyes. The main outcome variable studied was absolute prediction error (APE). APE was calculated as absolute difference between target refraction and postoperative refraction obtained at 6 weeks.

Results: A total of 92 eyes of 90 patients were included. Mean age at surgery was 12.04 ± 3.33 years. Mean axial length was $24.85 \pm 1.94 \text{ mm}$. Mean APE using SRK -II formula was 1.29 ± 1.12 for the entire group. However, 35% of the eyes had APE between 0 and 0.5. APE was affected by axial length ($P = 0.007$).

Discussion: IOL power calculation remains challenging in children specially in younger children. Various studies over the years have reported mean prediction error involving several different formulae. Kunnaya et al reported mean APE of 2.27 with SRK 2. There is no such data available for Vitrectomized eyes. Children in this study were older as well as had longer axial lengths compared to that in literature.

Conclusions: Prediction error using SRK-II formula in vitrectomized pediatric eyes is comparable to that in nonvitrectomized eyes as reported in literature.

044 Effect of oral propranolol in astigmatism-induced orbital hemangioma. Nora Aldohayan, Yasser Al-Thnayan

Introduction: Capillary hemangiomas are the most common orbital tumor in infancy, with a course of rapid proliferation followed by spontaneous regression. It may cause astigmatism or amblyopia