



The antimetabolite 3-bromopyruvate selectively inhibits *Staphylococcus aureus*

Paolo Visca^a, Federica Pisa^{a,b}, Francesco Imperi^{b,*}

^a Department of Sciences, University 'Roma Tre', Rome, Italy

^b Department of Biology and Biotechnology Charles Darwin, Sapienza University of Rome, Laboratory affiliated to Istituto Pasteur Italia – Fondazione Cenci Bolognetti, Rome, Italy



ARTICLE INFO

Article history:

Received 21 September 2018

Accepted 17 November 2018

Editor: Stefania Stefani

Keywords:

Bactericidal activity

Biofilm

Catabolism

Glutathione

ESKAPE

Pyruvate

ABSTRACT

Increased antibacterial resistance jeopardizes current therapeutic strategies to control infections, soliciting the development of novel antibacterial drugs with new mechanisms of action. This study reports the discovery of potent and selective antistaphylococcal activity of 3-bromopyruvate (3BP), an antimetabolite in preclinical development as an anticancer drug. 3BP showed bactericidal activity against *Staphylococcus aureus*, with active concentrations comparable with those reported to be effective against cancer cells. In contrast, no relevant activity was observed against other ESKAPE bacteria (*Enterococcus faecium*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* spp.). The anti-staphylococcal activity of 3BP was confirmed using a panel of human and veterinary strains, including multi-drug-resistant isolates. 3BP showed highest antibacterial activity under conditions that increase its stability (acidic pH) or promote *S. aureus* fermentative metabolism (anaerobiosis), although 3BP was also able to kill metabolically inactive cells. 3BP showed synergism with gentamicin, and also disrupted pre-formed *S. aureus* biofilms at concentrations only slightly higher than those inhibiting planktonic cells. This study unravels novel antibacterial and antibiofilm activities for the anticancer drug 3BP, paving the way for further preclinical studies.

© 2018 Elsevier B.V. and International Society of Chemotherapy. All rights reserved.

1. Introduction

Antibiotic resistance is a serious public health problem at a global level. Antibiotics have saved millions of lives, but are progressively losing their efficacy against many bacterial pathogens. While very few new antibiotics are being developed by the pharmaceutical industry, mainly due to the inherent low reward and high risk of antibiotic research, the rapid spread of antibiotic-resistant pathogens in hospitals and the community calls for new investments in antibacterial drug discovery. However, to be effective, these investments should support experimental approaches able to make the process of antibacterial drug discovery more rapid and economically sustainable.

Searching for side activities in drugs already approved for use in humans or at an advanced phase of preclinical development represents a potential shortcut to obtain new antibacterials [1]. Compared with de-novo drug discovery, the drug repurposing approach has a higher probability of yielding bioavailable and safe hit com-

pounds, which can move directly into clinical trials or serve as leads for drug optimization programmes. Moreover, drug repurposing is expected to reduce the time and costs of the conventional drug discovery process, as pharmacological/toxicological information is available for repurposing candidates. Notably, UK and US funders have recently launched programmes to re-evaluate de-prioritized drugs for new therapeutic uses [1].

In the last years, an increasing number of studies have identified some antibacterial activity in several drugs approved for various purposes, including anticancer, antifungal and cardiovascular therapies [2,3], strongly supporting the research on drug repurposing as a strategy to identify novel antibacterials that can be delivered rapidly to the clinical phase.

3-Bromopyruvate (3BP) is a chemically synthesized halogenated derivative of pyruvate that has been used by biochemists for several decades. Indeed, alkylating properties have made 3BP suitable for use in enzymatic studies *in vitro*, with preferential targets focused on metabolic enzymes. Approximately 15 years ago, 3BP was also proven to act as a powerful anticancer agent [4]. 3BP treatment abolishes ATP production in cancer cells, mainly by inhibition of glycolysis [5], inducing rapid cancer cell death both *in vitro* and in several animal tumour models, including hepatocarcinoma, colon, breast and lung cancers [6]. Irrespective

* Corresponding author. Address: Department of Biology and Biotechnology Charles Darwin, Sapienza University of Rome, Laboratory affiliated to Istituto Pasteur Italia – Fondazione Cenci Bolognetti, Rome, Italy.

E-mail address: francesco.imperi@uniroma1.it (F. Imperi).

of the strong alkylating activity of 3BP *in vitro*, this compound showed outstanding selectivity towards cancer cells in animal models, which appears to be related to: (i) the much higher expression of the main 3BP transporter (monocarboxylate transporter 1) in tumour cells compared with normal cells; and (ii) the metabolic differences between cancer cells and healthy cells. Indeed, while glycolytic metabolism is exacerbated in cancer cells, and these cells display high rates of aerobic glycolysis even in the presence of oxygen (i.e. the 'Warburg effect'), normal cells mainly rely on mitochondrial oxidative phosphorylation for energy production, and are therefore barely sensitive to the antiglycolytic effect of 3BP [6,7]. The tumour-eradicating activity of 3BP has been documented in different animal models. While the majority of *in vivo* studies did not report relevant adverse effects of 3BP treatment on healthy cells or tissues at therapeutically active doses, some liver toxicity was observed when very high dosing regimens and systemic delivery were used [6]. A few studies have described the use of 3BP in volunteer cancer patients [8,9], and 3BP appears to be used as a last-resort treatment in some cancer clinics worldwide [10, <https://www.cancertreatmentsresearch.com/3-bromopyruvate/>]. However, while the US company PreScience Labs stated that the Food and Drug Administration approved a phase I clinical evaluation of 3BP in liver cancer patients in 2013, 3BP has not yet undergone formal clinical trials, possibly because of lack of funding and/or the existence of competing patent applications [10]. A recent case of 3BP misuse by an unlicensed practitioner, who was accused of causing three deaths in a German alternative medicine clinic in 2016, may have complicated the road of this candidate drug to the clinic [10].

Regarding the possible activity of 3BP on micro-organisms, two studies highlighted the capability of 3BP to inhibit the growth of eukaryotic pathogens, such as the fungus *Cryptococcus neoformans* and the microalgae *Prototheca* spp. [11,12]. Although a patent application for use of antiglycolytic compounds, including 3BP, to prevent or treat bacterial infections was submitted in 2011 [13], the antibacterial activity of 3BP has not been investigated experimentally, with the exception of a recent study showing mild growth inhibition of *Mycobacterium tuberculosis* exposed to millimolar concentrations of 3BP [14].

This study aimed to assess the possibility of 3BP repurposing as an antibacterial agent against pathogens of the ESKAPE group (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Enterobacter* spp.) for which new drugs are urgently needed. The working hypothesis originated from the following: (i) bacterial pathogens possess analogues of eukaryotic glycolytic enzymes; (ii) pyruvate is an intermediate of energy metabolism in many pathogenic bacteria; and (iii) 3BP inhibits some bacterial glycolytic enzymes *in vitro* [12]. The results revealed that 3BP has bactericidal and antibiofilm activity against *S. aureus*, including multi-drug-resistant isolates, while it does not inhibit all the other ESKAPE bacteria, and suggest that the specificity of 3BP towards *S. aureus* is likely due to unique metabolic features of this species.

2. Materials and methods

2.1. Bacterial strains and growth conditions

Bacterial strains were routinely maintained in Luria-Bertani (LB) medium supplemented with 1.5% agar at 37°C. Mueller-Hinton (MH) broth was used for antibacterial assays. When indicated, MH was adjusted to different pH values by the addition of HCl and sterilized by filtration, or supplemented with glucose or pyruvate at 5 or 50 mM concentration. Anaerobic conditions were obtained by incubating plates in a GENBox Jar 2.5 L in the presence of a GENbox anaerobic generator (bioMérieux, Marcy l'Etoile, France).

To support anaerobic growth of *S. aureus*, MH broth was supplemented with 5 g/L glucose. Tryptic soy broth (TSB) supplemented with 2% glucose and 2% NaCl (TSB-GN) was used for biofilm assays [15]. 3BP was purchased from Sigma-Aldrich (St Louis, MO, USA), and the stock solution was prepared in water and stored at -20°C before use.

2.2. Minimum inhibitory concentration and minimum bactericidal concentration assays

The minimum inhibitory concentration (MIC) of 3BP and antibiotics was determined using the broth microdilution method according to the Clinical Laboratory Standards Institute (CLSI) guidelines [16], after 24 h of incubation at 37°C under static conditions. The minimum bactericidal concentration (MBC) of 3BP for *S. aureus* isolates was determined by plating dilutions from each well showing no bacterial growth on LB agar plates. Colonies were counted after 24 h of incubation at 37°C. The MBC was defined as the lowest concentration of 3BP required to kill 99.9% of the initial inoculum [17].

2.3. Time-kill assays

Time-kill assays were performed using the CLSI M26-A standard method [17] with some modifications. Briefly, *S. aureus* ATCC25923 was cultured in MH broth for 6–8 h and then diluted in MH broth at a final optical density at a wavelength of 600 nm (OD₆₀₀) of 0.01 [corresponding to c. 10⁶ colony-forming units (CFU)/mL] in the absence or presence of 3BP and/or gentamicin. Bacterial cultures were incubated at 37°C under vigorous shaking (200 rpm). At different time points, serial dilutions were prepared in saline (0.9% NaCl), and aliquots of each dilution were plated on LB agar plates to determine the number of viable cells (CFU).

2.4. Checkerboard assay

The checkerboard broth microdilution method [18] was performed to investigate the interaction of 3BP with ampicillin, gentamicin, tetracycline, rifampicin, chloramphenicol and ofloxacin. Two-fold serial dilutions of 3BP and each antibiotic in MH broth were perpendicularly dispensed in 96-well microtitre plates, and each well was inoculated with *S. aureus* ATCC25923 at c. 5 × 10⁵ CFU/mL. Microtitre plates were incubated at 37°C under static conditions for 24 h, and bacterial growth was assessed visually. Results were expressed as the fractional inhibitory concentration index (FICI), which corresponds to the sum of the FICs of the two antibacterial compounds, where the FIC for each drug is defined as the MIC of the drugs in combination divided by the MIC of the drugs used alone. The FICI was interpreted as follows: ≤0.5, synergy; >0.5–4, indifference; and >4, antagonism [18].

2.5. Biofilm assays

Biofilm formation was evaluated in TSB-GN as described previously, with a few modifications [15]. *S. aureus* strains were cultured in TSB at 37°C for 6–8 h, refreshed in TSB-GN at OD₆₀₀ of 0.01, and 100-μL aliquots were dispensed in the wells of tissue-culture-treated 96-well microtitre plates. After 24 h at 37°C under static conditions, planktonic cells were removed and microtitre plates were washed three times with distilled water to remove non-adherent cells. Microtitre plates were air-dried, and adherent biofilms were fixed with 100 μL of 95% ethanol for 10 min. Ethanol was discarded, microtitre plates were air-dried, and fixed biofilm was stained with 100 μL of 1% crystal violet for 10 min. Unbound crystal violet was removed and microtitre plates were

Table 1
3BP activity against ESKAPE bacteria.

Species	Strain	Reference/source	MIC ($\mu\text{g/mL}$)	MBC ($\mu\text{g/mL}$)
<i>Enterococcus faecium</i>	3137	Rossolini's laboratory collection	>320	NA
<i>Enterococcus faecalis</i>	ATCC29212	ATCC	>320	NA
<i>Staphylococcus aureus</i>	ATCC25923	ATCC	20	40
	ATCC43300 (MRSA)	ATCC	40	80
<i>Klebsiella pneumoniae</i>	ATCC27736	ATCC	>320	NA
	17830 (MDR)	[20]	>320	NA
<i>Acinetobacter baumannii</i>	ATCC19606	ATCC	>320	NA
	ACICU (MDR)	[21]	>320	NA
<i>Pseudomonas aeruginosa</i>	ATCC15692	ATCC	>320	NA
	SP1 (MDR)	[22]	>320	NA
<i>Enterobacter cloacae</i>	ATCC13047	ATCC	>320	NA
<i>Enterobacter aerogenes</i>	ATCC13048	ATCC	>320	NA

MIC, minimum inhibitory concentration; MBC, minimum bactericidal concentration; MDR, multi-drug resistant; MRSA, methicillin-resistant *Staphylococcus aureus*; ATCC, American Type Culture Collection; NA, not assessed.

washed three times with distilled water and air-dried. The biofilm-bound dye was solubilized with 100 μL of 33% glacial acetic acid and, after 10 min, quantified by measuring OD_{600} in a Wallac 1420 Victor3V plate reader (Perkin Elmer, Waltham, MA, USA). To assess the antibiofilm activity of 3BP, 24-h-old biofilms were incubated under static conditions at 37°C for a further 24 h in TSB-GN supplemented or not with 3BP at 1 \times , 2 \times , 4 \times or 8 \times MIC. Biofilms were fixed, stained and quantified as described above.

2.6. Construction of the deletion mutant *P. aeruginosa* PAO1 ΔgshA

Deletion mutagenesis in *P. aeruginosa* was performed as described previously [19] using the suicide vector pDM4 ΔgshA , which was generated by cloning c. 450-bp DNA fragments corresponding to the upstream and downstream genomic regions of *gshA* (PA5203) into the plasmid pDM4. Plasmids and primers used for cloning are listed in Table S1 (see online supplementary material).

2.7. Statistical analysis

Statistical analysis was performed using GraphPad InStat with one-way analysis of variance.

3. Results

3.1. Antibacterial activity of 3BP against ESKAPE pathogens

The MIC of 3BP was determined on a small selection of bacterial strains belonging to the ESKAPE group. 3BP showed antimicrobial activity against the two *S. aureus* strains, ATCC25923 and methicillin-resistant ATCC43300, with MIC values of 20 and 40 $\mu\text{g/mL}$, respectively. In contrast, all the other ESKAPE species were insensitive to 3BP up to 320 $\mu\text{g/mL}$ (Table 1). The anti-*S. aureus* activity of 3BP was further evaluated on 20 clinical and veterinary isolates, including some cystic fibrosis isolates and several antibiotic-resistant strains (Table S2, see online supplementary material). 3BP was found to inhibit the growth of all *S. aureus* isolates tested, with MIC values ranging between 20 and 80 $\mu\text{g/mL}$ (Table S2).

To further investigate the antistaphylococcal activity of 3BP, the MBC was also determined for each isolate. Notably, the MBC of 3BP was identical to the MIC for 77% of the strains (17 out of 22), and only 2 \times MIC for the remaining strains (Tables 1 and S1, see online supplementary material), suggesting that 3BP is primarily endowed with bactericidal activity against *S. aureus*. To confirm this, a time-kill assay was performed using *S. aureus* ATCC25923 as the reference strain and 3BP at concentrations corresponding to 1 \times , 2 \times or 4 \times MIC. 3BP was able to kill *S. aureus* cells in a dose-dependent




Fig. 1. Time-kill curves of *Staphylococcus aureus* ATCC25923 exposed to 3-bromopyruvate (3BP). *S. aureus* ATCC25923 was cultured in Mueller-Hinton (MH) broth, adjusted to c. 10^6 colony-forming units (CFU)/mL in MH broth containing 3BP at 20, 40 or 80 $\mu\text{g/mL}$, and incubated at 37°C with vigorous agitation. At the indicated time points (0, 1, 2 and 4 h), cell viability was assessed by CFU counting on agar plates. The results are mean (\pm standard deviation) of at least four independent assays.

manner with very fast bactericidal activity, as 97.3, 98.7 and 99.7% of cells died within 1 h of 3BP treatment at 1 \times , 2 \times and 4 \times MIC, respectively (Fig. 1). Under these experimental conditions, the 3 \log_{10} reduction in the number of CFU, which is required to define an antibiotic as bactericidal [17], was obtained with 80 $\mu\text{g/mL}$ of 3BP after ≥ 2 h of treatment (Fig. 1).

3.2. Effect of culture conditions on the antistaphylococcal activity of 3BP

It has been reported that the stability of 3BP in aqueous solutions is significantly higher at acidic pH compared with neutral (physiological) or basic pH [23]. To investigate any pH dependence of antibacterial activity, the MIC of 3BP was determined for the reference strain *S. aureus* ATCC25923 in MH broth adjusted to different pH values (range 5–8; Table 2). While the different pHs did not notably affect the ability of *S. aureus* ATCC25923 to grow under the tested conditions, the growth rate was reduced slightly at pH 5 compared with other pHs (Fig. S1, see online supplementary material). Notably, a significant decrease in the MIC of 3BP was observed from basic to acidic pH. Indeed, the MIC of 3BP was four-fold lower at pH 5 and eight-fold higher at pH 8 with respect to the MIC at pH 7 (Table 2), indicating that an acidic pH significantly enhances the antibacterial activity of 3BP, probably as a result of increased stability of the compound.

Notably, exogenous pyruvate protected *S. aureus* from the antibacterial activity of 3BP in a dose-dependent manner, while




Fig. 2. Killing activity of 3-bromopyruvate (3BP) on metabolically inactive cells. *Staphylococcus aureus* ATCC25923 was cultured in Mueller-Hinton (MH) broth, adjusted to $c. 10^6$ colony-forming units (CFU)/mL in saline, and (A) immediately treated with 3BP at $4 \times$ minimum inhibitory concentration (80 µg/mL) in saline for 2 h at 37°C, or (B) incubated at 4°C for 16 h and then treated with 3BP (80 µg/mL) in saline for 2 h at 37°C. Results are reported as percentage of CFU with respect to the untreated controls (100%), and represent the mean (\pm standard deviation) of three independent assays.

Table 2

Effect of pH and anaerobiosis on minimum inhibitory concentration (MIC) of 3-bromopyruvate for *Staphylococcus aureus* ATCC25923.

Condition	pH	Supplement	MIC (µg/mL)
Aerobiosis	5		5
	6		10
	6.5		10
	7		20
	7.5		40–80
	8		160
	7	Glucose (50 mM)	20
	7	Glucose (5 mM)	20
Anaerobiosis	7	Pyruvate (50 mM)	320
	7	Pyruvate (5 mM)	80
	7		5
	7		5

equivalent concentrations of glucose did not (Table 2), suggesting that pyruvate and 3BP compete for transporter(s) and/or specific intracellular target(s).

3BP activity against *S. aureus* was also assessed during anaerobic growth, which was reported to increase the rate of glycolysis in this bacterium [24,25]. The MIC of 3BP for ATCC25923 was four-fold lower than that obtained under aerobiosis (Table 2), implying that *S. aureus* cells are more sensitive to 3BP under anaerobic conditions.

To verify whether active metabolism is essential for the antibacterial activity of 3BP, the bactericidal activity of $4 \times$ MIC 3BP (80 µg/mL) in saline was tested against *S. aureus* ATCC25923 pre-incubated or not at 4°C for 16 h in saline to reduce metabolic activity. No increase in the resistance to 3BP-mediated killing was observed in metabolically inactive (or poorly active) cells (Fig. 2), suggesting that 3BP treatment could also be effective against dormant *S. aureus* cells.

3.3. Interaction between 3BP and antibiotics

To assess 3BP interaction(s) with currently available antibacterial drugs, checkerboard assays with 3BP and several antibiotics characterized by different modes of action were conducted using *S. aureus* ATCC25923 as the test strain. 3BP had a synergistic effect only with gentamicin (Table 3). Although indifference was observed with all other antibiotics, ampicillin–3BP and tetracycline–3BP combinations showed an FICI close to 0.5 (Table 3), suggestive of some additive effect. Notably, no drug pair showed antagonistic effects (FICI >4) (Table 3).

The synergy between 3BP and gentamicin was confirmed by time-kill assays; *S. aureus* ATCC25923 was treated with $0.5 \times$ MIC

Table 3

Interaction between 3-bromopyruvate and selected antibiotics against *Staphylococcus aureus* ATCC25923.

Antibiotic	MIC (µg/mL)	FICI ^a	Interpretation
Gentamicin	0.312	0.438	Synergy
Ampicillin	0.156	0.625	Indifference
Tetracycline	0.25	0.563	Indifference
Rifampicin	0.004	2	Indifference
Chloramphenicol	8	1	Indifference
Ofloxacin	0.25	2	Indifference

MIC, minimum inhibitory concentration; FICI, fractional inhibitory concentration index.

^a <0.5, synergy; 0.5–4, indifference; >4, antagonism.




Fig. 3. Time-kill curves of *Staphylococcus aureus* ATCC25923 exposed to 3-bromopyruvate (3BP) and gentamicin (Gm) alone or in combination. *S. aureus* ATCC25923 was cultured in Mueller-Hinton (MH) broth, adjusted to $c. 10^6$ colony-forming units (CFU)/mL in MH broth containing either 3BP or gentamicin or both at $0.5 \times$ minimum inhibitory concentration (10 µg/mL and 0.156 µg/mL, respectively), and incubated at 37°C with vigorous agitation. At the indicated time points (0, 1, 2, 4 and 20 h), cell viability was assessed by CFU counting on agar plates. The results are the mean (\pm standard deviation) of three independent assays.

of 3BP (10 µg/mL), gentamicin (0.156 µg/mL) or both compounds, and the number of CFU/mL was counted over time (Fig. 3). Notably, the number of CFU/mL decreased in the first 4 h of treatment with both compounds, either alone or in combination. However, while *S. aureus* ATCC25923 treated with sub-MIC concentrations of 3BP or gentamicin alone attained growth levels almost comparable to the untreated control (20-h incubation), the combination of 3BP with gentamicin at $0.5 \times$ MIC completely abrogated regrowth. Since synergism is defined as a ≥ 2 log₁₀ decrease in CFU/mL for the drug combination compared with each drug alone at 20 h [26], this experiment confirmed that 3BP and gentamicin exert a synergistic effect on *S. aureus*.




Fig. 4. 3-Bromopyruvate (3BP)-mediated disruption of *Staphylococcus aureus* biofilms. Biofilms of *S. aureus* strains ATCC25923, ATCC43300, BG2 and BG7 were developed in microtitre plates containing tryptic soy broth supplemented with 2% glucose and 2% NaCl (TSB-GN). After 24 h, supernatants were discarded, and fresh TSB-GN containing increasing 3BP concentrations was added (1 \times , 2 \times , 4 \times or 8 \times minimum inhibitory concentration). Plates were incubated for a further 24 h. Biofilm biomass was quantified as described in the 'Materials and methods' section. The results are the mean (\pm standard deviation) of four independent assays. Black histograms represent the biomass of 24-h-old biofilms before treatment. Grey and black asterisks denote significant differences with respect to 24-h-old biofilms or untreated 48-h-old biofilms, respectively. * P <0.05; ** P <0.01; *** P <0.001 (one-way analysis of variance).

3.4. Antibiofilm activity of 3BP

Biofilm formation during infection is a primary cause of antibiotic treatment failure, as bacterial cells encased in biofilms are generally more resistant to antibiotics than their planktonic counterparts [27]. To assess whether 3BP retains its antibacterial activity on biofilm cells, preformed biofilms of *S. aureus* ATCC25923 were treated for 24 h with increasing concentrations of 3BP, corresponding to 1 \times , 2 \times , 4 \times and 8 \times MIC. Notably, complete disruption of ATCC25923 biofilm was observed in the presence of 3BP concentrations $\geq 2 \times$ MIC (Fig. 4). No significant differences in biomass were observed between 24- or 48-h-old untreated biofilms and biofilms treated with 3BP at 1 \times MIC (Fig. 4).

To verify whether the antibiofilm properties of 3BP are conserved in different *S. aureus* strains, the same assay was performed on the strains ATCC43300, BG2 and BG7, endowed with different levels of sensitivity to 3BP in the MIC and MBC assays (Table S1, see online supplementary material). Overall, 3BP showed biofilm-disrupting activity against all strains, although some differences were observed. For instance, ATCC43300 biofilms were almost completely disrupted at $\geq 2 \times$ MIC (similarly to ATCC25923), while 3BP at 1 \times MIC only inhibited biofilm growth without causing biofilm disruption (Fig. 4). Biofilm disruption at $\geq 2 \times$ MIC was also observed for BG7, although the effect was weaker than that observed with ATCC25923 and ATCC43300 (Fig. 4). Finally, BG2 was slightly more resistant to the antibiofilm activity of 3BP, as significant biofilm disruption was only observed at $\geq 4 \times$ MIC (Fig. 4). This experiment demonstrated that 3BP does disrupt *S. aureus* biofilms, although at concentrations slightly higher than those active against planktonic cells.

4. Discussion

Increased resistance among bacterial pathogens leads to a growing need for novel antibacterials. This study assessed the antibacterial activity of a well-known antimetabolite, 3BP, which is in preclinical development as an antitumour drug. 3BP was found

to have bactericidal activity against *S. aureus*, with MIC and MBC values for different strains comparable to the concentrations active against cancer cell lines [6,28] or fungal cells [11] (Tables 1 and S1, see online supplementary material). In contrast, 3BP does not inhibit the growth of any other ESKAPE species (Table 1), either Gram-negative or Gram-positive, suggesting that the antibacterial activity of 3BP is specific to *S. aureus* (and possibly closely related species). The biological and molecular basis of such specificity is unknown. Considering that 3BP susceptibility in eukaryotic cells: (i) positively correlates with 3BP uptake rates; and (ii) negatively correlates with intracellular levels of glutathione [11], it is tempting to discuss whether these two factors could somehow justify the observed selectivity of 3BP towards *S. aureus*. Regarding 3BP uptake, at least two of the species found to be highly resistant to 3BP (i.e. *A. baumannii* and *P. aeruginosa*) are able to grow and/or obtain energy in minimal media containing lactate or pyruvate as the sole carbon source [29–32], indicating that these species possess transporter(s) for lactate/pyruvate uptake, which could also mediate internalization of the 3BP analogue. However, considering that enzymes and transporters involved in catabolism are often tightly regulated in bacteria [33], it cannot be excluded that the expression of these systems in these two species is somehow repressed during growth in the nutritionally rich MH broth. Unfortunately, knowledge on lactate and pyruvate transport systems in bacteria is still scarce [34], and thus the presence and/or expression of these transporters cannot be simply predicted by genomics and/or transcriptomic analyses. 3BP uptake was measured in fungal cells using [14C]-labelled 3BP [11]; however, this radioactive compound is not commercially available, hampering the direct assessment of 3BP internalization by bacteria in the present work. Thus, it cannot be ruled out that some specific feature(s) of the cell envelope can somehow influence the uptake of 3BP and, thus, its antibacterial activity in different bacteria.

Interestingly, a putative correlation can be predicted between glutathione and 3BP antibacterial activity. Indeed, among ESKAPE bacteria, *S. aureus* is the only pathogen that does not produce glutathione [35,36]. In both cancer and fungal cells, glutathione has

been proposed to protect against 3BP toxicity by preventing oxidative stress [11,37] and/or by directly interacting with and titrating 3BP, ultimately reducing its toxic levels in the cytoplasm [38]. It can be hypothesized that the higher sensitivity of *S. aureus* to 3BP could, at least partially, be related to the lack of glutathione-based detoxification mechanism(s). To test whether endogenous glutathione production can *per se* account for 3BP resistance, the *gshA* gene – which is essential for glutathione biosynthesis – was deleted in *P. aeruginosa* PAO1 [39]. Notably, the MIC of 3BP for the glutathione-deficient *P. aeruginosa* $\Delta gshA$ mutant was found to be 320 $\mu\text{g/mL}$, i.e. just two-fold lower than that of the wild type (640 $\mu\text{g/mL}$), and still four- to 16-fold higher than that for *S. aureus* strains (Table S1, see online supplementary material). This strongly suggests that the inability to produce glutathione does not represent the only reason for the specificity of 3BP towards *S. aureus*.

Finally, the anti-*S. aureus* activity of 3BP could be related to specific metabolic features of this bacterium. The ESKAPE group contains a strictly aerobic bacterium (*A. baumannii*) and five facultative anaerobes, which can use either anaerobic respiration and fermentation (*K. pneumoniae*, *Enterobacter* spp., *P. aeruginosa*), or fermentation alone (*S. aureus*, *Enterococcus* spp.) for energy production during anaerobic growth. Thus, the susceptibility of *S. aureus* to 3BP does not appear to be related to a distinctive catabolic potential of this bacterium. However, it cannot be excluded that the presence of unique metabolic pathway(s) and/or possible differences in the sequence or level of expression/activity of specific metabolic enzymes could account for the higher sensitivity of *S. aureus* to 3BP compared with other bacteria. Experiments are in progress to tentatively identify the transport mechanism(s) and intracellular target(s) of 3BP in *S. aureus*, and to define its mechanism of action as an antibacterial agent.

In addition to its bactericidal activity on planktonic cells, 3BP has also shown potent biofilm-disrupting properties, causing almost complete detachment of *S. aureus* biofilms at relatively low concentrations (only two- to four-fold higher than the MIC) (Fig. 4). This result indicates that, differently from the majority of antibiotics which are much less effective against biofilms relative to planktonic cells, 3BP retains its anti-*S. aureus* activity against biofilm-forming cells. The high tolerance of biofilm-forming cells to antibiotics is generally ascribed to many factors, such as expression of specific resistance genes, low growth rates, poor diffusion of antibiotics across the biofilm matrix, and lower antibiotic activity under low pH and/or low oxygen conditions, which are often present in bacterial biofilms [40–42]. In this view, the high antibiofilm activity of 3BP could be justified by: (i) its low molecular weight and high water solubility, which likely promote diffusion across the exopolysaccharide matrix and/or water channels of the biofilm; (ii) its high stability at low pH [23]; and (iii) its activity against cells that adopt a fermentative metabolism and/or metabolically inactive cells (Table 2 and Fig. 2), such as those present in the deepest layers of biofilm.

In conclusion, this study describes the promising in-vitro activity of the antimetabolite 3BP against both planktonic and biofilm *S. aureus* cells. This compound is currently in preclinical development for anticancer therapy. Although many studies have reported its efficacy and safety in several animal tumour models involving different administration routes, such as intravenous, intraperitoneal, inhalation or intratumoral delivery [6], clinical trials on 3BP have not been conducted to date and detailed pharmacokinetics studies are not available, either in animals or in humans. The concentrations of 3BP active against *S. aureus* are fairly high with respect to conventional antibiotics (Tables 1 and S1, see online supplementary material), but are similar to those active against cancer cell lines [6,28], suggesting that 3BP administration could attain in-vivo 3BP concentrations that are also effective against *S. aureus* cells, at least in animal models. Moreover, considering the

high antibiofilm activity of this compound and its increased activity under anaerobic and acidic conditions, which are common in bacterial biofilms [40–42], 3BP could have potential for the topical treatment of surface-exposed biofilm-related infections, such as chronic wound and ocular *S. aureus* infections [43], which would allow administration of relatively high doses of 3BP with lower risk of systemic toxicity. In this view, the in-vitro results could pave the way for the evaluation of 3BP as a narrow-spectrum antibacterial agent for the treatment of *S. aureus* in acute and/or chronic infection models. More relevantly, characterization of the mechanism(s) of action of this antimetabolite in *S. aureus* could highlight novel molecular target(s) and, therefore, drive the rational design of more active compounds against this important human pathogen.

Acknowledgements

The authors wish to thank Prof. Gian Maria Rossolini (University of Florence, Italy) for providing *E. faecium* strain 3137.

Funding

This work was supported by the Sapienza University of Rome (Ateneo 2017), the Regione Lazio (prot. 85-2017-13763), the Grant of Excellence Departments (MIUR-Italy) and the Pasteur Institute-Cenci Bolognetti Foundation. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Competing interests

None declared.

Ethical approval

Not required.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.ijantimicag.2018.11.008.

References

- [1] Mullard A. Drug repurposing programmes get lift off. *Nat Rev Drug Discov* 2012;11:505–6.
- [2] Rangel-Vega A, Bernstein LR, Mandujano-Tinoco EA, García-Contreras SJ, García-Contreras R. Drug repurposing as an alternative for the treatment of recalcitrant bacterial infections. *Front Microbiol* 2015;6:282.
- [3] Rampioni G, Visca P, Leoni L, Imperi F. Drug repurposing for antivirulence therapy against opportunistic bacterial pathogens. *Emerg Top Life Sci* 2017;1:13–22.
- [4] Geschwind JF, Ko YH, Torbenson MS, Magee C, Pedersen PL. Novel therapy for liver cancer: direct intraarterial injection of a potent inhibitor of ATP production. *Cancer Res* 2002;62:3909–13.
- [5] Cardaci S, Desideri E, Ciriolo MR. Targeting aerobic glycolysis: 3-bromopyruvate as a promising anticancer drug. *J Bioenerg Biomembr* 2012;44:17–29.
- [6] Azevedo-Silva J, Queirós O, Baltazar F, Ulaszewski S, Goffeau A, Ko YH, et al. The anticancer agent 3-bromopyruvate: a simple but powerful molecule taken from the lab to the bedside. *J Bioenerg Biomembr* 2016;48:349–62.
- [7] Ngo H, Tortorella SM, Ververis K, Karagiannis TC. The Warburg effect: molecular aspects and therapeutic possibilities. *Mol Biol Rep* 2015;42:825–34.
- [8] Ko YH, Verhoeven HA, Lee MJ, Corbin DJ, Vogl TJ, Pedersen PL. A translational study ‘case report’ on the small molecule ‘energy blocker’ 3-bromopyruvate (3BP) as a potent anticancer agent: from bench side to bedside. *J Bioenerg Biomembr* 2012;44:163–70.
- [9] El Sayed SM, Mohamed WG, Seddik MA, Ahmed AS, Mahmoud AG, Amer WH, et al. Safety and outcome of treatment of metastatic melanoma using 3-bromopyruvate: a concise literature review and case study. *Chin J Cancer* 2014;33:356–64.

- [10] Feldwisch-Drentrup H. Candidate cancer drug suspected after death of three patients at an alternative medicine clinic. *Science* August 2016;12. Available at <http://www.sciencemag.org/news/2016/08/candidate-cancer-drug-suspected-after-death-three-patients-alternative-medicine-clinic> accessed 11 February 2019.
- [11] Dyląg M, Lis P, Niedźwiecka K, Ko YH, Pedersen PL, Goffeau A, et al. 3-Bromopyruvate: a novel antifungal agent against the human pathogen *Cryptococcus neoformans*. *Biochem Biophys Res Commun* 2013;434:322–7.
- [12] Jagielski T, Niedźwiecka K, Roeske K, Dyląg M. 3-Bromopyruvate as an alternative option for the treatment of protothecosis. *Front Pharmacol* 2018;9:375.
- [13] Geschwind JF, Vali M. Methods of treatment using 3-bromopyruvate and other selective inhibitors of ATP production. Patent WO2011127200A2. October 13, 2011.
- [14] Singh KS, Singh SK. The *Mycobacterium tuberculosis* H37Ra gene MRA_1916 causes growth defects upon down-regulation. *Sci Rep* 2015;5:16131.
- [15] Elkhatib WF, Khairalla AS, Ashour HM. Evaluation of different microtiter plate-based methods for the quantitative assessment of *Staphylococcus aureus* biofilms. *Future Microbiol* 2014;9:725–35.
- [16] Clinical and Laboratory Standards Institute. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard. 9th ed. Wayne, PA: CLSI; 2012. CLSI M07-A9.
- [17] Clinical and Laboratory Standards Institute. Methods for determining bactericidal activity of antimicrobial agents; approved guideline, Wayne, PA: CLSI; 1999. CLSI M26-A.
- [18] White RL, Burgess DS, Manduru M, Bosso JA. Comparison of three different in vitro methods of detecting synergy: time-kill, checkerboard, and E test. *Antimicrob Agents Chemother* 1996;40:1914–18.
- [19] Fernández-Piñar R, Lo Sciuto A, Rossi A, Ranucci S, Bragonzi A, Imperi F. In vitro and in vivo screening for novel essential cell-envelope proteins in *Pseudomonas aeruginosa*. *Sci Rep* 2015;5:17593.
- [20] García-Fernández A, Miriagou V, Papagiannitsis CC, Giordano A, Venditti M, Mancini C, et al. An ertapenem-resistant extended-spectrum-beta-lactamase-producing *Klebsiella pneumoniae* clone carries a novel OmpK36 porin variant. *Antimicrob Agents Chemother* 2010;54:4178–84.
- [21] Iacono M, Villa L, Fortini D, Bordoni R, Imperi F, Bonnal RJ, et al. Whole-genome pyrosequencing of an epidemic multidrug-resistant *Acinetobacter baumannii* strain belonging to the European clone II group. *Antimicrob Agents Chemother* 2008;52:2616–25.
- [22] Bonchi C, Frangipani E, Imperi F, Visca P. Pyoverdine and proteases affect the response of *Pseudomonas aeruginosa* to gallium in human serum. *Antimicrob Agents Chemother* 2015;59:5641–6.
- [23] Glick M, Biddle P, Jantzi J, Weaver S, Schirch D. The antitumor agent 3-bromopyruvate has a short half-life at physiological conditions. *Biochem Biophys Res Commun* 2014;452:170–3.
- [24] Fuchs S, Pané-Farré J, Kohler C, Hecker M, Engelmann S. Anaerobic gene expression in *Staphylococcus aureus*. *J Bacteriol* 2007;189:4275–89.
- [25] Ferreira MT, Manso AS, Gaspar P, Pinho MG, Neves AR. Effect of oxygen on glucose metabolism: utilization of lactate in *Staphylococcus aureus* as revealed by in vivo NMR studies. *PLoS One* 2013;8:e58277.
- [26] European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) EUCAST Definitive Document E.Def 1.2, May 2000: Terminology relating to methods for the determination of susceptibility of bacteria to antimicrobial agents. *Clin Microbiol Infect* 2000;6:503–8.
- [27] Hall CW, Mah TF. Molecular mechanisms of biofilm-based antibiotic resistance and tolerance in pathogenic bacteria. *FEMS Microbiol Rev* 2017;41:276–301.
- [28] Zhang Q, Pan J, North PE, Yang S, Lubet RA, Wang Y, et al. Aerosolized 3-bromopyruvate inhibits lung tumorigenesis without causing liver toxicity. *Cancer Prev Res* 2012;5:717–25.
- [29] Bouvet PJM, Patrick AD. Grimont taxonomy of the genus *Acinetobacter* with the recognition of *Acinetobacter baumannii* sp. nov., *Acinetobacter haemolyticus* sp. nov., *Acinetobacter johnsonii* sp. nov., and *Acinetobacter junii* sp. nov. and emended descriptions of *Acinetobacter calcoaceticus* and *Acinetobacter lwoffii*. *Int J Syst Evol Microbiol* 1986;36:228–40.
- [30] Lin GH, Chen HP, Huang JH, Liu TT, Lin TK, Wang SJ, et al. Identification and characterization of an indigo-producing oxygenase involved in indole 3-acetic acid utilization by *Acinetobacter baumannii*. *Antonie Van Leeuwenhoek* 2012;101:881–90.
- [31] Brown PR, Tata R. Glycollate inhibition of growth of *Pseudomonas aeruginosa* on lactate medium. *J Gen Microbiol* 1987;133:1521–6.
- [32] Eschbach M, Schreiber K, Trunk K, Buer J, Jahn D, Schobert M. Long-term anaerobic survival of the opportunistic pathogen *Pseudomonas aeruginosa* via pyruvate fermentation. *J Bacteriol* 2004;186:4596–604.
- [33] Görke B, Stülke J. Carbon catabolite repression in bacteria: many ways to make the most out of nutrients. *Nat Rev Microbiol* 2008;6:613–24.
- [34] Kreth J, Lengeler JW, Jahreis K. Characterization of pyruvate uptake in *Escherichia coli* K-12. *PLoS One* 2013;8:e67125.
- [35] Smirnova GV, Oktyabrsky ON. Glutathione in bacteria. *Biochemistry (Mosc)* 2005;70:1199–211.
- [36] Masip L, Veeravalli K, Georgiou G. The many faces of glutathione in bacteria. *Antioxid Redox Signal* 2006;8:753–62.
- [37] Qin JZ, Xin H, Nickloff BJ. 3-Bromopyruvate induces necrotic cell death in sensitive melanoma cell lines. *Biochem Biophys Res Commun* 2010;396:495–500.
- [38] El Sayed SM, Baghdadi H, Zolaly M, Almarahy HH, Ayat M, Donki JG. The promising anticancer drug 3-bromopyruvate is metabolized through glutathione conjugation which affects chemoresistance and clinical practice: an evidence-based view. *Med Hypotheses* 2017;100:67–77.
- [39] Van Laar TA, Esani S, Birges TJ, Hazen B, Thomas JM, Rawat M. *Pseudomonas aeruginosa* *gshA* mutant is defective in biofilm formation, swarming, and pyocyanin production. *mSphere* 2018;3 pii:e00155–18.
- [40] Lewis K. Multidrug tolerance of biofilms and persister cells. *Curr Top Microbiol Immunol* 2008;322:107–31.
- [41] Vroom JM, De Grauw KJ, Gerritsen HC, Bradshaw DJ, Marsh PD, Watson GK, et al. Depth penetration and detection of pH gradients in biofilms by two-photon excitation microscopy. *Appl Environ Microbiol* 1999;65:3502–11.
- [42] Kharidia R, Liang JF. The activity of a small lytic peptide PTP-7 on *Staphylococcus aureus* biofilms. *J Microbiol* 2011;49:663–8.
- [43] Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: properties, regulation, and roles in human disease. *Virulence* 2011;2:445–59.